Using genetic data to identify transmission risk factors: statistical assessment and application to tuberculosis transmission

Isaac H. Goldstein[1], Damon Bayer[1], Ivan Barilar[2], Balladiah Kizito[3], Ogopotse Matsiri[3], Chawangwa Modongo[3], Nicola M. Zetola[4], Stefan Niemann[2*], Volodymyr M. Minin[1*], Sanghyuk S. Shin[5*],

1 Department of Statistics, University of California, Irvine, California
2 German Center for Infection Research, Research Center Borstel, Germany
3 Victus Global Botswana Organisation, Gaborone, Botswana
4 Augusta University, Augusta, Georgia
5 Sue & Bill Gross School of Nursing, University of California, Irvine, California

* Co-corresponding authors: stniemann@yahoo.de, vminin@uci.edu, ssshin2@hs.uci.edu

Abstract

Identifying host factors that influence infectious disease transmission is an important step toward developing interventions to reduce disease incidence. Recent advances in methods for reconstructing infectious disease transmission events using pathogen genomic and epidemiological data open the door for investigation of host factors that affect onward transmission. While most transmission reconstruction methods are designed to work with densely sampled outbreaks, these methods are making their way into surveillance studies, where the fraction of sampled cases with sequenced pathogens could be relatively low. Surveillance studies that use transmission event reconstruction then use the reconstructed events as response variables (i.e., infection source status of each sampled case) and use host characteristics as predictors (e.g., presence of HIV infection) in regression models. We use simulations to study estimation of the effect of a host factor on probability of being an infection source via this multi-step inferential procedure. Using `TransPhylo` — a widely-used method for Bayesian estimation of infectious disease transmission events — and logistic regression, we find that low sensitivity of identifying infection sources leads to dilution of the signal, biasing logistic regression coefficients toward zero. We show that increasing the proportion of sampled cases improves sensitivity and some, but not all properties of the logistic regression inference. Application of these approaches to real world data from a population-based TB study in Botswana fails to detect an association between HIV infection and probability of being a TB infection source. We conclude that application of a pipeline, where one first uses `TransPhylo` and sparsely sampled surveillance data to infer transmission events and then estimates effects of host characteristics on probabilities of these events, should be accompanied by a realistic simulation study to better understand biases stemming from imprecise transmission event inference.

Author summary

Factors that affect infectious disease transmission are poorly understood, which impede efforts to prevent the spread of infectious diseases. Recently, software packages have

been developed to infer transmission histories of infectious disease outbreaks using data from infectious disease genetics and epidemiology. These software packages have been used as part of methods to identify individual characteristics that affect infectious disease transmission. We used computer simulation to explore whether a statistical pipeline using the software package `TransPhylo` can successfully identify individual risk factors for being an infection source in a realistic public health setting where only a small proportion of pathogens are sequenced. We simulated tuberculosis (TB) outbreaks with different odds of being an infection source for TB transmission between people living with and without HIV. We found that the `TransPhylo`-based pipeline consistently underestimated the odds ratio for the association between HIV and being an infection source for TB transmission. We then applied this method to data from a TB study from Botswana and found no evidence of an association between HIV and being an infection source for TB transmission. Identification of transmission risk factors may be difficult in settings with low sampling proportion for genetic data.

Introduction

Better understanding of risk factors for transmitting infectious disease can help improve public health interventions. For example, HIV infection is associated with reduced bacterial load and lower probability of cavitary disease among people with tuberculosis (TB) [1]. This has led to the hypothesis that HIV infection may reduce the probability of onward transmission of *M. tuberculosis* among people with TB. [1,2] Improved understanding of the effect of HIV on *M. tuberculosis* transmission could inform prioritization of scarce public health resources for contact tracing. Moreover, identifying host factors for transmission could provide valuable insights into the pathophysiology of the infectious disease.

In practice, it is rare to have perfect knowledge of who infected whom in an infectious disease outbreak, making study of risk factors for transmission difficult. Advances in methods for pathogen genomics have led to enhanced understanding of infectious disease transmission. [3,4] Sequencing data can be compared among infected hosts to rule out direct transmission when substantial genomic differences exist. More recently, several methods have been developed for using genomic and epidemiological data to reconstruct transmission events and identify a putative infection source for each transmission event [5–12]. In general, these tools were originally designed for outbreak investigations, where capturing most, if not all, cases was feasible. Recently, researchers have used such methods on data collected in the context of passive disease surveillance as part of a statistical pipeline to determine the association between potential risk factors and being an infection source (transmitting an infection to at least one other host) [13–15]. **Fig 1** illustrates this pipeline. Researchers first infer a timed phylogenetic tree from genetic data, use the phylogenetic tree to infer infection source status labels for study participants, and finally infer the odds ratio of interest using the inferred infection source labels as observed response variables.

Fig 1. Visualization of statistical pipeline evaluated in this paper. Genomic data are used to construct timed phylogenetic trees, which are then used as inputs to infer transmission trees. Based on transmission trees, probability of being an infection source is derived, and then used as a response variable in an analysis of host factor associations.

While passive disease surveillance could capture a significant portion of infectious disease incidence, we expect a meaningfully smaller proportion of cases will be sampled as compared to an outbreak investigation study. It is worth pointing out that we expect the proportion of cases sampled to heavily depend on the resources of a public health system, as well as the prevalence of the disease being studied. In well resourced public health systems, the difference between passive surveillance and outbreak investigation may be small. Still, the performance of many of the transmission reconstruction methods in the context of passive surveillance is not at all clear. And while statistical pipelines relying on these transmission reconstruction methods have been used in practice, their ability to provide useful estimates of the associations between being an infection source and risk factors of interest remains largely untested. The goal of this study is to better understand the use of these pipelines in disease surveillance contexts by evaluating the performance of a set of statistical pipelines which use the commonly used `TransPhylo` software to reconstruct transmission events [5]. We evaluate bias and precision of the estimators of the odds ratios for probability of being an infection source. We also evaluate the power of statistical tests based on these estimators to reject a null hypothesis that the odds ratio is one, and coverage of corresponding 95% confidence and credible intervals. In particular, we evaluate the properties of using `TransPhylo` generated infection source labels and logistic regression, and `TransPhylo` infection source labels with two different statistical models that take into account measurement error. We accomplish this goal by simulating TB outbreaks with two classes of hosts (representing those living with and without HIV) with different probabilities of being an infection source where the underlying true odds ratios are known, and evaluating pipeline performance under a variety of true odds ratios and sampling designs. Finally, we apply our pipelines to real data to investigate the association between HIV infection and being an infection source for *M. tuberculosis* transmission.

Materials and methods

`TransPhylo` Overview

`TransPhylo` is an R package which uses a statistical model and Markov Chain Monte Carlo to produce a posterior distribution of transmission trees, graphs which describe who infected whom in an infectious disease outbreak [5]. `TransPhylo` uses timed phylogenetic trees of sampled cases in order to produce this posterior. One advantage of using `TransPhylo` for this task is that it is designed to allow for within-host evolution. It does this by "coloring" branches of the phylogenetic tree, where colors represent hosts, and allowing for color changes to occur anywhere on a branch, as opposed to just at bifurcations of the tree. Another advantage of `TransPhylo` is that it explicitly allows for outbreaks where not all individuals have been sampled. Using the posterior distribution of transmission trees, we can create a probability of being an infectious source for each sampled individual by calculating the proportion of posterior trees in which the individual transmitted the disease to another individual (regardless of whether the individual was sampled).

Simulation Set Up

All simulated data sets were generated in R version 4.0.2 or 4.0.4 using the `nosoi` package version 1.0.3 [16, 17]. `nosoi` simulates an infectious disease outbreak by simulating individual cases which generate new cases in discrete units of time until either a specified time limit is reached or a specified total number of cases is reached. The behavior of cases is controlled through three parameters: the probability of the recipient ceasing to be infectious at the beginning of a time step, the number of susceptible persons an individual case encounters in a time step, and finally, the probability that, given a contact with a susceptible individual, an individual case will transmit the pathogen. For all simulations in this study, we assumed discrete time steps represented one month of real time, and chose model parameters to mimic a plausible *M. tuberculosis* outbreak (see Appendix A.1.1 for simulation parameters). In brief, R_0 was set to be 1.18, the mean latent period was 9 months among hosts that became infectious, and the mean infectious period was 3 months. All outbreaks used in this study lasted 8 years. In order to control the number of sampled recipients, we generated simulations at random, but only used outbreaks with a total number of between fifty and two thousand cases. This meant that many potential outbreaks were rejected as part of our sampling simulation process. While this rejection sampling has the potential to change the true odds ratio being estimated, we have found empirically this is not an issue for outbreaks lasting 8 years. In our simulation set-up, each outbreak represents a cluster of *M. tuberculosis* infected individuals to be analyzed jointly using `TransPhylo`, mimicking the common real world practice of clustering using SNPs by some threshold and generating separate timed phylogenies for each cluster. Henceforth we will refer to these outbreaks as clusters.

We simulated clusters with two types of individuals representing people living with and without HIV co-infected with *M. tuberculosis*. Based on data collected for a population-based epidemiology study of *M. tuberculosis* in Botswana (discussed below), we specified that on average 53 percent of cases would be people living with HIV. We had five primary simulation settings where the only changing parameter was the ratio of the probabilities of transmission given contact for two classes of cases. The settings were: probability of transmission given contact was 3.02 times higher among people living without HIV compared to that among people with HIV, 1.75 times higher, equal, 1.75 times lower, or 3.33 times lower. All other parameters were identical for the two classes of cases. For the 3.02 setting, we will round down and refer to this as the 3

setting for the duration of the paper. 102

A single simulation consisted of fifty simulated clusters, where each cluster lasted 103
eight years. From a single cluster, individuals were sampled from the last three years of 104
the outbreak. Recipients were only sampled during their infectious periods, with the 105
sampling time equally likely at any point in this time frame. For each cluster, we 106
sampled 16% of eligible recipients for each of the last three years of the outbreak. Only 107
simulations where all fifty clusters had at least two sampled individuals were accepted. 108
In addition, because individuals could be active in multiple years, there were cases 109
where there were not enough individuals in a particular year left to reach the percentage 110
of 16%, simulations were likewise rejected in this scenario. 111

We chose this sampling scheme to reflect real world sampling conditions in our study 112
setting, where data were collected as part of ongoing surveillance of an endemic disease 113
over the course of multiple years, and where there was little chance of sampling 114
recipients during their latent period. For example, due to under-diagnosis, the World 115
Health Organization estimated that only 54% of people with new TB episodes in 116
Botswana were reported in 2019 [18]. Using this proportion, we divided the number of 117
M. tuberculosis sequences available by the number of estimated TB episodes (reported + 118
unreported) to calculate the sampling proportion for our study. In more detail: we 119
divided the number of diagnosed incidents of TB during the Kopanyo study (N = 4,331) 120
by 0.538 to determine the total number of reported and unreported TB episodes (N = 121
8049). We then divided the number of sequences initially available for analysis (N = 122
1306) by 8049 to determine the sampling proportion of 16%. We continued to use this 123
conservative sampling proportion estimate after increasing our sequenced samples to 124
1426. Thus, 16% reflects a reasonable sampling proportion one would expect when data 125
come from passive surveillance of TB. 126

Once sampled, we generated timed phylogenetic trees for each cluster using `nosoi`. 127
The trees from separate clusters represent the trees from separate transmission clusters 128
which would be generated as part of a real data analysis. These timed phylogenetic 129
trees served as the input for `TransPhylo` (version 1.4.4). For each simulation setting, 130
we generated one hundred simulations. 131

In addition to our primary simulations, we conducted a number of secondary 132
simulations to further investigate the operating characteristics of our statistical pipeline. 133
For all secondary simulations, people living without HIV had a probability of 134
transmission given contact that was set to be 1.75 times higher than people with HIV. 135
We then varied the sampling proportion, increased total sample size, and decreased time 136
from start of epidemic to start of sampling. To investigate the impact of increasing the 137
sampling density, we increased the sampling proportion from 16% to 32% while keeping 138
the sample size approximately the same by halving the number of clusters in a 139
simulation (from fifty to twenty-five). To further investigate the consequences of 140
increasing sampling density, we then increased the sampling proportion from 16% to 141
64%, again keeping the sample size approximately similar using only 13 clusters. To 142
investigate the impact of an increase in sample size, we kept the sampling proportion at 143
16%, but doubled the number of clusters from fifty to one hundred. To investigate the 144
impact of shortening the time from beginning of epidemic to beginning of sampling, we 145
kept the sampling proportion at 16%, but increased the sampling window from the last 146
three years to the last seven years of the simulation, while decreasing the number of 147
clusters from fifty to thirty-five to approximately maintain the original sample size. All 148
simulation settings are summarised in Appendix Table B1. 149

Using `TransPhylo` on Simulated Data Sets 150

We used the multiple cluster functionality of `TransPhylo` to analyze all simulated trees 151
from a single simulation simultaneously. The sampling proportion π was allowed to vary 152

among clusters, while all other parameters were shared across all trees. Based on empirical results from early simulations, for the primary simulations, the prior for the sampling proportion was set as a beta distribution with alpha equal to 1 and beta equal to 19. The other priors in `TransPhylo` cannot be changed by the user. `TransPhylo` assumes the number of individuals an infectious person will infect has a negative binomial distribution governed by two parameters r and p. These parameters also describe the reproduction number, and so their priors control the prior for the reproduction number. r has an exponential prior with rate parameter one, p has a uniform prior over the interval zero to one. We found in practice that p was not identifiable and fixed it at 0.5 for our analyses. Finally, the within-host effective population size also has an exponential prior with rate parameter one. The generation time and sampling time distribution were identical, with shape parameter 10, scale parameter 0.1. Parameters and their priors are summarised in Appendix Table B2. We sampled 100,000 draws from the posterior distribution of transmission trees, discarding the first 50,000 as burn-in, and saving only every tenth sample. Probability of being an infection source was calculated as the proportion of posterior draws in which a particular case transmitted the disease. We considered a case to be an infection source if their inferred posterior probability of being an infection source was greater than 0.6. This same cutoff probability of 0.6 was used for the real data analysis as well. We have found the exact value of this cutoff makes little difference, as the distribution of probabilities tends to be extremely bi-modal, clustered around one and zero (see appendix B.3).

Inferring Odds Ratios using Infection Source Labels

All statistical inference was conducted in R (either version 4.0.2 or 4.0.4). We estimated odds ratios through logistic regression via the `glm` function in R. The mean of the response variable was the probability of being an infection source and the covariate was HIV status, where one indicates a person living without HIV, and zero indicates a person with HIV. That is:

$$\log \left(\frac{P(Y = 1)}{1 - P(Y = 1)} \right) = \beta_0 + HIV^- \beta_1,$$

where $Y = 1$ when a case is labeled as an infection source and $Y = 0$ when they are not. For simulations, we used logistic regression using both the true infection source labels and the labels inferred by `TransPhylo`. True odds ratios for each simulation setting were calculated using Monte-Carlo simulations (see appendix B.2.1). Our results suggested logistic regression using the `TransPhylo` inferred labels yielded sub-optimal inference due to mislabeled cases (i.e. `TransPhylo` inferred a case was not an infection source when in fact they were). This is a well-known phenomenon in statistics called measurement error, see Neuhaus (1999) for a clear introduction to the problem [19]. Many methods have been developed to adjust for this misclassification, we chose to include two as potential remedies to our statistical pipeline.

The first method is available in the `SAMBA` package by Beesley and Mukherjee (2020) (we used version 0.9.0) [20]. The `SAMBA` misclassification model assumes no misclassification of negative results (specificity is 1) and first infers the sensitivity of the response variable, then incorporates this sensitivity in the final logistic regression. To infer sensitivity, the user is required to input a fixed unconditional probability of the response variable being one, in our setting, this is the unconditioned probability that a case is an infection source. For simulations, the true unconditional probability from the sampled data was used. For the real data analysis, the mean probability of being an infection source in the last three years of the cluster from one of the primary simulation settings was used.

The second method is a Bayesian approach we created to account for the case when specificity is not 1. For this model we assume the sensitivity and specificity for the response labels are known. In the model below Y represents the observed response label whereas Z represents the unobserved true response label.

$$\beta_0 \sim N(0,1),$$
$$\beta_1 \sim N(0,1),$$
$$\log \frac{P(Z_i = 1)}{1 - P(Z_i = 1)} = \beta_0 + HIV_i^- \beta_1,$$
$$Y_i \mid Z_i = 0 \sim \text{Bernoulli}(p = 1 - \text{Specificity}),$$
$$Y_i \mid Z_i = 1 \sim \text{Bernoulli}(p = \text{Sensitivity}).$$

For the simulations, Monte-Carlo estimated sensitivity and specificity were used, rounded to two significant digits. For the real data analysis, the Monte-Carlo estimated sensitivity and specificity from one of the primary simulations was used. There was little difference in estimated sensitivity and specificity between the different settings (**Figures B1, B2**). We used Hamiltonian Monte Carlo (HMC) to conduct Bayesian inference with this model using Stan through the `rstan` package (version 2.21.2). We generated 2000 posterior draws and discarded the first 1000 as burn-in. We used the median of the posterior distribution for point estimates and 95% Bayesian credible intervals as measures of uncertainty.

Data Collection

The real data used in this study was originally collected as part of a molecular epidemiology study of *M. tuberculosis* transmission in two regions of Botswana. Botswana has a high prevalence of HIV as well as *M. tuberculosis*. In 2017, 19.9% of people aged 15–49 were living with HIV, 300 of 100,000 people had TB, 48% of people with TB also had HIV [21]. All patients who were diagnosed with *M. tuberculosis* between 2013 and 2016 in the two regions were eligible for enrollment in the study. Of the 4331 enrolled participants, 2159 had culture-confirmed TB. Any participants who had unknown HIV status or who had had a negative result more than a year prior to sampling were offered HIV testing as part of the study protocol [22, 23].

Sequencing

Of the *M. tuberculosis* DNA extracted from cultured isolates, 1671 (77%) were of sufficient quality to undergo sequencing using Nextera XT libraries. We performed WGS using Illumina Technology (MiSeq, NextSeq 500) generating 2x151bp paired end reads. All genomes were analysed with the MTBseq pipeline using the standard input values [24]. In short, reads were mapped to the *M. tuberculosis* H37Rv genome (GenBank ID: NC_000962.3) using BWA-mem and Samtools. Base call recalibration and realignment of reads around insertions and deletions was performed using GATKv3. Finally, variant calling was performed with Samtools mpileup and in-house scripts that employed thresholds for coverage and base quality. After sequencing, 1426 libraries were of high enough quality, defined by at least 50x coverage and 95% of the reference genome covered, to be used for further analysis. Average coverage of the good quality samples was 120x.

Concatenated sequence alignment of variants was produced by including those genome positions that fulfilled the aforementioned criteria for coverage and variant frequency in 95% of all samples. Repetitive regions (PPE/PE-PGRS genes), InDels,

consecutive SNPs in 12bp windows and genes implicated in antibiotic resistance were excluded. Phylogenetically informative SNPs were taken from existing literature. We excluded sequences with heterogeneous reads along a base position for 45 - 50 SNPs or greater, which is indicative of mixed infections, leaving 1347 sequences for analysis.

Analysis of Real Data

Within each lineage, we calculated raw single nucleotide polymorphism (SNP) distances between all sequences and clustered sequences based on two cutoffs, either sequences with a SNP distance of five or fewer, or ten or fewer were clustered together. For our analysis, we excluded clusters with less than four sequences. We also excluded clusters with little genetic variation by excluding clusters where there were less than two times the number of sequences in the cluster minus two varying sites. For the five SNP analysis, we had 147 sequences from 21 clusters. For the ten SNP analysis, we had 388 sequences from 46 clusters to analyze.

We used BEAST 2 version 2.6.3 to create posterior distributions of timed phylogenetic trees with 10 million posterior draws for each of our clusters [25]. We then calculated maximum clade credibility trees to use as the inputs into `TransPhylo`, discarding the first ten percent of Markov chain Monte Carlo iterations. We used an HKY model with strict molecular clock and a constant effective population size model, and used an ascertainment bias correction for sequences of unknown length to account for the fact that analyzed sequences included only varying sites [26]. We used a uniform prior for the molecular clock with a lower bound 1×10^{-8} and upper bound 5×10^{-7} scaled by 4.2 million divided by the number of sites in the data for each cluster. The scaling was done to account for the fact that we used BEAST2 on SNPs, not the full genome. The upper and lower bounds were chosen based on work by Menardo et. al, and were chosen to be suitable for all TB lineages [27]. Priors on the frequency parameter, kappa and population size were all log normal distributions with parameter M set to 1.0 and parameters S set to 1.25.

For `TransPhylo`, shape and scale parameters for the generation time and sampling time distributions were fixed at 10 and 0.1 respectively. The multitree function was used to conduct inference simultaneously on all clusters generated by the same SNP threshold, with all parameters but the sampling proportion shared among all clusters. We used a beta distribution with alpha set to 1 and beta set to 19 as a prior for the sampling distribution. We drew 100,000 posterior samples, discarding the first 50,000 samples as burn-in and saving every tenth sample. We calculated the odds ratio for people living without HIV vs with HIV using the inferred `TransPhylo` labels using logistic regression, using the SAMBA package, and using the Bayesian model as described in a previous subsection.

Ethics Statement

The real data were collected for the Kopanyo Study, which received approval from the Health Research and Development Committee of the Botswana Ministry of Health and Wellness, the U.S. Center for Disease Control and Prevention Institutional Review Board (IRB) (6291), and the University of Pennsylvania IRB. Informed consent was obtained from all participants.

Reproducibility

All code and data needed to recreate the results of this study are available at https://github.com/igoldsteinh/kopanyo_tp_code.

Results 276

Simulated Data 277

Table 1 shows summary statistics of the simulated clusters used in the primary 278
simulation setting where people living with HIV were 1.75 times less likely to transmit 279
infection given contact compared to people without HIV. Median cluster size was 116, 280
with 95% of clusters ranging from 53 to 393 hosts. Latent periods for cases had a 281
median of 9 months, while infectious periods lasted a median of 3 months. Median tree 282
height was 5.9 years. 283

Metric	Median	[2.5%Q, 97.5% Q]
Cluster Size	116	[53, 393]
Samples per Cluster	9	[3,34]
Latent Period (Months)	9	[4,17]
Infectious Period (Months)	3	[1,15]
$P(IS^*)$	0.69	[0.64, 0.73]
$P(HIV^+)$	0.53	[0.49,0.56]
$P(IS|HIV^-)$	0.77	[0.71, 0.81]
$P(IS|HIV^+)$	0.61	[0.57, 0.67]
$P(HIV^+|IS = N)$	0.65	[0.58, 0.71]
$P(HIV^+|IS = Y)$	0.48	[0.43, 0.52]
Tree Height (Years)	5.92	[2.59, 7.42]
% TP Labeled IS	19.47	[17.11, 22.35]

Table 1. Simulation Summary Statistics for 8 Year Simulation. Summary
statistics of simulated infectious disease outbreaks from 100 simulations where
probability of transmission given contact for hosts without HIV was 1.75 times larger
than for hosts with HIV. *Infection Source - transmitting an infection to at least one
other host

Operating Characteristics of Statistical Pipeline 284

For each simulation setting, we calculated the sensitivity and specificity of `TransPhylo` 285
for identification of an individual as an infection source. Point estimates for each 286
simulation setting are displayed in **Figures B1, B2**. In general, specificity was high, 287
around 0.97, while sensitivity was low, approximately 0.28. Sensitivity increased when 288
either the sampling density or the sampling window was increased. 289

Then, for each version of our pipeline and each simulation setting, we calculated 290
frequentist coverage (proportion of simulations for which a 95% confidence/credible 291
interval contained the true parameter–ideally it should be 0.95), proportion reject: the 292
proportion of times that a null hypothesis that the odds ratio was one was rejected 293
(with significance level 5%), percent bias (bias divided by the true odds ratio), and 294
mean confidence interval width of 95% confidence intervals. **Figure 2** shows the results 295
from the primary set of simulations. The four pipelines tested for each simulation were: 296
a reference pipeline where true infection source labels were known and used as inputs 297
into a generalized linear model (Truth + GLM), a pipeline where `TransPhylo` labels 298
were used as input into a generalized linear model (TP + GLM), a pipeline where 299
`TransPhylo` labels were used as input into SAMBA (TP + ME1) for misclassification 300
adjustment and a pipeline where `TransPhylo` labels were used as inputs in our Bayesian 301
model for misclassification adjustment (TP + ME2). Across simulation settings, the 302
Truth + GLM pipeline had coverage around 0.95, proportion reject above 0.95 (except 303
in the case when the odds ratio was 1), percent bias near 0 and confidence interval 304
width ranging from 0.17 to 2.7. For the TP + GLM pipeline, coverage varied depending 305

on the simulation setting, ranging from 0.62 to 0.93. When the true odds ratio was not 306
one, TP + GLM had proportion reject ranging from 0.54 to 1. Percent bias ranged from 307
-16.76% to 48.34%, mean credible interval width was 0.33 to 2.28. The TP + ME1 308
pipeline had coverage and proportion reject similar to TP + GLM, with percent bias 309
ranging from -55.83% to 195.66% and MCIW between 0.25 and 23.94. Finally, the TP + 310
ME2 pipeline had coverage ranging from 0.79 to 0.98, proportion reject ranging from 311
0.44 to 1, percent bias from -30.52% to 118.51% and MCIW from 0.36 to 22.. 312

Fig 2. Operating characteristics of statistical pipelines in primary simulation settings.
Truth + GLM is a reference pipeline where the true infection source labels are used as
response variables in a generalized linear model. TP + GLM is a pipeline where
infection source labels generated by `TransPhylo` are used as response variables in a
generalized linear model. TP + ME1 is a pipeline where infection source labels are
generated by `TransPhylo` and used as an input into a model from the `SAMBA` package,
which allows for false positives. TP + ME2 again uses labels generated by `TransPhylo`
as response variables into a model which allows for both false positives and false
negatives. The settings denote different simulation settings, each value describes the
true ratio of the probability of transmission given contact for cases living without HIV
to cases with HIV. I.E., 0.57 means that the probability of transmission given contact
for hosts without HIV was 1.75 times as small as the probability of transmission given
contact for hosts with HIV. Coverage refers to proportion of simulations where 95%
confidence intervals captured the true odds ratio. Prop. Reject refers to the proportion
of simulations where a null hypothesis that the true odds ratio was one would be
rejected, assuming significance level of 5%. Percent bias is bias divided by the true odds
ratio, MCIW is mean confidence interval width.

The results of our secondary set of simulations (along with the original results from 313

the 1.75 setting) are displayed in **Figure** 3. For the TP + GLM pipeline, increasing the ⁣ ⁣314
percent of hosts sampled from 16% to 32% (Double Sampling Density) led to increased ⁣ ⁣315
coverage and increased proportion of scenarios where a null hypothesis that the odds ⁣ ⁣316
ratio was one was rejected. Quadrupling the percent of hosts sampled to 64% (Quad ⁣ ⁣317
Sampling Density) led to a decrease in coverage compared to both the default and the ⁣ ⁣318
Double Sampling Density simulation, although the proportion reject remained ⁣ ⁣319
comparable with Double Sampling Density simulation, and the bias decreased. ⁣ ⁣320
Increasing the sampling window to all but the first year of the outbreak slightly ⁣ ⁣321
decreased coverage and increased proportion reject. Doubling the number of clusters ⁣ ⁣322
analyzed from fifty to one hundred (Increase Sample Size) led to the lowest coverage of ⁣ ⁣323
all settings but the highest proportion reject. For the two measurement error pipelines, ⁣ ⁣324
the proportion reject improved in a similar manner as in the TP + GLM pipeline. ⁣ ⁣325
However, the coverage changed in a different manner, with both the Increase Sample ⁣ ⁣326
Sizes and the Increase Sampling Window having substantially lower coverage than the ⁣ ⁣327
default setting. ⁣ ⁣328

Fig 3. Operating characteristics of statistical pipelines in secondary simulation settings.
See Figure 2 for x-axis and y-axis labels. The settings denote different simulation
settings. For all simulations the probability of transmission given contact for cases
living without HIV was 1.75 times as large as the probability of transmission given
contact for cases living with HIV. Default refers to the simulation settings used in the
primary simulation setting. In the Double Sampling Density setting percent of active
cases sampled was doubled from 16% to 32%. In the Quad Sampling Density setting
percent active cases sampled was quadrupled to 64%. In the Increase Sample Size
setting the number of clusters sampled from was doubled from 50 to 100. In the
Increase Sampling Window the sampling window was changed from the last three years
of the simulation to the last seven years.

Real Data Results

Study participant characteristics from the Botswana study are summarized in **Table 2**. The vast majority of sampled TB bacteria belonged to lineage 4. More than half of study participants were male, and most participants were living with HIV.

	Count	Percent
TB Lineage		
Lineage 1	83	6.4
Lineage 2	69	5.3
Lineage 3	12	0.9
Lineage 4	1142	87.4
Sex		
Male	719	55.1
Female	587	44.9
Age		
Under 18	80	6.1
18 to 35	693	53.1
35 to 55	466	35.7
55 to 75	67	5.1
75 and older	10	0.8
HIV Status		
Without HIV	590	45.2
With HIV	716	54.8

Table 2. Study participant characteristics Summary statistics of participant characteristics from the study of *M. tuberculosis* in Botswana.

The four largest phylogenetic trees from each of our real data analyses (using five SNPs and ten SNPs as clustering cutoffs) produced by our statistical pipelines are displayed in **Figure 4**. From our genetic samples, we used BEAST2 to create a phylogenetic tree, then used `TransPhylo` to label tips of the tree as being infection sources or not infection sources and compare with their recorded HIV status. The bottom row have tips labeled by ID numbers, demonstrating the way trees change when inclusion criteria change. All participants from the five SNP tree (right) remain in the same ten SNP tree (left), with additional participants included because of the relaxed inclusion criteria.

Fig 4. Timed Phylogeny with tips labeled by HIV status and inferred infection source status using `TransPhylo`. Timed Phylogenies are generated in `BEAST 2`. Before timed phylogenies are generated, TB genomes are first clustered by SNP distance based on a cutoff of either five or ten SNPs. The trees generated under each cutoff are different, but related to each other, with larger trees overall using a cutoff of ten SNPs as opposed to five. The plot was made using R package `ggtree` [28].

Table 3 shows results from our real data analysis using clusters generated under a five SNP cutoff rule and a ten SNP cutoff rule. Mean cluster size and mean tree height increased marginally depending on which cutoff was used (increasing from 7 to 8.43 participants and from 6.86 years to 8.27 years), but both are similar to cluster size and tree height in simulated data (**Table 1**). Overall sample size was smaller than in simulation settings. In both analyses, between 11% to 12% of participants were identified as infection sources. Using any version of our pipeline, we fail to reject (at a 5% significance level) a null hypothesis that the odds ratio is one. Using either measurement error pipeline, the confidence intervals are larger than those produced by the TP + GLM pipeline. For the five SNP Analysis TP + GLM had confidence interval width of 2.39, while TP + ME1 had confidence interval width 17.36, and TP + ME2 had confidence interval width of 2.70. Using data from the 10 SNP analyses, all

pipelines had shorter confidence intervals. The TP + GLM pipeline had a confidence   354
interval width of 1.40, TP + ME1 had confidence interval width of 6.58, and TP + ME2   355
had confidence interval width of 2.15.   356

	5 SNP Analysis	10 SNP Analysis
Participants	145	377
Clusters	21	46
Mean Samples per Cluster	7	8.43
Mean Tree Height (Years)	6.86	8.27
Infection Sources	16	43
OR and 95% CI (TP + GLM)	0.96 [0.33, 2.72]	1.03 [0.55, 1.95]
OR and 95% CI (TP + ME1)	0.88 [0.04, 17.4]	1.10 [0.18, 6.76]
Mean OR and 95% CI (TP + ME2)	0.80 [0.20, 2.90]	0.96 [0.36, 2.51]

Table 3. Summary of analyses using five and ten SNP cutoffs Summary of
analyses using five and ten SNP cutoffs. Mean tree height refers to the height of timed
phylogenetic trees in years. TP + GLM refers to a statistical pipeline where infection
source labels are first generated using `TransPhylo` and then used as response variables
in a generalized linear model to calculate an odds ratio. The odds ratio is the odds ratio
for the probability of being an infection source given the case is living without HIV as
compared to cases living with HIV.

In addition, we used a generalized linear model with additional covariates for   357
participant gender, smoking status, past history of TB infection and excessive alcohol   358
use. The model had no covariates for which a null hypothesis the odds ratio was one   359
could be rejected, and the point estimates and confidence intervals for the odds ratio for   360
HIV status were similar to those reported in Table 3.   361

Discussion   362

In this study, we used simulations under realistic scenarios for passive TB surveillance   363
to show that the statistical pipeline using `TransPhylo`-generated labels as response   364
variables in a generalized linear model (TP + GLM) had poor frequentist properties for   365
estimating the association between host characteristic and being an infection source.   366
When the difference in probability of infection given contact was high between host   367
categories, as in the 0.3 and 3 settings, TP + GLM results would lead one to correctly   368
reject a null hypothesis the odds ratio was one more than 95% of the time, but due to   369
bias and overly narrow confidence intervals, often failed to capture the true odds ratio.   370
In the case of the 0.57 and 1.75 settings, TP + GLM results would lead to rejecting the   371
null hypothesis less than 65% of the time. This poor performance is partially explained   372
by the sensitivity and specificity of `TransPhylo` for generating infection source labels.   373
When used in our study setting, `TransPhylo` did not accurately label infection sources,   374
leading to substantial under-counts of the true number of infection sources (**Figures**   375
B1, **B2**). This measurement error resulted in poor estimation of the odds ratio of   376
interest. The two measurement error models tested as part of a pipeline did not lead to   377
improved coverage or higher rates of correctly rejecting the null hypothesis.   378
In all scenarios tested, the bias of using `TransPhylo` labels was such that the   379
estimated odds ratio was always closer to one than the true odds ratio (**Figure 2**).   380
When the true odds ratio was one, the pipeline performed well, with low bias, coverage   381
near 0.95, and the probability of falsely rejecting null hypothesis near 0.05. This   382
suggests that TP + GLM is a conservative pipeline, with bias towards failing to detect   383
a significant effect, unless the effect size is large. We note that the simulations used in   384
these studies created true timed phylogenies which were used as inputs into   385

`TransPhylo`. In practice, these phylogenies must also be estimated from sequence data. [386] Depending on the genetic diversity of available sequences, there may also be errors when [387] creating the timed phylogeny. The errors in reconstruction of phylogenies could [388] propagate further down the statistical pipeline, reducing its overall performance. [389]

Our findings are consistent with a recent benchmarking study of computational [390] methods for reconstructing TB transmission, which included `TransPhylo` and five other [391] methods. [29] That study used simulations to show low sensitivity for correctly [392] identifying transmission events under realistic low-TB burden scenarios for all methods [393] tested. Another recent study used simulations to evaluate the use of genomic data and [394] GLM for identifying risk factors for TB transmission, defined as genetic closeness (<2 [395] SNPs). [30] That study found a consistent underestimation of the true odds ratio under [396] multiple scenarios in low-TB burden settings. Our findings complement these prior [397] studies by providing further insights into the limitations of using pathogen genomic [398] data for understanding infectious disease transmission, with simulations and data based [399] on a high TB burden setting. [400]

The TP + GLM pipeline has now been implemented in multiple published studies [401] using surveillance data of *M. tuberculosis* [13–15]. These studies assumed higher [402] sampling proportions than those considered in our study. For example, Xu et. al. study [403] of TB in Spain used a prior for the sampling proportion with plausible values ranging [404] from 48% to 99% [13]. Depending on the surveillance setting, such high sampling [405] density may indeed be plausible. In our study of TB in Botswana, based on the World [406] Health Organizations estimates of the reporting rate for TB, we assumed a much lower [407] sampling proportion [18]. We recommend carefully considering both the sampling [408] scheme of any future studies using surveillance data when conducting research using [409] this pipeline, as well as the plausible effect sizes researchers wish to detect. Simulation [410] studies with pre-specified effect sizes could be used as part of study design in order to [411] help researchers decide what sampling protocol they should aim for in future studies. [412] The code used in this study could be used as a starting point for conducting such [413] simulation studies. [414]

We found that changes to the sampling scheme could improve frequentist properties [415] of the TP + GLM pipeline. For example, doubling the number of clusters under the [416] 1.75 setting increased the proportion of null hypotheses rejected to 0.94 (**Figure 3**). [417] However, just increasing the overall sample size did not improve coverage of confidence [418] interval. While increased sample size decreased variance, and thus confidence interval [419] width, it did not affect bias. In contrast, doubling the sampling density and increasing [420] the sampling window both decreased bias and mean confidence interval width by a [421] small amount. This led to improvements in proportion of null hypotheses rejected, and [422] marginal improvements in coverage (increased sampling density) or only marginal losses [423] in coverage (increased sampling window). The reduction in bias likely follows from [424] improvements in the sensitivity of `TransPhylo` under these settings(**Figures B1, B2**). [425] Less intuitively, we found that quadrupling the sampling density actually decreased the [426] coverage due to narrower confidence intervals, even though the sensitivity of [427] `Transphylo` improved. The bias decreased, as we would expect, but the mean [428] confidence interval width decreased so much that coverage dropped. We speculate this [429] is a consequence of the proportion of individuals labeled as infection sources. As this [430] proportion nears 50%, we expect the estimated standard errors to narrow, because it is [431] easier to estimate probabilities closer to 0.5. Our particular result is almost certainly [432] dependent on the true effect size, that is, we would not expect that for all uses of this [433] pipeline, a sampling proportion of 0.64 would decrease coverage as compared to a [434] sampling proportion of 0.32. Our results suggest the behavior of this pipeline is [435] unpredictable, a higher sampling density will not automatically result in improved [436] coverage. When exact estimation of the effect size is important, we urge great caution [437]

when using this pipeline.   438

The improved performance under the increased sampling window suggests there may   439
be room for improvement in the statistical model being used. Currently, `TransPhylo`   440
assumes that the sampling proportion is constant throughout the period of the   441
outbreak, while in our simulation scenarios sampling only took place in a fixed number 442
of years of an outbreak (as it often would in a real life surveillance study). Future work 443
could focus on changing this assumption in a `TransPhylo`-like model which would   444
better match the the model likelihood to the data generation process. Though, again,   445
we would urge caution, as improving the sensitivity of `TransPhylo` can lead to   446
non-intuitive pipeline performance.   447

Applying the `TransPhylo`-based inference of who-infected-whom to a   448
population-based TB study in Botswana, we failed to detect an association between   449
HIV status and being an infection source for *M. tuberculosis* transmission. TB is the   450
leading cause of death among people living with HIV [18]. The pathophysiology of   451
HIV-TB co-infection is complicated as HIV-associated immunosuppression reduces *M.*   452
tuberculosis bacterial load in HIV-infected hosts. This observation has led to the   453
hypothesis that people living with HIV have a lower risk of onward TB transmission   454
than people without HIV. [2] Epidemiologic studies investigating this hypothesis have   455
produced conflicting results [15, 31, 32]. In light of our simulation study results, our   456
failure to reject the null hypothesis may be due to inadequate statistical power for   457
detecting an association of low to moderate effect sizes when sampling proportion is low. 458
Additional studies using more densely sampled sequences are needed to provider   459
stronger evidence regarding this putative association.   460

This study had several limitations which should be acknowledged. First, our   461
simulations made several simplifying assumptions which make them imperfect   462
representations of *M. tuberculosis* outbreaks. In particular, we assumed latent and   463
infectious periods were the same for hosts with and without HIV, and further assumed   464
that sampling had no impact on infectiousness, whereas in the real world sampling   465
would presumably be followed by treatment which would impact infectiousness. In our   466
real data analysis, *M. tuberculosis* sequencing data were missing for majority of the   467
enrolled participants. We assumed these data were missing completely at random.   468
Moreover, the number of infection sources inferred by `TransPhylo` was small, which   469
limited the ability to detect associations using GLM. Finally, our study assessed   470
pipeline performance in the best case scenario: when timed phylogenetic trees are   471
perfectly accurate. However, in the real world, phylogenetic trees must first be   472
constructed from sequence data, a process which may affect pipeline performance.   473
Conducting assessments of these pipelines using simulation engines which simulate   474
sequence data is a useful area of future research.   475

Our findings provide new insights into the use of statistical pipelines that combine   476
pathogen genomic and epidemiologic data to investigate host factors associated with   477
onward transmission of infectious diseases. We recommend careful consideration of   478
potential biases due to imprecise inference of transmission events when using these   479
pipelines. Simulation studies based on the data collection context may improve the   480
interpretation of the results of studies using these pipelines.   481

Acknowledgements   482

We thank Xavier Didelot for helping us resolve issues using `TransPhylo` and Paul
Bastide and Sebastian Lequime for their help using `nosoi`. We also thank Caroline
Colijn, Jessica Stockdale and Vijay Jeevanantham Naidu for sharing their code using
`TransPhylo` and `BEAST2`. Research reported in this publication was supported by the
National Institute of Allergy And Infectious Diseases of the National Institutes of

Health under Award Number R01AI147336 and R01AI097045. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Peters JS, Andrews JR, Hatherill M, Hermans S, Martinez L, Schurr E, et al. Advances in the understanding of Mycobacterium tuberculosis transmission in HIV-endemic settings. The Lancet Infectious diseases. 2019;19(3):e65—-e76.

2. Yates TA, Abubakar I, Tanser F. HIV infection and the transmission of tuberculosis. J Infect Dis. 2015;211(9):1510.

3. Gardy JL, Johnston JC, Sui SJH, Cook VJ, Shah L, Brodkin E, et al. Whole-genome sequencing and social-network analysis of a tuberculosis outbreak. New England Journal of Medicine. 2011;364(8):730–739.

4. Hall M, Woolhouse M, Rambaut A. Using genomics data to reconstruct transmission trees during disease outbreaks. Revue scientifique et technique (International Office of Epizootics). 2016;35(1):287–296.

5. Didelot X, Fraser C, Gardy J, Colijn C. Genomic infectious disease epidemiology in partially sampled and ongoing outbreaks. Molecular Biology and Evolution. 2017;34(4):997—1007.

6. Wymant C, Hall M, Ratmann O, Bonsall D, Golubchik T, de Cesare M, et al. PHYLOSCANNER: inferring transmission from within-and between-host pathogen genetic diversity. Molecular Biology and Evolution. 2018;35(3):719—-733.

7. Volz EM, Frost SD. Inferring the source of transmission with phylogenetic data. PLoS Computational Biology. 2013;9(12):e1003397.

8. Hall M, Woolhouse M, Rambaut A. Epidemic reconstruction in a phylogenetics framework: transmission trees as partitions of the node set. PLoS Computational Biology. 2015;11(12):e1004613.

9. Campbell F, Cori A, Ferguson N, Jombart T. Bayesian inference of transmission chains using timing of symptoms, pathogen genomes and contact data. PLoS Computational Biology. 2019;15(3):e1006930.

10. Campbell F, Didelot X, Fitzjohn R, Ferguson N, Cori A, Jombart T. outbreaker2: a modular platform for outbreak reconstruction. BMC Bioinformatics. 2018;19(11).

11. Sledzieski S, Zhang C, Mandoiu I, Bansal MS. TreeFix-TP: phylogenetic error-correction for infectious disease transmission network inference. In: BIOCOMPUTING 2021: Proceedings of the Pacific Symposium. World Scientific; 2020.

12. Gallagher SK, Follmann D. Branching process models to identify risk factors for infectious disease transmission. Journal of Computational and Graphical Statistics. 2021;In Press.

13. Xu Y, Cancino-Muñoz I, Torres-Puente M, Villamayor LM, Borrás R, Borrás-Máñez M, et al. High-resolution mapping of tuberculosis transmission: Whole genome sequencing and phylogenetic modelling of a cohort from Valencia Region, Spain. PLoS Medicine. 2019;16(10):e1002961.

14. Yang C, Lu L, Warren JL, Wu J, Jiang Q, Zuo T, et al. Internal migration and transmission dynamics of tuberculosis in Shanghai, China: an epidemiological, spatial, genomic analysis. The Lancet Infectious Diseases. 2018;18(7):788—-795.

15. Sobkowiak B, Banda L, Mzembe T, Crampin AC, Glynn JR, Clark TG. Bayesian reconstruction of Mycobacterium tuberculosis transmission networks in a high incidence area over two decades in Malawi reveals associated risk factors and genomic variants. Microbial Genomics. 2020;6(4). doi:https://doi.org/10.1099/mgen.0.000361.

16. R Core Team. R: A Language and Environment for Statistical Computing; 2020. Available from: https://www.R-project.org/.

17. Lequime S, Bastide P, Dellicour S, Lemey P, Baele G. nosoi: A stochastic agent-based transmission chain simulation framework in R. Methods in Ecology and Evolution. 2020;11(8):1002–1007.

18. Organization WH. Global Tuberculosis Report 2020;. https://www.who.int/publications-detail-redirect/9789240013131.

19. Neuhaus JM. Bias and efficiency loss due to misclassified responses in binary regression. Biometrika. 1999;86(4):843–855.

20. Beesley LJ, Mukherjee B. Statistical inference for association studies using electronic health records: handling both selection bias and outcome misclassification. Biometrics. 2022;78:214–226.

21. Botswana Country Profile;. https://www.cdc.gov/globalhivtb/where-we-work/botswana/botswana.html.

22. Zetola N, Modongo C, Moonan P, Click E, Oeltmann J, Shepherd J, et al. Protocol for a population-based molecular epidemiology study of tuberculosis transmission in a high HIV-burden setting: the Botswana Kopanyo study. BMJ Open. 2016;6(5):e010046.

23. Zetola NM, Moonan PK, Click E, Oeltmann JE, Basotli J, Wen XJ, et al. Population-Based Geospatial and Molecular Epidemiologic Study of Tuberculosis Transmission Dynamics, Botswana, 2012–2016. Emerging Infectious Diseases. 2021;27(3):835—-844.

24. Kohl TA, Utpatel C, Schleusener V, De Filippo MR, Beckert P, Cirillo DM, et al. MTBseq: a comprehensive pipeline for whole genome sequence analysis of Mycobacterium tuberculosis complex isolates. PeerJ. 2018;6:e5895.

25. Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, et al. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. PLoS Computational Biology. 2019;15(4):e1006650.

26. Bouackert, Remco. Ascertainment correction/Sampling bias; 2019. Available from: https://www.beast2.org/2019/07/18/ascertainment-correction.html.

27. Menardo F, Duchêne S, Brites D, Gagneux S. The molecular clock of Mycobacterium tuberculosis. PLoS Pathogens. 2019;15(9):e1008067.

28. Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. ggtree: an R package for visualization and annotation of phylogenetic trees with their covariates and other associated data. Methods in Ecology and Evolution. 2017;8(1):28–36.

29. Sobkowiak B, Romanowski K, Sekirov I, Gardy JL, Johnston J. Comparing transmission reconstruction models with Mycobacterium tuberculosis whole genome sequence data. bioRxiv. 2022;doi:https://doi.org/10.1101/2022.01.07.475333.

30. Leavitt SV, Horsburgh CR, Lee RS, Tibbs AM, White LF, Jenkins HE. What Can Genetic Relatedness Tell Us About Risk Factors for Tuberculosis Transmission? Epidemiology. 2021;33(1):55–64.

31. Winter JR, Smith CJ, Davidson JA, Lalor MK, Delpech V, Abubakar I, et al. The impact of HIV infection on tuberculosis transmission in a country with low tuberculosis incidence: a national retrospective study using molecular epidemiology. BMC Medicine. 2020;18(1).

32. Middelkoop K, Mathema B, Myer L, Shashkina E, Whitelaw A, Kaplan G, et al. Transmission of tuberculosis in a South African community with a high prevalence of HIV infection. The Journal of Infectious Diseases. 2015;211(1):53—-61.

33. Svensson A. A note on generation times in epidemic models. Mathematical Biosciences. 2007;208(1):300–311.

34. Champredon D, Dushoff J. Intrinsic and realized generation intervals in infectious-disease transmission. Proceedings of the Royal Society B: Biological Sciences. 2015;282(1821):2015–2026. doi:10.1098/rspb.2015.2026.

35. Ma Y, Horsburgh CR, White LF, Jenkins HE. Quantifying TB transmission: a systematic review of reproduction number and serial interval estimates for tuberculosis. 2018;146(12):1478–1494. doi:10.1017/S0950268818001760.

36. Behr MA, Edelstein PH, Ramakrishnan L. Revisiting the timetable of tuberculosis. Bmj. 2018;362.

37. Molebatsi K, Wang Q, Dima M, Ho-Foster A, Modongo C, Zetola N, et al. Depression and delayed tuberculosis treatment initiation among newly diagnosed patients in Botswana;16(7):1088–1098. doi:10.1080/17441692.2020.1826049.

38. Cox H, Dickson-Hall L, Ndjeka N, Hoog Av, Grant A, Cobelens F, et al. Delays and loss to follow-up before treatment of drug-resistant tuberculosis following implementation of Xpert MTB/RIF in South Africa: A retrospective cohort study. 2017;14(2):e1002238. doi:10.1371/journal.pmed.1002238.

39. Dodd PJ, Looker C, Plumb ID, Bond V, Schaap A, Shanaube K, et al. Age-and sex-specific social contact patterns and incidence of Mycobacterium tuberculosis infection. American journal of epidemiology. 2016;183(2):156–166.

40. Mousa A, Winskill P, Watson OJ, Ratmann O, Monod M, Ajelli M, et al. Social contact patterns and implications for infectious disease transmission–a systematic review and meta-analysis of contact surveys. ELife. 2021;10:e70294.

41. Davison AC, Hinkley DV. Bootstrap methods and their application. 1. Cambridge University Press; 1997.