Impact Of an Immune Modulator Mycobacterium-w On Adaptive Natural Killer Cells

and Protection Against COVID-19

Sarita Rani Jaiswal [1,2,5], Jaganath Arunachalam[1], Ashraf Saifullah[2], Rohit Lakhchaura[2],

Dhanir Tailor [6], Anupama Mehta[2], Gitali Bhagawati[3], Hemamalini Aiyer[3], Bakulesh

Khamar[4], Sanjay V. Malhotra[6], Suparno Chakrabarti[1,2*]

Cellular Therapy and Immunology, Manashi Chakrabarti Foundation, New Delhi[1],

Department of Blood and Marrow Transplantation[2] and Department of Pathology and

Microbiology, Dharamshila Narayana Super-speciality Hospital[3], Research & Development,

Cadila Pharmaceuticals Ltd, Ahmedabad[4], Amity Institute of Molecular Medicine and Stem

Cell Research, Amity University, Noida, Uttar Pradesh[5], India, Department of Cell,

Development & Cancer Biology and Center for Experimental Therapeutics, Knight Cancer

Institute, Oregon Health & Science University, Portland, OR, USA[6].

Running Title: Mw for COVID-19

Keywords: *Mycobacterium w* (Mw), COVID-19, SARS-CoV-2, innate immunity, NKG2C,

Adaptive NK cells, NKG2A, ADCC

*CORRESPONDENCE:

Suparno Chakrabarti

Cellular Therapy and Immunology,

Manashi Chakrabarti Foundation, Vasundhara Enclave, New Delhi-110096, India

Email: foundationforcure@gmail.com

ABSTRACT

The kinetics of NKG2C$^+$ adaptive natural killer (ANK) cells and NKG2A$^+$inhibitory NK

(iNK) cells with respect to the incidence of SARS-CoV-2 infection were studied for 6 months

in a cohort of health-care workers following administration of heat killed *Mycobacterium w*

(Mw group) in comparison to a control group. In both groups, COVID-19 correlated with a

lower NKG2C$^+$ANK cells at baseline. There was a significant upregulation of NKG2C

expression and IFN-γ release in Mw group (p=0.0009), particularly in those with lower

baseline NKG2C expression, along with downregulation of iNK cells (p<0.0001). This

translated to a significant reduction in incidence and severity of COVID-19 in the Mw group

(IRR-0.15, p=0.0004). RNA-seq analysis at 6 months showed an upregulation of ANK

pathway genes and an enhanced ANK mediated ADCC signature. Thus, Mw was observed to

have a salutary impact on the ANK cell profile and a long-term upregulation of ANK-ADCC

pathways, which could have provided protection against COVID-19 in a non-immune high-

risk population.

INTRODUCTION

The rapid explosion of the novel coronavirus, SARS-CoV-2, in early 2020, across the globe overwhelmed even the most prepared health infrastructures[1] and exposed the healthcare workers to an unforeseen situation, where they remained at the greatest risk of exposure to the highest viral load, in the absence of a prevention or a cure. Despite a very high incidence of infections, witnessed in the Indian population as well, there was a surprising sparing of the urban slum-dwellers[2]. We hypothesized the role of a bolstered innate immune system secondary to chronic pathogen exposure as a plausible reason for this phenomenon. We reasoned that due to ubiquitous exposure to cytomegalovirus (CMV) in early childhood, followed by exposure to a multitude of pathogens subsequently, there might be stronger repertoire of NKG2C expressing adaptive natural killer (ANK) cells in this population [3].

Early CD56^{bright} NK cells have very high expression of NKG2A, which functions as an inhibitory check-point in the process of functional maturation of NK cells[4]. Both NKG2C and NKG2A bind to the same ligand, HLA-E, but the latter binds with several-fold greater affinity compared to NKG2C [5]. Unlike somatic mutations witnessed in adaptive immune cells to produce precise and clonal antigen specificity, NK cells express a plethora of germline encoded activating and inhibitory receptors. The regulation of NK cell function, which is termed as 'licensing', occurs with stochastic expression of killer-immunoglobulin-like receptors (KIR), which bind to self- HLA-Class 1 antigens with biallelic specificity[6]. Expression of KIRs for which appropriate self-HLA antigens exist enable continued inhibition of NK cells preventing autologous cytotoxicity. Hence, NKG2A$^+$inhibitory NK cells (iNK) are key to the prevention of autoreactivity of NK cells, prior to the KIR-driven process of licensing. In a subset of mature and licensed NK cells, exposure to CMV leads to the expression of a C-lectin type activating receptor, NKG2C, which is encoded by the

KLRC2 gene[7]. These cells are characterized by upregulation of NKG2C and

downregulation of the inhibitory counterpart, NKG2A[8]. This subset of NK cells, now

called NKG2C+ANK cells, exhibits the classic adaptive features, such as, clonal expansion,

persistence and recall memory more akin to memory cytotoxic T cells, than canonical NK

cells[9].

While the major subset of ANK cells express NKG2C, which is the defining phenotype,

several myeloid (FCεRG, PLZF) and B lymphoid (SYK, EAT-2) genes are downregulated,

and certain T lymphoid genes (CD3ζ, BCL11B) are upregulated in ANK cells[10]. The

alterations in these gene expressions in ANK cells favour an augmentation of antibody-

dependent cellular cytoxicity (ADCC). In a small subset of ANK cells, the adaptive functions

might be demonstrable with epigenetic distribution of myeloid and lymphoid associated

genes as described above, even without upregulation of NKG2C expression[10; 11]. Hence,

for the sake of clarity, the ANK cells described in this study are NKG2C+ ANK cells.

In the context of haploidentical hematopoietic cell transplantation (HCT), NKG2C+ANK

cells were found to afford protection, not only against leukemia, but also a range of viral

infections, other than CMV. It was suggested that high NKG2C+ ANK cells was essential in

maintaining a non-inflammatory milieu without compromising anti-tumor effect post-

HCT[12; 13; 14; 15]. Based on these seminal findings along with the existing evidence that

certain natural infections, such as Hantavirus, as well as Influenza and BCG vaccinations are

capable of upregulating NKG2C+ANK cells in CMV-exposed populations[16; 17; 18], we

hypothesized that a novel heat killed *Mycobacterium w* (Mw), also known as *Mycobacterium*

indicus pranii, an approved immunomodulator in India [19], might upregulate NKG2C+ANK

cells and offer protection against SARS-CoV-2 (Severe Acute Respiratory Syndrome

Coronavirus-2) infections in the process. We studied the impact of Mw on the incidence of

COVID-19 (Corona Virus Disease 2019 caused by SARS-CoV-2) in a cohort-control study

in front line health care workers and its impact on the kinetics and repertoire of

NKG2C$^+$ANK cells in the context of KLRC2 genotype.

MATERIALS AND METHODS

In a single-center non-randomised cohort control study, 50 HCWs from a single department

in the hospital were administered 0.1 ml *Mw* (Sepsivac, Cadila Pharmaceuticals, India)

intradermally in each arm on day 1 of the study (Mw group) and 50 randomly selected HCWs

from the rest of the institution were enrolled in a Control group. In addition, in the Mw

group, those without any local site reaction, who consented for second dose, were

administered an additional dose of 0.1 ml Mw, 30 days after the first dose. The observation

period of the study was 15 days after administration of the first dose of Mw to 180 days

(September 2020 to February 2021).

All HCW included in the study had nasopharyngeal swab evaluated for SARS-CoV-2 by

reverse transcriptase-polymerase chain reaction (RT-PCR), on development of symptoms

suggestive of COVID-19 or following contact with a patient, HCW or a family member in

contact with COVID-19 patient. Body temperature, pulse rate, oxygen saturation and self-

reporting of symptoms was evaluated before and after each working day[20]. COVID-19 was

diagnosed and its severity was graded as per established criteria. The duration of observation

was 6 months from September 2020 to February 2021. In addition, blood was collected, at

baseline, days 30, 60, and 100 from all the subjects, for evaluating the kinetics of

NKG2C$^+$ANK cells and T cells. All subjects of this study also underwent evaluation for

CMV status (seropositive or seronegative).

All subjects provided written informed consent for participating in the study. The study was

approved by institutional ethics committee and registered with Clinical Trials Registry of

India (CTRI/2020/10/028326).

Real Time Reverse Transcription Polymerase Chain Reaction (RT-PCR) for Detection

of SARS-CoV-2

All Covid tests were done by Truenat Real Time Reverse Transcriptase Polymerase chain

reaction (RT-PCR) test. Nasopharyngeal swabs were collected using standard nylon flocked

swab and inserted into the viral transport medium (VTM) provided from the same company

(Molbio diagnostics Pvt. Ltd. Goa, India). Samples were transported immediately to the

laboratory maintaining proper temperature and processed as per manufacturer's guidelines.

[Truenat Beta CoV Chip-based Real time PCR test for Beta Coronavirus, Molbio diagnostics

Pvt. Ltd. Goa, India]. The target sequence for this assay is E gene of Sarbeco virus and

human RNaseP (internal positive control). Confirmatory genes were RdRP gene and ORF1A

gene.

Immunological monitoring

Peripheral blood mononuclear cells (PBMC) were isolated from whole blood samples, by

density gradient centrifugations using HiSep™ LSM 1077 media. For surface staining, $0.5 \times$

10^6 cells were washed with phosphate-buffered saline (PBS) and stained with the following

antibodies which were used for phenotypic analysis: CD3(APC-H7, SK-7) CD16 (PE-Cy7,

B73.1), CD56 (APC R700, NCAM16.2), CD57 (BV605, NK-1), NKG2A (PE-Cy7, Z199),

CD4 (APC-H7), CD8 (Per-CP Cy), CD45RA (FITC), CD45RO (BV605), from BD

Biosciences, (San Jose, CA) and NKG2C (PE, REA205) from Miltenyi Biotec, Germany.

Cells were then incubated for 30 minutes. Viability was assessed with 7-AAD viability dye

(Beckman Coulter). For intracellular staining, cells were stained for IFN- gamma using

monoclonal antibodies for interferon-gamma (IFN-γ) (4S.B3) and perforin (Alexa647, DG9)

(BD Biosciences) after fixation and permeabilization with appropriate buffer (BD

Biosciences and e-biosciences, San Diego, CA, USA). Flow Cytometry was performed using

10 colour flow cytometry (BD FACS Lyrics) and the flow cytometry data was analyzed using

FlowJo software (v10.6.2, FlowJo). Unstained, single stained (one antibody/sample) as well

as fluorescence-minus-one (FMO) samples were used as controls for the acquisition as well

as the subsequent analysis. Statistical divergences were determined by the GraphPad Prism

software.

The gating strategy has been described earlier[15]. NKG2C$^+$ANK cells were defined as

CD56^{dim}NKG2C$^+$NKG2A$^-$CD57$^+$ subset of NK cells. NKG2C/NKG2A ratio was calculated

as relative % of NKG2C$^+$NKG2A$^-$ ANK cells / NKG2C$^-$NKG2A$^+$ iNK cells (Fig. S4).

KLRC2 (NKG2C) genotyping

KLRC2 gene encodes for NKG2C and is located in chromosome 12p13. For categorizing

subjects KLRC2 wildtype homozygous (*Wt$^+$/ Wt$^+$*), KLRC2 deletion homogyzous *(Del$^+$/*

Del$^+$) and KLRC2 heterozygous (*Wt$^+$/ Del$^+$*), PCR-based KLRC2 genotyping was carried

out. DNA was isolated from the peripheral blood using QIAGEN QIAamp@ DNA blood

mini kit method (Hilden, Germany). PCR amplification was carried out with forward and

reverse primer sequences as previously described[21]. PCR amplification was carried out in

20μl volume, containing 1x PCR master mix which has premixed taq polymerase, dNTPs,

PCR buffer (Thermo fisher scientific, Waltham, MA 02451, United States), 1.65 pmol

forward primers and reverse primers for KLRC2 deletion and wild type KLRC2 genes (see

Table S2), 100pg-1μg genomic DNA. Amplification was performed using a T100 thermal

cycler (Bio-Rad, Hercules, CA). Cycling temperature profiles were adopted from a previous

study[21] with minor modifications.

Briefly, the reaction mixture was subjected to one cycle of denaturation at 95°C for 10 min

followed immediately by 29 cycles of 95°C for 20 s, 50°C for 30 s, 72 $^\circ$C for 40 s; and a final

extension at 72°C for 10 min before cooling to 4°C. A non-template control was included in

each batch of PCR reactions. PCR products were identified by running the entire PCR

product on a 2% agarose gel for 60 minutes at 70 Volts. The size of the amplicons was

determined by comparison against the migration of a 100-bp DNA ladder (GeneDireX, Inc,

Taoyuan County, Taiwan). Agarose gels were visualized and documented using the Gel doc

XR+ gel documentation system (Bio-Rad, Hercules, CA). An illustration of the PCR assay is

shown in supplementary figure (Fig. S5).

RNA-seq Analysis

This was carried out on both Mw and Control group at 6 months following exposure to Mw.

Four subjects from each group without COVID-19 were selected for the study.

RNA isolation using Trizol method and polyA RNA selection:

Peripheral blood mononuclear cells (PBMC) were isolated from whole blood samples by

density gradient centrifugations using HiSep™ LSM 1077 media (Himedia, Mumbai, India).

5×10^6 PBMCs were used for RNA isolation using the Trizol method and followed by DNase

treatment for RNA purification. Purified RNA was quantified using a Qubit 4.0 fluorometer.

Take 5µg of total RNA used for the polyA RNA selection using NEB NEXT oligo d(T)$_{25}$

beads (NEB, MA, USA). PolyA RNA was quantified and integrity assessed using Qubit 4.0

fluorometer (Invitrogen, Waltham, Massachusetts, United States) according to manufacturer

instructions.

cDNA library preparation and whole transcriptome sequencing using MinION 2.0-

Oxford Nanopore Technologies (ONT):

For Direct cDNA Native Barcoding Sequencing (SQK-DCS109 with EXP-NBD104, oxford

nanopore technologies, Oxford, UK), 100ng of polyA RNA was used for the library

preparation. Using H minus Reverse Transcriptase (Invitrogen, Waltham, Massachusetts,

United States), complementary DNA (cDNA) was synthesised, followed by RNA

degradation and second-strand synthesis of cDNA. Double-stranded cDNA was used for the

end preparation, followed by native barcoding and adaptor ligation (all steps were followed,

according to the manufacturer's instructions). Ligated cDNA was loaded on the flow cell

(R9) in MinION Libraries were sequenced specifying 72 hours on the ONT MinION using

R9.4.1 flow cells and MinKNOW (v21.06.10, Microsoft Windows OS based) to generate

FAST5 files. FAST5 files were base-called with CPU based Guppy basecaller (v.5.0.11)

(ONT) to create summary text files and FASTQ files of the reads for further downstream

analysis.

Sample pre-processing and Quality assessment:

Demultiplexing of the pooled samples and adapter removal was carried out using inbuilt

algorithm of Minknow. Linux Long Time Support (v.20.04) Operating System was used for

all the analysis. A comprehensive report of the sequencing was generated by NanoComp

(https://github.com/NanoComp/h5utils) tool and sequencing quality assessment was done by

FastQc tool (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Sample reads

were subjected to a minimum phred quality score of 9 and reads with lower quality were

filtered out using NanoFilt (https://github.com/wdecoster/nanofilt) tool. Some initial reads are

usually prone to low base-calling quality. Hence, 50bp of each initial reads from every

sample were filtered out for quality maintenance using NanoFilt tool only. All the samples

were subjected to quality assessment before and after quality filtering.

Differential Gene Expression analysis:

Differential Gene Expression analysis (DGE) was done using "pipeline-transcriptome-de"

(https://github.com/nanoporetech/pipeline-transcriptome-de) pipeline. This pipeline from

nanopore tech uses snakemake, minimap2, salmon, edgeR, DEXSeq and stageR to automate

DGE workflows on long read data. Pipeline was set to make only reads aligned to minimum

3 samples to be considered for analysis. A separate conda (https://docs.conda.io/en/latest/#)

environment was created on Linux OS to host this pipeline. Quantification and DGE was

done by R language based (https://www.r-project.org/) tool edgeR

(https://bioconductor.org/packages/release/bioc/html/edgeR.html) employing gene-wise

negative binomial regression model and normalisation factor (Transcript Mean of M-value)

for each sequence library. Differentially expressed genes (DEGs) with log2Fold Change

(log2Fc) ≥ 0.5, ≤ -0.5 and associated p-value <0.05 were selected as significant for further

analysis. Annotation of DEGs was fetched from ENSEMBL database

(https://asia.ensembl.org/index.html) using R language biomaRt package

(https://bioconductor.org/packages/release/bioc/html/biomaRt.html). All the file compilation

was ultimately done using Microsoft Excel and Libre Office calc.

Hierarchical clustering analyses was performed between all four samples of each group to

generate Heatmap from normalised log2 counts per million expression values using web-

based START(Shiny Transcriptome Analysis Resource Tool,

https://kcvi.shinyapps.io/START/) tool. Comparative gene expression boxplot on the basis of

log2 Counts per million between Mw and Control group were generated from START tool.

Volcano plot involving all DEGs were created using R language based ggplot2

(https://ggplot2.tidyverse.org/) package.

Gene Ontology (GO) Pathway Analysis:

Pathway enrichment analysis was done for significant DEGs using g:Profiler

(https://biit.cs.ut.ee/gprofiler/gost) using Gene Ontology. Pathways related to adaptive

Natural killer (ANK), ADCC and innate immune inflammatory pathways were considered for

focussed analysis. Functional enrichment was done by R based clusterProfiler (v. 4.2.1)

https://bioconductor.org/packages/release/bioc/html/clusterProfiler.html tool. An adjusted p-

value threshold of ≤0.05 was considered for this study.

Statistics

Lymphocyte subsets have been represented as % of the parent population. Binary variables

were compared between the groups using chi square test. The continuous variables were

analysed using independent sample t-test considering Levene's test for equality of variances

and non-parametric tests (Mann-Whitney U test). P value < 0.05 was considered to be

significant. GraphPad Prism (version 8.0 for Windows, GraphPad Software, La Jolla, CA,

USA) was used for the statistical assessment (unpaired low-parametric Mann–Whitney or

Kruskal–Wallis test and Spearman correlation). Recursive partitioning analysis was carried

out using *rpart* package (https://cran.r-project.org/web/packages/rpart/index.html) in R

(https://cran.r-project.org/) to generate optimal cut off for ANK cells at baseline.

The efficacy of Mw in reducing the incidence of COVID-19 was calculated in terms of attack

rates incidence risk ratio (IRR), absolute risk reduction (ARR) and intervention efficacy (see

Table S3). This was calculated in terms of infections occurring at 2 weeks after

administration of Mw is customary in prophylactic studies as well as those occurring any

time between during the study period.

RESULTS

Mw and COVID-19

The characteristics and outcomes of the Mw and control groups are detailed in Table 1.

All subjects were seropositive for CMV.

Twenty (3 in Mw group vs 17 in the control group) out of 100 subjects in the entire cohort

had symptomatic COVID-19 during the study period (15 days-6 months). All infections on

the Mw group were mild, while 6 of 17 in control group had moderate disease and required

hospital admission (p=0.01). No deaths were recorded in either group. In Mw group all three

infections occurred beyond 150 days. In the control group, 17 subjects had COVID-19 at a

median of 56 days (range, 18–135). Additionally, three mild symptomatic COVID-19 were

documented within the first 7 days of Mw administration. The incidence rates (IR) of

COVID-19 were 3.57 and 24.05 /10,000 person-days in Mw and control groups respectively

(IRR-0.15, 95% CI, 0.04-0.5, p=0.0004) with an efficacy rate for Mw being 82.35%, (95%

CI, 50-96 %) (Fig. 1).

Twenty-four subjects in the Mw group were randomised 1:1 to receive the second dose, 30

days after the first dose. Impact of the second dose on protection could not be analysed as

none of the randomised subjects developed COVID-19 during the study period.

Safety Profile of Mw

Mw was safe with no systemic adverse effects. Only one subject had mild fever (37.5º C) for

less than two hours, 24 hours after administration, which was self-limiting. However, 14%

had pain at the local sites, with 12% developing pain and erythema lasting for more than 72

hours. In 8% of the subjects, ulceration was noted at the local site with long-term scars

similar to that observed with BCG vaccination. Those with severe local reactions were all

older (35 years and above), compared to the median age of the group (28 years).

Baseline NKG2C$^+$ANK cells correlated with increased risk of COVID-19 in both Mw

and Control groups

Baseline data on immune parameters was available in 80 subjects in the overall cohort, 30 in

control and 50 in Mw group. The NKG2C$^+$ANK cells in those who got infected (n=16) was

7.9 ± 6.1, (includes 3 who got infected within 7 days of receiving Mw) compared to 20.9 ±

13.8 in those who remained uninfected (p= 0.0005).

Amongst the 30 subjects in the control group, who were tested at baseline for NKG2C$^+$ANK

cells, 10 developed COVID-19. The NKG2C$^+$ANK cells at baseline was 9.7 ± 6.6 in those

with infection compared to 17.7 ± 5.56 in those without infection (p=0.0016). Those who

acquired the infection in the Mw group, also had a lower baseline NKG2C$^+$ANK (4.76 ± 3.9

vs 22.3 ± 16.1, Fig. 5, p=0.01). Interestingly, 2 of the 3 subjects, experiencing COVID-19 in

the Mw group, did not show any increment in NKG2C$^+$ANK levels or NKG2C/NKG2A

ratios at any of the time-points. We did not find any correlation between expression of

NKG2A as well as NKG2C/NKG2A ratio with development of symptomatic COVID-19 in

this study.

Mw and ANK cells

Thirty subjects from the Mw group and 15 from the control group, who did not develop

COVID-19 or any other significant infection during the study period were included in the

final longitudinal analysis of immunological impact of Mw.

Baseline ANK levels did not differ, but ANK Kinetics at 60 days were different in the

Mw group

There was no difference in overall NK cells between the two groups at baseline or at

subsequent time-points. NKG2C$^+$ANK and NKG2A$^+$ iNK cells expression were also similar

at baseline in both groups (Fig. 2, A & B). The data from control group shown in further

comparisons only pertain to the baseline and day 60, as the values in the control group for

both NKG2C$^+$ ANK cells and NKG2A$^+$ iNK cells did not change significantly at any of the

time-points.

In the Mw group on the other hand, (Fig. 2, A- D), NKG2C$^+$ANK cells increased from

baseline (23.0 ± 22.9) to peak at day 60 (40.9 ± 16.5, day 60 p=0.0009, Fig. 2 C), while

NKG2A$^+$ iNK cells reduced from 36.23 ± 14.4 to 26.22 ± 14.04 on day 30 and 6.17 ± 6.01 on

day 60 (p<0.0001, Fig. 2 D). The absolute values as well as the log two-fold change (log2FC)

were significantly different for both NKG2C$^+$ANK cells (1.5 ± 1.9 vs -0.62 ± 0.89, p=0.0002,

Fig. 2 E) as well as NKG2A$^+$ iNK cells (-2.9 ± 1.2 vs 0.17 ± 0.29, p<0.0001, Fig. 2 E) cells at

day 60, compared to the control group.

The impact of Mw on NKG2C upregulation was only observed in those with low ANK

cells at baseline

A cut-off of 15% was established for NKG2C$^+$ANK cells based on recursive partitioning for

evaluating effect of base line value on NKG2C upregulation. Upregulation of NKG2C$^+$ ANK

cells was seen in those with NKG2C$^+$ ANK levels below 15% (Low ANK group). From a

baseline value of 4.7 ± 2.2, this increased to 8.9 ± 4.9 on day 30 (p=0.005,) and 43.3 ± 19.12

on day 60 (p<0.0001, Fig. 3 A). This was also reflected in the log2FC values.

In those with NKG2C$^+$ANK levels above 15% (High NKG2C$^+$ANK), there was no

significant change in absolute values of NKG2C$^+$ANK cells at either day 30 or day 60(Fig. 3

B), but a positive impact on log2FC was noted (p=0.04, Fig. 3 E).

The impact on NKG2A⁺iNK downregulation was irrespective of baseline NKG2C⁺ANK expression

The downregulation of NKG2A⁺iNK cells was noted in the Mw group, irrespective of baseline NKG2C expression [Fig. 3, C & D]. The downregulation of NKG2A⁺iNK cells was significant at day 60 in the Mw group, irrespective of baseline NKG2C expression (-2.99 ± 1.21 vs 0.23 ± 0.23, p<0.0001 for Low ANK group and -2.84 ± 1.28 vs 0.09 ± 0.35, p<0.0001 for High ANK group) [Fig. 3 F].

Mw had a sustained effect on NKG2C+ANK/ NKG2A+iNK ratio until 100 days

NKG2C⁺ ANK/NKG2A⁺ iNK ratio was similar between two groups at baseline. However, this increased in the Mw group at day 60 (12.3 ± 9.2, p<0.0001) and persisted until day 100 (7.44 ± 13.55, p=0.01). This was witnessed for both Low and High ANK group in the Mw cohort but not in the control cohort. [Fig. 4, A- D].

IFN-γ release was higher in Mw group at 60 days

IFN-γ release potential of the NKG2C⁺ANK cells was studied at baseline and at 60 days. This was similar between the groups at the baseline (mean-6.7 vs 6.4). However, this was significantly increased in the Mw group at day 60, compared to the control group (mean-27.96 vs 9.9, p=0.01, Fig. 4 E).

KLRC2 and kinetics of ANK cells

KLRC2 deletion was studied in the Mw group only. KLRC2 *Wt ⁺/del⁺* was detected in 36% of the subjects in the Mw group. This was not associated with any significant decrease in the baseline NKG2C⁺ANK levels (15.1 ± 17.67 vs 20.4 ± 17.96, p=0.46). There was no difference in the effect of Mw between the KLRC2 *Wt⁺/ Wt⁺* and W*t⁺/Del⁺* groups at day 60 either (p=0.26), although the log2FC increase tended to be higher in the W*t⁺/Del⁺* group

(p=0.12) [Fig. S1, A & B]. KLRC2 deletion was detected in 2 out of 6 with COVID-19,

compared to 16 out of 44 without COVID-19 (p=0.6).

Mw did not influence kinetics of naïve and memory T cell subsets

CD4 and CD8 cells remained unchanged, as were the CD45RA and CD45RO subsets at days

30 and 60 in the Mw group [Fig. S2, A & B]. The same was witnessed in the control group

(data not shown).

NKG2A expression on CD3$^+$ T cells was examined at baseline and subsequent timelines.

Despite a significant downregulation of NKG2A observed in NK cells, no such effect was

evident in the T cells. [Fig. S2 C]

Impact of the second dose of Mw on NKG2C$^+$ANK cells

There was no difference between in NKG2C$^+$ANK cells at day 60 and day 100 (p=0.28) in

double vs single doses of Mw (Fig. 6 A). However, NKG2A$^+$ iNK cell expression was

significantly reduced at day 100 (7.7 ± 16.77 vs 24.85 ± 19.43, p=0.04) in double dose group

(Fig. 6 B). The NKG2C/NKG2A ratio was also significantly higher in double dose group at

day 100 (10.42 ± 7.19 vs 2.08 ± 3.26, p <0.001) [Fig. 6 C]. There was no difference in CD4$^+$

& CD8$^+$ T cells and memory or naïve subsets at day 60 and 100 between the randomized

groups (data not shown).

RNAseq Analysis at 6 months

Sequencing and mapping metrics along with Quality scores are shown in Supplement 1

(Table S1)

Identification of differentially expressed genes

Based on criteria of p-value < 0.05, a total of 3,603 out of 16,058 were found to be

differentially expressed genes (DEGs) between Mw and Control group, including 1568 genes

that were upregulated and 2035 genes that were downregulated in Mw group. Those DEGs

with a log2 fold change of at least 0.5 were considered for further analysis. DEGs related to

392 Adaptive Natural Killer Cells and ADCC were selected out to analyse the differential

393 expression pattern between two groups.

394 **ANK and ADCC genes were upregulated in Mw group**

395 Nineteen DEGs were associated with ANK and ADCC pathways (Fig. 7, A & B). 11 genes

396 were upregulated and 8 were found to be downregulated in Mw group. KLRC2 (NKG2C),

397 BCL11B, ARID5B, B3GAT1 (CD57), KLRC4 were upregulated and KLRC1(NKG2A),

398 ZBTB16 (PLZF1), KIT and SH2D1B (EAT-2) were downregulated in relation to ANK

399 pathway (Fig. 7 C). In the ADCC pathway, CD247 (CD3ζ), FCGR1A (CD64),

400 FCGR2A(CD32a), FCGR2C (CD32c) were upregulated and FCER1G was downregulated

401 (Fig. 7 C), favouring ANK mediated ADCC.

402 **Gene Ontology (GO) Pathway Analysis**

403 Functional enrichment of differentially expressed genes were carried out on the selected

404 genes focusing on ANK and ADCC pathways. Selected 19 DEGs were analysed using

405 goProfiler and clusterProfiler for Gene Ontology (GO). GO analysis of the selected DEGs

406 demonstrated that immune response-regulating cell surface receptor signalling pathway

407 (GO:0002768) and positive regulation of natural killer cells mediated toxicity (GO:0045954)

408 were significantly upregulated (Fig. 7 D).

410 **DISCUSSION**

411 The identification of NK cells with ability to persist and mount recall responses has

412 challenged the traditional compartmentalization of adaptive and immune responses[22; 23].

413 The NKG2C[+]ANK cells explored in this study are known to be induced by primary exposure

414 to CMV[24; 25]. However, in CMV exposed patients, these NKG2C[+]ANK cells have been

415 found to proliferate in response to a variety of viral pathogens, including HIV, HCV,

416 Hantavirus, influenza, and pox viruses[16; 26; 27; 28; 29]. The unique nature of this non-

antigen specific recall response against a wide variety of pathogens, puts NKG2C[+]ANK cells

at the forefront in the battle against novel pathogens such as SARS-CoV-2 [3; 30; 31; 32; 33;

34], where antigen-specific adaptive response generated from T and B cells are absent at the

outset of the pandemic.

BCG vaccine has been shown to induce long-term heterologous memory response of NK

cells[31]. This has been studied, both in the context of infections as well as cancer[35].

Influenza vaccination has been shown to induce proliferation of NKG2C[+]ANK cells with

potential for enhanced cytokine release[36]. These considerations prompted us to explore this

novel immunomodulator, Mw, which has been extensively studied as adjuvant or prophylaxis

in other infections[19; 37; 38], and some forms of cancers including BCG resistant bladder

cancer [39].

Our study demonstrated that prophylaxis with Mw was associated with a six-fold reduction in

the incidence of symptomatic COVID-19 in this high-risk cohort over a 6-month period. This

protective efficacy was accompanied by a sharp increase in NKG2C[+]ANK cells between 30-

and 60-days following exposure to Mw. A critical observation was a steep decline in the

expression of the inhibitory counterpart, NKG2A[+]iNK cells, which continued through 100

days. This resulted in an increasing NKG2C/NKG2A ratio through the study period.

Exposure to a second dose of Mw resulted in a continued downregulation of NKG2A[+] iNK

cell and a further increase in NKG2C/NKG2A ratios. The importance of the relative

proportion of NKG2C and NKG2A expressing NK cells could be appreciated better in the

context of the fact that the inhibitory impact of NKG2A overrides that of NKG2C, as it binds

to the ligand HLA-E with several fold greater affinity than NKG2C[5]. In the context of

SARS-CoV-2 infection, in-vitro studies have demonstrated that the viral spike protein-1 (SP-

1) upregulates NKG2A on NK cells and HLA-E on the infected lung epithelial cells, causing

a strong inhibition of cytotoxicity of NK cells[40]. Our group has demonstrated a correlation

between high expression of NKG2A, suppression of NKG2C, and adverse outcome following

severe COVID-19 lung disease, despite viral clearance[34]. It is possible that the adverse

effect of KLRC2 deletion genotype as reported earlier[30], was mitigated by Mw. Hence, it

might be inferred that if any intervention can achieve the reverse, i.e., downregulation of

NKG2A and upregulation of NKG2C expression, this might offer innate protection against

COVID-19.

In our study, a lower baseline NKG2C$^+$ANK cells predisposed to COVID-19 in the control

group. The same in the Mw group was associated with COVID-19 in the first 2 weeks, but

not thereafter until 150 days. It is also worth noting that COVID-19 occurring in the Mw

group beyond 150 days was associated with failure in modulation of the adverse

NKG2C$^+$ANK cell profile in 2 out of 3 subjects. Apart from phenotypic alteration, Mw

seemed to influence the cytokine release potential as well, as evidenced by an increase in

IFN-γ release at day 60 in the Mw group, compared to both baseline as well as the control

group. Furthermore, upregulation of NKG2C was dominantly noted in those with a lower

baseline NKG2C$^+$ANK cell. Upregulation of NKG2C expression might be more tightly

controlled, unlike downregulation of NKG2A, in mature and licensed NK cells, where an

inhibitory regulation mediated by NKG2A is physiologically redundant due to KIR-mediated

inhibitory control. More importantly, this also demonstrates that downregulation of

checkpoint receptor NKG2A is an important determinant of a favorable ANK profile.

RNA-seq analysis confirmed that upregulation of ANK pathway was evident at 6 months in

the Mw group. Apart from upregulation of KLRC2 and B3GAT1 and downregulation of

KLRC1, the key transcription factor in the ANK pathway, BCL11b, was persistently

upregulated[41]. Downregulation of EAT-2 and PLZF further corroborated the classic gene

expression signature of ANK cells. Moreover, increased expression of AT-rich interaction

domain 5B (ARID5B), as demonstrated in the Mw group plays an important role in enhanced

metabolism in ANK cells as well as increased IFN-γ release and survival[42]. DGE analysis

also revealed an enhancement of ANK mediated ADCC pathway, with significant

upregulation of CD247 along with downregulation of FCER1G, which is a typical signature

of ANK-ADCC[43]. Both CD247 and FCER1G are adapter molecules for FCGRIIIA (CD16)

with CD247 possessing 3 ITAMs against one ITAM of FCER1G, increasing the ADCC

several folds[44]. It is possible that Mw induced augmentation of NK-ADCC might

potentiate the efficacy of SARS-CoV2 vaccines as well[45]. Prior studies on Mw have

explored its effect on TLRs via the monocyte/macrophage pathway[46; 47; 48] and not on

NK cells. The suggested mechanistic pathway as to how Mw might be favorably influencing

ANK mediated protection against COVID-19 has been depicted in Fig. 8.

In conclusion, our study shows that Mw offered protection against COVID-19 in a high-risk

population at the peak of the pandemic in the absence of antigen-specific immunity. This

could be through a sustained upregulation of NKG2C$^+$ANK cell and ADCC pathway, along

with simultaneous downregulation of NKG2A$^+$iNK pathways. If borne out in a randomized

setting, these findings might usher a novel approach to bolster heterologous immunity in the

current and future pandemics and identify ANK pathway in the immunological vulnerability

for developing COVID-19.

Acknowledgement: We thank all the nursing and technical staff of the department who have assisted in carrying out this study. This study was supported by a grant from Indo-US Science and Technology Forum (IUSSTF/VN-COVID/049/2020).

Authorship Contribution: SRJ, BK and SC designed the study. AM, RL, GB and HM performed the study. SRJ, AS, AM, GB and HM collected the data. SRJ, AS, JA, SVM, DT and SC analyzed the data; SRJ and SC wrote the manuscript. All the co-authors reviewed and approved the manuscript.

Conflicts of Interest Statement: BK and SVM are employed by Cadila Pharmaceuticals Ltd. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding: This study was supported by grant from Indo-US Science and Technology Forum (IUSSTF/VN-COVID/049/2020).

Data Availability Statement: The RNA-seq data generated from the experiment related to study have been uploaded in 'Sequence Read Archive' (SRA)

REFERENCE

[1] D. Blumenthal, E.J. Fowler, M. Abrams, and S.R. Collins, Covid-19 - Implications for the Health Care System. N Engl J Med 383 (2020) 1483-1488.

[2] J. Kaushal, and P. Mahajan, Asia's largest urban slum-Dharavi: A global model for management of COVID-19. Cities 111 (2021) 103097.

[3] S.R. Jaiswal, P. Malhotra, D.K. Mitra, and S. Chakrabarti, Focusing On A Unique Innate Memory Cell Population Of Natural Killer Cells In The Fight Against COVID-19: Harnessing The Ubiquity Of Cytomegalovirus Exposure. Mediterr J Hematol Infect Dis 12 (2020) e2020047.

[4] L. Moretta, F. Locatelli, D. Pende, S. Sivori, M. Falco, C. Bottino, M.C. Mingari, and A. Moretta, Human NK receptors: from the molecules to the therapy of high risk leukemias. FEBS Lett 585 (2011) 1563-7.

[5] V. Beziat, B. Hervier, A. Achour, D. Boutolleau, A. Marfain-Koka, and V. Vieillard, Human NKG2A overrides NKG2C effector functions to prevent autoreactivity of NK cells. Blood 117 (2011) 4394-6.

[6] V. Beziat, B. Descours, C. Parizot, P. Debre, and V. Vieillard, NK cell terminal differentiation: correlated stepwise decrease of NKG2A and acquisition of KIRs. PLoS. One 5 (2010) e11966.

[7] S. Lopez-Verges, J.M. Milush, B.S. Schwartz, M.J. Pando, J. Jarjoura, V.A. York, J.P. Houchins, S. Miller, S.M. Kang, P.J. Norris, D.F. Nixon, and L.L. Lanier, Expansion of a unique CD57(+)NKG2Chi natural killer cell subset during acute human cytomegalovirus infection. Proc Natl Acad Sci U S A 108 (2011) 14725-32.

[8] M. Della Chiesa, S. Sivori, S. Carlomagno, L. Moretta, and A. Moretta, Activating KIRs and NKG2C in Viral Infections: Toward NK Cell Memory? Front Immunol 6 (2015) 573.

[9] J.C. Sun, and L.L. Lanier, Natural killer cells remember: an evolutionary bridge between innate and adaptive immunity? Eur J Immunol 39 (2009) 2059-64.

[10] H. Schlums, F. Cichocki, B. Tesi, J. Theorell, V. Beziat, T.D. Holmes, H. Han, S.C. Chiang, B. Foley, K. Mattsson, S. Larsson, M. Schaffer, K.J. Malmberg, H.G. Ljunggren, J.S. Miller, and Y.T. Bryceson, Cytomegalovirus infection drives adaptive epigenetic diversification of NK cells with altered signaling and effector function. Immunity 42 (2015) 443-56.

[11] L. Muccio, M. Falco, A. Bertaina, F. Locatelli, F. Frassoni, S. Sivori, L. Moretta, A. Moretta, and M. Della Chiesa, Late Development of FcepsilonRgamma(neg) Adaptive Natural Killer Cells Upon Human Cytomegalovirus Reactivation in Umbilical Cord Blood Transplantation Recipients. Front Immunol 9 (2018) 1050.

[12] S.R. Jaiswal, P. Bhakuni, G. Bhagawati, H.M. Aiyer, M. Soni, N. Sharma, R.R. Jaiswal, A. Chakrabarti, and S. Chakrabarti, CTLA4Ig-primed donor lymphocyte infusions following haploidentical transplantation improve outcome with a distinct pattern of early immune reconstitution as compared to conventional donor lymphocyte infusions in advanced hematological malignancies. Bone Marrow Transplant (2020).

[13] S.R. Jaiswal, P. Bhakuni, G. Bhagwati, H.M. Aiyar, A. Chakrabarti, and S. Chakrabarti, Alterations in NKG2A and NKG2C Subsets of Natural Killer Cells Following Epstein-Barr Virus Reactivation in CTLA4Ig-based Haploidentical Transplantation Is Associated With Increased Chronic Graft-Versus-Host Disease. Transplantation 104 (2020) e23-e30.

[14] S.R. Jaiswal, and S. Chakrabarti, Natural killer cell-based immunotherapy with CTLA4Ig-primed donor lymphocytes following haploidentical transplantation. Immunotherapy 11 (2019) 1221-1230.

[15] S.R. Jaiswal, S. Chakraborty, R. Lakhchaura, P. Shashi, A. Mehta, M. Soni, and S. Chakrabarti, Early and Sustained Expansion of Adaptive Natural Killer Cells Following Haploidentical Transplantation and CTLA4Ig-Primed Donor Lymphocyte Infusions Dissociate Graft-versus-Leukemia and Graft-versus-Host Effects. Transplantation and Cellular Therapy 27 (2021) 144-151.

[16] N.K. Bjorkstrom, T. Lindgren, M. Stoltz, C. Fauriat, M. Braun, M. Evander, J. Michaelsson, K.J. Malmberg, J. Klingstrom, C. Ahlm, and H.G. Ljunggren, Rapid expansion and long-term persistence of elevated NK cell numbers in humans infected with hantavirus. J Exp Med 208 (2011) 13-21.

[17] J.C. Sun, and L.L. Lanier, Is There Natural Killer Cell Memory and Can It Be Harnessed by Vaccination? NK Cell Memory and Immunization Strategies against Infectious Diseases and Cancer. Cold Spring Harb Perspect Biol 10 (2018).

[18] F. Abebe, Immunological basis of early clearance of Mycobacterium tuberculosis infection: the role of natural killer cells. Clin Exp Immunol 204 (2021) 32-40.

[19] I.S. Sehgal, R. Agarwal, A.N. Aggarwal, and S.K. Jindal, A randomized trial of Mycobacterium w in severe sepsis. J Crit Care 30 (2015) 85-9.

[20] Z. Wu, and J.M. McGoogan, Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. JAMA (2020).

[21] R. Miyashita, N. Tsuchiya, K. Hikami, K. Kuroki, T. Fukazawa, M. Bijl, C.G. Kallenberg, H. Hashimoto, T. Yabe, and K. Tokunaga, Molecular genetic analyses of human NKG2C (KLRC2) gene deletion. Int Immunol 16 (2004) 163-8.

[22] J.G. O'Leary, M. Goodarzi, D.L. Drayton, and U.H. von Andrian, T cell- and B cell-independent adaptive immunity mediated by natural killer cells. Nat Immunol 7 (2006) 507-16.

[23] L.L. Lanier, Shades of grey — the blurring view of innate and adaptive immunity. Nature Reviews Immunology 13 (2013) 73-74.

[24] F. Cichocki, S. Cooley, Z. Davis, T.E. DeFor, H. Schlums, B. Zhang, C.G. Brunstein, B.R. Blazar, J. Wagner, D.J. Diamond, M.R. Verneris, Y.T. Bryceson, D.J. Weisdorf, and J.S. Miller, CD56dimCD57+NKG2C+ NK cell expansion is associated with reduced leukemia relapse after reduced intensity HCT. Leukemia 30 (2016) 456-63.

[25] A. Rolle, M. Meyer, S. Calderazzo, D. Jager, and F. Momburg, Distinct HLA-E Peptide Complexes Modify Antibody-Driven Effector Functions of Adaptive NK Cells. Cell Rep 24 (2018) 1967-1976 e4.

[26] J. Nattermann, H.D. Nischalke, V. Hofmeister, B. Kupfer, G. Ahlenstiel, G. Feldmann, J. Rockstroh, E.H. Weiss, T. Sauerbruch, and U. Spengler, HIV-1 infection leads to increased HLA-E expression resulting in impaired function of natural killer cells. Antivir Ther 10 (2005) 95-107.

[27] R. Thomas, H.Z. Low, K. Kniesch, R. Jacobs, R.E. Schmidt, and T. Witte, NKG2C deletion is a risk factor of HIV infection. AIDS Res Hum Retroviruses 28 (2012) 844-51.

[28] R. Zhang, J. Xu, K. Hong, L. Yuan, H. Peng, H. Tang, P. Ma, Y. Zhang, H. Xing, Y.
Ruan, and Y. Shao, Increased NKG2A found in cytotoxic natural killer subset in
HIV-1 patients with advanced clinical status. AIDS 21 Suppl 8 (2007) S9-17.
[29] D. Mele, B. Oliviero, S. Mantovani, S. Ludovisi, A. Lombardi, F. Genco, R. Gulminetti,
S. Novati, M.U. Mondelli, and S. Varchetta, Adaptive Natural Killer Cell Functional
Recovery in Hepatitis C Virus Cured Patients. Hepatology 73 (2021) 79-90.
[30] H. Vietzen, A. Zoufaly, M. Traugott, J. Aberle, S.W. Aberle, and E. Puchhammer-
Stockl, Deletion of the NKG2C receptor encoding KLRC2 gene and HLA-E variants
are risk factors for severe COVID-19. Genet Med 23 (2021) 963-967.
[31] C. Covian, A. Fernandez-Fierro, A. Retamal-Diaz, F.E. Diaz, A.E. Vasquez, M.K. Lay,
C.A. Riedel, P.A. Gonzalez, S.M. Bueno, and A.M. Kalergis, BCG-Induced Cross-
Protection and Development of Trained Immunity: Implication for Vaccine Design.
Front Immunol 10 (2019) 2806.
[32] G.T. Hart, T.M. Tran, J. Theorell, H. Schlums, G. Arora, S. Rajagopalan, A.D.J.
Sangala, K.J. Welsh, B. Traore, S.K. Pierce, P.D. Crompton, Y.T. Bryceson, and E.O.
Long, Adaptive NK cells in people exposed to Plasmodium falciparum correlate with
protection from malaria. J Exp Med 216 (2019) 1280-1290.
[33] J. Kleinnijenhuis, J. Quintin, F. Preijers, L.A. Joosten, C. Jacobs, R.J. Xavier, J.W. van
der Meer, R. van Crevel, and M.G. Netea, BCG-induced trained immunity in NK
cells: Role for non-specific protection to infection. Clin Immunol 155 (2014) 213-9.
[34] S.R. Jaiswal, J. Arunachalam, A. Bhardwaj, A. Saifullah, R. Lakhchaura, M. Soni, G.
Bhagawati, and S. Chakrabarti, Impact of adaptive natural killer cells, KLRC2
genotype and cytomegalovirus reactivation on late mortality in patients with severe
COVID-19 lung disease. Clin Transl Immunology 11 (2022) e1359.
[35] G. Esteso, N. Aguilo, E. Julian, O. Ashiru, M.M. Ho, C. Martin, and M. Vales-Gomez,
Natural Killer Anti-Tumor Activity Can Be Achieved by In Vitro Incubation With
Heat-Killed BCG. Front Immunol 12 (2021) 622995.
[36] P. Riese, S. Trittel, R.D. Pathirana, F. Klawonn, R.J. Cox, and C.A. Guzman,
Responsiveness to Influenza Vaccination Correlates with NKG2C-Expression on NK
Cells. Vaccines (Basel) 8 (2020).
[37] R. Kamal, V. Pathak, A. Kumari, M. Natrajan, K. Katoch, and H.K. Kar, Addition of
Mycobacterium indicus pranii vaccine as an immunotherapeutic to standard
chemotherapy in borderline leprosy: a double-blind study to assess clinical
improvement (preliminary report). Br J Dermatol 176 (2017) 1388-1389.
[38] A. Gupta, F.J. Ahmad, F. Ahmad, U.D. Gupta, M. Natarajan, V. Katoch, and S. Bhaskar,
Efficacy of Mycobacterium indicus pranii immunotherapy as an adjunct to
chemotherapy for tuberculosis and underlying immune responses in the lung. PLoS
One 7 (2012) e39215.
[39] M. Subramaniam, L.L. In, A. Kumar, N. Ahmed, and N.H. Nagoor, Cytotoxic and
apoptotic effects of heat killed Mycobacterium indicus pranii (MIP) on various human
cancer cell lines. Sci Rep 6 (2016) 19833.
[40] D. Bortolotti, V. Gentili, S. Rizzo, A. Rotola, and R. Rizzo, SARS-CoV-2 Spike 1
Protein Controls Natural Killer Cell Activation via the HLA-E/NKG2A Pathway.
Cells 9 (2020).
[41] T.D. Holmes, R.V. Pandey, E.Y. Helm, H. Schlums, H. Han, T.M. Campbell, T.T.
Drashansky, S. Chiang, C.Y. Wu, C. Tao, M. Shoukier, E. Tolosa, S. Von
Hardenberg, M. Sun, C. Klemann, R.A. Marsh, C.M. Lau, Y. Lin, J.C. Sun, R.
Mansson, F. Cichocki, D. Avram, and Y.T. Bryceson, The transcription factor Bcl11b
promotes both canonical and adaptive NK cell differentiation. Sci Immunol 6 (2021).

[42] F. Cichocki, C.Y. Wu, B. Zhang, M. Felices, B. Tesi, K. Tuininga, P. Dougherty, E. Taras, P. Hinderlie, B.R. Blazar, Y.T. Bryceson, and J.S. Miller, ARID5B regulates metabolic programming in human adaptive NK cells. J Exp Med 215 (2018) 2379-2395.

[43] I. Hwang, T. Zhang, J.M. Scott, A.R. Kim, T. Lee, T. Kakarla, A. Kim, J.B. Sunwoo, and S. Kim, Identification of human NK cells that are deficient for signaling adaptor FcRgamma and specialized for antibody-dependent immune functions. Int Immunol 24 (2012) 793-802.

[44] W. Liu, J.M. Scott, E. Langguth, H. Chang, P.H. Park, and S. Kim, FcRgamma Gene Editing Reprograms Conventional NK Cells to Display Key Features of Adaptive Human NK Cells. iScience 23 (2020) 101709.

[45] X. Chen, C.A. Rostad, L.J. Anderson, H.Y. Sun, S.A. Lapp, K. Stephens, L. Hussaini, T. Gibson, N. Rouphael, and E.J. Anderson, The development and kinetics of functional antibody-dependent cell-mediated cytotoxicity (ADCC) to SARS-CoV-2 spike protein. Virology 559 (2021) 1-9.

[46] R.K. Pandey, A. Sodhi, S.K. Biswas, Y. Dahiya, and M.K. Dhillon, Mycobacterium indicus pranii mediates macrophage activation through TLR2 and NOD2 in a MyD88 dependent manner. Vaccine 30 (2012) 5748-54.

[47] P. Kumar, V. John, S. Marathe, G. Das, and S. Bhaskar, Mycobacterium indicus pranii induces dendritic cell activation, survival, and Th1/Th17 polarization potential in a TLR-dependent manner. J Leukoc Biol 97 (2015) 511-20.

[48] S. Das, B.P. Chowdhury, A. Goswami, S. Parveen, J. Jawed, N. Pal, and S. Majumdar, Mycobacterium indicus pranii (MIP) mediated host protective intracellular mechanisms against tuberculosis infection: Involvement of TLR-4 mediated signaling. Tuberculosis (Edinb) 101 (2016) 201-209.

 LEGEND OF FIGURES

Figure 1: Impact of Mw vaccination on COVID-19 compared to a control group: Points

and connecting line plot show the infection trend in Mw group (n=50) and the Control group

(n=50). The x- axis shows the time in weeks and y- axis shows the Cumulative Incidence (CI)

in %. Red and black shaded circles represent Mw group and Control group respectively.

Figure 2: Impact of Mw on both Adaptive NK (ANK) cells and inhibitory NK (iNK)

cells:

A) Scatter dot with bar plot showing the expression of NKG2C$^+$ ANK cells (Mw group -

n=30 and control group- n=15) and B) NKG2A$^+$ iNK cells (Mw group-n=30 and control

group- n=15) with and without of Mw vaccine at baseline and day 60. C) Scatter dot with bar

plot showing kinetics of NKG2C$^+$ ANK cells, and D) NKG2A$^+$ iNK cells expression in Mw

group (n=30) at Baseline, day 30, day 60 and day 100. E) log2FC expression of NKG2C$^+$ and

NKG2A$^+$ in both Mw group (n=30) and Control group (n=15) at day 60 after normalization

with baseline. Red upside and downside shaded triangles represent NKG2C$^+$ ANK and

NKG2A$^+$ iNK respectively for Mw group. Black upside and downside shaded triangles

represent NKG2C$^+$ ANK and NKG2A$^+$ iNK respectively for Control group. ****- p <

0.0001, **- p <0.01, ***- p <0.001 and ns= p value not significant.

Figure 3: Impact of Mw on upregulation of NKG2C$^+$ ANK cells with respect to NKG2C

expression at baseline: Scatter dot with bar plot showing the kinetics of A) NKG2C$^+$ ANK

cells (n=15) and C) NKG2A$^+$ iNK (n=15) cells expression in Mw group at baseline, day 30

and 60 in respect to <15% NKG2C at baseline. B) NKG2C$^+$ ANK (n=15) cells and D)

NKG2A$^+$ iNK (n=15) cells expression in Mw group at baseline, day 30 and 60 in respect to

>15% NKG2C at baseline. Log2FC expression of E) NKG2C$^+$ ANK and F) NKG2A$^+$ iNK

cells at day 60 after normalization with baseline value in Mw group (>/<15 %, n=15) and

control group. Red upside and downside shaded triangles represent NKG2C$^+$ ANK and

NKG2A[+] iNK respectively for Mw group. Black upside and downside shaded triangles

represent NKG2C[+] ANK and NKG2A[+] iNK respectively for Control group. ****- p <

0.0001, **- p < 0.01, *- p < 0.05 and ns= p value not significant.

Figure 4: Mw showed sustained effect on NKG2C/NKG2A ratios until 100 days and

Impact of Mw intracellular Cytokine (IFN-γ) release: Scatter dot with bar plot showing

A) NKG2C/NKG2A ratio of Control group (n=15) and Mw group (n=30) at baseline and day

60. B) Kinetics of NKG2C/NKG2A ratio at baseline, day 30, 60 and 100 in Mw group. C) &

D) Kinetics of NKG2C/NKG2A ratio at baseline and day 60 on the basis of >/< 15 %

NKG2C at baseline in Mw group and control group respectively. E) intracellular IFN-γ

release in both control (n=5) and Mw (n=5) group at baseline and day 60. Red and black

shaded circles represent Mw group and Control group respectively. ****- p < 0.0001, ***- p

< 0.001, *- p < 0.05 and ns= p value not significant.

Figure 5: Relationship between NKG2C[+]ANK cells and COVID-19: Scatter dot with bar

plot showing expression of NKG2C[+] ANK cells at baseline in the overall cohort (no

infection, n=64 & infection, n=16), Control group (no infection, n=20 & infection, n=10) and

Mw group (no infection, n=44 & infection, n=6) with respect to SARS-CoV-2 infection.

Black and red upside shaded triangles represent NKG2C[+] ANK for Control group and Mw

group respectively and unshaded upside triangles represent NKG2C[+] ANK of both groups

combined. ***- p < 0.001, **- p < 0.01 and * - P < 0.05.

Figure 6: Impact of the Second dose of Mw on Adaptive and Inhibitory NK cells: Scatter

dot with bar plot showing expression of A) NKG2C[+] ANK (day 60, single dose; n=17, double

dose; n=12 and day 100, single dose; n=17, double dose; n=12), B) NKG2A[+] iNK cells (day

60, single dose; n=12, double dose; n=12 and day 100, single dose; n=17, double dose; n=12)

and C) NKG2C/NKG2A (day 60, single dose; n=17, double dose; n=12 and day 100, single

dose; n=17, double dose; n=12) ratio with respect to single and double dose of Mw vaccine at

day 60 and 100. Red upside and downside shaded triangles represent NKG2C$^+$ ANK and

NKG2A$^+$ iNK respectively for Mw group. Red shaded circle represents Mw group.

 ****- $p < 0.0001$, **- $p < 0.01$, *- $p < 0.05$ and ns= p value not significant.

Figure 7: Differential gene expression and network analysis in Mw and Control group:

A) Volcano plot showing genes differentially expressed in Mw as compared to control group.

Genes with a p-value <0.05 and log2 Fold Change >=0.5 <= -0.5 were considered significant.

Selected genes are highlighted (Red: Upregulated in Mw group, Blue: Upregulated in Control

group). B) Heatmap of RNA-seq expression data showing selected DEGs related to this

study. Hierarchical cluster analysis were performed between all samples of Mw and Control

group. Gene expression is shown in normalized log2 counts per million. C) Boxplots showing

expression levels of 19 selected genes in Mw (green) as compared to control group (orange).

Relative gene expression is shown in normalized log2 counts per million (log2CPM) for I)

ANK pathway in the top panel, II) ADCC pathway in the middle panel. (*p-value < 0.05,

**p-value < 0.01, ***p-value < 0.001, ****p-value<0.0001). D) GO network analysis of the

top 5 enriched GO terms in the differentially expressed genes for Adaptive NK cell and

Antibody-dependent Cellular Cytotoxicity (Fisher's exact test using enrichGO function in R

package clusterProfiler, multiple test correction by Benjamini- Hochberg method, adj. p-

value <0.05).

Figure 8: The suggested mechanistic pathway of the impact of Mw on ANK cells and

protection against COVID-19.

	Control group (N=50)	Mw group (N=50)	p-value
Age at vaccination, Median (Range), Years	28 (21-55)	28 (22-56)	0.11
<45	45	42	
>45	5	8	
Gender			0.3
Male	34	28	
Female	16	22	
SARS-CoV-2 Infection	17	3	0.0012
Mild	11	3	0.02
Moderate	6	0	0.01
Severe	0	0	1.0
Incidence rate/10000 person days	24.05	3.57	
Time to Infection, Median (Range), Mean, Days	56 (18-135) 66.4 ± 39.1	156 (151-159) 155.3 ± 4.0	<0.0001

760 **Table 1: Characteristics and Outcomes**

762 **Table 2: Incidence Rate Ratio of SARS-CoV-2 infections and Mw efficacy.**

Parameter	Estimate	*p*-value	95% CI
Incidence Rate Ratio (IRR)	0.15	0.0004	0.04 to 0.5
Attack rate in Mw control (ARU)	0.34		
Attack rate in Mw treated (ARV)	0.06		
Mw efficacy (%)	82.35%		50 to 96 %
Absolute Risk Reduction (ARR)	0.28		0.13 to 0.43
Number Needed to Treat (NNT)	3.6		2.3 to 7.5

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

(A)

(B)

(C)

Low NKG2C   High NKG2C   Low NKG2C   High NKG2C

NKG2C⁺ ANK cells

Monocyte   NKG2A⁺ iNK cells   NKG2A⁺ iNK cells

Target cell

IFN-γ
Perforin

Cytokine mediated cytotoxicity   Viral destruction   ADCC (Antibody dependent cell cytotoxicity)