A Cloth Facemask Causes No Major Respiratory or Cardiovascular Perturbations

during Moderate to Heavy Exercise

Natália Mendes Guardieiro[1,2], Gabriel Barreto[1], Felipe Miguel Marticorena[1], Tamires Nunes

Oliveira[1], Luana Farias de Oliveira[1], Ana Lucia de Sá Pinto[2,3], Danilo Marcelo Leite do

Prado[1], Bryan Saunders*[1,4], Bruno Gualano*[1,3,5]

*both authors equally contributed to this study

1 – Applied Physiology and Nutrition Research Group, School of Physical Education and

Sport; Rheumatology Division; Faculdade de Medicina FMUSP, Universidade de São Paulo,

São Paulo, SP, BR, University of São Paulo, SP, Brazil.

2 – Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo

3 - Laboratory of Assessment and Conditioning in Rheumatology (LACRE), Rheumatology

Division, Universidade de São Paulo, São Paulo, Brazil

4 – Institute of Orthopaedics and Traumatology, Faculty of Medicine FMUSP, University of

São Paulo, Brazil.

5 – Food Research Center, University of São Paulo, São Paulo, Brazil.

Correspondence:

Dr Bruno Gualano - Applied Physiology & Nutrition Research Group, Rheumatology

Division, Faculty of Medicine FMUSP

Av. Dr. Arnaldo, 455 - Cerqueira César - CEP: 01246903 - University of São Paulo,

São Paulo, SP, Brazil. Phone: +55 11 3061-8789

E-mail: gualano@usp.br

Abstract

Objectives: To investigate whether wearing a cloth facemask could affect physiological and perceptual responses to exercise at distinct exercise intensities in non-trained men and women.

Methods: In a crossover design**,** participants (17 men and 18 women) underwent a progressive square-wave test at four intensities (i. at 80% of the ventilatory anerobic threshold [80%VAT]; ii. at VAT; iii. at the respiratory compensation point [RCP]; iv. at exercise peak [Peak] to exhaustion), with or without a triple-layered cloth mask (Mask or No-Mask). Several physiological, metabolic and perceptual measures were analyzed.

Results: Mask reduced inspiratory capacity at all exercise intensities vs. No-Mask (p<0.0001), irrespective of sex. Mask reduced respiratory frequency vs. No-Mask (p=0.001) at Peak (-8.3 breaths·min^{-1}; CI: -5.8, -10.8), RCP (-6.9 breaths·min^{-1}; CI: -4.6, -9.2) and VAT (-6.5 breaths·min^{-1}; CI: -4.1, -8.8), but not at Baseline or at 80%VAT. Mask also reduced tidal volume (p<0.0001) at both RCP (-0.5L; CI: -0.3, -0.6) and Peak (-0.8L; CI: -0.6, -0.9), but not at Baseline, 80%VAT or VAT. Shallow breathing index was increased with Mask at Peak compared to No-Mask (11.3; CI: 7.5, 15.1), but not at any other intensities. Mask did not change heart rate, lactate, ratings of perceived exertion, blood pressure or oxygen saturation.

Conclusions: Wearing a cloth facemask during exercise at moderate to heavy intensities is unlikely to incur significant respiratory or cardiovascular changes, irrespective of sex. These data can inform new exercise recommendations for health during the COVID-19 pandemic and debunk unfounded allegations of harmful effects of masks during exercise. ClinicalTrials.gov: NCT04887714

Key-words: physical activity; mask; COVID-19 pandemic; oxygen saturation; lactate.

What are the new findings?

✓ Using a progressive square-wave test, we showed that wearing a cloth facemask during exercise increased breathing difficulty, but this was dependent upon the exercise intensity.

✓ Respiratory variables (e.g., inspiratory capacity, respiratory frequency, shallow breathing index) were affected at higher rather than lower intensities.

✓ Mask wearing did not change heart rate, lactate, ratings of perceived exertion, blood pressure or oxygen saturation at any exercise intensity.

✓ There were no substantial sex differences on the effects of mask wearing during exercise.

How might it impact on clinical practice in the future?

✓ These data can debunk unfounded allegations on harmful effects of masks during exercise, and help inform new exercise recommendations for health during the COVID-19 pandemic, particularly where facemasks remain necessary.

Introduction

The use of face masks is one of the most effective non-pharmacological strategy to prevent SARS-CoV-2 infections [1, 2]. The recent resurgence of cases and deaths worldwide has led to some decision-makers to re-issue mask orders to contain the disease, suggesting that this safety tool will remain important as long as the pandemic is not fully mitigated [3], and possibly after it is under control, as in several cultures employing facemasks as a routine practice to protect against several health threats before COVID-19 outbreak [4]. Wearing a facemask is recommended even during exercise, particularly at indoor fitness facilities and gyms, where COVID-19 outbreaks have been reported [5]. Nevertheless, the physiological impact of facemasks during exercise remains underexplored.

A facemask may reduce the ability to breathe comfortably during exercise, which has been confirmed by some [6, 7], but not all [8] studies. It is possible to conjecture that the effects of wearing a mask on cardiorespiratory responses may manifest during exhaustive high-intensity exercise, but not (or less) during low- to moderate-intensity exercise. In fact, Driver et al. [6] provide preliminary evidence that the effect of facemasks is dependent on exercise intensity, with differences in oxygen saturation, ratings of perceived exertion (RPE) and dyspnoea occurring at different stages of an incremental cardiopulmonary test as exercise intensity increased. However, short-stage maximal incremental tests do not normalize the physiological responses to exercise in relation to the gas exchange or blood acid-base profiles [9 10], since %VO_{2max} at the ventilatory thresholds largely varies between individuals [9], hampering accurate determination of exercise intensities and ultimately confounding data interpretation [10, 11]. To overcome this limitation, constant-load tests based on the dynamic behaviour of the pulmonary gas exchange and blood acid-base status have been recommended to accurately determine exercise intensity domains (i.e., moderate, heavy,

severe, and extreme) [10]. This approach allows determination of whether facemasks affect

physiological and perceptual parameters at different, well-defined exercise intensities,

helping tailor exercise prescription for health that can minimize any negative effects of

wearing a mask on cardiorespiratory responses.

Another remaining question is whether women and men respond differently to mask wearing

during exercise. Generally, women have smaller lungs and airways, which limits their ability

to generate expiratory flow [12, 13], which results in reduced ventilation during exercise

compared to men. Women also have lower oxygen (O_2) carrying capacity, maximum cardiac

output, and arteriovenous O_2 difference [14]. Considering the number of physiological and

morphological sex differences to exercise, one could speculate that any physiological

perturbations brought about wearing a mask during exercise could be greater in women, since

men have an overall higher cardiorespiratory reserve.

This study aimed to investigate whether wearing a cloth facemask could affect physiological

and perceptual responses to exercise at distinct exercise intensities in non-trained individuals.

A secondary aim was to test whether sex differences influenced these responses.

Methods

Ethics statement

The protocol was approved by the institutional ethics committee. Written informed consent was obtained before participants' enrollment.

Study design and setting

This was a crossover study (clinicaltrials.gov: NCT04887714) performed at an intrahospital, exercise physiology laboratory in São Paulo, Brazil. Data collection took place between April and November 2021.

Participants

Men and women not engaged in competitive sports (*i.e.,* non-trained) were eligible for this study. Exclusion criteria included any cardiac, pulmonary, and rheumatologic diseases, musculoskeletal limitations, and BMI <18.5 or >30 kg·m^2. A total of 18 men and 20 women entered the study, although 3 dropped out for reasons unrelated to the study. Thirty-five individuals (17 men and 18 women) completed all main sessions (age: women: 28 ± 5 y, men: 30 ± 4 y; BMI: women: 22.9 ± 2.0 kg·m^2, men: 24.5 ± 2.6 kg·m^2) and were analyzed. According to the short form of the International Physical Activity Questionnaire (IPAQ) instrument [15], 31 participants were physically active, whereas 4 were inactive.

Experimental design

Participants attended the laboratory on three separate occasions, separated by a minimum of 48 h, at the same time of day to account for circadian variation [16]. The first visit consisted of an incremental cardiopulmonary running test to exhaustion to determine peak oxygen uptake (V$\Box$O$_{2peak}$) and ventilatory thresholds. The remaining two main visits consisted of a

running progressive square-wave test (PSWT), performed with or without the use of a triple-layered cloth facemask (Fashion Masks, São Paulo, Brazil). This facemask was chosen because it is widely accessible, recommended to the general public by the WHO and appropriate for exercise (https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/cloth-face-cover-guidance.html). The outer layer was waterproof polyester fabric, the middle layer a polypropylene filter, and the inner layer was absorbable cotton. The size was "one size fits all" and was thus identical for all participants. Athletes were required to keep the mask firmly in place over the nose, mouth, and chin during the entire session. The order of sessions was determined by an individual not involved in data collection. Blocks of two individuals were allocated to the two possible orders (Mask–No Mask; No Mask–Mask) using a random number generator (https://www.randomizer.org/) to ensure the study was counterbalanced. Since the investigators could see that the participants were wearing a cloth facemask or not, the session order was provided directly to the research team. All participants were habituated to wear a mask during their daily routines due to mandates, but not specifically during exercise. Participants were requested to refrain from strenuous exercise, caffeine and alcohol, and replicated their diet, in the 24 h prior to each visit.

Cardiorespiratory Exercise Test

Immediately prior to the cardiorespiratory exercise test, participants performed a pulmonary function test according to recommendations [17]. The cardiorespiratory exercise test was performed on a motorized treadmill (Centurion 300, Micromed, Brazil). For men, the test started at 5 km·h^{-1} and increased speed (1 km·h^{-1}·min^{-1}) up to a maximum velocity of 14 km·h^{-1}. For women, the test started at 4 km·h^{-1} and increased speed (1 km·h^{-1}·min^{-1}) up to 13 km·h^{-1}. For those participants who reached these maximal speeds, there was a subsequent increase in inclination (2%·min^{-1}) until exhaustion. Ventilatory and gas exchange

measurements were recorded continuously throughout the test using a breath-by-breath system (MetaLyzer 3B, Cortex, Germany), as was heart rate (HR; ergo PC elite, Micromed, Brazil). Maximal effort was determined according to published criteria [18] and individual $\dot{V}O_{2peak}$ was determined as the $\dot{V}O_2$ averaged over the final 30 s.

Progressive square-wave test (PSWT)

Data from the cardiorespiratory exercise test was used to determine exercise workload for the square wave treadmill test according to the ventilatory anaerobic threshold (VAT) and the respiratory compensation point (RCP). All thresholds were determined by the same respiratory physiologist with clinical experience in the area. The PSWT protocol was performed on a motorized treadmill (Centurion, model 200, Micromed, Brazil) and consisted of three 5-min stages at workloads equivalent to 1) 80%VAT, 2) VAT, and 3) RCP. These stages represented moderate, heavy and severe domains [10] and corresponded to 41±9%, 53±9% and 81±8% of $\dot{V}O_{2peak}$ of the volunteers. Participants then completed a final stage to exhaustion at a running speed equivalent to the maximum achieved during the cardiorespiratory exercise test (Peak). Ventilatory and gas exchange measurements were recorded continuously throughout, with the spirometer mask placed over the cloth facemask.

To determine the effect of the mask on pattern of change in operating lung volumes we evaluated end-expiratory volume to functional vital capacity ratio (EELV/FVC). Inspiratory capacity was determined at rest and at the end of each exercise stage during the PSWT. Ventilatory constraint was evaluated as the difference between inspiratory capacity at rest and at each exercise workload [19]. Ventilatory efficiency was determined using the ventilatory equivalent for carbon dioxide ($\dot{V}_E/\dot{V}CO_2$) and end-tidal carbo dioxide pressure (PetCO$_2$)

during each stage. Breathing pattern was evaluated during each stage using the breathing

frequency to tidal volume ratio (BF/TV) ratio [20].

RPE were assessed at the end of each stage with participants pointing to a chart using the 6-

to 20-point Borg scale [21]. Heart rate was monitored continuously throughout (ergo PC elite,

Micromed, Brazil). A fingertip blood sample (20 μL) was collected at baseline, at the end of

each stage and 4-min post-exhaustion for the subsequent analysis of lactate. Blood was

homogenized in the same volume of 2% NaF, centrifuged at 2000 g for 5 min before plasma

was removed and stored at $-20^{o}C$ until analysis. Plasma lactate was determined

spectrophotometrically using an enzymatic-colorimetric method (Katal, Interteck, Brazil).

Subjective Perception of Discomfort Questionnaire

Participants completed a questionnaire [22] following the completion of the PSWT to rate

their perception of ten sensations of discomfort related to the facemask: humidity, heat,

breathing resistance, itchiness, tightness, saltiness, feeling unfit, odor and fatigue. They were

also required to rate their overall feeling of discomfort related to the facemask on a scale

from 0 to 10, with 0–3 representing "Comfortable", >3–7 representing "Uncomfortable" and

>7 representing "Extremely uncomfortable".

Statistical analyses

Lactate data from 2 individuals (1 male, 1 female) who did not complete the third stage were

excluded from the RCP and Peak analyses. Furthermore, 1 male and 1 female reported

extreme discomfort with the PSWT mask and stopped exercising before volitional

exhaustion; and were excluded from the Peak analysis for all outcomes. The same 4

individuals were excluded from the TTE analysis. Lactate data were log10 transformed

before mixed model analysis, turning the model into an exponential data mixed model, and

back transformed through exponentiation for the final reporting of data. Whenever outlying

data points were considered improbable (e.g., a value of 50 mmHg for systolic blood

pressure), they were considered measurement or transcription error and were excluded.

Repeated measures Mixed Model ANOVAs were performed with condition (Mask, No-

Mask), sex and exercise intensity (Baseline [except RPE], 80%VAT, VAT, RCP, Peak) as

fixed factors and individuals as random factors. Exceptions were Spirometry, TTE and

Questionnaire related outcomes, which were not repeated measures and, therefore, timing

was not included as a fixed factor. For TTE, the model was corrected by treatment order,

since participants were not familiarized to the protocol. When a significant main effect or

interaction was detected (accepted at $p \leq 0.05$), post-hoc pairwise comparisons were performed

with Tukey's adjustment. All analyses were performed with the RStudio software (Rstudio

1.4.11003, PBC, MA) using the "lmer" function of the lmerTest package and "emmeans"

function of the emmeans package. Data are expressed as estimated differences and 95%

confidence intervals (CI), and data in figures are represented as mean±1standard deviation.

Patient and Public Involvement statement

Participants were not involved in the design or in performing the study.

Results

Cardiorespiratory Exercise Test

Participants' characteristics and respiratory data at the ventilatory thresholds and calculated

stages are presented in Table 1.

Table 1. Participants' characteristics and cardiorespiratory data from the cardiopulmonary exercise test.

VE/MVV: minute ventilation/maximum voluntary ventilation ratio.

	Women	Men
	Mean ± SD	**Mean ± SD**
Age (years)	28 ± 5	30 ± 4
Weight (kg)	60.9 ± 9.0	73.96 ± 8.1
Height (m)	1.63 ± 0.07	1.74 ± 0.07
80%VAT		
Heart rate (bpm)	115 ± 12	109 ± 20
Respiratory exchange ratio	0.79 ± 0.11	0.83 ± 0.07
Minute ventilation (L/min)	25.4 ± 5.5	34.4 ± 10.7
VO_2 absolute (L/min)	1.01 ± 0.19	1.43 ± 0.4
VO_2 relative (ml/kg/min)	16.2 ± 3.2	19.4 ± 6
VO_2 %peak (%)	42.1 ± 7.9	39.8 ± 10.5
VAT		
Heart rate (bpm)	128 ± 13	122 ± 17
Respiratory exchange ratio	0.85 ± 0.09	0.89 ± 0.06
Minute ventilation (L/min)	32.6 ± 6.4	44.7 ± 10.2
VO_2 absolute (L/min)	1.25 ± 0.21	1.83 ± 0.35
VO_2 relative (ml/kg/min)	20.7 ± 3.5	24.9 ± 5.5
VO_2 %peak (%)	53.6 ± 7.5	51.4 ± 9.8
RCP		
Heart rate (bpm)	168 ± 9	161 ± 10
Respiratory exchange ratio	1.02 ± 0.05	1.04 ± 0.05

Minute ventilation (L/min)	64.3 ± 10.9	83.5 ± 14.8
VO$_2$ absolute (L/min)	2 ± 0.24	2.75 ± 0.35
VO$_2$ relative (ml/kg/min)	33.1 ± 3.6	37.2 ± 4.9
VO$_2$ %peak (%)	86.1 ± 8.4	77.0 ± 8.0
Peak		
Heart rate (bpm)	183 ± 6	183 ± 6
Respiratory exchange ratio	1.18 ± 0.06	1.22 ± 0.07
Minute ventilation (L/min)	95.3 ± 15.3	139.4 ± 25.1
VE/MVV	0.73 ± 0.13	0.77 ± 0.16
VO$_2$ absolute (L/min)	2.33 ± 0.26	3.59 ± 0.45
VO$_2$ relative (ml/kg/min)	38.7 ± 4.6	48.6 ± 6.3

Progressive square-wave test (PSWT)

Inspiratory capacity (IC)

Mask reduced IC at all exercise intensities (interaction effect condition*timing: F=8.6,

p<0.0001, Figure 1, Panel A) compared to No-Mask irrespective of sex (80%VAT: -0.4 L;

CI: -0.2, -0.6; VAT: -0.5 L; CI: -0.4, -0.7; RCP: -0.7 L; CI: -0.5, -0.9; Peak: -1.0 L; CI: -0.8, -

1.2), except at Baseline (-0.2 L; CI: 0.0, -0.4).

End expiratory lung volume (EELV) and EELV/Forced Vital capacity (EELV/FVC)

Mask did not influence EELV or EELV/FVC, irrespective of exercise intensities and sex (all

p≥0.1, Figure 1, Panel B and C).

Respiratory Frequency (Rf)

Mask reduced Rf vs. No-Mask (interaction effect of condition*timing: F=4.6, p=0.001,

Figure 1, Panel D) at Peak (-8.3 breaths·min^{-1}; CI: -5.8, -10.8), RCP (-6.9 breaths·min^{-1}; CI: -

4.6, -9.2) and VAT (-6.5 breaths·min^{-1}; CI: -4.1, -8.8), but not at Baseline or at 80%VAT

(both p≥0.06). Rf was reduced similarly in men (-7.5 breaths·min^{-1}; CI: -6.0, -9.0) and

women (-3.4 breaths·min^{-1}; CI: -1.9, -4.9) with Mask (Figure 1, Panel D).

Tidal Volume (VT)

Mask reduced VT (interaction effect of condition*intensity, F=18.3, p<0.0001, Figure 1,

Panel E) at both RCP (-0.5L; CI: -0.3, -0.6) and Peak (-0.8L; CI: -0.6, -0.9), but not at

Baseline, 80%VAT or VAT (all p≥0.97). Sex had no influence on the effects of Mask on VT

(p=0.053).

Tobin index (Rf/VT or shallow breathing index)

Mask increased the Tobin index at Peak compared to No-Mask (+11.3; CI: 7.5, 15.1), but not

at any other intensity (all p≥0.4, interaction effect of condition*intensity: F=7.3, p<0.0001).

Rf/VT in men was not affected by Mask, whereas it was increased in Mask vs. No-Mask for

women (interaction effect of condition*sex: F=25.1, p<0.0001; men: +1.1; CI: -1.1, 3.3;

women: +6.9; CI: 4.7, 9.1; Figure 1, Panel F).

PetCO$_2$

Mask increased PetCO$_2$ at both RCP (+4.0 mmHg; CI: 2.8, 5.3, Figure 1, Panel G) and Peak

(+4.9 mmHg; CI: 3.5, 6.3) compared to No-Mask (interaction effect condition*intensity: F =

6.8, p<0.0001), but had no effect at Baseline, 80%VAT or VAT (all p≥0.09). The effect of

Mask on PetCO$_2$ increases was comparable in men (+3.4 mmHg; CI: 2.5, 4.2) and women

(+1.9 mmHg; CI: 1.1, 2.7).

$V\square_E/V\square CO_2$

Mask reduced VE/VCO_2 at the three highest intensities compared to No-Mask (interaction

effect of condition*intensity, F=3.7, p=0.006): VAT (-2.2; CI: -0.9, -3.5), RCP (-3.2; CI:

-1.9, -4.5), and Peak (-4.4; CI: -3.0, -5.8) (Figure 1, Panel H), irrespective of sex.

Heart rate during exercise (HR)

No effects of Mask at any exercise intensity were seen for HR. Despite an interaction effect

of condition*sex for HR (F=4.6, p=0.03), post-hoc comparisons did not show any significant

differences (all p≥0.4). (Figure 2, Panel A).

Lactate

Mask did not affect lactate measures at any exercise intensities, irrespective of sex (Figure 2,

Panel B).

Ratings of Perceived Exertion (RPE)

Mask did not influence RPE at any exercise intensity. However, Mask increased RPE for

men, but not women (interaction effect of condition*sex, F=6.2, p=0.01, men: +1.4; CI: 0.9,

2.0; women: +0.4; CI: -0.1, 1.0; Figure 2, Panel C).

Blood pressure

Mask did not affect both systolic or diastolic blood pressure at any exercise intensity (Figure

2, Panel D and E), regardless of sex.

Oxygen saturation ($SatO_2$)

Mask did not affect $SatO_2$ at any exercise intensity (Figure 2, Panel F), regardless of sex.

Time-to-exhaustion (TTE)

Mask reduced TTE compared to No-Mask (-34.5 s; CI: -17.0, -52.1; main effect of condition:

F=14.9, p=0.0007, Figure 3), with no condition*sex interaction.

Forced Vital Capacity at rest (FVC)

Mask reduced FVC compared to No-Mask (-1.8L; CI: -1.1, -1.5; condition: F=117.7,

p<0.0001), with no condition*sex interaction (Table 2).

Forced expiratory volume in 1 second at rest (FEV1)

Mask reduced FEV1 compared to No-Mask (-1.2; CI: -1.0, -1.4; main effect of condition:

F=156.2, p<0.0001), with no condition*sex interaction (Table 2).

FVC/FEV1 at rest (FVC by FEV1 ratio)

Mask did not influence the FVC/FEV1 ratio, irrespective of sex (Table 2).

Peak expiratory flow at rest (PEF)

Mask reduced PEF compared to No-Mask (-3.4; CI: -2.8, -4.0, main effect of condition:

F=122.1, p<0.0001), independently of sex (Table 2).

Subjective perception questionnaire

Mask increased the subjective feelings of Heat, Misfitting, Discomfort, Fatigue, Resistance,

Saltiness and Humidity (all p≤0.01), but did not affect feelings of Saltiness, Tightness or

Itchiness (all p≥0.1). No interaction between condition and sex was detected (Table 2).

Table 2. Spirometry at rest and subjective questionnaire outcomes.

	Women		Men	
	No-Mask	**Mask**	**No-Mask**	**Mask**
	Mean ± SD	**Mean ± SD**	**Mean ± SD**	**Mean ± SD**
Spirometry				
FEV1 (L)[$]!	3.4 ± 0.4	2.3 ± 0.5	4.5 ± 0.5	3.2 ± 0.6
FVC (L)[$]!	3.8 ± 0.6	2.6 ± 0.6	5 ± 0.7	3.7 ± 0.7
FVC/FEV1(%)	90.7 ± 7.7	90.5 ± 7.2	90.8 ± 6.7	87.3 ± 8.4
PEF (L/s)[$]!	7.7 ± 1.8	4.4 ± 1.1	9.2 ± 1.9	5.7 ± 1.6
Questionnaire				
Tightness	3.4 ± 2.4	4.2 ± 3.4	2.5 ± 2.2	3.4 ± 2.5
Heat[$]	3.6 ± 2.1	6.2 ± 2.5	3.3 ± 2.2	5.9 ± 2.3
Itchiness	0.8 ± 1.1	1.1 ± 2.2	1.6 ± 2.7	2.8 ± 3.1
Misfitting[$]	1.8 ± 2.3	2.6 ± 2.5	1.6 ± 2.1	4.1 ± 2.7
Discomfort[$]	4.5 ± 1.4	7.9 ± 2.3	4.2 ± 2.9	7.1 ± 2.6
Fatigue[$]	5.2 ± 3.1	8.2 ± 1.8	5.0 ± 2.6	8.6 ± 2.4
Odour	1.2 ± 2.4	0.9 ± 1.8	0.9 ± 1.3	1.7 ± 2.4
Resistance[$]	3.4 ± 2.3	8.6 ± 1.2	4.0 ± 2.8	8.4 ± 1.3
Saltiness[$]!	0.3 ± 0.5	0.7 ± 1.6	1.0 ± 1.8	2.7 ± 3.5
Humidity[$]	3.7 ± 2.6	5.3 ± 2.3	4.1 ± 1.9	6.6 ± 2.4

322 FEV1, forced expiratory volume in 1 second; FVC, forced vital capacity; FVCbyFEV1,

323 forced vital capacity by FEV1 ratio; PEF, peak expiratory flow.

324 Significance labels: [$], main effect of condition; [!], main effect of sex; [&], interaction effect

Figure 1. Inspiratory capacity, end expiratory lung volume (EELV), EELV by forced vital capacity ratio (EELV/FVC), respiratory frequency, tidal volume, respiratory frequency by tidal volume, CO_2 partial pressure (PetCO$_2$) and ventilatory equivalent for carbon dioxide (V_E/VCO$_2$) data expressed as mean ± 1 standard deviation.

$^\$$ main effect of Mask; $^!$ main effect of intensity; $^\&$ main effect of sex; * condition*intensity interaction; $^\male$ condition*intensity interaction for men; $^\female$ condition*intensity interaction for women.

Figure 2. Heart rate, lactate, rating of perceived effort (RPE), systolic (SBP) and diastolic (DBP), and oxygen saturation (SatO$_2$) expressed as mean ± 1 standard deviation.

$ main effect of Mask; ! main effect of intensity; & main effect of sex; ♂ condition*intensity interaction for men.

Figure 3. Time-to-exhaustion during the final stage. Dashed lines connect individual
performance data between No-Mask and Mask condition.

$ main effect of Mask

Discussion

This study showed that breathing difficulty with a cloth facemask is dependent upon the exercise intensity, with lower distress at less severe intensities. Mask wearing did not substantially affect physiological or metabolic variables during exercise, regardless of sex and intensity. From a practical perspective, these data suggest that use of a cloth facemask for protecting individuals from SARS-CoV-2 infections should not be a barrier to the engagement in moderate-to-heavy physical activity.

The main novelty of the current study is that we assessed the influence of wearing a mask on respiratory and cardiovascular variables across several exercise intensities, spanning moderate to severe domains [10]. Wearing a mask did not modify most respiratory variables in the moderate to heavy exercise domains, with only inspiratory capacity reduced at 80%VAT with the mask. Our findings suggest that reductions in inspiratory capacity early in exercise are reflective of a decrease in contractile power of the inspiratory muscles. This inspiratory distress may place a greater strain on the respiratory muscles to maintain breathing requirements during exercise. This was not seen in the moderate to heavy domains but was manifested in an inability to maintain the physiological increases in respiratory frequency and tidal volume at the higher intensities. For example, during VAT, although breathing frequency was reduced, this was likely compensated by increases in tidal volume as evidenced by no differences between conditions, suggesting that the cloth facemask did not negatively inhibit the ability of the respiratory system to work at moderate, heavy and severe exercise intensities. Indeed, this might explain the reduced V_E/VCO_2 with a mask. At higher intensities, however, both respiratory frequency and tidal volume were reduced with a mask, which may have led to an inability to maintain respiratory homeostasis and affected subsequent performance. Our data are in agreement with previous studies showing a reduced

exercise capacity with different facemasks [6 22], though others have shown no negative

impact [8, 23]. It is possible that different types of facemasks and respirators (e.g., FFP2/N95,

surgical, cloth facemasks) and participants' fitness level may have contributed to these

conflicting findings.

Despite some changes in respiratory variables, no cardiovascular measure was affected using

a cloth facemask. Even during the highest exercise intensity domains, there were no changes

in heart rate, or systolic and diastolic blood pressure when wearing a cloth facemask.

Moreover, the slight overall reduction in blood oxygen saturation (not detected at individual

exercise intensities) was not clinically meaningful, supporting previous studies showing no

effect of wearing a facemask on oxygen saturation [24, 25]. These findings collectively can

counteract misinformation during the COVID-19 pandemic, particularly relating to the use of

masks during exercise and its supposed negative effects on cardiac overload, acid-base

balance, and oxygen saturation [28].

We had speculated that any potential physiological effects associated with wearing a mask

during exercise could be greater in women, who have an overall lower higher

cardiorespiratory reserve than men, owing to classically described morphological and

physiological sex-differences (e.g., smaller lungs, lower O_2 carrying capacity and maximum

cardiac output). This hypothesis was not confirmed, as the effects of the cloth facemask on

physiological measures were mostly similar between women and men irrespective of exercise

intensities. Exercise capacity was reduced by 23.9% in women and 17.8% in men when using

a cloth facemask, with no differences between sexes. This suggests that the stress imposed by

wearing a mask does not constitute a greater physiological or metabolic burden to women vs.

men, despite the well-known sex-differences during exercise.

This study has several limitations. Although the measurement of respiratory variables during the PSWT provide novel information regarding the respiratory response during different intensities, participants were required to wear a facemask for breath-by-breath measures over the cloth facemask. This may have increased the discomfort felt by the participants and may also have led to some inaccuracies in measurements due to air escaping. The current data cannot be directly extrapolated to trained individuals; however, we felt it important to investigate this matter among a non-trained population, as there has been intense debate on the physiological repercussions and potential adverse effects of face masks in non-trained individuals. Since sufficient levels of physical activity prevent morbidities and mortality [29-31] and improve vaccine immunogenicity [32], it is important that mask mandates do not lead to a reduction in physical activity. In this regard, the present data provide relevant information that wearing a cloth facemask will not have a negative impact during exercise at moderate-to-heavy intensities, which is associated with a plethora of health-related benefits [33, 34]. Whether the negative perpetual feelings related to the use of masks may result in less adherence to exercise remains to be examined. Furthermore, the influence of mask wearing during exercise in clinical populations warrants investigation.

In conclusion, although wearing a cloth facemask may reduce maximal exercise capacity, it did not impose significant respiratory or cardiovascular changes during exercise performed at moderate-to-heavy intensities. These data have important practical implications as they can debunk unfounded allegations of harmful effects of facemasks during exercise and help inform new exercise recommendations for health during the COVID-19 pandemic.

Acknowledgments

The authors would like to thank all of the athletes for taking part in this research. The authors received no specific funding for this work. B.G (2017/13552-2), G.B. (2020/12036-3), T.N.O. (2020/04368-6) and B.S. (2016/50438-0) have been financially supported by Fundação de Amparo à Pesquisa do Estado de São Paulo. B.S. has also received a grant from Faculdade de Medicina da Universidade de São Paulo (2020.1.362.5.2).

References

References

1. Clase CM, Fu EL, Joseph M, et al. Cloth Masks May Prevent Transmission of COVID-19:
An Evidence-Based, Risk-Based Approach. Annals of internal medicine
2020;**173**(6):489-91 doi: 10.7326/M20-2567[published Online First: Epub Date]|.

2. Leung NHL, Chu DKW, Shiu EYC, et al. Respiratory virus shedding in exhaled breath and
efficacy of face masks. Nature medicine 2020;**26**(5):676-80 doi: 10.1038/s41591-020-
0843-2[published Online First: Epub Date]|.

3. Stephenson J. CDC Studies Underscore Continued Importance of Masks to Prevent
Coronavirus Spread. JAMA Health Forum 2021;**2**(2):e210207-e07 doi:
10.1001/jamahealthforum.2021.0207[published Online First: Epub Date]|.

4. Burgess A, Horii M. Risk, ritual and health responsibilisation: Japan's 'safety blanket' of
surgical face mask-wearing. Sociology of health & illness 2012;**34**(8):1184-98 doi:
10.1111/j.1467-9566.2012.01466.x[published Online First: Epub Date]|.

5. Groves LM, Usagawa L, Elm J, et al. Community Transmission of SARS-CoV-2 at Three
Fitness Facilities — Hawaii, June–July 2020. MMWR and Morbidity and Mortality
Weekly Report 2021;**70**:316-20

6. Driver S, Reynolds M, Brown K, et al. Effects of wearing a cloth face mask on
performance, physiological and perceptual responses during a graded treadmill
running exercise test. Br J Sports Med 2021 doi: 10.1136/bjsports-2020-
103758[published Online First: Epub Date]|.

7. Fikenzer S, Uhe T, Lavall D, et al. Effects of surgical and FFP2/N95 face masks on
cardiopulmonary exercise capacity. Clinical research in cardiology : official journal of
the German Cardiac Society 2020;**109**(12):1522-30 doi: 10.1007/s00392-020-01704-
y[published Online First: Epub Date]|.

8. Shaw K, Butcher S, Ko J, et al. Wearing of Cloth or Disposable Surgical Face Masks has
no Effect on Vigorous Exercise Performance in Healthy Individuals. Int J Environ
Res Public Health 2020;**17**(21) doi: 10.3390/ijerph17218110[published Online First:
Epub Date]|.

9. Coyle EF, Coggan AR, Hopper MK, et al. Determinants of endurance in well-trained
cyclists. J Appl Physiol (1985) 1988;**64**(6):2622-30 doi:
10.1152/jappl.1988.64.6.2622[published Online First: Epub Date]|.

10. Burnley M, Jones AM. Oxygen uptake kinetics as a determinant of sports performance.
Eur J Sport Sci 2007;**7**(2):63-79 doi: 10.1080/17461390701456148[published Online
First: Epub Date]|.

11. Whipp BJ, Ward SA, Rossiter HB. Pulmonary O2 uptake during exercise: conflating
muscular and cardiovascular responses. Med Sci Sports Exerc 2005;**37**(9):1574-85
doi: 10.1249/01.mss.0000177476.63356.22[published Online First: Epub Date]|.

12. Duke JW. Sex Hormones and Their Impact on the Ventilatory Responses to Exercise and
the Environment. In: Hackney AC, ed. Sex Hormones, Exercise and Women:
Scientific and Clinical Aspects. Cham: Springer International Publishing, 2017:19-34.

13. Freedson P, Katch VL, Sady S, et al. Cardiac output differences in males and females
during mild cycle ergometer exercise. Medicine and science in sports 1979;**11**(1):16-9

14. Lewis DA, Kamon E, Hodgson JL. Physiological differences between genders.
Implications for sports conditioning. Sports Med 1986;**3**(5):357-69 doi:
10.2165/00007256-198603050-00005[published Online First: Epub Date]|.

15. Hagstromer M, Oja P, Sjostrom M. The International Physical Activity Questionnaire
(IPAQ): a study of concurrent and construct validity. Public health nutrition
2006;**9**(6):755-62 doi: 10.1079/phn2005898[published Online First: Epub Date]|.

16. Atkinson G, Reilly T. Circadian variation in sports performance. Sports Med
1996;**21**(4):292-312 doi: 10.2165/00007256-199621040-00005[published Online
First: Epub Date]|.

17. Graham BL, Steenbruggen I, Miller MR, et al. Standardization of Spirometry 2019
Update. An Official American Thoracic Society and European Respiratory Society
Technical Statement. American journal of respiratory and critical care medicine
2019;**200**(8):e70-e88 doi: 10.1164/rccm.201908-1590ST[published Online First:
Epub Date]|.

18. Howley ET, Bassett DR, Jr., Welch HG. Criteria for maximal oxygen uptake: review and
commentary. Med Sci Sports Exerc 1995;**27**(9):1292-301

19. Guenette JA, Chin RC, Cory JM, et al. Inspiratory Capacity during Exercise:
Measurement, Analysis, and Interpretation. Pulmonary medicine 2013;**2013**:956081
doi: 10.1155/2013/956081[published Online First: Epub Date]|.

20. Neder JA, Andreoni S, Lerario MC, et al. Reference values for lung function tests: II.
Maximal respiratory pressures and voluntary ventilation. Brazilian Journal of Medical
and Biological Research [online] 1999;**32**(6):719-27 doi: 10.1590/S0100-
879X1999000600007[published Online First: Epub Date]|.

21. Borg GA. Psychophysical bases of perceived exertion. Medicine and science in sports and
exercise 1982;**14**(5):377-81

22. Li Y, Tokura H, Guo YP, et al. Effects of wearing N95 and surgical facemasks on heart
rate, thermal stress and subjective sensations. International archives of occupational
and environmental health 2005;**78**(6):501-9 doi: 10.1007/s00420-004-0584-
4[published Online First: Epub Date]|.

23. Epstein D, Korytny A, Isenberg Y, et al. Return to training in the COVID-19 era: The
physiological effects of face masks during exercise. Scand J Med Sci Sports
2021;**31**(1):70-75 doi: 10.1111/sms.13832[published Online First: Epub Date]|.

24. Shaw KA, Zello GA, Butcher SJ, et al. The impact of face masks on performance and
physiological outcomes during exercise: a systematic review and meta-analysis. Appl
Physiol Nutr Metab 2021:1-11 doi: 10.1139/apnm-2021-0143[published Online First:
Epub Date]|.

25. Fukushi I, Nakamura M, Kuwana S-i. Effects of wearing facemasks on the sensation of
exertional dyspnea and exercise capacity in healthy subjects. PloS one
2021;**16**(9):e0258104 doi: 10.1371/journal.pone.0258104[published Online First:
Epub Date]|.

26. Kouzy R, Abi Jaoude J, Kraitem A, et al. Coronavirus Goes Viral: Quantifying the
COVID-19 Misinformation Epidemic on Twitter. Cureus 2020;**12**(3):e7255 doi:
10.7759/cureus.7255[published Online First: Epub Date]|.

27. Roozenbeek J, Schneider CR. Susceptibility to misinformation about COVID-19 around
the world. 2020;**7**(10):201199 doi: 10.1098/rsos.201199[published Online First: Epub
Date]|.

28. Chandrasekaran B, Fernandes S. "Exercise with facemask; Are we handling a devil's
sword?" – A physiological hypothesis. Medical Hypotheses 2020;**144**:110002 doi:
https://doi.org/10.1016/j.mehy.2020.110002[published Online First: Epub Date]|.

29. Lavie CJ, Ozemek C, Carbone S, et al. Sedentary Behavior, Exercise, and Cardiovascular
Health. Circulation research 2019;**124**(5):799-815 doi:
10.1161/circresaha.118.312669[published Online First: Epub Date]|.

30. Fiuza-Luces C, Garatachea N, Berger NA, et al. Exercise is the real polypill. Physiology
(Bethesda, Md) 2013;**28**(5):330-58 doi: 10.1152/physiol.00019.2013[published
Online First: Epub Date]|.

31. Mok A, Khaw KT, Luben R, et al. Physical activity trajectories and mortality: population
based cohort study. 2019;**365**:l2323 doi: 10.1136/bmj.l2323[published Online First:
Epub Date]|.

32. Gualano B, Lemes I, Silva R, et al. Physical activity associates with enhanced
immunogenicity of an inactivated virus vaccine against SARS-CoV-2 in patients with
autoimmune rheumatic diseases: Research Square, 2021.
33. Gebel K, Ding D, Chey T, et al. Effect of Moderate to Vigorous Physical Activity on All-
Cause Mortality in Middle-aged and Older Australians. JAMA Internal Medicine
2015;**175**(6):970-77 doi: 10.1001/jamainternmed.2015.0541[published Online First:
Epub Date]|.
34. Hupin D, Roche F, Gremeaux V, et al. Even a low-dose of moderate-to-vigorous physical
activity reduces mortality by 22% in adults aged ≥60 years: a systematic review and
meta-analysis. Br J Sports Med 2015;**49**(19):1262-67 doi: 10.1136/bjsports-2014-
094306[published Online First: Epub Date]|.