

Predictors of uncertainty and unwillingness to receive the COVID-19 booster vaccine in a sample of 22,139 fully vaccinated adults in the UK

Elise Paul, PhD^a & Daisy Fancourt, PhD^a

^a Department of Behavioural Science and Health, University College London, 1-19 Torrington Place, London, WC1E 7HB, United Kingdom

*Corresponding author: Dr Elise Paul, 1-19 Torrington Place, London, WC1E 7HB, United Kingdom, e.paul@ucl.ac.uk

Funding

The Nuffield Foundation [WEL/FR-000022583], the MARCH Mental Health Network funded by the Cross-Disciplinary Mental Health Network Plus initiative supported by UK Research and Innovation [ES/S002588/1], and the Wellcome Trust [221400/Z/20/Z and 205407/Z/16/Z].

Keywords

COVID-19, booster vaccine, vaccine refusal

Summary

Background The continued success of the COVID-19 vaccination programme in the UK will depend on widespread uptake of booster vaccines. However, there is evidence of hesitancy and unwillingness to receive the booster vaccine, even in fully vaccinated adults. Identifying factors associated with COVID-19 booster vaccine intentions specifically in this population is therefore critical.

Methods We used data from 22,139 fully vaccinated adults who took part in the UCL COVID-19 Social Study. Multinomial logistic regression examined longitudinal predictors of uncertainty and unwillingness (versus willingness) to receive a COVID-19 booster vaccine (measured 22 November 2021 to 6 December 2021), including (i) socio-demographic factors, (ii) COVID-19 related factors (e.g., having been infected with COVID-19), and (iii) initial intent to receive a COVID-19 vaccine in the four months following the announcement in the UK that the vaccines had been approved (2 December 2020 to 31 March 2021).

Findings 4% of the sample reported that they were uncertain about receiving a COVID-19 booster vaccine, and a further 4% unwilling. Initial uncertainty and unwillingness to accept the first COVID-19 vaccine in 2020-21 were each associated with over five times the risk of being uncertain about and unwilling to accept a booster vaccine. Healthy adults (those without a pre-existing physical health condition) were also more likely to be uncertain or unwilling to receive a booster vaccine. In addition, low levels of current stress about catching or becoming seriously ill from COVID-19, consistently low compliance with COVID-19 government guidelines during periods of strict restrictions (e.g., lockdowns), lower levels of educational qualification, lower socio-economic position and age below 45 years were all associated with uncertainty and unwillingness.

Interpretation Our findings highlight that there are a range of factors that predict booster intentions, with the strongest predictor being previous uncertainty and unwillingness. Two other concerning patterns also emerged from our results. First, administration of booster vaccinations may increase social inequalities in experiences of COVID-19 as adults from lower socio-economic backgrounds are also most likely to be uncertain or unwilling to accept a booster vaccine as well as most likely to be seriously affected by the virus. Second, some of those most likely to spread COVID-19 (i.e., those with poor compliance with guidelines) are most likely to be uncertain and unwilling. Public health messaging should be tailored specifically to these groups.

Introduction

Although highly effective in preventing severe disease and death,^{1,2} emerging evidence suggests that immunity from COVID-19 vaccines diminishes over time.(Falsey et al., 2021; Levine-Tiefenbrun et al., 2021) Licensed COVID-19 vaccines may also be less effective against new more transmissible variants^{5,6} and may therefore need to be updated.⁷ Widespread uptake of COVID-19 booster vaccines in the general population will therefore be critical for maintaining the progress that has been made in preventing deaths and serious illness due to COVID-19 and reducing the burden on the healthcare system in the UK.

Although COVID-19 vaccine uptake in the UK has, overall, been high,⁸ some groups have been more likely than others to refuse the vaccine.⁹ People from ethnic minority groups, those who have already been infected with COVID-19, people living in more deprived areas, people who are unemployed or not working, and smokers are less likely to have received at least one dose of a COVID-19 vaccine.⁹ Some of these same groups had also expressed hesitancy or unwillingness to receive a COVID-19 vaccine before they were available: people with lower levels of education,¹⁰⁻¹³ lower household incomes,^{10,11,14} and people from ethnic minority groups.¹² However, other demographic groups reporting COVID-19 vaccine hesitancy before the vaccines were available¹⁰ have not shown differential uptake in the UK (e.g., women).⁹

Recent data from the Opinions and Lifestyle Survey (OPN) in the UK suggest a considerable proportion of the general population who have received two doses of a COVID-19 vaccine are either unwilling (5%; very unlikely or fairly unlikely) or unsure (4%; neither likely nor unlikely) about accepting a booster vaccine.¹⁵ This proportion is much higher amongst adults who previously reported they or were unlikely or unsure about having a first dose of a COVID-19 vaccine (22%).¹⁶ Young adults (8%) and adults over the age of 70 (9%) were twice as likely as other age groups (4% of adults ages 30-49 and 3% of adults ages 50-69) to say they would be unwilling to accept the booster, whilst the proportion of men saying they would refuse is was twice that of women (6% vs 3%).¹⁵ But further research into predictors of booster hesitancy or unwillingness is lacking.

Therefore, the aims of this study were to identify longitudinal associations between socio-demographic and COVID-19 related factors (e.g., having been infected with COVID-19) and initial intent to receive the COVID-19 vaccine with uncertainty and unwillingness to receive a COVID-19 booster vaccine in adults who had been fully vaccinated.

Methods

Study design and participants

Data were drawn from the COVID-19 Social Study; a large panel study of the psychological and social experiences of over 75,000 adults (aged 18+) in the UK during the COVID-19 pandemic. The study commenced on 21 March 2020 and involves online weekly data collection from participants for the duration of the COVID-19 pandemic in the UK. The study is not random and therefore is not representative of the UK population. But it does contain a well-stratified sample that was recruited using three primary approaches. First, convenience sampling was used, including promoting the study through existing networks and mailing lists (including large databases of adults who had previously consented to be involved in health research across the UK), print and digital media coverage, and social media. Second, more targeted recruitment was undertaken focusing on groups who were anticipated to be less likely to take part in the research via our first strategy, including (i) individuals from a low-income background, (ii) individuals with no or few educational qualifications, and (iii) individuals who were unemployed. Third, the study was promoted via partnerships with

third sector organisations to vulnerable groups, including adults with pre-existing mental health conditions, older adults, carers, and people experiencing domestic violence or abuse. The study was approved by the UCL Research Ethics Committee [12467/005] and all participants gave informed consent. Participants were not compensated for participation.

In the current study, we included individuals who had (i) data on intent to receive a COVID-19 booster vaccine (collected 22 November to 6 December 2021, hereafter ‘follow-up’), (ii) data on initial intent to receive a first COVID-19 vaccine (collected 2 December 2020 to 31 March 2021), (iii) received at least two doses of a COVID-19 vaccine at follow-up, and (iv) complete data on other study variables.

A total of 32,096 participants met the first criterion, and of these, 26,619 also had data on initial COVID-19 vaccine intent, and 25,821 also met the third criterion of having had at least two doses of a COVID-19 vaccine at follow-up. 3,682 individuals were excluded due to missing data on other study variables, for example those who had selected “other/prefer not to say” in response to gender or “prefer not to say” on ethnicity or income, due to insufficient statistical power and an inability to create statistical weights for these individuals, leaving a total analytical sample of 22,139. See Supplemental Table S1 for a comparison of excluded and included participants on study variables.

Measures

Outcome variable

Intention to receive a COVID-19 booster vaccine was measured at follow-up with a study developed item (“How likely do you think you are to get a COVID-19 booster vaccine if/when you are offered one?”). Response options ranged from 1 “very unlikely” to 6 “very likely”, as well as 7 “I have already had one/accepted one”. An ordinal variable was coded: (0) intend to receive the booster vaccine (responses of 5-7), (1) unsure about whether to receive the booster vaccine (responses of 3-4), and (2) unwilling to receive the booster vaccine (responses of 1-2).

Predictor variables

Socio-demographics

Socio-demographic factors were collected at baseline and included gender (male vs female), age group (60+, 45-59, 30-44, and 18-29) ethnicity (white vs ethnic minority groups [i.e. Asian/Asian British, Black/Black British, White and Black/Black British, mixed race, Chinese/Chinese British, Middle Eastern/Middle Eastern British, or other ethnic group]), education (undergraduate degree or higher [further education after the age of 18], A- levels (equivalent to education to age 18) or vocational training, and up to GCSE (General Certificate of Secondary Education) [equivalent to education to age 16]), income (annual household income: >£90,000, £60,000-89,999, £30,000-59,999, £16,000-29,999, and < £16,000), smoking status (non-smoker vs former or current smoker), employment status (not employed [still at school, at university, unable to work due to disability, unemployed and seeking work, or retired vs employed [self-employed, part-time or full-time employed]), area of dwelling (urban [city, large town, small town] vs rural [village, hamlet, isolated dwelling]), living arrangement (living alone vs living with others but not including children vs living with others, including children), and government’s identified key worker status (not a key worker vs key worker). The latter included people with jobs deemed essential during the pandemic (e.g., health and social care, education, and childcare) and who were required to leave home to carry out this work during lockdowns.

Participants reported at baseline whether they had received a clinical diagnosis of a mental illness (e.g., depression, anxiety, or other psychiatric diagnosis) or chronic physical health condition (e.g., high blood pressure, diabetes, or other physical health condition) were used to create two binary

variables (yes/no) to indicate the presence or absence of pre-existing physical and mental health conditions.

COVID-19 related factors

Initial intent to receive a COVID-19 vaccine was measured with a study-developed item administered 2 December 2020 to 31 March 2021. Many participants had vaccine intention data for all four months during this period, but because we were interested in their first recorded intention to receive the vaccine after its authorisation was announced in the UK on 2 December 2020, we coded initial intent in this way. In a sensitivity analysis, participants' last recorded COVID-19 vaccine intention during the period 2 December 2020 to 31 March 2021 was used. See Supplemental Table S2 for a listing of all question wording for study-developed items.

Ever having had COVID-19 was operationalised from a study-developed item administered at follow-up which was coded as: yes, confirmed by a test (either a positive COVID-19 test or an antibody test), yes, suspected by either a doctor or the participant, and no.

Confidence in government and the health service to handle the pandemic were assessed with one question each at follow-up. Response options ranged from 1 (none at all) to 7 (lots). Two binary variables were created to indicate a lot of (5-7) versus low (1-4) confidence in the government and health system.

Responses to the question on compliance with government COVID-19 guidelines were on a scale from 1 (none at all) to 7 (very much so). We analyse this as an ordinal variable reflecting patterns of compliance whilst restrictions were in place across the pandemic. As most of our sample resides in England, we used dates for which there were strict COVID-19 restrictions in England (e.g., rather than Scotland or Northern Ireland). See Supplemental Table S3 for a listing of these dates. The ordinal variable was coded as: consistently high compliance (a mean score of 6-7 across all strict restriction periods) vs consistently low compliance (mean score of 1-5 across all strict restriction periods). Sensitivity analyses using i) only participants with compliance data from all three strict restriction periods and ii) a variable reflecting compliance across the entirety of the pandemic were conducted.

Knowledge of COVID-19 was measured at follow-up with one study-developed question and rated on a 7-point scale from 1 (very poor knowledge) to 7 (very good knowledge). Responses of 1-4 were categorised as low compared to high (5-7) COVID-19 knowledge. The presence or absence of worry about either contracting COVID-19 or becoming seriously ill from it were captured from two multiple choice questions asked at follow-up. A binary variable was created to indicate not having endorsed either as a source of stress vs having endorsed one or both.

Statistical analysis

Two multinomial regression models were fitted to examine associations between socio-demographic and COVID-19 related factors and uncertainty and unwillingness to receive a COVID-19 booster vaccine. The first model included only socio-demographic variables whilst the second additionally included COVID-19 related variables. The outcome variable in these two models was coded such that those who were uncertain or unwilling were compared to those who were very likely to receive the COVID-19 booster vaccine or had already accepted one. Multinomial regression coefficients were exponentiated and presented as relative risk ratios (RRR) with corresponding 95% confidence intervals (CI).

To account for the non-random nature of the sample and increase representativeness of the UK general population, all data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics.¹⁷ A multivariate reweighting method was implemented using the Stata user written command 'ebalance'.¹⁸ Analyses were conducted using Stata version 16.¹⁹

Results

Unweighted and weighted sample characteristics are presented in Table 1. Before weighting, the sample was disproportionately female, older (ages 60+), white, and had education levels of at least an undergraduate degree. But after weighting the sample aligned with expected population characteristics. 4% of the weighted sample expressed unwillingness to receive a COVID-19 booster vaccine (e.g., a score of 1-2 on a scale of 1 to 6), whilst 4% were uncertain (a score of 3-4) (Figure 1). Descriptive differences between adults who were unwilling or uncertain about receiving a COVID-19 booster vaccine vs those who were willing by several demographic characteristics are shown in Table S4.

Of those who were initially willing to accept a COVID-19 vaccine when it became available, 5% said they were either uncertain (3%) or unwilling (2%) to accept a booster (Table 2). 73% of those who were initially uncertain and 65% of those who were initially unwilling had become willing to accept a booster. But 12% of adults who were initially uncertain about accepting a COVID-19 vaccine were now unwilling to accept a booster, and 1 in 5 (20%) who were initially unwilling about a first dose remained unwilling to accept a booster vaccine.

Socio-demographic predictors of booster uncertainty and unwillingness

Results from the multinomial regression model predicting uncertainty and unwillingness to receive a COVID-19 booster vaccine from socio-demographic factors are shown in Table 3. The strongest predictor of booster unwillingness was younger age. Young adults were nearly six times as likely than older adults to be unwilling to receive a booster (RRR = 5.74; 95% CI: 2.82 to 11.68), and adults ages 30-44 were more than twice as likely (RRR = 2.56; 95% CI: 1.31 to 5.00). Education to A-levels or vocational and up to a GCSE were each associated with a 2.1-2.5 times higher relative risk of being unwilling to receive a COVID-19 booster vaccine relative to people who held to degrees (A-levels or vocational: RRR = 2.09; 95% CI: 1.15 to 3.79; up to GCSE: RRR = 2.50; 95% CI: 1.31 to 4.79). Additionally, healthy adults were also at increased risk of being unwilling (RRR 1.52; 95% CI: 1.00 to 2.30) to receive the booster compared to individuals with long-term physical health conditions. Ethnicity, living arrangement, smoking status, keyworker status, employment status, income, and having a mental health condition were not associated with uncertainty or unwillingness.

For booster uncertainty, the same three factors continued to be predictors. Relative to adults aged 60 and above, adults aged 18-29 were over 5 times more likely to be uncertain (RRR = 5.30; 95% CI: 2.87 to 9.77), adults aged 30-44 were over 4 times more likely (RRR = 4.31; 95% CI: 2.63 to 7.05), and even adults aged 45-59 were 1.6 times more likely to be uncertain (RRR = 1.60; 95% CI: 1.00 to 2.54). Education up to GCSE was additionally associated with a 1.7 times increased risk of being uncertain (RRR = 1.76; 95% CI: 1.09 to 2.84) relative to having a degree, but the association for adults with A-levels or vocational qualifications was not present. As for unwillingness, healthy adults were also at increased risk of being unwilling (RRR = 1.67; 95% CI: 1.17 to 2.40) to receive the booster compared to individuals with a long-term physical health condition. For uncertainty, two further socio-demographic factors emerged as predictors too. Adults who were employed (vs unemployed/retired/students/not working) were more than 3 times more likely to be uncertain (RRR = 3.25; 95% CI: 2.11 to 5.02). There were mixed results for income, with some evidence that household income levels below £60,000 and below £16,000 per year were associated with increased risk of being uncertain (£30,000-59,999: RRR = 2.43; 95% CI: 1.06 to 5.59; <£16,000: RRR = 3.48; 95% CI: 1.32 to 9.16) relative to higher income households. But no further socio-demographic factors were associated with uncertainty, including ethnicity, living arrangement, smoking status, keyworker status and having a mental health condition were not associated with uncertainty or unwillingness.

COVID-19 related predictors of booster uncertainty and unwillingness

The strongest predictor of booster uncertainty and unwillingness was original intentions to be vaccinated when the vaccines were first approved (Table 4). Adults who were initially uncertain about receiving a COVID-19 vaccine were five times more likely to say they were uncertain or unwilling to receive a booster (uncertain: RRR = 4.92; 95% CI: 2.98 to 8.11; unwilling: RRR = 5.29; 95% CI: 3.07 to 9.09) than those who were initially willing. Initial unwillingness associated with an over six times higher relative risk (RRR = 6.40; 95% CI: 3.94 to 10.41) and eleven times higher relative risk (RRR = 11.29; 95% CI: 6.79 to 18.78) of booster uncertainty and unwillingness, respectively.

The second highest predictor of unwillingness was consistently low compliance with COVID-19 guidelines during strict restriction periods (RRR = 2.45; 95% CI: 1.65 to 3.65) which also predicted uncertainty (RRR = 1.51; 95% CI: 1.04 to 2.19). Not having said that catching or falling seriously ill from COVID-19 was a source of stress in past week at follow-up also predicted unwillingness (RRR = 1.91; 95% CI: 1.19 to 3.07) and uncertainty (RRR = 1.80; 95% CI: 1.19 to 2.72) about boosters. Low self-reported knowledge of COVID-19 also predicted uncertainty but not unwillingness to get a booster (uncertainty: RRR = 1.78; 95% CI: 1.19 to 2.67; unwillingness: RRR = 1.19; 95% CI: 0.70 to 2.05). Confidence in government, confidence in the health system, and past experience of having COVID were not predictors of booster intentions.

Sensitivity analyses

Sensitivity analyses conducted using participants' last recorded (rather than first recorded) COVID-19 vaccine intention during the period 2 December 2020 to 31 March 2021 (Supplemental Table S5), from the model which included a variable indicating low vs high levels of compliance with guidelines across the entirety of the pandemic (rather than only during periods of strict restrictions) (Supplemental Table S6), and from the model which included only sample participants who had had exactly two doses of a COVID-19 vaccine at follow-up (Supplemental Table S7) resulted in substantively similar findings. However, amongst adults who had had exactly two doses of a COVID-19 vaccine at follow-up, age was no longer a predictor of booster uncertainty (likely because most older adults had already received boosters by then), whilst education up to A-levels or vocational and age 30-44 were no longer predictors of booster unwillingness (Supplemental Table S7).

Discussion

This is the first study, to our knowledge, to examine longitudinal socio-demographic and COVID-19 related factors, and initial intent to receive a COVID-19 vaccine with future uncertainty and unwillingness to receive a COVID-19 booster vaccine in fully vaccinated adults. The proportions in our sample who were uncertain about receiving (4%) and unwilling (4%) to receive a COVID-19 booster vaccine are similar to those in the OPN (4% and 5%, respectively).¹⁵ Our findings broadly align with the factors already identified as predicting initial vaccine intentions when the vaccine was first approved (e.g. age, income, employment, and education), showing that some of the same groups should remain the focus of public health campaigns to increase booster vaccine uptake.¹⁰ Underlining this finding, our results also highlight that previous uncertainty and unwillingness to be vaccinated remains the strongest predictor of intentions to receive a booster vaccine, even amongst adults who did get fully vaccinated despite initial concerns. Two other concerning patterns also emerged from our results. First, administration of booster vaccinations may increase social inequalities in experiences of COVID-19 as adults from lower socio-economic backgrounds are also most likely to be uncertain or unwilling to accept a booster vaccine as well as most likely to be seriously affected by the virus. Second, some of those most likely to spread COVID-19 (i.e., those with poor compliance with guidelines) are most likely to be uncertain and unwilling.

Previous vaccine attitudes and intentions are logical predictors of attitudes to booster vaccines, and previous research has highlighted a range of reasons for initial hesitancy, including high levels of mistrust in vaccines and concerns about unforeseen future side effects.¹⁰ However, the adults in our study had already received two vaccines, so their initial uncertainty or unwillingness had previously been overcome. Furthermore, not all of the fully vaccinated adults in our sample who were uncertain or unwilling to receive a booster vaccine were initially hesitant about receiving a first dose (Table 2); 5% of the initially willing were now either uncertain (3%) or unwilling (2%) to accept a booster vaccine. So there is a need to understand the reasons for uncertainty and unwillingness to accept a booster vaccine in more detail. One potential explanation has to do with misinformation about the length of protection provided by COVID-19 vaccines. In recent data from fully vaccinated adults in the OPN study, the most commonly cited reason for unwillingness to receive a booster vaccine (59%) was a belief that the first and second vaccines were enough to keep them safe, and nearly half (49%) thought that a booster would not offer them extra protection.¹⁵ Indeed, it is notable that healthy adults were 1.5 times more likely to be uncertain or unwilling to receive a booster vaccine, suggesting that the booster vaccine is held as less important than individuals with pre-existing health conditions. This indicates an urgent need for public health messaging to correct widespread misinformation on the length of protection offered from COVID-19 vaccines. A second potential explanation relates to recent publicised concerns that giving third doses to people in high income countries whilst many lower income countries do not have access to a first or second dose will increase inequities.²⁰ Indeed, recent findings from the OPN indicate that around 1 in 4 (22%) fully vaccinated adults who are unwilling to accept a booster vaccine said they were unwilling to do so because they thought the vaccine should be offered to someone else.¹⁵ Further, a study conducted by Imperial College London asked respondents specifically if they would be willing to donate their COVID-19 booster vaccine to people in low income countries if a test showed they did not need a booster dose, and the majority of respondents in the UK (78%) and the US (69%) said they would do so.²¹ Therefore, altruism may be a driver for booster hesitancy or unwillingness for some adults in our sample. Further research is needed to understand these new attitudes further in order that specific strategies to encourage booster uptake can be designed and implemented.

It is also concerning that some we found a socio-economic gradient in intentions to take up the booster vaccine, as some of the same socio-demographic groups most at risk of uncertainty and unwillingness to receive the COVID-19 booster vaccine are also at increased risk for severe illness and death from COVID-19, including those with low incomes and who have low levels of education.^{10,12,13,25} These groups have already experienced a disproportionately higher number of infections and deaths, relative to other groups and had lower uptake of initial vaccines.²⁶⁻²⁹ Consequently, our finding of reduced intentions to take up the booster in this group, even amongst individuals who have already received two vaccinations, is concerning as it may exacerbate existing inequalities in exposure to and experience of the virus. However, other socio-demographic groups did not show any differences in booster vaccine intentions (key worker status, smoking, ethnicity etc.) This provides direction as to which groups may require more specific public health strategies to encourage booster vaccinations, and such work may need to take account of specific logistical or financial barriers individuals may face in getting their booster vaccine.

Worryingly, some of those who are most likely to spread the virus are also the least likely to say they will accept a booster vaccine. First, the most important socio-demographic determinant of both uncertainty and unwillingness to receive a COVID-19 booster vaccine in our study was age, which has been found in prior work.^{12,15,25} Young adults were over 5 times as likely to be uncertain about accepting a booster vaccine compared to older adults, and nearly 6 times as likely to be unwilling. Young adults may be more hesitant about accepting a booster vaccine because they feel they are less likely to be severely affected by the disease. Other work has shown that young adults are less stressed by the virus³⁰ and less compliant with government guidelines,³¹ which were also factors identified as predictors of original COVID-19 vaccine unwillingness and hesitancy.^{10,25} Young people may also be less likely to be vaccinated because of concerns about side effects.¹⁰ However, in the

OPN study, young adults (and adolescents) who were initially uncertain about accepting a COVID-19 vaccine were the most likely to have later received at least one dose.¹⁶ Therefore, the younger adults age group may be most amenable to public health programmes to provide more information and clarity around the benefits of booster vaccination. We also found an association between uncertainty and unwillingness to receive a COVID-19 booster vaccine and poor compliance with government guidelines during lockdowns, which is congruent with other research.^{10,25} This is concerning, as individuals who are non-compliant with behavioural interventions such as mask wearing and isolation when infected or symptomatic, or after contact with someone who has been infected, are more likely to spread the virus.³² Hospitalisations may subsequently increase, further driving the need for more of the behavioural interventions these individuals fail to comply with. Interestingly, though, having been previously infected with COVID-19 was not a predictor of either unwillingness or uncertainty about accepting a booster vaccine, as has been found elsewhere, suggesting that individuals are not making a trade in terms of strategies to avoid infection.⁹

Strengths and limitations

Strengths of the current study are that it is a large sample well-stratified on major demographic characteristics and analyses were weighted to increase representativeness of the general UK population. Analyses were also longitudinal and included rich information on a range of socio-demographic and COVID-19 related factors. We focused on initial COVID-19 vaccine attitudes in the four months following the day their approval was announced in the UK in 2019 as well as gathering new data on booster vaccine attitudes to allow comparisons of attitudes over the past 12 months. However, there were some limitations. First, our sample is not nationally representative of the general UK population. We used a heterogeneous and large sample in our analyses and weighted our responses to population proportions, but it is possible that more extreme views towards booster vaccines may therefore not be represented. Second, our recruitment strategy could have resulted in clustering (i.e., individuals from the same households participating), and attitudes may be more similar within households. However, this would be considered a minimal risk given the large sample and we used robust standard errors. Third, due to small sub-group sample sizes we were unable to examine specific effects of different ethnicities on booster vaccine attitudes and we treated this socio-demographic characteristic as binary. This could have led to oversimplification of the diverse opinions and experiences held by different ethnicity subgroups.

Conclusions

Findings suggest initial uncertainty and unwillingness about receiving a COVID-19 booster vaccine will be a barrier to maintaining the progress that has been made thus far in controlling the COVID-19 pandemic through vaccines. There is therefore an urgent need for public health messaging emphasising the safety and importance of COVID-19 booster vaccines even for fully vaccinated adults. Focusing on people who were initially uncertain rather than unwilling may be more effective, as they may be easier to convince to vaccinate than the latter.³³ In the Vaccine Opinions Study in the UK, adults who were previously unsure about the COVID-19 vaccine were more likely to later have accepted at least one dose it than those who were initially unwilling (60% vs 40%).¹⁶ But more specific nuanced public health strategies are recommended based on the findings of this paper, in particular focusing on individuals from lower socio-economic backgrounds and individuals at greatest risk of spreading the virus. Overall, there is an urgent need for more research into ongoing attitudes to COVID-19 vaccines and boosters. This is particularly important in light of evidence showing that existing COVID-19 vaccines offer protection for limited time periods and therefore will likely need to be updated on a regular period moving forwards.

Role of the funding source

The funders had no role in the study design; in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the paper for publication. All researchers listed as authors are independent from the funders and all final decisions about the research were taken

by the investigators and were unrestricted. All authors had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Contributors

DF conceptualised and designed the study. DF also acquired funding, led the investigation, provided oversight on the methodology, administered the project, provided software and other resources, and supervised the project. Data were curated, validated, and formally analysed by EP. EP created visualisations, wrote the original manuscript draft with input from all authors. All authors reviewed and edited the manuscript. All authors approved the final version of the manuscript and had full access to and verified the data.

Declaration of interests

All authors declare no conflicts of interest.

Data sharing

The UCL COVID-19 Social Study documentation and codebook are available for download at <https://www.covidsocialstudy.org/>. Statistical code is available upon request from Elise Paul (e.paul@ucl.ac.uk).

Acknowledgements

The researchers are grateful for the support of a number of organisations with their recruitment efforts including: the UKRI Mental Health Networks, Find Out Now, UCL BioResource, SEO Works, FieldworkHub, and Optimal Workshop.

Ethics approval and consent to participate

Ethical approval for the COVID-19 Social Study was granted by the UCL Ethics Committee. All participants provided fully informed consent and the study is GDPR compliant.

References

1. Haas EJ, Angulo FJ, McLaughlin JM, Anis E, Singer SR, Khan F, et al. Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a nationwide vaccination campaign in Israel: an observational study using national surveillance data. *The Lancet*. 2021;397(10287):1819–29.
2. Mor V, Gutman R, Yang X, White EM, McConeghy KW, Feifer RA, et al. Short-term impact of nursing home SARS-CoV-2 vaccinations on new infections, hospitalizations, and deaths. *J Am Geriatr Soc*. 2021;
3. Falsey AR, Fenck Jr RW, Walsh EE, Kitchin N, Absalon J, Gurtman A, et al. SARS-CoV-2 neutralization with BNT162b2 vaccine dose 3. *N Engl J Med*. 2021;385(17):1627–9.
4. Levine-Tiefenbrun M, Yelin I, Alapi H, Katz R, Herzog E, Kuint J, et al. Viral loads of Delta-variant SARS-CoV2 breakthrough infections following vaccination and booster with the BNT162b2 vaccine. *medRxiv*. 2021;
5. Abu-Raddad LJ, Chemaitelly H, Butt AA. Effectiveness of the BNT162b2 Covid-19 Vaccine against the B. 1.1. 7 and B. 1.351 Variants. *N Engl J Med*. 2021;
6. Lopez Bernal J, Andrews N, Gower C, Gallagher E, Simmons R, Thelwall S, et al. Effectiveness of Covid-19 vaccines against the B. 1.617. 2 (Delta) variant. *N Engl J Med*. 2021;585–94.
7. Milne G, Hames T, Scotton C, Gent N, Johnsen A, Anderson RM, et al. Does infection with or vaccination against SARS-CoV-2 lead to lasting immunity? *Lancet Respir Med*. 2021 Dec 1;9(12):1450–66.
8. UK Government. Vaccinations in the UK | Coronavirus in the UK [Internet]. 2021 [cited 2021 Dec 6]. Available from: <https://coronavirus.data.gov.uk/details/vaccinations>
9. Office for National Statistics. Coronavirus (COVID-19) Infection Survey technical article: Analysis of characteristics associated with vaccination uptake [Internet]. 2021. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionsurveytechnicalarticleanalysisofcharacteristicsassociatedwithvaccinationuptake/2021-11-15>
10. Paul E, Steptoe A, Fancourt D. Attitudes towards vaccines and intention to vaccinate against COVID-19: Implications for public health communications. *Lancet Reg Health - Eur* [Internet]. 2021 Feb 1 [cited 2021 Feb 22];1(100012). Available from: [https://www.thelancet.com/journals/lanep/article/PIIS2666-7762\(20\)30012-0/abstract](https://www.thelancet.com/journals/lanep/article/PIIS2666-7762(20)30012-0/abstract)
11. Rhodes A, Hoq M, Measey M-A, Danchin M. Intention to vaccinate against COVID-19 in Australia. *Lancet Infect Dis*. 2020 Sep;S1473309920307246.
12. Robertson E, Reeve KS, Niedzwiedz CL, Moore J, Blake M, Green M, et al. Predictors of COVID-19 vaccine hesitancy in the UK household longitudinal study. *Brain Behav Immun*. 2021 May 1;94:41–50.
13. Sherman SM, Smith LE, Sim J, Amlôt R, Cutts M, Dasch H, et al. COVID-19 vaccination intention in the UK: results from the COVID-19 vaccination acceptability study (CoVAccS), a nationally representative cross-sectional survey. *Hum Vaccines Immunother*. 2021;17(6):1612–21.

14. Loomba S, de Figueiredo A, Piatek SJ, de Graaf K, Larson HJ. Measuring the impact of COVID-19 vaccine misinformation on vaccination intent in the UK and USA. *Nat Hum Behav*. 2021;5(3):337–48.
15. Office for National Statistics. Coronavirus and the social impacts on Great Britain- 3 December 2021 [Internet]. 2021 [cited 2021 Dec 6]. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthandwellbeing/bulletins/coronavirusandthesocialimpactsongreatbritain/latest>
16. Office for National Statistics. Coronavirus and changing attitudes towards vaccination, England [Internet]. 2021 [cited 2021 Dec 9]. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthandwellbeing/bulletins/coronavirusandchangingattitudestowardsvaccinationengland/7to16september2021#attitudes-towards-covid-19-booster-vaccines>
17. Office for National Statistics. Population estimates for the UK, England and Wales, Scotland and Northern Ireland [Internet]. 2020 May [cited 2020 Sep 30]. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/bulletins/annualmidyearpopulationestimates/mid2018>
18. Hainmueller J, Xu Y. Ebalance: A Stata package for entropy balancing. *J Stat Softw*. 2013;54(7).
19. StataCorp. Stata Statistical Software: Release 16. College Station, TX: StataCorp LP; 2019.
20. Dyer O. Covid-19: Rich countries' booster plans will impede global vaccination, experts say. *BMJ*. 2021;374:n2353.
21. Institute of Global Health Innovation. UK and USA attitudes towards COVID-19 booster vaccines [Internet]. Imperial College London; 2021. Available from: https://www.imperial.ac.uk/media/imperial-college/institute-of-global-health-innovation/UK_US-vaccine-insights_ICL-YouGov-Covid-19-Behaviour-Tracker_20210625_final.pdf
22. Singh AK, Gupta R, Ghosh A, Misra A. Diabetes in COVID-19: Prevalence, pathophysiology, prognosis and practical considerations. *Diabetes Metab Syndr Clin Res Rev*. 2020;14(4):303–10.
23. Wang B, Li R, Lu Z, Huang Y. Does comorbidity increase the risk of patients with COVID-19: evidence from meta-analysis. *Aging*. 2020 Apr 8;12(7):6049–57.
24. Ssentongo P, Ssentongo AE, Heilbrunn ES, Ba DM, Chinchilli VM. Association of cardiovascular disease and 10 other pre-existing comorbidities with COVID-19 mortality: A systematic review and meta-analysis. *PLOS ONE*. 2020 Aug 26;15(8):e0238215.
25. Czeisler MÉ, Rajaratnam SM, Howard ME, Czeisler CA. COVID-19 Vaccine Intentions in the United States—December 2020 to March 2021. *MedRxiv*. 2021;2021.05.16.21257290.
26. Abedi V, Olulana O, Avula V, Chaudhary D, Khan A, Shahjouei S, et al. Racial, Economic, and Health Inequality and COVID-19 Infection in the United States. *J Racial Ethn Health Disparities*. 2021;8(3):732–42.
27. Krieger N, Waterman PD, Chen JT. COVID-19 and Overall Mortality Inequities in the Surge in Death Rates by Zip Code Characteristics: Massachusetts, January 1 to May 19, 2020. *Am J Public Health*. 2020;110(12):1850–2.

28. Tan AX, Hinman JA, Abdel Magid HS, Nelson LM, Odden MC. Association Between Income Inequality and County-Level COVID-19 Cases and Deaths in the US. *JAMA Netw Open*. 2021 May 3;4(5):e218799.
29. Whittle RS, Diaz-Artiles A. An ecological study of socioeconomic predictors in detection of COVID-19 cases across neighborhoods in New York City. *BMC Med*. 2020 Sep 4;18(1):271.
30. Niño M, Harris C, Drawve G, Fitzpatrick KM. Race and ethnicity, gender, and age on perceived threats and fear of COVID-19: Evidence from two national data sources. *SSM - Popul Health*. 2021;13:100717.
31. Wright L, Fancourt D. Do predictors of adherence to pandemic guidelines change over time? A panel study of 22,000 UK adults during the COVID-19 pandemic. *Prev Med*. 2021 Dec 1;153:106713.
32. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ, et al. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *The Lancet*. 2020;395(10242):1973–87.
33. World Health Organization. Best practice guidance: How to respond to vocal vaccine deniers in public. 2016.




Figure 1. Changes in willingness to receive a COVID-19 vaccine (2 December 2020 to 31 March) and a COVID-19 booster vaccine (22 November to 6 December 2021) (weighted, $N = 22,139$)

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

Table 1. Unweighted and weighted sample characteristics (N = 22,139)

| | | Unweighted | | Weighted | |
|--|---|------------|------|----------|------|
| | | Mean | SD | Mean | SD |
| Gender | Male | 0.26 | 0.44 | 0.49 | 0.50 |
| | Female | 0.74 | 0.44 | 0.51 | 0.50 |
| Age | 60+ | 0.43 | 0.50 | 0.30 | 0.46 |
| | 45-59 | 0.34 | 0.47 | 0.27 | 0.44 |
| | 30-44 | 0.19 | 0.39 | 0.24 | 0.43 |
| | 18-29 | 0.04 | 0.19 | 0.20 | 0.40 |
| Ethnicity | White | 0.97 | 0.18 | 0.87 | 0.33 |
| | Ethnic minority groups | 0.03 | 0.18 | 0.13 | 0.33 |
| Education | Undergraduate degree or higher | 0.71 | 0.46 | 0.34 | 0.47 |
| | A-levels or vocational | 0.16 | 0.37 | 0.34 | 0.47 |
| | Up to GCSE | 0.13 | 0.34 | 0.32 | 0.47 |
| Income | >£90,000 | 0.10 | 0.30 | 0.07 | 0.26 |
| | £60,000-89,999 | 0.15 | 0.35 | 0.13 | 0.33 |
| | £30,000-59,999 | 0.35 | 0.48 | 0.33 | 0.47 |
| | £16,000-29,999 | 0.26 | 0.44 | 0.29 | 0.45 |
| | <£16,000 | 0.14 | 0.35 | 0.18 | 0.39 |
| Employment status | Not employed | 0.42 | 0.49 | 0.42 | 0.49 |
| | Employed | 0.58 | 0.49 | 0.58 | 0.49 |
| Living arrangement | Alone | 0.23 | 0.42 | 0.20 | 0.40 |
| | With others, not with children | 0.57 | 0.49 | 0.59 | 0.49 |
| | With, including children | 0.20 | 0.40 | 0.21 | 0.41 |
| Area of dwelling | Urban | 0.75 | 0.43 | 0.79 | 0.40 |
| | Rural | 0.25 | 0.43 | 0.21 | 0.40 |
| Smoking status | Former or non-smoker | 0.93 | 0.25 | 0.89 | 0.32 |
| | Current smoker | 0.07 | 0.25 | 0.11 | 0.32 |
| Key worker status | Not a key worker | 0.80 | 0.40 | 0.78 | 0.41 |
| | Key worker | 0.20 | 0.40 | 0.22 | 0.41 |
| Long-term physical health condition | No long-term physical health condition | 0.57 | 0.50 | 0.59 | 0.49 |
| | Long-term physical health condition | 0.43 | 0.50 | 0.41 | 0.49 |
| Long-term mental health condition | No long-term mental health condition | 0.85 | 0.36 | 0.81 | 0.40 |
| | Long-term mental health condition | 0.15 | 0.36 | 0.19 | 0.40 |
| Confidence in government to handle pandemic | Low confidence in government to handle pandemic | 0.33 | 0.47 | 0.36 | 0.48 |
| | High confidence in government to handle pandemic | 0.67 | 0.47 | 0.64 | 0.48 |
| Confidence in health system to handle pandemic | High confidence in health system to handle pandemic | 0.62 | 0.49 | 0.66 | 0.47 |
| | Low confidence in health system to handle pandemic | 0.38 | 0.49 | 0.34 | 0.47 |
| Knowledge of COVID 19 | High knowledge of COVID-19 | 0.87 | 0.34 | 0.83 | 0.38 |
| | Low knowledge of COVID-19 | 0.13 | 0.34 | 0.17 | 0.38 |

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

| | | | | |
|---|------|------|------|------|
| Compliance across strict restriction periods | | | | |
| Consistently high compliance with COVID-19 guidelines | 0.72 | 0.45 | 0.65 | 0.48 |
| Consistently low compliance with COVID-19 guidelines | 0.28 | 0.45 | 0.35 | 0.48 |
| Have had COVID-19 | | | | |
| Have not had COVID-19 | 0.79 | 0.41 | 0.77 | 0.42 |
| Have had COVID-19, suspected by self or doctor | 0.10 | 0.30 | 0.11 | 0.31 |
| Have had COVID-19, confirmed by test | 0.11 | 0.31 | 0.12 | 0.33 |
| Stressed in the past week about catching/becoming ill from COVID-19 | | | | |
| Stressed | 0.35 | 0.48 | 0.32 | 0.47 |
| Not stressed | 0.65 | 0.48 | 0.68 | 0.47 |
| Initial COVID-19 vaccine intention | | | | |
| Initially willing to accept or had accepted a COVID-19 vaccine | 0.92 | 0.27 | 0.90 | 0.30 |
| Initially uncertain about accepting a COVID-19 vaccine | 0.05 | 0.22 | 0.07 | 0.25 |
| Initially unwilling to accept a COVID-19 vaccine | 0.03 | 0.16 | 0.03 | 0.18 |
| COVID-19 booster vaccine intentions | | | | |
| Willing/accepted a COVID-19 booster vaccine | 0.95 | 0.21 | 0.92 | 0.27 |
| Uncertain about a COVID-19 booster vaccine | 0.02 | 0.16 | 0.04 | 0.20 |
| Unwilling to accept COVID-19 booster vaccine | 0.02 | 0.15 | 0.04 | 0.19 |

Note. Ethnic minority groups refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education. Data in the weighted sample were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. GCSE refers to General Certificate of Secondary Education.

It is made available under a [CC-BY 4.0 International license](#) .

Table 2. Proportion of adults in each COVID-19 booster vaccine intent category by initial COVID-19 vaccine intention (weighted, N = 22,319)

| | Initially willing to accept a COVID-19 vaccine | Initially uncertain about accepting a COVID-19 vaccine | Initially unwilling to accept a COVID-19 vaccine |
|--|--|--|---|
| | Prop. | Prop. | Prop. |
| COVID-19 booster vaccine intention | | | |
| Willing to accept or have accepted a booster | 94.77% | 73.48% | 64.80% |
| Uncertain about accepting a booster | 2.87% | 14.37% | 15.39% |
| Unwilling to accept a COVID-19 booster | 2.37% | 12.15% | 19.81% |

Note. Data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics

Table 3. Socio-demographic predictors of uncertainty and unwillingness to receive a COVID-19 booster vaccine using a multivariable multinomial regression (weighted, $N = 22,139$)

| | | Uncertain | | | Unwilling | | |
|---|--|----------------|--------|------|-----------|--------|-------|
| | | RRR | 95% CI | | RRR | 95% CI | |
| Gender (ref male) | Female | 1.06 | 0.74 | 1.52 | 0.96 | 0.61 | 1.50 |
| Age (ref 60+) | 45-59 | 1.60 | 1.00 | 2.54 | 1.57 | 0.91 | 2.72 |
| | 30-44 | 4.31 | 2.63 | 7.05 | 2.56 | 1.31 | 5.00 |
| | 18-29 | 5.30 | 2.87 | 9.77 | 5.74 | 2.82 | 11.68 |
| Ethnicity (ref white) | Ethnic minority groups | 1.07 | 0.58 | 1.95 | 1.50 | 0.74 | 3.05 |
| Education (ref undergraduate degree or higher) | A-levels or vocational | 1.51 | 0.95 | 2.39 | 2.09 | 1.15 | 3.79 |
| | Up to GCSE | 1.76 | 1.09 | 2.84 | 2.50 | 1.31 | 4.79 |
| | Income (ref >£90,000) | £60,000-89,999 | 0.50 | 0.21 | 1.22 | 0.35 | 0.11 |
| | £30,000-59,999 | 2.43 | 1.06 | 5.59 | 0.58 | 0.22 | 1.54 |
| | £16,000-29,999 | 2.25 | 0.88 | 5.78 | 1.38 | 0.49 | 3.87 |
| | <£16,000 | 3.48 | 1.32 | 9.16 | 2.20 | 0.80 | 6.03 |
| Employment status (ref not employed) | Employed | 3.25 | 2.11 | 5.02 | 1.57 | 0.92 | 2.68 |
| Living arrangement (ref alone) | Living with others, not including children | 0.77 | 0.43 | 1.39 | 0.77 | 0.42 | 1.39 |
| | Living with others, including children | 1.74 | 0.95 | 3.20 | 1.67 | 0.91 | 3.09 |
| Area of dwelling (ref urban) | Rural | 0.87 | 0.57 | 1.31 | 1.11 | 0.65 | 1.88 |
| Smoking status (ref former or non-smoker) | Current smoker | 1.61 | 0.96 | 2.72 | 1.08 | 0.57 | 2.05 |
| | Key worker status (ref not a key worker) | Key worker | 0.77 | 0.49 | 1.21 | 1.26 | 0.70 |
| Long-term physical health condition (ref condition) | No long-term physical health condition | 1.67 | 1.17 | 2.40 | 1.52 | 1.00 | 2.30 |
| Long-term mental health condition (ref condition) | No long-term mental health condition | 1.19 | 0.80 | 1.79 | 1.43 | 0.82 | 2.50 |

Note. 'Very likely' to accept a COVID-19 booster vaccine was the reference group in the multinomial regression model. Ethnic minority groups refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education. Data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

Table 4. COVID-19 related predictors of uncertainty and unwillingness to receive a COVID-19 booster vaccine using a multivariable multinomial regression (weighted, $N = 22,139$)

| | Uncertain | | | Unwilling | | |
|--|-----------|--------|-------|-----------|--------|-------|
| | RRR | 95% CI | | RRR | 95% CI | |
| Confidence in government to handle pandemic (ref high) | | | | | | |
| Low confidence in government to handle pandemic | 1.19 | 0.81 | 1.75 | 0.87 | 0.53 | 1.44 |
| Confidence in health system to handle pandemic (ref high) | | | | | | |
| Low confidence in health system to handle pandemic | 1.32 | 0.90 | 1.94 | 0.85 | 0.54 | 1.34 |
| Knowledge of COVID 19 (ref high) | | | | | | |
| Low knowledge of COVID-19 | 1.78 | 1.19 | 2.67 | 1.19 | 0.70 | 2.05 |
| Compliance across strict restriction periods (ref high) | | | | | | |
| Consistently low compliance with COVID-19 guidelines | 1.51 | 1.04 | 2.19 | 2.45 | 1.65 | 3.65 |
| Have had COVID-19 (ref have not had COVID-19) | | | | | | |
| Have had COVID-19, suspected by self or doctor | 1.38 | 0.89 | 2.15 | 1.24 | 0.76 | 2.03 |
| Have had COVID-19, confirmed by test | 1.11 | 0.66 | 1.88 | 1.06 | 0.59 | 1.91 |
| Stressed in the past week about catching/becoming ill from COVID-19 (stressed) | | | | | | |
| Not stressed | 1.80 | 1.19 | 2.72 | 1.91 | 1.19 | 3.07 |
| Initial COVID-19 vaccine intention (ref willing) | | | | | | |
| Initially uncertain about accepting a COVID-19 vaccine | 4.92 | 2.98 | 8.11 | 5.29 | 3.07 | 9.09 |
| Initially unwilling to accept a COVID-19 vaccine | 6.40 | 3.94 | 10.41 | 11.29 | 6.79 | 18.78 |

Note. 'Very likely' to accept a COVID-19 booster vaccine was the reference group in the multinomial regression model. Ethnic minority groups refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education. Data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Model includes all socio-demographic variables in Table 3.

Supplementary Material

Table S1. Comparison of excluded and included participants, unweighted

| | | Excluded | | Included | |
|--|---|----------|------|----------|------|
| | | Mean | SD | Mean | SD |
| Gender | | | | | |
| | Male | 0.23 | 0.42 | 0.26 | 0.44 |
| | Female | 0.77 | 0.42 | 0.74 | 0.44 |
| Age | | | | | |
| | 60+ | 0.36 | 0.48 | 0.43 | 0.50 |
| | 45-59 | 0.35 | 0.48 | 0.34 | 0.47 |
| | 30-44 | 0.22 | 0.42 | 0.19 | 0.39 |
| | 18-29 | 0.07 | 0.25 | 0.04 | 0.19 |
| Ethnicity | | | | | |
| | White | 0.95 | 0.21 | 0.97 | 0.18 |
| | Ethnic minority groups | 0.05 | 0.21 | 0.03 | 0.18 |
| Education | | | | | |
| | Undergraduate degree or higher | 0.64 | 0.48 | 0.71 | 0.46 |
| | A-levels or vocational | 0.19 | 0.39 | 0.16 | 0.37 |
| | Up to GCSE | 0.17 | 0.38 | 0.13 | 0.34 |
| Income | | | | | |
| | >£90,000 | 0.09 | 0.29 | 0.10 | 0.30 |
| | £60,000-89,999 | 0.15 | 0.35 | 0.15 | 0.35 |
| | £30,000-59,999 | 0.34 | 0.47 | 0.35 | 0.48 |
| | £16,000-29,999 | 0.25 | 0.43 | 0.26 | 0.44 |
| | <£16,000 | 0.17 | 0.38 | 0.14 | 0.35 |
| Employment status | | | | | |
| | Not employed | 0.41 | 0.49 | 0.42 | 0.49 |
| | Employed | 0.59 | 0.49 | 0.58 | 0.49 |
| Living arrangement | | | | | |
| | Alone | 0.18 | 0.38 | 0.23 | 0.42 |
| | With others, not with children | 0.56 | 0.50 | 0.57 | 0.49 |
| | With, including children | 0.26 | 0.44 | 0.20 | 0.40 |
| Area of dwelling | | | | | |
| | Urban | 0.74 | 0.44 | 0.75 | 0.43 |
| | Rural | 0.26 | 0.44 | 0.25 | 0.43 |
| Smoking status | | | | | |
| | Former or non-smoker | 0.90 | 0.30 | 0.93 | 0.25 |
| | Current smoker | 0.10 | 0.30 | 0.07 | 0.25 |
| Key worker status | | | | | |
| | Not a key worker | 0.76 | 0.43 | 0.80 | 0.40 |
| | Key worker | 0.24 | 0.43 | 0.20 | 0.40 |
| Long-term physical health condition | | | | | |
| | Long-term physical health condition | 0.57 | 0.50 | 0.57 | 0.50 |
| | No long-term physical health condition | 0.43 | 0.50 | 0.43 | 0.50 |
| Long-term mental health condition | | | | | |
| | Long-term mental health condition | 0.81 | 0.39 | 0.85 | 0.36 |
| | No long-term mental health condition | 0.19 | 0.39 | 0.15 | 0.36 |
| Confidence in government to handle pandemic | | | | | |
| | Low confidence in government to handle pandemic | 0.31 | 0.46 | 0.33 | 0.47 |
| | Low confidence in government to handle pandemic | 0.69 | 0.46 | 0.67 | 0.47 |
| Confidence in health system to handle pandemic | | | | | |
| | High confidence in health system to handle pandemic | 0.52 | 0.50 | 0.62 | 0.49 |
| | Low confidence in health system to handle pandemic | 0.48 | 0.50 | 0.38 | 0.49 |
| Knowledge of COVID 19 | | | | | |

It is made available under a [CC-BY 4.0 International license](#) .

| | | | | | |
|---|--|------|------|------|------|
| | High knowledge of COVID-19 | 0.86 | 0.35 | 0.87 | 0.34 |
| | Low knowledge of COVID-19 | 0.14 | 0.35 | 0.13 | 0.34 |
| Compliance across strict restriction periods | | | | | |
| | Consistently high compliance with COVID-19 guidelines | 0.75 | 0.44 | 0.72 | 0.45 |
| | Consistently low compliance with COVID-19 guidelines | 0.25 | 0.44 | 0.28 | 0.45 |
| Have had COVID-19 | | | | | |
| | Have not had COVID-19 | 0.74 | 0.44 | 0.79 | 0.41 |
| | Have had COVID-19, suspected by self or doctor | 0.12 | 0.32 | 0.10 | 0.30 |
| | Have had COVID-19, confirmed by test | 0.14 | 0.34 | 0.11 | 0.31 |
| Stressed in the past week about catching/becoming ill from COVID-19 | | | | | |
| | Stressed | 0.32 | 0.47 | 0.35 | 0.48 |
| | Not stressed | 0.68 | 0.47 | 0.65 | 0.48 |
| Initial COVID-19 vaccine intention | | | | | |
| | Initially willing to accept or had accepted a COVID-19 vaccine | 0.80 | 0.40 | 0.92 | 0.27 |
| | Initially uncertain about accepting a COVID-19 vaccine | 0.08 | 0.26 | 0.05 | 0.22 |
| | Initially unwilling to accept a COVID-19 vaccine | 0.12 | 0.33 | 0.03 | 0.16 |
| COVID-19 booster vaccine intentions | | | | | |
| | Willing/accepted a COVID-19 booster vaccine | 0.85 | 0.36 | 0.95 | 0.21 |
| | Uncertain about a COVID-19 booster vaccine | 0.04 | 0.19 | 0.02 | 0.16 |
| | Unwilling to accept COVID-19 booster vaccine | 0.11 | 0.32 | 0.02 | 0.15 |

Note: 'Very likely' to accept a COVID-19 booster vaccine was the reference group in the multinomial regression model. Ethnic minority groups refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education.

Table S2. Wording of study-developed items

| Variable | Question wording | Response options |
|---|---|--|
| Intention to receive a COVID-19 booster vaccine | We are interested in your thoughts on the BOOSTER VACCINATION. How likely do you think you are to get a COVID-19 booster vaccine if/when you are offered one? | 1- Very unlikely 6- very likely I have already had one/accepted one |
| Have ever had COVID-19 | Overall, do you believe you have ever had Covid-19? | Yes, confirmed by a positive Covid-19 test at the time I felt ill Yes, confirmed by a positive antibody test since Yes, suspected by a doctor but not tested Yes, my own suspicions No, not that I know of |
| Confidence in government to handle pandemic | To what extent do you have confidence in the UK Government's ability to handle the Coronavirus situation as it continues to develop? | 1- none at all to 7- lots |
| Confidence in health system to handle pandemic | How much confidence do you have that the UK health service can cope during Covid-19? If you live in a devolved nation, we ask you to focus on the health service within your country (e.g., NHS Health Scotland / NHS Wales / HSCNI)? | 1- none at all to 7- lots |
| Compliance with COVID-19 guidelines | Are you following the recommendations from authorities to prevent spread of COVID-19? | 1- none at all to 7- very much so |
| Knowledge of COVID-19 | How would you rate your knowledge of COVID-19? | 1- very poor knowledge to 7- very good knowledge |

It is made available under a [CC-BY 4.0 International license](#) .

Table S3. Dates of strict restrictions in England

| Restriction period | Dates |
|--------------------|-------------------------------|
| First | 23 March to 9 May 2020 |
| Second | 2 November to 3 December 2020 |
| Third | 5 January to 12 April 2021 |

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Table S4. Comparison of sample participants by COVID-19 booster vaccine intention (weighted, N = 22,139)

| | | Willing | | Uncertain | | Unwilling | |
|--|---|---------|------|-----------|------|-----------|------|
| | | Mean | SD | Mean | SD | Mean | SD |
| Gender | Male | 0.50 | 0.50 | 0.42 | 0.49 | 0.43 | 0.50 |
| | Female | 0.50 | 0.50 | 0.58 | 0.49 | 0.57 | 0.50 |
| Age | 60+ | 0.31 | 0.46 | 0.07 | 0.26 | 0.13 | 0.34 |
| | 45-59 | 0.27 | 0.44 | 0.20 | 0.40 | 0.22 | 0.42 |
| | 30-44 | 0.23 | 0.42 | 0.44 | 0.50 | 0.27 | 0.44 |
| | 18-29 | 0.19 | 0.39 | 0.29 | 0.45 | 0.38 | 0.49 |
| Ethnicity | White | 0.88 | 0.33 | 0.85 | 0.36 | 0.80 | 0.40 |
| | Ethnic minority groups | 0.12 | 0.33 | 0.15 | 0.36 | 0.20 | 0.40 |
| Education | Undergraduate degree or higher | 0.35 | 0.48 | 0.31 | 0.46 | 0.22 | 0.42 |
| | A-levels or vocational | 0.33 | 0.47 | 0.40 | 0.49 | 0.44 | 0.50 |
| | Up to GCSE | 0.32 | 0.47 | 0.29 | 0.45 | 0.33 | 0.47 |
| Income | >£90,000 | 0.07 | 0.26 | 0.05 | 0.21 | 0.08 | 0.27 |
| | £60,000-89,999 | 0.13 | 0.34 | 0.05 | 0.22 | 0.06 | 0.24 |
| | £30,000-59,999 | 0.33 | 0.47 | 0.45 | 0.50 | 0.20 | 0.40 |
| | £16,000-29,999 | 0.28 | 0.45 | 0.27 | 0.44 | 0.37 | 0.48 |
| | <£16,000 | 0.18 | 0.38 | 0.18 | 0.39 | 0.29 | 0.45 |
| Employment status | Not employed | 0.43 | 0.49 | 0.18 | 0.38 | 0.33 | 0.47 |
| | Employed | 0.57 | 0.49 | 0.82 | 0.38 | 0.67 | 0.47 |
| Living arrangement | Alone | 0.20 | 0.40 | 0.18 | 0.39 | 0.20 | 0.40 |
| | With others, not with children | 0.60 | 0.49 | 0.42 | 0.49 | 0.46 | 0.50 |
| | With, including children | 0.20 | 0.40 | 0.40 | 0.49 | 0.34 | 0.47 |
| Area of dwelling | Urban | 0.79 | 0.41 | 0.84 | 0.37 | 0.81 | 0.40 |
| | Rural | 0.21 | 0.41 | 0.16 | 0.37 | 0.19 | 0.40 |
| Smoking status | Former or non-smoker | 0.89 | 0.31 | 0.81 | 0.39 | 0.86 | 0.35 |
| | Current smoker | 0.11 | 0.31 | 0.19 | 0.39 | 0.14 | 0.35 |
| Key worker status | Not a key worker | 0.79 | 0.41 | 0.72 | 0.45 | 0.69 | 0.46 |
| | Key worker | 0.21 | 0.41 | 0.28 | 0.45 | 0.31 | 0.46 |
| Long-term physical health condition | No long-term physical health condition | 0.42 | 0.49 | 0.23 | 0.42 | 0.28 | 0.45 |
| | Long-term physical health condition | 0.58 | 0.49 | 0.77 | 0.42 | 0.72 | 0.45 |
| Long-term mental health condition | No long-term mental health condition | 0.19 | 0.39 | 0.20 | 0.40 | 0.21 | 0.41 |
| | Long-term mental health condition | 0.81 | 0.39 | 0.80 | 0.40 | 0.79 | 0.41 |
| Confidence in government to handle pandemic | Low confidence in government to handle pandemic | 0.36 | 0.48 | 0.24 | 0.43 | 0.34 | 0.47 |
| | High confidence in government to handle pandemic | 0.64 | 0.48 | 0.76 | 0.43 | 0.66 | 0.47 |
| Confidence in health system to handle pandemic | High confidence in health system to handle pandemic | 0.67 | 0.47 | 0.53 | 0.50 | 0.65 | 0.48 |
| | Low confidence in health system to handle pandemic | 0.33 | 0.47 | 0.47 | 0.50 | 0.35 | 0.48 |
| Knowledge of COVID 19 | High knowledge of COVID-19 | 0.84 | 0.37 | 0.63 | 0.48 | 0.67 | 0.47 |

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

| | | | | | | |
|---|------|------|------|------|------|------|
| Low knowledge of COVID-19 | 0.16 | 0.37 | 0.37 | 0.48 | 0.33 | 0.47 |
| Compliance across strict restriction periods | | | | | | |
| Consistently high compliance with COVID-19 guidelines | 0.67 | 0.47 | 0.45 | 0.50 | 0.35 | 0.48 |
| Consistently low compliance with COVID-19 guidelines | 0.33 | 0.47 | 0.55 | 0.50 | 0.65 | 0.48 |
| Have had COVID-19 | | | | | | |
| Have not had COVID-19 | 0.78 | 0.42 | 0.68 | 0.47 | 0.69 | 0.46 |
| Have had COVID-19, suspected by self or doctor | 0.10 | 0.31 | 0.14 | 0.35 | 0.12 | 0.32 |
| Have had COVID-19, confirmed by test | 0.12 | 0.32 | 0.18 | 0.38 | 0.19 | 0.39 |
| Stressed in the past week about catching/becoming ill from COVID-19 | | | | | | |
| Stressed | 0.33 | 0.47 | 0.22 | 0.42 | 0.18 | 0.38 |
| Not stressed | 0.67 | 0.47 | 0.78 | 0.42 | 0.82 | 0.38 |
| Initial COVID-19 vaccine intention | | | | | | |
| Initially willing to accept or had accepted a COVID-19 vaccine | 0.92 | 0.27 | 0.63 | 0.48 | 0.59 | 0.49 |
| Initially uncertain about accepting a COVID-19 vaccine | 0.05 | 0.23 | 0.24 | 0.43 | 0.23 | 0.42 |
| Initially unwilling to accept a COVID-19 vaccine | 0.02 | 0.15 | 0.13 | 0.33 | 0.19 | 0.39 |

Note. 'Very likely' to accept a COVID-19 booster vaccine was the reference group in the multinomial regression model. Ethnic minority groups refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education. Data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

Table S5. Sensitivity analysis: Socio-demographic, COVID-19 related, and initial COVID-19 vaccine intent predictors of uncertainty and unwillingness to receive a COVID-19 booster vaccine using a multivariable multinomial regression and last recorded initial COVID-19 vaccine intent (weighted, $N = 22,319$)

| | | Uncertain | | | Unwilling | | |
|--|--|-----------|--------|-------|-----------|--------|-------|
| | | RRR | 95% CI | | RRR | 95% CI | |
| Gender (ref male) | Female | 1.12 | 0.77 | 1.62 | 1.06 | 0.68 | 1.65 |
| Age (ref 60+) | 45-59 | 1.43 | 0.90 | 2.25 | 1.33 | 0.77 | 2.28 |
| | 30-44 | 3.09 | 1.82 | 5.25 | 1.61 | 0.81 | 3.21 |
| | 18-29 | 3.20 | 1.76 | 5.83 | 3.24 | 1.46 | 7.23 |
| Ethnicity (ref white) | Ethnic minority groups | 0.80 | 0.42 | 1.50 | 1.13 | 0.53 | 2.42 |
| Education (ref undergraduate degree or higher) | A-levels or vocational | 1.39 | 0.87 | 2.20 | 1.69 | 0.98 | 2.92 |
| | Up to GCSE | 1.57 | 1.00 | 2.45 | 2.03 | 1.14 | 3.59 |
| Income (ref >£90,000) | £60,000-89,999 | 0.49 | 0.20 | 1.23 | 0.34 | 0.12 | 0.99 |
| | £30,000-59,999 | 2.54 | 1.10 | 5.87 | 0.69 | 0.29 | 1.66 |
| | £16,000-29,999 | 2.27 | 0.89 | 5.79 | 1.41 | 0.53 | 3.78 |
| | <£16,000 | 3.40 | 1.25 | 9.19 | 2.37 | 0.89 | 6.30 |
| Employment status (ref not employed) | Employed | 3.29 | 2.10 | 5.14 | 1.57 | 0.91 | 2.71 |
| Living arrangement (ref alone) | Living with others, not including children | 0.83 | 0.47 | 1.45 | 0.83 | 0.45 | 1.53 |
| | Living with others, including children | 1.76 | 0.95 | 3.25 | 1.66 | 0.87 | 3.16 |
| Area of dwelling (ref urban) | Rural | 0.86 | 0.56 | 1.34 | 1.01 | 0.55 | 1.87 |
| Smoking status (ref former or non-smoker) | Current smoker | 1.65 | 1.02 | 2.68 | 1.22 | 0.63 | 2.35 |
| Key worker status (ref not a key worker) | Key worker | 0.74 | 0.49 | 1.13 | 1.20 | 0.67 | 2.15 |
| Long-term physical health condition (ref condition) | No long-term physical health condition | 1.41 | 0.98 | 2.05 | 1.09 | 0.71 | 1.68 |
| Long-term mental health condition (ref condition) | No long-term mental health condition | 1.34 | 0.87 | 2.06 | 1.52 | 0.83 | 2.79 |
| Confidence in government to handle pandemic (ref high) | Low confidence in government to handle pandemic | 1.19 | 0.81 | 1.76 | 0.85 | 0.51 | 1.42 |
| Confidence in health system to handle pandemic (ref high) | Low confidence in health system to handle pandemic | 1.32 | 0.91 | 1.93 | 0.83 | 0.53 | 1.30 |
| Knowledge of COVID 19 (ref high) | Low knowledge of COVID-19 | 1.81 | 1.22 | 2.68 | 1.11 | 0.63 | 1.95 |
| Compliance across strict restriction periods (ref high) | Consistently low compliance with COVID-19 guidelines | 1.58 | 1.09 | 2.31 | 2.82 | 1.87 | 4.25 |
| Have had COVID-19 (ref have not had COVID-19) | Have had COVID-19, suspected by self or doctor | 1.37 | 0.87 | 2.14 | 1.30 | 0.80 | 2.14 |
| | Have had COVID-19, confirmed by test | 1.16 | 0.69 | 1.93 | 1.12 | 0.63 | 1.99 |
| Stressed in the past week about catching/becoming ill from COVID-19 (ref stressed) | Not stressed | 1.79 | 1.17 | 2.75 | 1.85 | 1.15 | 2.95 |
| Initial COVID-19 vaccine intention (ref willing) | Initially uncertain about accepting a COVID-19 vaccine | 4.41 | 2.08 | 9.35 | 10.86 | 5.18 | 22.78 |
| | Initially unwilling to accept a COVID-19 vaccine | 6.47 | 2.46 | 17.02 | 10.67 | 3.98 | 28.59 |

Note. 'Very likely' to accept a COVID-19 booster vaccine was the reference group in the multinomial regression model. Ethnic minority groups refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education. Data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Table S6. Sensitivity analysis: Socio-demographic, COVID-19 related, and initial COVID-19 vaccine intent predictors of uncertainty and unwillingness to receive a COVID-19 booster vaccine using a multivariable multinomial regression using average compliance with government guidelines across the entirety of the pandemic (weighted, $N = 22,139$)

| | | Uncertain | | | Unwilling | | |
|--|--|-----------|--------|-------|-----------|--------|-------|
| | | RRR | 95% CI | | RRR | 95% CI | |
| Gender (ref male) | Female | 1.05 | 0.72 | 1.53 | 0.98 | 0.64 | 1.50 |
| Age (ref 60+) | 45-59 | 1.45 | 0.91 | 2.31 | 1.46 | 0.83 | 2.56 |
| | 30-44 | 3.33 | 1.97 | 5.64 | 2.05 | 1.01 | 4.16 |
| | 18-29 | 3.91 | 2.14 | 7.12 | 4.73 | 2.12 | 10.52 |
| Ethnicity (ref white) | Ethnic minority groups | 0.77 | 0.40 | 1.48 | 1.18 | 0.59 | 2.35 |
| Education (ref undergraduate degree or higher) | A-levels or vocational | 1.33 | 0.84 | 2.13 | 1.78 | 1.04 | 3.06 |
| | Up to GCSE | 1.47 | 0.93 | 2.30 | 2.12 | 1.20 | 3.77 |
| Income (ref >£90,000) | £60,000-89,999 | 0.50 | 0.20 | 1.26 | 0.34 | 0.11 | 1.06 |
| | £30,000-59,999 | 2.52 | 1.08 | 5.85 | 0.70 | 0.28 | 1.74 |
| | £16,000-29,999 | 2.22 | 0.87 | 5.70 | 1.39 | 0.52 | 3.71 |
| | <£16,000 | 3.07 | 1.13 | 8.29 | 2.09 | 0.78 | 5.56 |
| Employment status (ref not employed) | Employed | 3.15 | 2.00 | 4.94 | 1.43 | 0.84 | 2.46 |
| Living arrangement (ref alone) | Living with others, not including children | 0.73 | 0.42 | 1.28 | 0.72 | 0.39 | 1.33 |
| | Living with others, including children | 1.67 | 0.89 | 3.12 | 1.61 | 0.85 | 3.08 |
| Area of dwelling (ref urban) | Rural | 0.87 | 0.55 | 1.36 | 1.01 | 0.56 | 1.81 |
| Smoking status (ref former or non-smoker) | Current smoker | 1.79 | 1.11 | 2.91 | 1.27 | 0.66 | 2.44 |
| Key worker status (ref not a key worker) | Key worker | 0.74 | 0.48 | 1.14 | 1.21 | 0.67 | 2.18 |
| Long-term physical health condition (ref condition) | No long-term physical health condition | 1.40 | 0.96 | 2.04 | 1.18 | 0.77 | 1.79 |
| Long-term mental health condition (ref condition) | No long-term mental health condition | 1.37 | 0.88 | 2.13 | 1.54 | 0.86 | 2.75 |
| Confidence in government to handle pandemic (ref high) | Low confidence in government to handle pandemic | 1.19 | 0.81 | 1.75 | 0.87 | 0.53 | 1.44 |
| Confidence in health system to handle pandemic (ref high) | Low confidence in health system to handle pandemic | 1.32 | 0.90 | 1.94 | 0.85 | 0.54 | 1.34 |
| Knowledge of COVID 19 (ref high) | Low knowledge of COVID-19 | 1.78 | 1.19 | 2.67 | 1.19 | 0.70 | 2.05 |
| Compliance across strict restriction periods (ref high) | Consistently low compliance with COVID-19 guidelines | 1.51 | 1.04 | 2.19 | 2.45 | 1.65 | 3.65 |
| Have had COVID-19 (ref have not had COVID-19) | Have had COVID-19, suspected by self or doctor | 1.38 | 0.89 | 2.15 | 1.24 | 0.76 | 2.03 |
| | Have had COVID-19, confirmed by test | 1.11 | 0.66 | 1.88 | 1.06 | 0.59 | 1.91 |
| Stressed in the past week about catching/becoming ill from COVID-19 (ref stressed) | Not stressed | 1.80 | 1.19 | 2.72 | 1.91 | 1.19 | 3.07 |
| Initial COVID-19 vaccine intention (ref willing) | Initially uncertain about accepting a COVID-19 vaccine | 4.92 | 2.98 | 8.11 | 5.29 | 3.07 | 9.09 |
| | Initially unwilling to accept a COVID-19 vaccine | 6.40 | 3.94 | 10.41 | 11.29 | 6.79 | 18.78 |

Note. 'Very likely' to accept a COVID-19 booster vaccine was the reference group in the multinomial regression model. Ethnic minority groups refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education. Data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

Table S7. Sensitivity analysis: Socio-demographic, COVID-19 related, and initial COVID-19 vaccine intent predictors of uncertainty and unwillingness to receive a COVID-19 booster vaccine using a multivariable multinomial regression including only participants who had had exactly two doses of a COVID-19 vaccine at follow-up (weighted, $N = 12,297$)

| | | Uncertain | | | Unwilling | | |
|--|--|-----------|--------|------|-----------|--------|-------|
| | | RRR | 95% CI | | RRR | 95% CI | |
| Gender (ref male) | Female | 1.12 | 0.76 | 1.63 | 1.04 | 0.67 | 1.60 |
| Age (ref 60+) | 45-59 | 0.82 | 0.50 | 1.34 | 1.06 | 0.59 | 1.91 |
| | 30-44 | 1.69 | 0.98 | 2.91 | 1.29 | 0.62 | 2.66 |
| | 18-29 | 1.85 | 0.97 | 3.50 | 2.69 | 1.15 | 6.30 |
| Ethnicity (ref white) | Ethnic minority groups | 0.85 | 0.44 | 1.64 | 1.15 | 0.57 | 2.31 |
| Education (ref undergraduate degree or higher) | A-levels or vocational | 1.24 | 0.77 | 1.99 | 1.70 | 0.98 | 2.95 |
| | Up to GCSE | 1.37 | 0.86 | 2.20 | 1.83 | 1.00 | 3.34 |
| Income (ref >£90,000) | £60,000-89,999 | 0.55 | 0.23 | 1.35 | 0.34 | 0.11 | 1.08 |
| | £30,000-59,999 | 2.49 | 1.08 | 5.75 | 0.71 | 0.28 | 1.79 |
| | £16,000-29,999 | 2.20 | 0.86 | 5.67 | 1.50 | 0.55 | 4.06 |
| | <£16,000 | 2.91 | 1.06 | 7.97 | 1.95 | 0.71 | 5.33 |
| Employment status (ref not employed) | Employed | 3.12 | 1.92 | 5.06 | 1.49 | 0.85 | 2.62 |
| Living arrangement (ref alone) | Living with others, not including children | 0.77 | 0.43 | 1.38 | 0.83 | 0.41 | 1.68 |
| | Living with others, including children | 1.69 | 0.89 | 3.23 | 1.77 | 0.86 | 3.61 |
| Area of dwelling (ref urban) | Rural | 0.85 | 0.53 | 1.34 | 1.03 | 0.54 | 1.94 |
| Smoking status (ref former or non-smoker) | Current smoker | 1.95 | 1.18 | 3.22 | 1.43 | 0.73 | 2.79 |
| Key worker status (ref not a key worker) | Key worker | 0.91 | 0.58 | 1.41 | 1.45 | 0.80 | 2.63 |
| Long-term physical health condition (ref condition) | No long-term physical health condition | 1.07 | 0.71 | 1.60 | 0.90 | 0.57 | 1.41 |
| Long-term mental health condition (ref condition) | No long-term mental health condition | 1.39 | 0.88 | 2.21 | 1.49 | 0.81 | 2.74 |
| Confidence in government to handle pandemic (ref high) | Low confidence in government to handle pandemic | 1.13 | 0.76 | 1.69 | 0.88 | 0.52 | 1.50 |
| Confidence in health system to handle pandemic (ref high) | Low confidence in health system to handle pandemic | 1.33 | 0.90 | 1.97 | 0.85 | 0.53 | 1.37 |
| Knowledge of COVID 19 (ref high) | Low knowledge of COVID-19 | 1.73 | 1.15 | 2.61 | 1.14 | 0.65 | 2.01 |
| Compliance across strict restriction periods (ref high) | Consistently low compliance with COVID-19 guidelines | 1.50 | 1.03 | 2.20 | 2.43 | 1.61 | 3.66 |
| Have had COVID-19 (ref have not had COVID-19) | Have had COVID-19, suspected by self or doctor | 1.23 | 0.78 | 1.94 | 1.12 | 0.67 | 1.87 |
| | Have had COVID-19, confirmed by test | 1.07 | 0.63 | 1.83 | 1.16 | 0.63 | 2.12 |
| Stressed in the past week about catching/becoming ill from COVID-19 (ref stressed) | Not stressed | 1.90 | 1.24 | 2.91 | 1.83 | 1.11 | 3.00 |
| Initial COVID-19 vaccine intention (ref willing) | Initially uncertain about accepting a COVID-19 vaccine | 4.62 | 2.77 | 7.70 | 5.31 | 3.05 | 9.25 |
| | Initially unwilling to accept a COVID-19 vaccine | 5.36 | 3.25 | 8.84 | 9.15 | 5.29 | 15.84 |

Note. 'Very likely' to accept a COVID-19 booster vaccine was the reference group in the multinomial regression model. Ethnic minority groups refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education. Data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics.