

1 **Activity of convalescent and vaccine serum against a B.1.1.529** 2 **variant SARS-CoV-2 isolate**

3 Juan Manuel Carreño¹, Hala Alshammary¹, Johnstone Tcheou¹, Gagandeep Singh¹, Ariel Raskin¹,
4 Hisaaki Kawabata¹, Levy Sominsky¹, Jordan Clark¹, Daniel C. Adelsberg¹, Dominika Bielak¹, Ana
5 Silvia Gonzalez-Reiche⁵, PSP/PARIS Study Group¹, Komal Srivastava¹, Emilia Mia Sordillo^{3*},
6 Goran Bajic^{1*}, Harm van Bakel^{5*}, Viviana Simon^{1,2,3,4*}, Florian Kramer^{1,3*}

7 *PSP/PARIS Study Group (in alphabetical order): A. Amoako, R. Banu, K. Beach, M.C. Bermúdez-*
8 *González, G. Cai, I. Ceglia, C. Cognigni, K. Farrugia, C. Gleason, A. van de Guchte, G. Kleiner,*
9 *Z. Khalil, N. Lyttle, W. Mendez, L.C.F. Mulder, A. Oostenink, A. Rooker, A.*
10 *Salimbangon, M. Saksena, A. Paniz-Mondolfi, J. Polanco*

11
12 ¹ Department of Microbiology, Icahn School of Medicine at Mount Sinai, NY (ISMMS)

13 ² Division of Infectious Diseases, Department of Medicine, ISMMS

14 ³ Department of Pathology, Molecular and Cell based Medicine, ISMMS

15 ⁴ The Global Health and Emerging Pathogens Institute, ISMMS

16 ⁵ Department of Genetics and Genomic Sciences, ISMMS

17 * shared senior authorship, correspondence to emilia.sordillo@mountsinai.org,

18 goran.bajic@mssm.edu, harm.vanbakel@mssm.edu, Viviana.simon@mssm.edu and

19 florian.kramer@mssm.edu

20

21 **The B.1.1.529 (Omicron) variant of severe acute respiratory syndrome coronavirus 2 (SARS-**
22 **CoV-2) was identified in November of 2021 in South Africa and Botswana as well as in a sample**
23 **of a traveler from South Africa in Hong Kong.^{1,2} Since then, B.1.1.529 has been detected in many**
24 **countries globally. This variant seems to be more infectious than B.1.617.2 (Delta), has already**
25 **caused super spreader events³ and has outcompeted Delta within weeks in several countries**
26 **and metropolitan areas. B.1.1.529 hosts an unprecedented number of mutations in its spike**
27 **gene and early reports have provided evidence for extensive immune escape and reduced**
28 **vaccine effectiveness.^{2,4-6} Here, we investigated the neutralizing and binding activity of sera**
29 **from convalescent, mRNA double vaccinated, mRNA boosted as well as convalescent double**
30 **vaccinated and convalescent boosted individuals against wild type, B.1.351 and B.1.1.529**
31 **SARS-CoV-2 isolates. Neutralizing activity of sera from convalescent and double vaccinated**
32 **participants was undetectable to very low against B.1.1.529 while neutralizing activity of sera**
33 **from individuals who had been exposed to spike three or four times was maintained, albeit at**
34 **strongly reduced levels. Binding to the B.1.1.529 receptor binding domain (RBD) and N-**
35 **terminal domain (NTD) was reduced in convalescent not vaccinated but was mostly retained in**
36 **vaccinated individuals.**

37 Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged in late 2019 in Wuhan,
38 China and has since then caused the coronavirus disease 2019 (COVID-19) pandemic. While SARS-
39 CoV-2 was antigenically relatively stable during its first few months of circulation, the first
40 antigenically distinct variants, B.1.1.7 (Alpha), B.1.351 (Beta) and P.1 (Gamma), emerged in late
41 2020. Other variants of interest (VOI) and variants of concern (VOCs) followed. So far, B.1.351

42 has shown the most antigenic drift in terms of reduction of *in vitro* neutralization rivaled only by
43 B.1.621 (Mu).⁷ B.1.617.2 (Delta), which emerged in early 2020 has been the most consequential
44 variant given the fact that it is more infectious than the viruses circulating in the beginning of the
45 pandemic and also partially escapes neutralization *in vitro*.⁸ B.1.1529 (Omicron) was first
46 detected in South Africa, Botswana and in a traveler from South Africa in Hong Kong.^{1,2} The
47 variant hosts a large number of mutations in its spike protein including at least 15 amino acid
48 changes in the receptor binding domain (RBD). These mutations are predicted to affect most
49 neutralizing antibody epitopes. In addition, B.1.1.529 seems to be fit and highly transmissible³
50 and it rapidly spread across the globe. In fact, within four weeks this variant has outcompeted
51 B.1.617.2 and is now the dominant circulating variant in several countries and urban areas.

52 Immunity to SARS-CoV-2 in human populations is highly variable and likely differs in individuals
53 with infection induced immunity, double vaccinated individuals, boosted individuals and
54 individuals with hybrid immunity due to the combination of infection followed by vaccination.
55 Understanding residual neutralizing and binding activity against highly antigenically distinct viral
56 variants such as B.1.1.529 in these distinct groups is essential to gauge the level of protection
57 that a specific community has against infection, mild or severe COVID-19.

58 To address these questions, we determined the loss of *in vitro* neutralizing and binding activity
59 for B.1.1529 in sera from individuals with different levels of immunity (infection, vaccine, hybrid).
60 We included samples from convalescent individuals (N=15), individuals vaccinated twice with
61 BNT162b2 (Pfizer/BioNTech mRNA vaccine, N=10), individuals vaccinated twice with mRNA-1273
62 (Moderna mRNA vaccine, N=10), individuals vaccinated three times with BNT162b2 (boosted,
63 N=10), individuals vaccinated three times with mRNA-1273 (boosted, N=10), convalescent
64 individuals who got 2 doses of BNT162b2 (N=10), convalescent individuals who got 2 doses of
65 mRNA-1273 (N=10) and finally convalescent individuals who got 3 doses of BNT162b2 (boosted,
66 N=10) (**Figure 1A** and **Supplementary Tables 1 and 2**). First, we tested the *in vitro* neutralizing
67 activity of the sera against wild type SARS-CoV-2 (as a reference for ancestral strains), B.1.351 (as
68 a reference for the most pronounced *in vitro* escape phenotype) and B.1.1.529 (isolated from
69 one of the first cases identified in New York City in late November 2021, **Supplementary Table**
70 **3**). The neutralization assay used mimics physiological conditions, since it is performed with
71 authentic SARS-CoV-2 in a multicycle replication setting in which serum/antibody is present at all
72 times akin to the situation in a seropositive individual. Across all 85 samples, the reduction in
73 neutralization for B.1.1.529 was greater than 14.5-fold (the actual fold reduction could not be
74 calculated since many samples were below the limit of detection) (**Figure 1B**). In comparison,
75 there was “only” a four-fold reduction against B.1.351 in the same sample set. In fact, 16.5% of
76 samples lost all neutralizing activity against B.1.1529. When looking at the different groups, we
77 noted that convalescent individuals had lower titers against wild type and B.1.351 with the
78 majority (73.3%) of samples having no measurable neutralizing activity for B.1.1529 (**Figure 1C**).
79 For samples from individuals double vaccinated with BNT162b2 and mRNA-1273, we observed a
80 more than 23-fold and a 42-fold reduction in neutralizing activity respectively (**Figure 1D and E**).
81 However, most individuals retained low but detectable neutralizing activity. Boosted individuals
82 experienced lower reduction with a 7.5-fold drop in neutralization for BNT162b2 boosted
83 individuals and a 16.7-fold reduction in mRNA-1273 boosted individuals (**Figure 1F and G**). Of

84 note, the lower fold change and the higher starting neutralization titers led to considerable
85 residual neutralizing activity in those groups. Convalescent individuals who received two
86 BNT162b2, two mRNA-1272 or three BNT162b2 vaccine doses showed 14-fold, 11-fold and 13-
87 fold drops in B.1.1.529 neutralization (**Figure 1H, I and J**). However all individuals in these groups
88 maintained relatively robust neutralization activity. These data indicate that convalescent
89 individuals greatly benefit from vaccination, an observation that is of significant public health
90 importance.

91 While *in vitro* neutralization is an important antibody function, antibody binding – even in the
92 absence of detectable neutralizing activity – can provide protection through Fc-mediated effector
93 functions. This type of protection has been described in detail for influenza virus⁹⁻¹¹ but binding
94 antibody titers also represent a correlate of protection for SARS-CoV-2.^{12,13} Furthermore,
95 retained binding to a highly mutated RBD or NTD, even if reduced, indicates that cognate B-cells
96 are present. These B-cells could likely be rapidly recalled during variant infection or variant
97 specific vaccination producing a strong plasmablast response leading to rapid control of viral
98 spread. In addition, B-cells with low affinity binding to antigenically drifted variant proteins may
99 enter lymph nodes and engage in germinal center reactions leading to antibodies that may regain
100 neutralizing activity through affinity maturation. To investigate the reduction in binding, we
101 expressed a recombinant RBD of B.1.1.529 and compared binding of this RBD with binding to wild
102 type and B.1.351 RBD (**Figure 2A**). Overall, reduction in binding to B.1.1.529 RBD was much less
103 pronounced than reduction in neutralization (**Figure 2B**). However, this reduction was
104 significantly higher than what we observed here and in the past for B.1.351.⁸ Reduction in binding
105 was most pronounced for convalescent individuals (**Figure 2C**) with a drop of more than 7.5-fold
106 and undetectable reactivity by enzyme linked immunosorbent assay (ELISA) in two thirds of the
107 convalescent individuals who were infected early in the pandemic prior to the circulation of viral
108 variants of concern. In all other groups binding was relatively well maintained with a reduction
109 in binding ranging from a 2.9-fold drop in individuals who had received two vaccinations with
110 mRNA-1273 to a 1.5-fold drop in individuals boosted with BNT162b2 (**Figures 2D to 2J**).

111 In addition to the RBD, the NTD is a prime target for B-cells after COVID-19 mRNA vaccination.¹⁴
112 The NTD also hosts neutralizing epitopes within and outside of the immunodominant ‘super site’.
113 The NTD of B.1.1.529 carries a large number of amino acid substitutions, three deletions and one
114 three amino acid long insertion which are, collectively, predicted to significantly change the
115 ‘super site’ as well as neutralizing epitopes outside of the ‘super site’. To determine if infection
116 induced and vaccine induced antibodies retain binding to the B.1.1.529 NTD, we expressed both
117 the wild type and variant NTDs to be probed by ELISA using the same 85 samples testing for
118 neutralization. Surprisingly, binding to the NTD was maintained with relatively minor reductions
119 (maximum 1.9-fold), suggesting either maintained binding (e.g. at lower affinity) to the ‘super
120 site’ or the presence of a large number of unchanged epitopes within this domain (**Figure 3**).

121 Our data aligns well with initial reports on the impact of B.1.1.529 on *in vitro* neutralizing activity
122 of convalescent and vaccine serum and expands on these initial reports by inclusion of
123 subcohorts with divergent SARS-CoV-2 exposure history including infection induced, primary
124 vaccine regimen as well as booster induced and hybrid immunity.^{2,5,6} We found that neutralizing
125 activity of B.1.1.529 is most impacted in unvaccinated, convalescent individuals and in naive

126 individuals who acquired immunity through two mRNA COVID-19 vaccine doses. Our findings
127 support recent reports describing significantly reduced protection from reinfection¹⁵ and
128 basically non-existing vaccine effectiveness against symptomatic disease after two BNT162b2
129 vaccinations.⁴ However, boosted individuals had, at least, within the short time after the booster
130 dose, significant protection against symptomatic disease in the range of 75%.⁴ While it is unclear
131 how long this protection lasts, we observe titers similar to those in boosted individuals in
132 convalescent vaccinated individuals, suggesting that those individuals may experience significant
133 protection.

134 This study also provides first insights into B.1.1.529 RBD and NTD specific binding changes.
135 Compared to the changes in neutralizing activities, binding was surprisingly well preserved
136 especially against NTD, in general, and against the RBD in vaccinated, boosted and convalescent
137 vaccinated individuals. It is conceivable, that these binding antibodies which often have non-
138 neutralizing phenotypes in cell culture, contribute to protection from disease as has been seen
139 for other viral infections.⁹⁻¹¹ In addition, the presence of strong binding antibodies suggest that,
140 while some antibodies may have lost affinity for the drifted epitopes, B-cells may be recalled
141 when encountering B.1.1.519 spike through infection or vaccination. This could lead to a strong
142 anamnestic response, which could positively impact COVID-19 progression. It could also lead to
143 the recruitment of these B-cells into germinal centers for further affinity maturation resulting in
144 potent, high affinity neutralizing antibodies against B.1.1.529.¹⁶ Importantly, our data adds to the
145 growing body of evidence suggesting that B.1.1.529 specific vaccines are urgently needed.

146

147 **Materials and methods**

148 **Human Serum Samples:** Convalescent and post-vaccine sera were collected from participants in
149 the longitudinal observational PARIS (Protection Associated with Rapid Immunity to SARS-CoV-2)
150 study.^{8,17} This cohort follows health care workers longitudinally since April 2020. The study was
151 reviewed and approved by the Mount Sinai Hospital Institutional Review Board (IRB-20-03374).
152 All participants signed written consent forms prior to sample and data collection. All participants
153 provided permission for sample banking and sharing.

154 For the antigenic characterization of the antigenically diverse B.1.1.529 variant, we selected 85
155 serum samples from 54 participants. 20/54 participants were seronegative prior to vaccination
156 while 34/54 had COVID-19 prior to vaccination (see **Supplemental Tables 1 and 2** for
157 demographics and vaccine information). All participants with pre-vaccination immunity were
158 infected in 2020 when only ancestral SARS-CoV-2 strains circulated in the New York metropolitan
159 area. Convalescent samples (N=15) were obtained within three months of SARS-CoV-2 infection
160 (average: 58 days, range: 23-87 days) while the post vaccinations samples were collected, on
161 average, 23 days (range: 14-39 days) after the second dose (N= 40, 20 Pfizer 2x and 20 Moderna
162 2x) or 19 days (range: 14-33 days) after the third booster (N= 30, 20 Pfizer 3x and 10 Moderna
163 3x) vaccine dose.

164 **Cells:** Vero-E6 cells expressing TMPRSS2 were cultured in Dulbecco's modified Eagles medium
165 (DMEM; Corning, #10-013-CV) containing 10% heat-inactivated fetal bovine serum (FBS);

166 GeminiBio, #100-106) and 1% minimum essential medium (MEM) amino acids solution (Gibco,
167 #11130051), supplemented with 100 U/ml penicillin and 100 µg/ml streptomycin (Gibco,
168 #15140122), 100 µg/ml normocin (InvivoGen, #ant-nr), and 3 µg/ml puromycin (InvivoGen, #ant-
169 pr). FreeStyle™ 293-F cells (Gibco, #R79007) were cultured in ESF-SFM medium (Expression
170 Systems, cat. no. 98-001) supplemented with 100 U/ml penicillin and 100 µg/ml streptomycin
171 (Gibco, #15140122). Expi293F™ Cells (Gibco, #A14527) were cultured in Expi293™ Expression
172 Medium (Gibco, #A1435102) supplemented with 100 U/ml penicillin and 100 µg/ml streptomycin
173 (Gibco, #15140122).

174 **Selection and culture of replication competent SARS-CoV-2 isolates.** The Mount Sinai Pathogen
175 Surveillance program (IRB approved, HS#13-00981) actively screens nasopharyngeal swab
176 specimens from patients seeking care at the Mount Sinai Health System for emerging viral
177 variants. After completion of the diagnostics, de-identified biospecimen were sequenced either
178 using an established complete virus genome sequencing approach¹⁸ (e.g., Beta isolate USA/NY-
179 MSHSPSP-PV27007/2021) or based on the spike S1 mutational profile determined by Spike-ID
180 (Omicron, manuscript in preparation). The B.1.1.529 isolate USA/NY-MSHSPSP-PV44488/2021
181 represents one of the first cases diagnosed in New York State (female, age bracket: 30-40 years,
182 mild COVID-19 symptoms, vaccinated and boosted) in late November 2021. The SARS-CoV-2 virus
183 USA-WA1/2020 was used as wild-type reference (BEI resources, NR-52281). **Supplemental Table**
184 **3** summarizes the amino acid substitutions, insertions and deletions in the spike region of each
185 of the three viral isolates

186 Viruses were grown by adding 200ul of viral transport media from the nasopharyngeal swabs to
187 Vero-E6-TMPRSS2 cells in culture media supplemented with 0.5 µg/ml amphotericin B (Gibco, #
188 15290-018). Cytopathic effects (CPE) appears within 4-6 days at which point the culture
189 supernatants was clarified by centrifugation at 4,000 g for 5 minutes. Expanded viral stocks used
190 were sequence-verified and titered by the 50% tissue culture infectious dose (TCID₅₀) method on
191 Vero-E6-TMPRSS2 cells prior to use in micro neutralization assays.

192 **Generation of recombinant variant RBD and NTD proteins:** The recombinant RBD proteins were
193 produced using Expi293F cells (Life Technologies). The proteins were cloned into a mammalian
194 expression vector, pCAGGS as described earlier^{19,20} and purified after transient transfections with
195 each respective plasmid. Six-hundred million Expi293F cells were transfected using the
196 ExpiFectamine 293 Transfection Kit and purified DNA. Supernatants were collected on day four
197 post transfection, centrifuged at 4,000 g for 20 minutes and finally filtered using a 0.22 µm filter.
198 Ni-nitrilotriacetic acid (Ni-NTA) agarose (Qiagen) was used to purify the proteins via gravity flow
199 and proteins were eluted as previously described.^{19,20} The buffer was exchanged using Amicon
200 centrifugal units (EMD Millipore) and all recombinant proteins were finally re-suspended in
201 phosphate buffered saline (PBS). Proteins were also run on a sodium dodecyl sulphate (SDS)
202 polyacrylamide gels (5–20% gradient; Bio-Rad) to check for purity.^{21,22} The NTD protein
203 constructs (residues 1-306) were cloned into pVRC8400 expression vector between *Sall* and *NotI*
204 endonuclease restriction sites yielding an NTD with an human rhinovirus (HRV) 3C protease-
205 cleavable C-terminal hexahistidine and a streptavidin-binding protein tags. The NTDs were
206 transiently expressed in FreeStyle™ 293-F cells. Four days post-transfection, supernatants were
207 harvested by centrifugation and further purified using immobilized metal affinity

208 chromatography (IMAC) with cobalt-TALON® resin (Takara) followed by Superdex 200 Increase
209 10/300 GL size exclusion column (GE Healthcare).

210 **Enzyme linked immunosorbent assay (ELISA):** Antibody titers in sera from convalescent
211 individuals and vaccinees were measured by a research grade ELISA using recombinant versions
212 of the RBD and NTD of wild type SARS-CoV-2 as well as the B.1.351 (Beta), and B.1.1.529
213 (Omicron) (see **Supplemental Table 4** for specific substitutions in each variant). All samples were
214 analyzed in a blinded manner. Briefly, 96-well microtiter plates (Corning, #353227) were coated
215 with 50 µl/well of recombinant protein (2 µg/ml) overnight at 4 °C. Plates were washed three
216 times with phosphate-buffered saline (PBS; Gibco, #10010-031) supplemented with 0.1% Tween-
217 20 (PBS-T; Fisher Scientific ref. 202666) using an automatic plate washer (BioTek 405TS
218 microplate washer). For blocking, PBS-T containing 3% milk powder (American Bio, #
219 AB1010901000) was used. After 1-hour incubation at room temperature (RT), blocking solution
220 was removed and initial dilutions (1:100) of heat-inactivated sera (in PBS-T 1%-milk powder) were
221 added to the plates, followed by 2-fold serial dilutions. After 2-hour incubation, plates were
222 washed three times with PBS-T and 50 µl/well of the pre-diluted secondary antibody anti-human
223 IgG (Fab-specific) horseradish peroxidase (HRP) antibody (produced in goat; Sigma-Aldrich, Cat#
224 A0293, RRID: AB_257875) diluted 1:3,000 in PBS-T containing 1% milk powder were added. After
225 1-hour incubation at RT, plates were washed three times with PBS-T and SigmaFast o-
226 phenylenediamine dihydrochloride (Sigmafast OPD; Sigma-Aldrich, Ref. P9187-50SET) was added
227 (100 µl/well) for 10min, followed by addition of 50 µl/well of 3 M hydrochloric acid (Thermo
228 Fisher, Ref. S25856) to stop the reaction. Optical density was measured at a wavelength of 490
229 nm using a plate reader (BioTek, SYNERGY H1 microplate reader). The area under the curve (AUC)
230 values were calculated and plotted using Prism 9 software (GraphPad).

231

232 **SARS-CoV-2 multi-cycle Microneutralization assay:** Sera from vaccinees were used to assess the
233 neutralization of wild type (WA1), B.1.351 (Beta) and B.1.1.529 (Omicron) SARS-CoV-2 isolates
234 (**Supplementary Table 3**). All procedures were performed in a biosafety level 3 (BSL-3) facility at
235 the Icahn School of Medicine at Mount Sinai following standard safety guidelines. Vero-E6-
236 TMPRSS2 cells were seeded in 96-well high binding cell culture plates (Costar, #07620009) at a
237 density of 20,000 cells/well in complete Dulbecco's modified Eagle medium (cDMEM) one day
238 prior to the infection. Heat inactivated serum samples (56°C for 1 hour) were serially diluted (3-
239 fold) in minimum essential media (MEM; Gibco, #11430-030) supplemented with 2 mM L-
240 glutamine (Gibco, #25030081), 0.1% sodium bicarbonate (w/v, HyClone, #SH30033.01), 10 mM
241 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES; Gibco, #15630080), 100 U/ml
242 penicillin, 100 µg/ml streptomycin (Gibco, #15140122) and 0.2% bovine serum albumin (BSA, MP
243 Biomedicals, Cat#. 810063) starting at 1:10. Remdesivir (Medkoo Bioscience inc., #329511) was
244 included to monitor assay variation. Serially diluted sera were incubated with 10,000 TCID₅₀ of
245 WT USA-WA1/2020 SARS-CoV-2, MSHSPSP-PV27007/2021 (B.1.351, Beta) or USA/NY-MSHSPSP-
246 PV44488/2021 (B.1.1.529, Omicron) for one hour at RT, followed by the transfer of 120µl of the
247 virus-sera mix to Vero-E6-TMPRSS2 plates. Infection proceeded for one hour at 37°C and
248 inoculum was removed. 100 µl/well of the corresponding antibody dilutions plus 100µl/well of
249 infection media supplemented with 2% fetal bovine serum (FBS; Gibco, #10082-147) were added

250 to the cells. Plates were incubated for 48h at 37°C followed by fixation overnight at 4°C in 200
251 µl/well of a 10% formaldehyde solution. For staining of the nucleoprotein, formaldehyde solution
252 was removed, and cells were washed with PBS (pH 7.4) (Gibco, #10010-031) and permeabilized
253 by adding 150 µl/well of PBS, 0.1% Triton X-100 (Fisher Bioreagents, #BP151-100) for 15 min at
254 RT. Permeabilization solution was removed, plates were washed with 200 µl/well of PBS (Gibco,
255 #10010-031) twice and blocked with PBS, 3% BSA for 1 hour at RT. During this time the primary
256 antibody was biotinylated according to manufacturer protocol (Thermo Scientific EZ-Link NHS-
257 PEG4-Biotin). Blocking solution was removed and 100 µl/well of biotinylated mAb 1C7C7²³, a
258 mouse anti-SARS nucleoprotein monoclonal antibody generated at the Center for Therapeutic
259 Antibody Development at The Icahn School of Medicine at Mount Sinai ISMMS (Millipore Sigma,
260 Cat# ZMS1075) at a concentration of 1µg/ml in PBS, 1% BSA was added for 1 hour at RT. Cells
261 were washed with 200 µl/well of PBS twice and 100 µl/well of HRP-conjugated streptavidin
262 (Thermo Fisher Scientific) diluted in PBS, 1% BSA were added at a 1:2,000 dilution for 1 hour at
263 RT. Cells were washed twice with PBS, and 100 µl/well of o-phenylenediamine dihydrochloride
264 (Sigmafast OPD; Sigma-Aldrich) were added for 10 min at RT, followed by addition of 50 µl/well
265 of a 3 M HCl solution (Thermo Fisher Scientific). Optical density (OD) was measured (490 nm)
266 using a microplate reader (Synergy H1; Biotek). Analysis was performed using Prism 7 software
267 (GraphPad). After subtraction of background and calculation of the percentage of neutralization
268 with respect to the “virus only” control, a nonlinear regression curve fit analysis was performed
269 to calculate the 50% inhibitory dilution (ID₅₀), with top and bottom constraints set to 100% and
270 0% respectively. All samples were analyzed in a blinded manner.

271 **Statistics:** A one-way ANOVA with Tukey’s multiple comparisons test was used to compare the
272 neutralization and RBD binding antibody titers. The exception is the 2x BNT162b2 RBD ELISA
273 group where a mixed effects model had to be used due to a missing data point. A student’s t test
274 was used for comparing wild type and B.1.1.529 NTD binding data. Statistical analyses were
275 performed using Prism 9 software (GraphPad).

276

277 **Acknowledgements**

278 We thank the study participants for their generosity and willingness to participate in longitudinal
279 COVID-19 research studies. None of this work would be possible without their contributions.

280 We would like to thank Dr. Randy A. Albrecht for oversight of the conventional BSL3
281 biocontainment facility, which makes our work with live SARS-CoV-2 possible. Furthermore, we
282 would like to thank Drs. Janine Kimpel and Dorothee von Laer for their willingness to speedily
283 share B.1.1.529 virus isolates (even though they could not included in this work). We are also
284 grateful for Mount Sinai's leadership during the COVID-19 pandemic. We want to especially thank
285 Drs. Peter Palese, Carlos Cordon-Cardo, Dennis Charney, David Reich and Kenneth Davis for their
286 support and Daniel Caughey for expert administrative assistance.

287 This work is part of the PARIS/SPARTA studies funded by the NIAID Collaborative Influenza
288 Vaccine Innovation Centers (CIVIC) contract 75N93019C00051. In addition, this work was also
289 partially funded by the NIAID Centers of Excellence for Influenza Research and Response (CEIRR)

290 contract and 75N93021C00014 by anonymous donors. This work is part of the NIAID SARS-CoV-
291 2 Assessment of Viral Evolution (SAVE) program.

292 **Conflict of interest statement**

293 The Icahn School of Medicine at Mount Sinai has filed patent applications relating to SARS-CoV-
294 2 serological assays and NDV-based SARS-CoV-2 vaccines which list Florian Krammer as co-
295 inventor. Viviana Simon is also listed on the serological assay patent application as co-inventor.
296 Mount Sinai has spun out a company, Kantaro, to market serological tests for SARS-CoV-2. Florian
297 Krammer has consulted for Merck and Pfizer (before 2020), and is currently consulting for Pfizer,
298 Third Rock Ventures, Seqirus and Avimex. The Krammer laboratory is also collaborating with
299 Pfizer on animal models of SARS-CoV-2.

300

301

302 **References**

- 303 1 Gu, H. *et al.* Probable Transmission of SARS-CoV-2 Omicron Variant in Quarantine Hotel,
304 Hong Kong, China, November 2021. *Emerg Infect Dis* **28**, doi:10.3201/eid2802.212422
305 (2021).
- 306 2 Cele, S. *et al.* SARS-CoV-2 Omicron has extensive but incomplete escape of Pfizer
307 BNT162b2 elicited neutralization and requires ACE2 for infection. *medRxiv*,
308 2021.2012.2008.21267417, doi:10.1101/2021.12.08.21267417 (2021).
- 309 3 Brandal, L. T. *et al.* Outbreak caused by the SARS-CoV-2 Omicron variant in Norway,
310 November to December 2021. *Eurosurveillance* **26**, 2101147,
311 doi:doi:<https://doi.org/10.2807/1560-7917.ES.2021.26.50.2101147> (2021).
- 312 4 Andrews, N. *et al.* Effectiveness of COVID-19 vaccines against the Omicron (B.1.1.529)
313 variant of concern. *medRxiv*, 2021.2012.2014.21267615,
314 doi:10.1101/2021.12.14.21267615 (2021).
- 315 5 Wilhelm, A. *et al.* Reduced Neutralization of SARS-CoV-2 Omicron Variant by Vaccine
316 Sera and Monoclonal Antibodies. *medRxiv*, 2021.2012.2007.21267432,
317 doi:10.1101/2021.12.07.21267432 (2021).
- 318 6 Rössler, A., Riepler, L., Bante, D., Laer, D. v. & Kimpel, J. SARS-CoV-2 B.1.1.529 variant
319 (Omicron) evades neutralization by sera from vaccinated and convalescent individuals.
320 *medRxiv*, 2021.2012.2008.21267491, doi:10.1101/2021.12.08.21267491 (2021).
- 321 7 Uriu, K. *et al.* Neutralization of the SARS-CoV-2 Mu Variant by Convalescent and Vaccine
322 Serum. *N Engl J Med* **385**, 2397-2399, doi:10.1056/NEJMc2114706 (2021).
- 323 8 Carreño, J. M. *et al.* Evidence for retained spike-binding and neutralizing activity against
324 emerging SARS-CoV-2 variants in serum of COVID-19 mRNA vaccine recipients.
325 *EBioMedicine* **73**, 103626, doi:10.1016/j.ebiom.2021.103626 (2021).
- 326 9 Dilillo, D. J., Tan, G. S., Palese, P. & Ravetch, J. V. Broadly neutralizing hemagglutinin
327 stalk-specific antibodies require FcγR interactions for protection against influenza virus
328 in vivo. *Nat Med*, doi:10.1038/nm.3443 (2014).
- 329 10 Asthagiri Arunkumar, G. *et al.* Broadly Cross-Reactive, Nonneutralizing Antibodies
330 against Influenza B Virus Hemagglutinin Demonstrate Effector Function-Dependent
331 Protection against Lethal Viral Challenge in Mice. *J Virol* **93**, doi:10.1128/JVI.01696-18
332 (2019).
- 333 11 Ng, S. *et al.* Novel correlates of protection against pandemic H1N1 influenza A virus
334 infection. *Nat Med* **25**, 962-967, doi:10.1038/s41591-019-0463-x (2019).
- 335 12 Gilbert, P. B. *et al.* Immune correlates analysis of the mRNA-1273 COVID-19 vaccine
336 efficacy clinical trial. *Science*, eab3435 (2021).
- 337 13 Earle, K. A. *et al.* Evidence for antibody as a protective correlate for COVID-19 vaccines.
338 *Vaccine*, doi:10.1016/j.vaccine.2021.05.063 (2021).
- 339 14 Amanat, F. *et al.* SARS-CoV-2 mRNA vaccination induces functionally diverse antibodies
340 to NTD, RBD, and S2. *Cell*, doi:10.1016/j.cell.2021.06.005 (2021).

- 341 15 Pulliam, J. R. C. *et al.* Increased risk of SARS-CoV-2 reinfection associated with
342 emergence of the Omicron variant in South Africa. *medRxiv*, 2021.2011.2011.21266068,
343 doi:10.1101/2021.11.11.21266068 (2021).
- 344 16 Turner, J. S. *et al.* SARS-CoV-2 mRNA vaccines induce persistent human germinal centre
345 responses. *Nature*, doi:10.1038/s41586-021-03738-2 (2021).
- 346 17 Krammer, F. *et al.* Antibody Responses in Seropositive Persons after a Single Dose of
347 SARS-CoV-2 mRNA Vaccine. *N Engl J Med*, doi:10.1056/NEJMc2101667 (2021).
- 348 18 Gonzalez-Reiche, A. S. *et al.* Introductions and early spread of SARS-CoV-2 in the New
349 York City area. *Science*, doi:10.1126/science.abc1917 (2020).
- 350 19 Amanat, F. *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans.
351 *Nat Med* **26**, 1033-1036, doi:10.1038/s41591-020-0913-5 (2020).
- 352 20 Stadlbauer, D. *et al.* SARS-CoV-2 Seroconversion in Humans: A Detailed Protocol for a
353 Serological Assay, Antigen Production, and Test Setup. *Curr Protoc Microbiol* **57**, e100,
354 doi:10.1002/cpmc.100 (2020).
- 355 21 Margine, I., Palese, P. & Krammer, F. Expression of functional recombinant
356 hemagglutinin and neuraminidase proteins from the novel H7N9 influenza virus using
357 the baculovirus expression system. *J Vis Exp*, e51112, doi:10.3791/51112 (2013).
- 358 22 Amanat, F. *et al.* Antibodies to the Glycoprotein GP2 Subunit Cross-React between Old
359 and New World Arenaviruses. *mSphere* **3**, doi:10.1128/mSphere.00189-18 (2018).
- 360 23 Amanat, F. *et al.* An In Vitro Microneutralization Assay for SARS-CoV-2 Serology and
361 Drug Screening. *Curr Protoc Microbiol* **58**, e108, doi:10.1002/cpmc.108 (2020).
- 362 24 Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to the
363 ACE2 receptor. *Nature* **581**, 215-220, doi:10.1038/s41586-020-2180-5 (2020).
- 364 25 Chi, X. *et al.* A neutralizing human antibody binds to the N-terminal domain of the Spike
365 protein of SARS-CoV-2. *Science* **369**, 650-655, doi:10.1126/science.abc6952 (2020).

366 Figure Legends

367 **Figure 1: Sera of convalescent and vaccinated individuals have strongly reduced neutralizing**
368 **activity against B.1.1.529 as compared to wild type SARS-CoV-2. A** Overview of different
369 exposure groups from whom samples were obtained. Further details are provided in
370 **Supplemental Table 1 and 2. B** shows absolute titers (left) and fold reduction (right) for the
371 combined samples, **C** to **J** shows the different groups. A one-way ANOVA with Tukey's multiple
372 comparisons test was used to compare the neutralization titers and significant p values (<0.05)
373 are indicated in the figure.

374 **Figure 2: Sera of vaccinated individuals mostly maintain binding to the B.1.1.529 RBD. A** shows
375 a model of the B.1.1.529 spike protein in complex with the angiotensin converting enzyme 2
376 (ACE2) receptor with B.1.1.529 specific mutations indicated. The figure is based on PDB 6M0J²⁴
377 and 7C2L²⁵ and was built in PyMol. **B** shows absolute titers (left) and fold reduction (right) for
378 the combined samples, **C** to **J** shows the different groups. A one-way ANOVA with Tukey's
379 multiple comparisons test was used to compare the neutralization titers and significant p values
380 (<0.05) are indicated in the figure. The exception is panel **D** where a mixed effects model had to
381 be used due to a missing data point.

382

383 **Figure 3: Serum of vaccinated individuals maintains binding to the B.1.1.529 NTD. A** shows
 384 absolute titers (left) and fold reduction (right) for the combined samples, **B to I** shows the
 385 different groups. A student's t test was used for comparing wild type and B.1.1.529 NTD binding
 386 data and significant p values (<0.05) are indicated in the figure.

387

388 **Supplemental Tables**

389 **Supplemental Table 1: Overall description of samples used.**

390

	COVID-19		Two vaccine doses				Three vaccine doses		
	Convalescent		BNT16 2b2		mRNA-1273		BNT16 2b2		mRNA-1273
Total N	15		naïve	convalescent	naïve	convalescent	naïve	convalescent	naïve
			10	10	10	10	10	10	10
Sex									
Female	10		8	7	8	5	8	7	8
Male	5		2	3	2	5	2	3	2
Days since infection									
Average (days)	58		no infection		no infection		no infection		no infection
Range (days)	23-87		no infection		no infection		no infection		no infection
Days since vaccine dose									
Average (days)	no vaccine		18	26	26	20	19	20	19
Range (days)	no vaccine		14-21	15-39	14-36	15-28	14-28	14-30	14-33

391

392 **Supplemental Table 2: Detailed description of samples used.**

393

Participant ID	Age Bracket	Sex	Ancestry	Time points included in this study	SARS-CoV-2 infection prior to vaccination	Vaccine Type
OMI-001	30-39	Female	Caucasian	Post-Vax, Post-Boost	No	Moderna
OMI-002	18-29	Female	Caucasian	Post-Vax, Post-Boost	No	Moderna
OMI-003	40-49	Male	Caucasian	Post-Vax, Post-Boost	No	Moderna
OMI-004	40-49	Female	Caucasian	Post-Vax, Post-Boost	No	Moderna
OMI-005	40-49	Female	Caucasian	Post-Vax, Post-Boost	No	Moderna
OMI-006	40-49	Male	Caucasian	Post-Vax, Post-Boost	No	Moderna
OMI-007	50-59	Female	Caucasian	Post-Vax, Post-Boost	No	Moderna
OMI-008	60-69	Female	Asian	Post-Vax, Post-Boost	No	Moderna
OMI-009	30-39	Female	Caucasian	Post-Vax, Post-Boost	No	Moderna
OMI-010	18-29	Female	Caucasian	Post-Vax, Post-Boost	No	Moderna
OMI-011	40-49	Female	Other	Post-Vax, Post-Boost	No	Pfizer
OMI-012	18-29	Female	Asian	Post-Vax, Post-Boost	No	Pfizer
OMI-013	60-69	Female	Caucasian	Post-Vax, Post-Boost	No	Pfizer

OMI-014	50-59	Female	Caucasian	Post-Vax, Post-Boost	No	Pfizer
OMI-015	18-29	Female	Caucasian	Post-Vax, Post-Boost	No	Pfizer
OMI-016	70-79	Male	Caucasian	Post-Vax, Post-Boost	No	Pfizer
OMI-017	30-39	Female	Asian Indian	Post-Vax, Post-Boost	No	Pfizer
OMI-018	30-39	Female	Caucasian	Post-Vax, Post-Boost	No	Pfizer
OMI-019	40-49	Male	Caucasian	Post-Vax, Post-Boost	No	Pfizer
OMI-020	40-49	Female	n.a.	Post-Vax, Post-Boost	No	Pfizer
OMI-021	40-49	Female	Asian	Post-Infection, Post-Vax	Yes	Pfizer
OMI-022	30-39	Female	Caucasian	Post-Infection, Post-Vax	Yes	Pfizer
OMI-023	30-39	Female	Asian	Post-Infection, Post-Vax	Yes	Pfizer
OMI-024	30-39	Male	Caucasian	Post-Infection, Post-Vax	Yes	Pfizer
OMI-028	18-29	Female	Caucasian	Post-Infection, Post-Vax	Yes	Moderna
OMI-029	30-39	Female	Caucasian	Post-Infection, Post-Vax	Yes	Pfizer
OMI-032	50-59	Male	Caucasian	Post-Infection, Post-Vax	Yes	Pfizer
OMI-027	40-49	Female	Caucasian	Post-Infection, Post-Vax	Yes	Pfizer
OMI-033	30-39	Female	Caucasian	Post-Infection, Post-Vax	Yes	Pfizer
OMI-034	40-49	Male	Latino	Post-Infection, Post-Vax	Yes	Pfizer
OMI-035	30-39	Female	Caucasian	Post-Infection, Post-Vax	Yes	Pfizer
OMI-025	50-59	Male	Latino	Post-Infection	Yes	no vax
OMI-026	30-39	Male	Caucasian	Post-Infection	Yes	no vax
OMI-030	40-49	Female	Other	Post-Infection	Yes	no vax
OMI-031	30-39	Female	Mexican	Post-Infection	Yes	no vax
OMI-036	50-59	Male	Asian	Post-Vax	Yes	Moderna
OMI-037	30-39	Male	Caucasian	Post-Vax	Yes	Moderna
OMI-038	18-29	Male	Caucasian	Post-Vax	Yes	Moderna
OMI-039	30-39	Female	Caucasian	Post-Vax	Yes	Moderna
OMI-040	60-69	Female	African American	Post-Vax	Yes	Moderna
OMI-041	40-49	Female	Caucasian	Post-Vax	Yes	Moderna
OMI-042	40-49	Female	Caucasian	Post-Vax	Yes	Moderna
OMI-043	60-69	Male	Caucasian	Post-Vax	Yes	Moderna
OMI-044	40-49	Male	n.a.	Post-Vax	Yes	Moderna
OMI-045	30-39	Male	Caucasian	Post-Boost	Yes	Pfizer
OMI-046	30-39	Female	Caucasian	Post-Boost	Yes	Pfizer
OMI-047	50-59	Female	n.a.	Post-Boost	Yes	Pfizer
OMI-048	60-69	Male	Caucasian	Post-Boost	Yes	Pfizer
OMI-049	60-69	Male	Caucasian	Post-Boost	Yes	Pfizer
OMI-050	30-39	Female	African American	Post-Boost	Yes	Pfizer
OMI-051	50-59	Female	Caucasian	Post-Boost	Yes	Pfizer
OMI-052	50-59	Female	Caucasian	Post-Boost	Yes	Pfizer
OMI-053	18-29	Female	Caucasian	Post-Boost	Yes	Pfizer
OMI-054	30-39	Female	Caucasian	Post-Boost	Yes	Pfizer

394 **Table 2 Legend:** Summary of the metadata of the 54 PARIS participants from whom a total of 85 serum samples
 395 were analyzed. For 31 PARIS participants, samples from two different time points were included (Post-Vax, Post-
 396 Boost or Post-Infection, Post-Vax).

397 The following abbreviations are used in the table:
 398 Post-infection: serum collected 23 to 87 days after SARS-CoV-2 diagnosis
 399 Post-Vax: serum collected 14-39 days after the second dose of mRNA vaccine
 400 Post-Boost: serum collected 14-30 days after the booster vaccine dose
 401 no vax: participant was not vaccinated at the time of sample collection
 402 n.a: data not available

403

404

405 **Supplementary Table 3:** Information on the viral isolates used in neutralization assays.

Isolate #	GISAID ID	Isolate ID	Lineage	RBD substitution s	NTD deletion /insertio n	NTD substitution s	Spike substitutions
NR-52281	EPI_ISL_404895	USA-WA1/2020	A	None	None	None	None
PV27007	EPI_ISL_1708926	USA/NY-MSHSPSP-PV27007/2021	B.1.351	K417N, E484K, N501Y	ΔL241-A243	D80A, D215G	D80A, D215G, K417, E484K, N501Y, D614G, A701V
PV44488	pending	USA/NY-MSHSPSP-PV44488/2021	B.1.1.529	G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H	ΔH69-V70, ΔG142-Y144, ΔN211, ins214EP E	A67V, T95I, Y145D, L212I	G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H, T547K, D614G,

							H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H, N969K, L981F
--	--	--	--	--	--	--	---



















407

408 **Supplemental Table 4:** Overview of the mutations encoded in the RBD and NTD proteins used
















409 for the binding assays.

Mutations in the sequence of recombinant RBD and spike proteins generated for this study					
Protein	Lineage	RBD substitutions	NTD deletions	NTD insertions	NTD substitutions
RBD	WA1	None	NA	NA	NA
RBD	B.1.351	K417K, E484K, N501Y	NA	NA	NA
RBD	B.1.1.529	G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, G496S, Q498R, N501Y, Y505H	NA	NA	NA
NTD	WA1	NA	None	None	None
NTD	B.1.1.529	NA	69-70, 143-145, 211	214EPE	A67V, T95I, G142D, L212I



















410



medRxiv preprint doi: <https://doi.org/10.1101/2021.12.20.21264134>; this version posted December 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

A**B****C****G****D****H****E****I****F****J**

medRxiv preprint doi: <https://doi.org/10.1101/2021.12.20.21268134>; this version posted December 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).



medRxiv preprint doi: <https://doi.org/10.1101/2021.12.20.21268134>; this version posted December 21, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).