Wastewater sequencing uncovers early, cryptic SARS-CoV-2 variant transmission

Smruthi Karthikeyan[1,#], Joshua I Levy[2,#], Peter De Hoff[3,9,21], Greg Humphrey[1], Amanda
Birmingham[4], Kristen Jepsen[5], Sawyer Farmer[1], Helena M. Tubb[1], Tommy Valles[1], Caitlin E
Tribelhorn[1], Rebecca Tsai[1], Stefan Aigner[3], Shashank Sathe[3], Niema Moshiri[6], Benjamin
Henson[5], Adam M. Mark[4], Abbas Hakim[3,9,21], Nathan A Baer[3], Tom Barber[3], Pedro Belda-
Ferre[3], Marisol Chacón[3], Willi Cheung[3,9,21], Evelyn S Cresini[3], Emily R Eisner,[3] Alma L
Lastrella[3], Elijah S Lawrence[3], Clarisse A Marotz[3], Toan T Ngo[3], Tyler Ostrander[3], Ashley
Plascencia[3], Rodolfo A Salido[3], Phoebe Seaver[3], Elizabeth W Smoot[3], Daniel McDonald[1],
Robert M Neuhard[7,12], Angela L Scioscia[8,9], Alysson M. Satterlund[10], Elizabeth H Simmons[11] ,
Dismas B. Abelman[12], David Brenner[12], Judith C. Bruner[12], Anne Buckley[12], Michael Ellison[12],
Jeffrey Gattas[12], Steven L. Gonias[13], Matt Hale[12], Faith Hawkins[12], Lydia Ikeda[12], Hemlata
Jhaveri[12], Ted Johnson[12], Vince Kellen[12], Brendan Kremer[12], Gary Matthews[12], Ronald W.
McLawhon[12], Pierre Ouillet[12], Daniel Park[12], Allorah Pradenas[12], Sharon Reed[12], Lindsay
Riggs[12], Alison Sanders[12], Bradley Sollenberger[12], Angela Song[7,12], Benjamin White[12], Terri
Winbush[12], Christine M Aceves[2], Catelyn Anderson[2], Karthik Gangavarapu[2], Emory Hufbauer[2],
Ezra Kurzban[2], Justin Lee[2], Nathaniel L Matteson[2], Edyth Parker[2], Sarah A Perkins[2], Karthik S
Ramesh[2], Refugio Robles-Sikisaka[2], Madison A Schwab[2], Emily Spencer[2], Shirlee Wohl[2], Laura
Nicholson[2], Ian H Mchardy[2], David P Dimmock[15], Charlotte A Hobbs[15], Omid Bakhtar[16], Aaron
Harding[16], Art Mendoza[16], Alexandre Bolze[17], David Becker[17], Elizabeth T Cirulli[17], Magnus
Isaksson[17], Kelly M Schiabor Barrett[17], Nicole L Washington[17], John D Malone[18], Ashleigh
Murphy Schafer[18], Nikos Gurfield[18], Sarah Stous[18], Rebecca Fielding-Miller[19,20], Richard S.
Garfein[19], Tommi Gaines[20], Cheryl Anderson[19], Natasha K. Martin[19], Robert Schooley[19] , Brett
Austin[16], Duncan R. MacCannell[22], Stephen F Kingsmore[15] , William Lee[17], Seema Shah[18], Eric
McDonald[18], Alexander T. Yu[21], Mark Zeller[2], Kathleen M Fisch[4,9], Christopher Longhurst[1,23],
Patty Maysent[24], David Pride[25], Pradeep K. Khosla[6], Louise C. Laurent[3,9,26], Gene W Yeo[3,26,27],
Kristian G Andersen[2,*], Rob Knight[1,6,28,*]

[#]equal contribution
*Senior author

[1] Department of Pediatrics, University of California San Diego, La Jolla, CA, USA
[2] Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA,
USA
[3] Expedited COVID Identification Environment (EXCITE) Laboratory, Department of Pediatrics,
University of California San Diego, La Jolla, CA, USA
[4] Center for Computational Biology and Bioinformatics, University of California San Diego, La
Jolla, CA, USA
[5] Institute for Genomic Medicine, University of California San Diego, La Jolla, CA, USA
[6] Department of Computer Science and Engineering, University of California San Diego, La
Jolla, CA, USA
[7] Operational Strategic Initiatives, University of California San Diego, La Jolla, CA, USA
[8] Student Health and Well-Being, University of California San Diego, La Jolla, CA, USA
[9] Department of Obstetrics, Gynecology, and Reproductive Sciences, University of California
San Diego, La Jolla, CA, USA

[10] Student Affairs, University of California San Diego, La Jolla, CA, USA
[11] Academic Affairs, University of California San Diego, La Jolla, CA, USA
[12] Return to Learn, University of California San Diego, La Jolla, CA, USA
[13] Department of Pathology, University of California San Diego, La Jolla, CA, USA
[14] Scripps Health, San Diego, La Jolla, CA, USA
[15] Rady Children's Institute for Genomic Medicine, San Diego, CA, USA
[16] Sharp Healthcare, San Diego, CA, USA
[17] Helix, San Mateo, CA, USA
[18] County of San Diego Health and Human Services Agency, San Diego, CA, USA
[19] Herbert Wertheim School of Public Health and Human Longevity Science, University of
California San Diego, La Jolla, CA, USA
[20] Division of Infectious Disease and Global Public Health, University of California San Diego,
La Jolla, CA, USA
[21] COVID-19 Detection, Investigation, Surveillance, Clinical, and Outbreak Response, California
Department of Public Health, Richmond, CA, USA
[22] Office of Advanced Molecular Detection, Centers for Disease Control and Prevention, Atlanta,
GA, USA
[23] Department of Biomedical Informatics, University of California, San Diego, La Jolla,
California, USA
[24] Office of the UC San Diego Health CEO, University of California, San Diego
[25] Departments of Pathology and Medicine, University of California, San Diego, La Jolla, CA
[26] Sanford Consortium of Regenerative Medicine, University of California San Diego, La Jolla,
CA
[27] Department of Cellular and Molecular Medicine, University of California San Diego, La Jolla,
CA
[28] Department of Bioengineering, University of California San Diego, La Jolla, CA, USA
73
74 **Address correspondence to:**
75
Rob Knight
Department of Pediatrics
University of California San Diego
9500 Gilman Drive, MC 0763
La Jolla, CA 92093, USA
robknight@ucsd.edu
+1 858-246-1184

83
84
85 **Summary**
86

As SARS-CoV-2 continues to spread and evolve, detecting emerging variants early is critical for
public health interventions. Inferring lineage prevalence by clinical testing is infeasible at scale,
especially in areas with limited resources, participation, or testing/sequencing capacity, which
can also introduce biases. SARS-CoV-2 RNA concentration in wastewater successfully tracks
regional infection dynamics and provides less biased abundance estimates than clinical testing.
Tracking virus genomic sequences in wastewater would improve community prevalence

estimates and detect emerging variants. However, two factors limit wastewater-based genomic
surveillance: low-quality sequence data and inability to estimate relative lineage abundance in
mixed samples. Here, we resolve these critical issues to perform a high-resolution, 295-day
wastewater and clinical sequencing effort, in the controlled environment of a large university
campus and the broader context of the surrounding county. We develop and deploy improved
virus concentration protocols and deconvolution software that fully resolve multiple virus strains
from wastewater. We detect emerging variants of concern up to 14 days earlier in wastewater
samples, and identify multiple instances of virus spread not captured by clinical genomic
surveillance. Our study provides a scalable solution for wastewater genomic surveillance that
allows early detection of SARS-CoV-2 variants and identification of cryptic transmission.
103
104   **Introduction**
105
SARS-CoV-2 continues to evolve, producing diverse new lineages[1]. Emerging variants of
concern (VOCs) and variants of interest (VOIs) demonstrate increased transmissibility, disease
severity, and/or immune escape[2]. Timely and accurate quantification of local prevalence of
SARS-CoV-2 variants is thus essential for effective public health measures. However, existing
strategies for variant detection based on virus genome sequencing of biospecimens obtained from
clinical testing ("clinical genomic surveillance") are expensive, inefficient, and have sampling
bias because of systemic healthcare disparities, particularly in poor and underserved
communities[3–5].
114
In contrast, PCR-based wastewater surveillance of SARS-CoV-2 RNA is not subject to clinical
testing biases and can track temporal changes in overall SARS-CoV-2 prevalence in a region [6–8],
but cannot identify epidemiological transmission links or monitor virus lineage prevalence,
which require genome sequence information. Virus genome sequencing from wastewater
("wastewater genomic surveillance") has the potential to cost-effectively capture community
virus spread[9,10], acting as a surrogate to clinical surveillance in elucidating lineage geospatial
distributions and track emerging SARS-CoV-2 variants (including new variants for which
targeted assays do not yet exist), and provide genome sequence data needed for transmission
network analysis and interpretation[11].
124
However, wastewater genomic surveillance is technically challenging[10]. Low viral loads, heavily
fragmented RNA, and PCR inhibitors in complex environmental samples lead to poor
sequencing coverage[12,13]. Obtaining high quality sequences from samples with low viral load and
elevated levels of PCR inhibitors remains an outstanding technical challenge in implementation
of wastewater genomic surveillance at scale. Additionally, tools for SARS-CoV-2 lineage
classification, such as pangolin[14] and UShER[15], were designed for clinical samples containing a
single dominant variant, and cannot estimate relative abundances of multiple SARS-CoV-2
lineages in samples with virus mixtures such as wastewater.
133
Here, we report a high-resolution approach to study community virus transmission using
wastewater genomic surveillance, leveraging several technical advances in wastewater virus
concentration and nucleic acid sequencing, and a computational tool for resolving multiple
SARS-CoV-2 lineages in short-read sequence data from a mixed sample (lineage deconvolution).
We obtained near 95% genome coverage even for samples with low viral load, compared with

40% or below from previous studies[11-13], a key advance that allowed us to build a robust pipeline
to monitor virus lineage prevalence in community wastewater.
141
Because places of communal living, such as university campuses, are considered key sites for
virus spread and represent well-controlled and relatively isolated environments, they are ideal for
comparing the relative utility of clinical and wastewater genomic surveillance[16]. Accordingly, we
conducted a high-resolution, longitudinal wastewater genomic surveillance effort at the
University of California San Diego (UCSD) campus, in parallel with clinical genomic
surveillance from nasal swabs in the local community, from November 2020 to September 2021:
ten months that effectively capture the surges in the region caused by the three main VOCs (as
determined by US CDC) in the United States, Epsilon, Alpha and Delta[1]. In more recent San
Diego-wide data collected from September 2021 to February 2022, we studied ongoing
transmission of the Delta variant and the rapid spread of the Omicron variant and its sublineages.
152
Our wastewater genomic surveillance approach identified VOCs up to 2 weeks prior to detection
through clinical genomic surveillance, even though a large proportion of clinical SARS-CoV-2
samples are sequenced in San Diego relative to other cities in the United States. In addition to
providing a detailed history of community virus spread, wastewater genomic surveillance also
identified multiple instances of cryptic community transmission not observed through clinical
genomic surveillance. Matching wastewater and clinical genome sequences provided
epidemiological information identifying specific transmission events. Our results demonstrate
the viability of wastewater genomic surveillance at scale, enabling early detection and tracking
of virus lineages and guiding clinical genomic surveillance efforts. This work informed public
health guidance and interventions on the UCSD campus as well as San Diego county in real
time, and our data and analyses were disseminated to both public health officials as well as the
general public via custom dashboards (see **Data Availability** for links).
165
166 **Results**
167
To directly compare wastewater genomic surveillance to clinical surveillance, we conducted a
large-scale SARS-CoV-2 genome sequencing study from wastewater samples collected daily
from 131 wastewater samplers covering 360 campus buildings, in many cases reaching single
building-level resolution. To identify epidemiological transmission links and monitor lineages in
the population, we sequenced all SARS-CoV-2 positive clinical and wastewater samples from
campus using a miniaturized tiled-amplicon sequencing approach. During this period of this
study, we collected and analyzed 21,383 wastewater samples: 19,944 wastewater samples from
the UCSD campus, and, for comparison, 1,475 wastewater samples from the greater San Diego
area, including the Point Loma wastewater treatment plant (the primary wastewater treatment
plant for the county with a catchment size of 2.3 million people) and 17 public schools spanning
four San Diego school districts[17]. We compared sequencing of 600 campus wastewater samples
to 759 genomes obtained from campus clinical swabs (46.2% of all positive tests on campus), all
processed by the CALM and EXCITE CLIA labs at UCSD. In addition, we compared 31,149
genomes obtained from clinical genomic surveillance of the greater San Diego community to
sequencing of 837 wastewater samples collected from San Diego county (including those from
the UCSD campus) during the same period.
184

High-resolution spatial sampling reveals micro-scale community spread

We implemented a GIS (geographic information system)-enabled building-level wastewater surveillance system to cover 360 buildings on the UCSD campus (**Figure 1A**). During the period of daily wastewater sampling, approximately 10,000 students lived on campus and 25,000 individuals were on campus on a daily basis. We found that wastewater test positivity correlated strongly with the number of clinical positives (**Figure 1B** and **Extended Data Figure 1**), showing that wastewater effectively captures the community infection dynamics based on total viral load. This is also consistent with our past studies that showed SARS-CoV-2 RNA can be detected ~85% of the time downstream from buildings containing individuals known to be infected[9].

Unlike qPCR-based mutant surveillance, genomic surveillance using full-length virus genomes can detect which strains of SARS-CoV-2 are circulating in the population, and can identify potential transmission links between infected individuals[18,19]. While targeted qPCR mutant panels have the ability to detect specific lineages in wastewater, they only target a small set of mutations that must be known beforehand in addition to the development and validation time before implementation. Furthermore, they cannot provide sub-lineage resolution (for instance, BA.1 v. BA.2 sublineage of Omicron) and will fail altogether if a sublineage loses the specific mutation targeted by the qPCR assay. To test the utility of wastewater genomic surveillance for studying virus spread in the community, we obtained near complete virus genomes for wastewater samples with cycle quantification (Cq) values as high as 38 (median genome coverage: 96.49% [75.67% - 100.00%], **Extended Data Figure 2**). However, using two common metrics of virus diversity, Shannon entropy (a measure of the uncertainty associated with randomly sampling an allele) and richness (the number of single nucleotide variant, or SNV, sites)[20], we found that SARS-CoV-2 genetic diversity is significantly greater in wastewater samples than clinical samples (**Figure 1C**, Mann-Whitney U test, p<0.001 for each, with effect size r=0.99, 0.97 for Shannon Entropy and Richness, respectively). This suggests that multiple virus lineages, likely shed from different infected individuals, are often present in wastewater samples.

Figure 1: Campus sampling locations and SARS-CoV-2 testing statistics. A. Geospatial distribution of the 131 actively deployed wastewater autosamplers and the corresponding 360 university buildings on the campus sewer network. Building-specific data have been de-

221 identified in accordance with university reporting policies. B. Campus wastewater and diagnostic
222 testing statistics over the 295 day sampling period (WW = wastewater, positivity is the fraction
223 of WW samplers with a positive qPCR signal). C.Virus diversity in wastewater and clinical
224 samples: Boxplots of Shannon entropy (top) and richness (bottom) for each sample type.
225
226 **Sample deconvolution robustly recovers the abundance of SARS-CoV-2 lineages in mixed**
227 **samples**
228
229 Wastewater systems aggregate stool, urine, and other biological waste products carrying viruses
230 from multiple infected individuals in the community in a single location, allowing for sampling
231 of virus mixtures that are representative of local lineage prevalence. However, existing methods
232 for determining virus lineage from sequencing are intended for non-mixed clinical samples and
233 can only be used to identify a single (dominant) lineage per sample.
234
To fully capture the virus diversity in community biospecimens, we developed Freyja, a tool to
estimate the relative abundance of virus lineages in a mixed sample. Freyja uses a "barcode"
library of lineage-defining mutations to represent each SARS-CoV-2 lineage in the global
phylogeny[21](**Figure 2A**). To encode each sample, Freyja stores the SNV frequencies (proportion
of reads at a site that contain the SNV) for each of the lineage-defining mutations (**Figure 2B,**
top). Since SNV frequencies at positions with greater sequencing depth more accurately estimate
the true mutation frequency, Freyja recovers relative lineage abundance by solving a depth-
weighted least absolute deviation regression problem, a mixed sample analog of minimizing the
edit distance between sequences and a reference (**Figure 2B, bottom**). To ensure results are
meaningful, Freyja constrains the solution space such that each lineage abundance value is non-
negative, and overall lineage abundance sums to one. Importantly, Freyja performs site-specific
weighting to account for non-constant variance in measured SNV frequency across sites,
enabling prioritization of information at each site as a function of sequencing depth. Read depths
are log-transformed, providing robustness to common attributes of real sequencing data such as
heavily skewed read depth across amplicons.
250
To validate Freyja, we sequenced "spike-in" synthetic mixtures from five key SARS-CoV-2
lineages (Lineage A, Beta, Delta, Epsilon, and Gamma) at proportions ranging from 5% to 100%
in each sample, with between 1 and 5 different lineages per mixture (**Figure 2C**, and see **Table**
1). We found that Freyja robustly recovered the expected lineage abundances for all mixtures,
even for lineages at 5% abundance (**Figure 2D**, and see **Extended Data Figure 3** for lineage
specific predictions).To further validate Freyja, we used wastewater samples from the UCSD
isolation dorms as well as Point Loma wastewater treatment plant, collection sites likely to
contain mixed-lineage samples, to compare Freyja-detected lineages with qPCR testing for 8
mutations associated with different variants of concern (N501Y, DelHV69/70, DelY144, K417N,
K417T, E484Q, P681R and L452R, **Figure 2E**). We found that Freyja consistently identified the
same lineages as qPCR testing, but, as expected, also identified additional lineages with SNVs
not included in our qPCR panel that were known to be circulating in San Diego at the time of
collection. Combined, these results show that Freyja robustly estimates viral lineage abundance
from samples containing a mixture of lineages, including synthetic virus mixtures and field
wastewater collections.
266

To compare Freyja with other wastewater analysis pipelines, we tested the performance of other
wastewater deconvolution methods including the method from Baaijens et al.[12], cojac[22], and LCS
[23] using the spike-in mixtures (**Extended Data Figure 4)**. We found that Freyja greatly
outperforms other methods in terms of accuracy, false positive rate, and computational
efficiency. The method from Baaijens et al. required greater than ten times more computation
time per sample relative to Freyja (~13.2 minutes vs ~1.1 minutes per sample, respectively).
Although cojac was fast, the small amplicon length used for the spike-in mixtures caused cojac
to fail to identify most of the variants entirely, while LCS failed to return estimates within two
days.
276

277
278
Figure 2: Sample deconvolution robustly recovers relative virus abundance. A. Subset of
lineage defining mutation "barcode" matrix. Each row represents one lineage (out of >1000
lineages included in the UShER global phylogenetic tree), and individual nucleotide mutations
are represented as columns. B. Single nucleotide variant frequencies obtained from iVar used for
recovering relative abundance of each lineage. C. Schematic of the spike-in validation
experiment. D. Depth-weighted de-mixing estimates of the virus abundance versus
expected/known abundance. Details on lineage specific predictions are provided in **Extended
Data Figure 3**. E. Comparison of wastewater sample deconvolution with VOC qPCR panel, with
lookup table (bottom) showing amino acid mutations corresponding to each variant.
288
289 **Detection of early and cryptic community transmission in wastewater**
290
SARS-CoV-2 RNA concentrations in wastewater have been shown to be an early indicator of
rising COVID-19 community incidence[9,24] (and see **Extended Data Figure 5A**), but whether

wastewater can be used to detect emerging variants, including VOCs and VOIs, prior to their
observation in clinical surveillance is unknown. To test if wastewater can enable early detection
of emerging lineages, we applied Freyja to our wastewater sequencing data and compared the
collection date of VOC positive samples from wastewater with the collection dates of samples
from clinical genomic surveillance (**Figure 3A**). With only 2.6% as many sequenced wastewater
samples as sequenced clinical samples, we detected the Alpha and Delta VOC lineages in
wastewater genomic surveillance up to 14 days prior to their first detection in genomic clinical
surveillance (Epsilon was circulating at the start of wastewater collection, and thus could not be
detected early). To further quantify our uncertainty in prevalence estimates, we used a fast
bootstrapping approach (**Extended Data Figure 6)** and found that the resampled distributions
did not include zero abundance. Since emerging VOC lineages may evade immune responses or
lessen the effectiveness of public health interventions[18], this early detection provides additional
time to make necessary adjustments to existing countermeasures.
306
To test if wastewater genomic surveillance can identify changes in the abundance of circulating
lineages, we compared VOC detection rates in clinical and wastewater sequencing over time. We
found that both wastewater and clinical genomic surveillance tracked changes in lineage
abundance, but increases in lineage detection frequency were generally observed first in
wastewater surveillance. For example, for the Epsilon variant, which was first detected in San
Diego in September of 2020, we observed increases in detection frequency in wastewater
approximately 5 days prior to the corresponding increase in clinical genomic surveillance data
(**Figure 3A,** see **Methods**). We noticed varying periods of ongoing lineage detection across
VOCs relative to clinical surveillance, possibly due to different virus shedding characteristics
across lineages [25]. For Epsilon specifically, elevated sampling density on the UCSD campus
relative to elsewhere in the county early on in the experiment may have biased San Diego wide
detection trends towards campus trends, particularly during the end of the wave. We also
observed clear signatures of times with elevated travel, as seen in the pulsing of Alpha detections
in wastewater around the end of holidays and school breaks. During these periods as well as
other times of mass student arrival, students were mandated to test immediately upon arrival
before they moved into their respective on-campus housing. In late March of 2021 following the
university break, mandated clinical testing identified spread of the Alpha variant exclusively in
off-campus residents (see **Figure 1B)**, suggesting that campus mitigation protocols kept the
Alpha outbreak from spreading on campus during this period.
326
To study the effectiveness of wastewater genomic surveillance at a smaller community scale, we
restricted our analysis to samples from the UCSD campus. We found that wastewater genomic
surveillance consistently identified the three major VOCs (Epsilon, Alpha, and Delta) throughout
their period of occurrence, despite detection gaps of one month or longer in clinical surveillance
that included regular asymptomatic testing, longer than the expected signal due to extended virus
shedding [26–28] (**Figure 3B**). During these gaps, positive samples were collected from multiple
distinct locations, with most locations not repeated, suggesting that this continued detection in
wastewater was not simply due to extended shedding. From mid-December to late-March, the
Alpha variant was detected more than once per week on average in wastewater but was not
detected by clinical surveillance. Similarly, wastewater surveillance detected continued Delta
transmission from mid-April to mid-June, but no cases were identified by clinical surveillance.
This explains in part the long tails of wastewater positivity on campus relative to clinical

surveillance on campus (**Figure 1B**), in which we control for extended shedding by excluding samples from campus isolation dorms (see **Methods** for details). The high wastewater positivity level in February-March 2020 extends beyond the expected duration of extended shedding, indicating that cryptic transmission likely played a significant role in campus virus spread during this period.

Figure 3: Freyja recovers early and cryptic transmission of SARS-CoV-2 variants of concern A. Timeline and normalized epidemiological curves for VOC detection in both wastewater and clinical sequences from San Diego County for the 3 major VOCs in circulation during the sampling period. Both Alpha and Delta are detected first in wastewater before clinical samples. Markers for clinical detections correspond to the ceiling of the daily detection count divided by 30 (e.g. 1-30 samples= one marker, 31-60 = two markers) , while wastewater markers correspond to a single detection. B. Timeline and epidemiological curves for VOC detection in the campus samples. Markers correspond to a single detection event for both clinical and

wastewater surveillance. All wastewater detections correspond to an estimated VOC prevalence
of at least 10%.

To study the effectiveness of wastewater surveillance in detecting and tracking other emerging
variants, we aggregated all wastewater sequencing data to estimate the temporal profile of
community lineage prevalence. We found that estimates of lineage abundance using wastewater
enable early identification of other VOCs/VOIs, even for lineages that are rarely observed in
clinical surveillance (**Figure 4**). For example, we detected the Mu (B.1.621) variant via
wastewater genomic surveillance on July 27th, nearly four weeks prior to its first detection
through clinical genomic surveillance on campus, on August 23rd (**Figure 4A,C**). However,
despite persistent Mu detection in campus wastewater throughout July and early August, we did
not detect the Mu variant in clinical or wastewater genomic surveillance on campus in
September, suggesting that local community transmission did not continue.

To test if Freyja continues to provide representative estimates of lineage prevalence for mixtures
containing closely related lineages, we analyzed the rise of the Delta variant (B.1.617.2) and its
sublineages (AY.*) in San Diego, from June-September 2021 (**Extended Data Figure 5B,C**). At
both the UCSD campus and the Point Loma wastewater treatment plant, we identified the rapid
emergence of B.1.617.2 and its sublineages (AY.*), along with low but persistent levels of the
P.1 (Gamma) variant. The relative abundances of each of the variants were within 2-fold of
prevalence estimates observed in clinical nasal swab data, suggesting that Freyja effectively
identifies prevalence even for closely related lineages, both at the university and county-scale.

378
Figure 4: Deconvolution recovers a fine-grained estimate of virus population dynamics. A.
Prevalence of SARS-CoV-2 variants in UCSD clinical surveillance, and B. Variant prevalence in
all clinical samples collected in San Diego County. C,D. Variant prevalence in wastewater at
UCSD as well as the greater San Diego County (includes wastewater samples collected from
Point Loma wastewater treatment plant as well as public schools in the San Diego districts).
Further analysis of Point Loma wastewater samples is shown in **Extended Data Figure 5**. All
curves show rolling average, window ±10 days. "Other" contains all lineages not designated as
VOCs. Bottom panels show number of sequenced samples per day.
387
In more recent data from Point Loma wastewater treatment plant, we identified the Omicron
variant (B.1.1.529 and descendants) at an abundance of near 1.7 % on November 27th, more than
10 days prior to the first clinical detection in San Diego on December 8th (**Figure 5A-B**). To
confirm these findings, we applied our VOC qPCR panel to the same samples and consistently
detected two mutations associated with the Omicron variant (DelHV69/70 and N501Y) in
samples detected after November 27th, while neither was detected in samples from earlier in
November (**Extended Data Table 3,** P681R was included to confirm the presence of Delta**).**
395

To visualize the dynamics of competition between the Delta and Omicron variants, we analyzed
wastewater collected at Point Loma from late September through early February. We found that
upon introduction to the community, Omicron rapidly rose to dominance and reached roughly
95% prevalence by December 26th. During the same period, the estimates for 95% Omicron
abundance in clinical samples tracked via S-gene target failures (SGTFs) was January 7th,
further suggesting wastewater genomic surveillance is a leading indicator of lineage dynamics
for emerging variants (**Figure 5A, Extended Data Figure 7**). To understand the magnitude of
lineage abundance, we scaled each sample by the measured virus RNA concentration of the
sample (**Figure 5B**). We observed that the absolute amount of circulating Delta variant remained
largely constant upon the introduction of Omicron, even as it appeared to decrease to a small
fraction of all viruses in the community.
407
To study the contribution of individual virus lineages to virus RNA concentration, we further
analyzed the growth dynamics of Delta and Omicron sub-lineages (**Figure 5C-D**). We found that
the many Delta lineages circulating in October and November were rapidly displaced by the
BA.1 Omicron lineage, which was soon after displaced by the BA.1.1 lineage, suggesting a
significant growth advantage over BA.1 and B.1.1.529. We did not observe significant levels of
any other Omicron sublineages.
414
415

Figure 5: Community wastewater enables early Omicron detection and reveals lineage dynamics. A. Prevalence of SARS-CoV-2 VOCs in wastewater collected from the Point Loma wastewater treatment plant from late September 2021 to early February 2022. B. Estimated VOC concentrations, prevalence estimates scaled by normalized viral load in wastewater. C,D. Lineage-specific estimates of prevalence and concentration. All curves show an adaptive rolling average calculated using a local linear approximation (Savitzky-Golay filter) of virus copies/L, with window size ± 1 sampling date.

Wastewater identifies both known and unknown history of campus infections

Phylogenetic analysis of virus genomes can be used to identify fine-scale spatial and temporal transmission networks, but it is unknown if wastewater can be used to further refine possible sites of transmission, elucidate transmission networks ("who-infected-whom"), or identify specific infected individuals[19]. To investigate the scale, structure, and timing of SARS-CoV-2 spread on campus, we reconstructed a maximum likelihood phylogenetic tree for each of the

major VOCs using all high-quality consensus genomes (see **Methods** for details) obtained from
the UCSD campus, as well as reference sequences for each lineage obtained elsewhere in the
United States (**Figure 6A-C**). In each tree, we identified many independent introductions, some
of which led to extended transmission on campus. The resulting virus diversity among the VOCs
present on campus enables ruling out of most transmission links and suggests campus virus
spread consisted of many separate, small outbreaks.
438
To analyze the spatial structure of virus spread, we identified collection sites for wastewater
sequences connected to transmission chains on campus, with building-specific resolution
(**Figure 6 A-C**). We observed multiple small, linked outbreaks clustered in nearby buildings.
Campus isolation protocol required students in congregate living to relocate to an isolation room
and linkages in the wastewater samples from buildings used for isolation reflected this co-
location. We also found multiple instances of successive exactly matching sequences from
wastewater collected from a single building, possibly due to continued viral shedding from the
same infected individuals from extended shedding in stool[26–28] or a transmission chain in the
building leading to multiple infections by genetically identical viruses.
448
To study the temporal delay between clinical and wastewater lineage detection, we compared
collection times of sequences from campus wastewater that match sequences from campus
clinical surveillance (including non-VOC lineages). We found 20 exact sequence matches and
103 near-matches (SNP distance of 3 or less) but did not observe any overall bias towards earlier
or later detection in wastewater (**Figure 6D**), suggesting that on average, wastewater and clinical
genomic surveillance identify a similar timing of individual detection events. However, despite
current technical difficulties with isolating haplotypes from diverse virus mixtures, more than
half of the clinical-wastewater sequence pairs demonstrate earlier detection in wastewater or are
from the same date. Importantly, since detection is often delayed or missed by clinical
surveillance, detections occur first in wastewater (despite a loss of sequences due to limited
haplotype recovery), further suggesting that wastewater genomic surveillance can reveal the
presence of specific genome sequences prior to clinical surveillance.
461
462

463
Figure 6: Wastewater identifies clinically known and unknown virus transmission. A-C.
Maximum likelihood phylogenetic trees for each of the dominant variants of concern using high
quality samples obtained at UCSD, as well as a representative set of sequences from the entire
United States. Wastewater sequences from the same sampler that differ by 1 or fewer SNPs are
denoted with a red asterisk. For all sequences, consensus bases were called at sites with >50%
nucleotide frequency. Location information is provided for select outbreaks. D. Pairwise
comparison of collection date for matching and near-matching wastewater and nasal swab

471 samples obtained at UCSD. Positive values indicate earlier collection in nasal swabs, and
472 negative values indicate earlier detection in wastewater.
473
474 **Discussion**
475
We show that improved virus concentration from wastewater, coupled with a method for
resolving multiple lineages from mixed samples, captures community virus lineage prevalence
and enables early detection of emerging variants, often before observation in clinical
surveillance. By sequencing both clinical and wastewater samples from the UCSD campus, we
detect VOCs persistently in wastewater even when their appearance in clinical samples is
intermittent. However, we also found occasions when rarer lineages, like B.1.1.318, were
detected in clinical samples but not in wastewater. This is not unexpected on campus since many
students living off-campus did not contribute to campus wastewater but were still clinically
tested as part of testing mandates and policies. In the larger San Diego community context, this
suggests that we may not be able to identify lineages circulating at low prevalence (< 1%) using
a single wastewater collection site. In addition, we note that clinical sequences identified from
the community may not be observable in the contributing catchment, as precise geolocation of all
clinical samples was not possible. On the other hand, we also observed rare lineages in
wastewater not seen in clinical samples from campus or the community. Since campus testing
mandates are unable to capture all cases (e.g. fully vaccinated individuals were not required to
test and not all community samples were sequenced), rare lineages can be missed.
492
The considerable benefits of wastewater surveillance may stem from biases in clinical testing,
including population testing availability and compliance, university quarantine policies, and
asymptomatic transmission, which may distort estimates of virus lineage prevalence from
clinical samples. Wastewater offers less biased and more consistent viral lineage prevalence
estimates, especially in areas with limited access and/or higher testing hesitancy rates, where
limited clinical surveillance can delay detection of emerging variants. Since it requires
considerably fewer samples, it is also more cost-effective than clinical testing, and could serve as
a long-term passive surveillance tool. This is particularly important for developing public health
interventions in low-resource and underserved communities, where widespread clinical genomic
surveillance for SARS-CoV-2 remains limited.
503
Wastewater is an information-dense resource for estimating the prevalence of specific viral
lineages, providing a community wide-snapshot not only of overall infection dynamics but of the
rise and fall of specific VOCs. Our method, Freyja, deconvolutes these information-rich mixtures
of virus lineages. For a large catchment area, such as San Diego's Point Loma wastewater
treatment plant, which covers over 2 million residents, even limited sampling may accurately
estimate lineage prevalence in the population and provide an early warning indicator of the rise
of new VOCs (as evidenced by the detection of Omicron at just over 1% abundance 11 days
ahead of the first local clinical observation). In addition, wastewater genomic surveillance with
building-level resolution provides a detailed description of the structure and dynamics of
community virus transmission, and can identify transmission links. It can be used to better direct
public health interventions, and can do so in real-time when combined with fast-turnaround
sequencing technologies. This high-resolution approach is of particular utility in community
gathering and transit sites, such as schools and airports, as well as sites with highly vulnerable

individuals, such as nursing homes and hospitals, where spatially resolved monitoring for
directing public health interventions is of great importance.
519
As SARS-CoV-2 continues to evolve, the risk of new VOCs remains high and there is a growing
need to identify these viruses ahead of their proliferation in the community. Accordingly,
development of technologies that are cost-effective, reduce biases, and provide leading rather
than trailing indicators of infection are essential to removing "blind spots" in our understanding
of local virus dynamics. Although technical issues have made wastewater sequencing difficult to
perform at scale, our key advances in virus concentration and sample deconvolution provide
evidence that this approach is now viable. Continued improvements to sequencing turnaround
speeds, lineage barcoding, and haplotype recovery from mixed samples will further accelerate
efforts to achieve earlier identification of emerging variants and improve the precision and
effectiveness of interventions.
530
531 **Methods**
532
533 **Wastewater sampling**
534
High-resolution spatial sampling at the campus level
131 wastewater autosamplers collecting 24h time-weighted composites were deployed across
manholes or sewer cleanouts of 360 campus buildings. GIS (geographic information systems)
informed analyses as well as agent-based network modeling of SARS-CoV-2 transmission on the
UCSD campus enabled identification of most optimal locations for wastewater sampling. During
the pilot phase (November 23-Dec 29th 2020), 68 samplers were prioritized to cover 239
residential buildings identified as the highest risk areas for large outbreaks on campus as a part of
an observational study of wastewater monitoring in high-density buildings [29]. This was based on
preliminary dynamic modeling which showed the largest potential outbreaks to occur within the
largest residential buildings [9]. In addition to the observational study of wastewater monitoring in
these high-density buildings, a cluster randomized study was also performed concurrently. This
included a randomized modified version of a stepped wedge crossover design, in which there
was random assignment of manholes for wastewater sampling. Clusters of manholes associated
with residential buildings were randomized to receive wastewater monitors at one of two-time
steps to evaluate the impact of wastewater monitoring on outbreak size in the associated
buildings. During the same time period, all students in these residences were mandated to
undergo weekly diagnostic testing which was used to validate the utility of building-level
wastewater monitoring. Furthermore, on-campus residences were initially focused due to the
relatively static nature of the population which enabled a more robust cross-validation of the
sensitivity and efficacy of the wastewater surveillance. The coverage of wastewater surveillance
was then increased to cover the rest of the campus buildings (including non-residential buildings
on campus) from January 2021. Four of the deployed wastewater samplers covered the
designated isolation and quarantine buildings on campus.
558
Wastewater composites were collected from the 131 samplers every day for the on-campus
residence buildings and Monday through Friday for the nonresidential campus buildings. 19,944
wastewater samples were collected and analyzed for the presence of SARS-CoV-2 RNA via RT-
qPCR between November 23rd 2020 and September 20th 2021. During this time, 9700 students

lived in campus residences and 25,000 worked on campus on a daily basis. Between October
2020 to January 1st 2021, all on-campus residents were mandated to test on a bi-weekly (once
every 2 weeks) basis and on a weekly basis from January 2nd 2021 (start of the Winter term).
However, fully vaccinated individuals were not mandated to test on a regular basis. Campus
protocols required SARS-CoV-2 positive students living in congregate housing to relocate to
designated isolation housing. Accordingly, our analysis of wastewater positivity (**Figure 1B)** did
not include isolation housing samplers, in order to control - as best as possible, a small number
of students in non-congregate housing spaces were allowed to isolate "in-place", for example -
for possible repeat detection due to extended shedding from infected individuals. Automated,
localized wastewater-triggered notifications were sent to the residents/employees of buildings
associated with a positive wastewater signal which further led to a surge in testing uptake rates
by 2 to 40-fold in the associated buildings.
575
576 *Wastewater sampling at the county level*
577 24h flow-weighted composites were collected thrice a week from the main pump station for the
578 Point Loma wastewater treatment plant, the primary treatment plant serving the greater San
579 Diego county with a catchment size of approximately 2.3 million. 132 wastewater samples were
580 collected between February 24th 2021 to February 7th, 2022.
581
582 **Wastewater sample processing and viral genome sequencing**
583
Sample processing
SARS-CoV-2 RNA was concentrated from 10ml of raw sewage and processed as described
elsewhere[7]. In brief, the viral RNA was concentrated using an automated affinity capture
magnetic hydrogel particle (Ceres Nanosciences Inc., USA) based concentration method after
which the nucleic acid was extracted and sample eluted in 50uL of elution buffer. The extracted
RNA was then screened for SARS-CoV-2 RNA via real-time RT-qPCR for 3 gene targets (N1,
N2 and E-gene). PMMoV (pepper mild mottle virus) was also screened to adjust for changes in
load. Positive wastewater samples were sequenced within 1-2 weeks of collection, comparable to
the delay for clinical samples. To cross-validate the ability of the deconvolution tool in reliably
resolving mixtures of strains in wastewater, the wastewater samples from the county as well as
the ones from the isolation dorms on campus (where multiple infected individuals were isolating)
were also run through a PCR panel targeting 8 mutations associated with the strains designated
as VOCs. The mutations screened for in wastewater using RT-qPCR included N501Y,
DelHV69/70, DelY144, K417N, K417T, E484Q, P681R and L452R (Promega Corp. Cat#
CS3174B02).
599
Miniaturized wastewater SARS-CoV-2 amplicon sequencing
The Swift Normalase® Amplicon Panels (SNAP) kit (PN: SN-5X296 (core) COVG1V2-96
(amplicon primers), Integrated DNA Technologies, Coralville, IA) was used on RNA from
wastewater samples that were positive for SARS-CoV-2 RNA to prepare the multiplex NGS
amplicon libraries and indexed using the SN91384 series of dual indexing oligos, yielding up to
1536 index pairs per pool. A miniaturized version of the protocol was used with the following
modifications: the Superscript IV VILO (Thermo Fisher, Carlsbad, CA) cDNA synthesis
reaction was scaled down to ~1/12 the normal reaction volume with 0.333uL of enzyme mix and
1.333uL of RNA being used. The multiplex amplicon amplification and Ampure XP bead

purification steps were scaled down ~1/6 the normal reaction volume. The Index adapter PCR
reaction and Ampure XP bead purification steps were scaled down to ~2/13 the normal reaction
volume. The final library resuspension volume was 29uL. 1uL of each library was pooled for an
initial shallow NGS run on a MiSeq (Illumina, San Diego, CA) using a Nano flow cell. This
equal volume pool was used to estimate the differential volumes required for similar read depths
across samples using a NovaSeq SP or S4 flow cell (Illumina, San Diego, CA). Between 5uL and
0.2uL of library material, depending on the data provided from the MiSeq Nano run, was
pipetted into a single pool for the NovaSeq run. Transfer volumes were capped at 5uL to reduce
pipetting time and because these types of "high volume" samples typically contained a higher
proportion of likely adapter dimers that inhibit flow cell performance for all samples. A
Dragonfly Discovery (SPT Labtech, UK) was used to dispense reaction master mixes or water
depending on the step. A BlueWasher (BlueCatBio, MA) was used for high throughput
centrifugal 384-well plate washing during the AmpureXP bead reaction cleanup steps. An IKA
MS3 Control linear plate mixer (IKA Works Inc, Wilmington, NC) set to 2600 RPM for 5' was
used to resuspend the AmpureXP beads during the rehydration steps. A Mosquito Genomics HV
16 channel robotic liquid handler (SPT Labtech, UK) was used to dispense the RNA, the reaction
master mixes, and prepare the equal volume pools for the initial MiSeq Nano (Illumina, San
Diego, CA) balancing runs. A Mosquito X1 single channel "hit picker" robotic liquid handler
(SPT Labtech, UK) was used for the final library balancing for the NovaSeq (Illumina, San
Diego, CA) NGS lanes.
629
630 Sequencing data were analyzed using the C-VIEW (COVID-19 VIral Epidemiology Workflow)
631 platform for initial QC and SARS-CoV-2 lineage assignment and phylogenetics. In brief,
632 sequencing reads are aligned with minimap2[30], and primer sequences trimming and quality
633 filtering is applied using the iVar trim method[20]. Sequencing depth and single nucleotide variant
634 (SNV) calls are obtained using samtools mpileup[31] and the iVar variants method[20].
635
Controls were included at all stages of sample processing (viral concentration, extraction, qPCR
and sequencing) to assess potential inhibition and cross-contamination. Most of the sample
processing steps were performed by liquid handling robots for consistency and to minimize
human error. Replicates were included for all wastewater samples. If any of the controls failed or
indicated cross-contamination, the entire batch was rerun. The clinical samples and wastewater
samples were processed separately for sequencing due to significant differences in viral load
between the two sample types.
643
644 **Virus diversity**
645 As reported previously[20], virus SNVs were used to characterize the populations derived from
646 wastewater and clinical samples. Richness was defined as the total number of SNV sites, and
647 mean Shannon entropy $H(p)$ was defined as
648

$$H(p) = \frac{1}{N} \sum_{i=1}^{N} -p_i \log_2(p_i) - (1 - p_i) \log_2(1 - p_i).$$

649
where p_i is the SNV frequency of at the i-th site, of N total sites. For statistical testing, a Mann-
Whitney U test was performed using all wastewater samples that were not sampled from the
same source within a 10 day period in order to ensure independence across samples, as well as
all clinical samples. Effect size was calculated using the rank-biserial correlation,

$r = 2U/(n_{WW}n_{CS}) - 1$ where U is the Mann-Whitney test statistic and n_{WW} and n_{CS} are the
numbers of wastewater and clinical samples, respectively.
656
Wastewater sample deconvolution

657
658
659   To infer relative abundance within a wastewater sample, we used a "barcode" matrix containing
660   the lineage defining mutations for each known virus lineage,
661

$$A = \begin{bmatrix} a_{1,1} & \dots & a_{1,N} \\ \vdots & \ddots & \vdots \\ a_{M,1} & \dots & a_{M,N} \end{bmatrix}$$

662
663
where $a_{i,j}$ denotes the i-th lineage, at mutation j. Lineage defining mutations were obtained from
the UShER global phylogenetic tree using the matUtils package[15]. Similarly, we let b and d
encode the frequency of each mutation and the corresponding sequencing depth (using the log-
transform $d_i = log_2(\text{depth}_i + 1)$ to adjust for large differences in depth across amplicons, which
we use to control for heteroskedasticity and down-weight the importance of sites with little or no
sequencing depth),
670

$$b = \begin{bmatrix} b_1 \\ \vdots \\ b_N \end{bmatrix}, d = \begin{bmatrix} d_1 \\ \vdots \\ d_N \end{bmatrix}.$$

671
672
673   We were then able to write this as a constrained (weighted) least absolute deviations problem

$$\hat{x} = \operatorname*{argmin}_{\substack{x \geq 0 \\ \sum x = 1}} ||A^T x - b||_{1W}, \quad \text{where} \quad ||\mu||_{1W} = \sum_{i=1}^{N} d_i |\mu_i|$$

674
which yields the "demixing" vector $\hat{x} = [\hat{x}_1 \dots \hat{x}_M]$ that specifies the relative abundances of
each of the known haplotypes. Analysis was only performed on samples with greater than 70%
coverage, with the exception of March samples from UCSD for which all samples with greater
than 50% coverage were used. Constrained minimization was performed in Python using the
cvxpy convex optimization package[32,33]. Mapping of lineages to variant WHO lineages (VOCs,
VUMs, etc.) was performed using curated lineage data from outbreak.info[1]. We note that the
Epsilon variant received different maximum escalation levels at CDC and WHO, which assigned
VOC and VOI status, respectively. Since the Epsilon variant was widespread in California and
much of the United States, we use the more "local" CDC designation.
684
Fast-bootstrapping method

685
Bootstrapping was performed at the nucleotide level by resampling each site based on a
multinomial distribution of read depth across all sites, where the event probabilities are
determined by the fraction of the total sample reads found at each site, followed by a secondary
resampling at each site according to a multinomial distribution (i.e. binomial when there was
only one SNV at a site), where event probabilities were determined by the frequencies of each
base at the site, and the number of trials is given by the sequencing depth. 1000 resamplings and
demixings were performed for all samples.

693
694 **Spike-in mixture experiment**
695 RNA was isolated from supernatants of a mammalian cell culture infected with one of five
696 strains of SARS-CoV-2. (A, B.1.1.7, B.1.351, P.1, or B.1.617.2).
697
RNA concentration standardization
Virus concentration was quantified by the UCSD EXCITE COVID testing laboratory using the
Thermo COVID-19 Test kit (PN:A47814, Thermo Scientific Corporation, Carlsbad, CA). The
median Cq values (N-gene, Orf1ab, & S-gene (where applicable)) was calculated and used to
determine how much the RNA needed to be diluted with water to reach a Cq value of 23. A post
dilution RT-qPCR reaction was performed and used to calculate the final dilution of the more
concentrated samples to the new target value of Cq 23.296. The number of freeze thaw cycles
between RNA samples was kept the same.
706
Virus Mixing
RNA standardized in the prior section was used to make a volumetric mixing array (final volume
10uL) using a Mosquito X1 HV robotic liquid handler (SPT Labtech, UK). Pairwise mixes of
5:95, 10:90, 20:80, 60:40, and 50:50 were made for each virus lineage and in both directions.
Equal mixes (20%) for each of the five test strains were made. 25% mixes and 33% mixes were
made for a subset of possible combinations and controls of 100:0 were prepared. See **Extended**
Data Table 1 for complete array. Corrected estimates of the fraction of each virus lineage based
were performed using the final measured Cq values for each pure virus lineage sample to control
for issues encountered during the dilution step (repeat Cq measurements had a coefficient of
variation of 0.007, **Extended Data Table 2**). Across all 95 mixtures, we observed a coefficient
of variation of 0.016. Since initial virus concentrations are controlled for using measured Cq
values, we expect remaining lineage specific bias (see **Extended Data Figure 3**) is likely due to
experimental inconsistencies encountered during mixture creation.
720
Deconvolution method performance comparison
A subset of the spike-in mixtures (1 of each type, for a total of 95 mixtures) was used to compare
Freyja, cojac (using VOC definitions from the public cojac github repository; Lineage A and
Epsilon definitions were created manually), the Kallisto-based method from Baaijens et al. 2021,
and LCS. Kallisto was run using 10 cores (with no bootstrapping), and LCS was run using 16
cores, both on an Intel Xeon processor (2.2GHz). LCS was run for 48 hours, but failed to
complete. Timing was performed using the "time" command, and included all steps after
alignment, trimming, and sorting. Times correspond to total CPU time.
729
730
731 **Estimation of delay in detection frequency**
732 Estimation of the lag time between epidemiological curves for wastewater and clinical
733 surveillance of the Epsilon variant in San Diego was performed by identifying the shift with
734 maximal cross-correlation. All time points leading up to the time of initial peak in detection
735 frequency were included for both wastewater and clinical data.
736
737 **Phylogenetic analyses**

Reconstruction of maximum likelihood trees was performed on all SARS-CoV-2 VOC genomes with 10x (10 reads or greater per site) genome coverage >95% and quality score >20 obtained from UCSD campus sampling, using IQtree[34]. This analysis included 150 (112 clinical, 38 wastewater) Epsilon, 49 (37 clinical, 12 wastewater) Alpha, and 160 (136 clinical, 24 wastewater) Delta lineage genomes from UCSD, in addition to 60 Epsilon, 20 Alpha, and 39 Delta randomly selected genomes from elsewhere in the United States. We used iVar [20] to identify consensus sequences for all San Diego samples. Bases were only included in the sequence if there was a consensus base at the site (>50% nucleotide frequency). We also masked known homoplasic sites prior to tree reconstruction[35]. Analysis of temporal comparison was performed on 608 samples (443 clinical, 165 wastewater, all lineages were included) with 10x genome coverage >95% and quality score >20 from UCSD. Sample collection SNP distances were calculated without considering ambiguous bases and gaps.

Code availability

Freyja is hosted publicly on github (https://github.com/andersen-lab/Freyja) and is available under a BSD-2-Clause License. Freyja is accessible as a package via bioconda (https://bioconda.github.io/recipes/freyja/README.html) in container form via dockerhub (https://hub.docker.com/r/andersenlabapps/freyja). COVID-19 VIral Epidemiology Workflow (C-VIEW) is available at https://github.com/ucsd-ccbb/C-VIEW as an open-source, end-to-end workflow for viral epidemiology focused on SARS-CoV-2 lineage assignment and phylogenetics.

Data Availability

All raw wastewater sequencing data is available via the NCBI Sequence Read Archive under the BioProject ID PRJNA819090. Consensus sequences from clinical and wastewater surveillance are all available on GISAID. Spike-in sequencing data is available via google cloud (https://console.cloud.google.com/storage/browser/search-reference_data). The UCSD campus dashboard can be accessed at https://returntolearn.ucsd.edu/dashboard/ . The county wastewater data from Point Loma are available through the public dashboard that can be accessed at https://searchcovid.info/dashboards/wastewater-surveillance/. The SEARCH genomic surveillance dashboard is available at https://searchcovid.info/dashboards/sequencing-statistics/.

Acknowledgements

We thank Laralyn Asato and the Microbiology Lab at the SD Public Utilities Department for providing us with county wastewater samples. We thank UC San Diego's Return to Learn (RTL) program for funding the campus-wide wastewater surveillance efforts. We also thank Jason Kayne, Rich Cota, Jesus Ortiz, and the Facilities management team (FM) at UCSD, Joseph Mayer from the Center for Aerosol Impacts on Chemistry of the Environment (CAICE) and Luke Arnold of the Campus Research Machine Shop (CRMS) for assistance with the installation and operation of the autosamplers; Robbie Jacobs, Shawn Knepple and their team at UCSD Logistics for assisting with our daily sampling efforts; Brett Pollak and the UCSD Information Technology Services team for assisting with the daily notifications; Office of Academic Affairs for contact tracing and targeted campus messaging assistance; Jana Severson, Patrick Hochstein, the UCSD HDH team, and the UCSD Environmental Health and Safety (EHS) personnel; Jack Gilbert and the Microbiome Sample Processing Core at UCSD for access to qPCR equipment. We also thank the CDC SPHERES consortium, SEARCH (San Diego Epidemiology and

784 Research for COVID Health) Alliance, and members of the Andersen lab for discussion and help
785 with logistics. We thank the healthcare workers, frontline workers, and patients who made the
786 collection of this SARS-CoV-2 dataset possible and all those who made genomic data available
787 for analysis via GISAID.
788

Funding
This work has been funded by CDC BAA contracts 75D30121P10258 (Helix) and
75D30120C09795 (G.W.Y., R.K., L.C.L., and K.G.A.), NIH NIAID 3U19AI135995-03S2
(K.G.A.), U19AI135995 (K.G.A.), U01AI151812 (K.G.A.), NIH NCATS UL1TR002550
(K.G.A.), the Conrad Prebys Foundation (K.G.A.), NIH 5T32AI007244-38 (J.I.L.), NIH Pioneer
Grant 1DP1AT010885 (R.K), NSF RAPID 2029069 (R.K.), San Diego County Health and
Human Services Agency (R.F.M), NIH S10OD026929 (K.J.). The findings and conclusions in
this report are those of the authors and do not necessarily represent the official position of the
Centers for Disease Control and Prevention. Use of trade names is for identification only and
does not imply endorsement by the Centers for Disease Control and Prevention.

Ethics declarations
The University of California San Diego Institutional Review Boards (IRB) provided human
subject protection oversight of the of the data obtained by the EXCITE lab for the campus
clinical samples (IRB approval #210699, #200477). All necessary patient/participant consent has
been obtained and the appropriate institutional forms have been archived, and any sample
identifiers included were de-identified. The wastewater component of this project was discussed
with our Institutional Review Board, and was not deemed to be human subject research, as it did
not record personally identifiable information.
808

Author Contributions
Conceptualization: RK, KGA
Methodology: SK, JIL, NKM, PDH, AK, SS, KMF, AB, LCL, GWY, KGA, RK
Software: JIL, DM, NM, KMF, AB, BH, SS, KG, NLM, KSR, CMA, EH, AMM
Formal Analysis: SK, JIL
Investigation: SK, JIL, NM, SF, HMT, TV, CET, RT, NAB, TB, MC, WC, ESC, ERE, AH, GH,
ALL, EL, TTN, TO, AP, RAS, PS, PBF, EWS, SA, PDH, CAM, LCL, GWY, CA, EK, MAS,
SAP, JL, EP, MZ, ES, RFK, TG, RG, KGA, RK
Resources: CA, NKM, RMN, RS, EHS, AMS, SFK, DPD, CAH, AM, SS, BA, SS, NG, JDM,
EM, IAM, AH, OB, AM, AB, KMSB, ETC, NLW, WL, MI, DB, LN, SW, MZ, RRS, RFM, TG,
RG, DBA, DB, JCB, AB, ME, JG, SLG, MH, FKH, LI, HJ, TJ, VK, BK, LR, CAH, GCM, PM,
RM, PO, DP, AP, AMS, BS, AS, BW, TW, SR, PKK ATY, DMC
Data curation: SK, JIL, PDH, GH, SF, HMT, CET, RT, TV, PDH, AB, NM, AMM, KMF
Writing – Original Draft: SK, JIL, KGA, RK
Writing – Review and editing: all authors
Visualization: SK, JIL
Supervision: RMN, NKM, RS, ALS, EHS, AMS, PDH, LCL, DP, GWY, KGA, RK
Project administration: RMN, NKM, RS, ALS, EHS, AMS, PDH, LCL, GWY, KGA, RK
Funding acquisition: RK, KGA

References

1. Julia L. Mullen, Ginger Tsueng, Alaa Abdel Latif, Manar Alkuzweny, Marco Cano, Emily Haag, Jerry Zhou, Mark Zeller, Emory Hufbauer, Nate Matteson, Kristian G. Andersen, Chunlei Wu, Andrew I. Su, Karthik Gangavarapu, Laura D. Hughes, and the Center for Viral Systems Biology. outbreak.info. *outbreak.info* https://outbreak.info/ (2021).

2. Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune escape. *Nat. Rev. Microbiol.* **19**, 409–424 (2021).

3. Reitsma, M. B. *et al.* Racial/Ethnic Disparities In COVID-19 Exposure Risk, Testing, And Cases At The Subcounty Level In California. *Health Aff.* **40**, 870–878 (2021).

4. Lieberman-Cribbin, W., Tuminello, S., Flores, R. M. & Taioli, E. Disparities in COVID-19 Testing and Positivity in New York City. *Am. J. Prev. Med.* **59**, 326–332 (2020).

5. Brito, A. F. *et al.* Global disparities in SARS-CoV-2 genomic surveillance. *medRxiv* (2021) doi:10.1101/2021.08.21.21262393.

6. Hata, A., Hara-Yamamura, H., Meuchi, Y., Imai, S. & Honda, R. Detection of SARS-CoV-2 in wastewater in Japan during a COVID-19 outbreak. *Sci. Total Environ.* **758**, 143578 (2021).

7. Karthikeyan, S. *et al.* High-Throughput Wastewater SARS-CoV-2 Detection Enables Forecasting of Community Infection Dynamics in San Diego County. *mSystems* **6**, (2021).

8. Randazzo, W. *et al.* SARS-CoV-2 RNA in wastewater anticipated COVID-19 occurrence in a low prevalence area. *Water Res.* **181**, 115942 (2020).

9. Karthikeyan, S. *et al.* Rapid, Large-Scale Wastewater Surveillance and Automated Reporting System Enable Early Detection of Nearly 85% of COVID-19 Cases on a University Campus. *mSystems* **6**, e0079321 (2021).

10. Mercer, T. R. & Salit, M. Testing at scale during the COVID-19 pandemic. *Nat. Rev. Genet.* **22**, 415–426 (2021).

11. Crits-Christoph, A. *et al.* Genome Sequencing of Sewage Detects Regionally Prevalent SARS-CoV-2 Variants. *MBio* **12**, (2021).

12. Baaijens, J. A. *et al.* Variant abundance estimation for SARS-CoV-2 in wastewater using RNA-Seq quantification. *medRxiv* (2021) doi:10.1101/2021.08.31.21262938.

13. Amman, F. *et al.* National-scale surveillance of emerging SARS-CoV-2 variants in wastewater. *medRxiv* 2022.01.14.21267633 (2022).

14. Rambaut, A. *et al.* A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nat Microbiol* **5**, 1403–1407 (2020).

15. Turakhia, Y. *et al.* Ultrafast Sample placement on Existing tRees (UShER) enables real-time phylogenetics for the SARS-CoV-2 pandemic. *Nat. Genet.* **53**, 809–816 (2021).

16. Walke, H. T., Honein, M. A. & Redfield, R. R. Preventing and Responding to COVID-19 on College Campuses. *JAMA* **324**, 1727–1728 (2020).

17. Fielding-Miller, R. K. *et al.* Wastewater and surface monitoring to detect COVID-19 in elementary school settings: The Safer at School Early Alert project. (2021) doi:10.1101/2021.10.19.21265226.

18. Ladner, J. T., Grubaugh, N. D., Pybus, O. G. & Andersen, K. G. Precision epidemiology for infectious disease control. *Nat. Med.* **25**, 206–211 (2019).

19. Grubaugh, N. D. *et al.* Tracking virus outbreaks in the twenty-first century. *Nat Microbiol* **4**, 10–19 (2019).

20. Grubaugh, N. D. *et al.* An amplicon-based sequencing framework for accurately measuring intrahost virus diversity using PrimalSeq and iVar. *Genome Biol.* **20**, 8 (2019).

21. McBroome, J. *et al.* A daily-updated database and tools for comprehensive SARS-CoV-2 mutation-annotated trees. *Mol. Biol. Evol.* (2021) doi:10.1093/molbev/msab264.

22. Jahn, K. *et al.* Detection and surveillance of SARS-CoV-2 genomic variants in wastewater. *bioRxiv* (2021) doi:10.1101/2021.01.08.21249379.

23. Valieris, R. *et al.* A mixture model for determining SARS-Cov-2 variant composition in pooled samples. *Bioinformatics* (2022) doi:10.1093/bioinformatics/btac047.

24. Peccia, J. *et al.* Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. *Nat. Biotechnol.* **38**, 1164–1167 (2020).

25. Singanayagam, A. *et al.* Community transmission and viral load kinetics of the SARS-CoV-2 delta (B.1.617.2) variant in vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study. *Lancet Infect. Dis.* **22**, 183–195 (2022).

26. Cevik, M. *et al.* SARS-CoV-2, SARS-CoV, and MERS-CoV viral load dynamics, duration of viral shedding, and infectiousness: a systematic review and meta-analysis. *Lancet Microbe* **2**, e13–e22 (2021).

27. Wu, Y. *et al.* Prolonged presence of SARS-CoV-2 viral RNA in faecal samples. *Lancet Gastroenterol Hepatol* **5**, 434–435 (2020).

28. Xu, Y. *et al.* Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat. Med.* **26**, 502–505 (2020).

29. Goyal, R., Hotchkiss, J., Schooley, R. T., De Gruttola, V. & Martin, N. K. Evaluation of SARS-CoV-2 transmission mitigation strategies on a university campus using an agent-based network model. *Clin. Infect. Dis.* (2021) doi:10.1093/cid/ciab037.

30. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100 (2018).

31. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

32. Diamond, S. & Boyd, S. CVXPY: A Python-Embedded Modeling Language for Convex Optimization. *J. Mach. Learn. Res.* **17**, (2016).

33. Agrawal, A., Verschueren, R., Diamond, S. & Boyd, S. A rewriting system for convex optimization problems. *Journal of Control and Decision* **5**, 42–60 (2018).

34. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

35. Issues with SARS-CoV-2 sequencing data. https://virological.org/t/issues-with-sars-cov-2-sequencing-data/473 (2020).

Extended Data:

Extended Data Table 1: Platemap of spike-in mixtures used for method validation

	1	2	3	4	5	6
A	5% Delta: 95% A	10% Delta: 90% A	20% Delta: 80% A	40% Delta: 60% A	50% Delta: 50% A	100% A
B	5% Delta: 95% Beta	10% Delta: 90% Beta	20% Delta: 80% Beta	40% Delta: 60% Beta	50% Delta: 50% Beta	100% Delta
C	5% Delta: 95% Gamma	10% Delta: 90% Gamma	20% Delta: 80% Gamma	40% Delta: 60% Gamma	50% Delta: 50% Gamma	100% Beta
D	5% Delta: 95% Alpha	10% Delta: 90% Alpha	20% Delta: 80% Alpha	40% Delta: 60% Alpha	50% Delta: 50% Alpha	100% Gamma
E	5% Beta: 95% A	10% Beta: 90% A	20% Beta: 80% A	40% Beta: 60% A	50% Beta: 50% A	100% Alpha
F	5% Beta: 95% Delta	10% Beta: 90% Delta	20% Beta: 80% Delta	40% Beta: 60% Delta	50% Beta: 50% Delta	20% A: 20% Delta: 20% Beta: 20% Gamma: 20% Alpha
G	5% Beta: 95% Gamma	10% Beta: 90% Gamma	20% Beta: 80% Gamma	40% Beta: 60% Gamma	50% Beta: 50% Gamma	25% Delta: 25% Beta : 25% Gamma: 25% Alpha
H	5% Beta: 95% Alpha	10% Beta: 90% Alpha	20% Beta: 80% Alpha	40% Beta: 60% Alpha	50% Beta: 50% Alpha	25% Delta: 25% Beta: 25% Gamma: 25% A
I	5% Gamma: 95% A	10% Gamma: 90% A	20% Gamma: 80% A	40% Gamma: 60% A	50% Gamma: 50% A	25% Delta: 25% Beta: 25% A: 25% Alpha
J	5% Gamma: 95% Delta	10% Gamma: 90% Delta	20% Gamma: 80% Delta	40% Gamma: 60% Delta	50% Gamma: 50% Delta	25% Delta: 25% A: 25% Gamma: 25% Alpha
K	5% Gamma: 95% Beta	10% Gamma: 90% Beta	20% Gamma: 80% Beta	40% Gamma: 60% Beta	50% Gamma: 50% Beta	25% A: 25% Beta: 25% Gamma: 25% Alpha
L	5% Gamma: 95% Alpha	10% Gamma: 90% Alpha	20% Gamma: 80% Alpha	40% Gamma: 60% Alpha	50% Gamma: 50% Alpha	33% Delta: 33% Beta: 33% Gamma
M	5% Alpha: 95% A	10% Alpha: 90% A	20% Alpha: 80% A	40% Alpha: 60% A	50% Alpha: 50% A	33% Delta: 33% Beta: 33% Alpha
N	5% Alpha: 95% Delta	10% Alpha: 90% Delta	20% Alpha: 80% Delta	40% Alpha: 60% Delta	50% Alpha: 50% Delta	33% Delta: 33% Alpha: 33% Gamma
O	5% Alpha: 95% Beta	10% Alpha: 90% Beta	20% Alpha: 80% Beta	40% Alpha: 60% Beta	50% Alpha: 50% Beta	33% Alpha: 33% Beta: 33% Gamma
P	5% Alpha: 95% Gamma	10% Alpha: 90% Gamma	20% Alpha: 80% Gamma	40% Alpha: 60% Gamma	50% Alpha: 50% Gamma	Neg

Extended Data Table 2: Consistency of Lineage A Cq values across repeated measurements

Replicate Number	Cq N Gene	Cq Orf1ab	Cq S Gene	Cq RNAseP	Average
1	31.228	30.807	30.045	29.581	30.693
2	30.783	29.77	29.546	29.49	30.033
3	31.201	30.622	29.733	29.745	30.519
4	30.621	30.953	29.578	28.925	30.384
5	31.188	30.073	29.366	28.745	30.209
6	30.604	29.788	29.829	28.797	30.074
7	30.308	30.335	29.573	29.149	30.072
8	30.738	30.36	29.711	28.79	30.269
9	31.144	29.97	30.045	28.79	30.386
10	31.122	30.822	29.566	29.671	30.503
11	31.825	29.763	29.833	29.134	30.474
12	31.434	30.18	29.773	29.133	30.462
13	31.209	29.793	29.402	29.559	30.135
14	30.641	30.181	29.816	29.833	30.213
15	30.744	29.371	29.695	29.257	29.937
16	30.396	29.728	29.441	28.428	29.855
17	30.957	29.449	29.913	28.242	30.107
18	30.791	30.113	29.601	29.277	30.169
19	31.561	29.839	29.943	29.06	30.448
20	31.434	29.711	29.568	28.864	30.238

Extended Data Table 3: Omicron surveillance at Point Loma Wastewater Treatment Plant

Collection Date	Avg. Estimated Omicron Abundance (%)	qPCR Detection		
		DelHV69/70	N501Y	P681R
10/04/21	0			x
10/06/21	0			x
10/10/21	0			x
10/11/21	0			x
10/13/21	0			x
10/17/21	0			x
10/18/21	0			x
10/20/21	0			x
11/12/21	0			x
11/22/21	0			x
11/27/21	1.726	x	x	x
11/28/21	1.967	x	x	x
12/1/21	2.439	x	x	x
12/5/21	17.11	x	x	x
12/6/21	19.764	x	x	x
12/12/21	50.65	x	x	x
12/16/21	67.14	x	x	x
12/20/21	79.135	x	x	x
12/21/21	80.567	x	x	x

Extended Data Figure 1: Relationship of daily UCSD campus wastewater sampler positivity and campus clinical positives. Black line indicates the linear fit to the data, with bootstrap 95% confidence interval shown in gray.

Extended Data Figure 2: Relationship between genome coverage and cycle quantification values. 10x genome coverage (fraction of sites with 10 reads or greater) remains high, even for Cq values of nearly 38. Points indicate median value in each bin, while error bars indicate the median absolute deviation.

Extended Data Figure 3: Lineage-specific prediction of variant abundance in spike-in validation samples. A. Schematic of "spike-in" sample design. B-F. Lineage specific prediction. Proportions of each lineage in the sample are shown as a pie chart marker (Grey = Lineage A, Orange = Alpha, Pink = Beta, Turquoise = Delta, and Purple = Gamma) with error bars indicating the standard deviation from the mean, across four replicates.

Extended Data Figure 4: Freyja more accurately estimates virus abundance, with fewer false positives. A-B. Estimated vs expected fraction of each lineage in the mixture. The Kallisto-based approach from Baaijens et. al shows a wider range of estimates for each known mix fraction, and generally underestimates the fraction. C. False positives with abundance greater than 0.5%.

Extended Data Figure 5: The rise of the Delta variant during Summer 2021. A. Mean SARS-CoV-2 viral gene copies/L of raw sewage (blue) collected from the Point Loma Wastewater Treatment Plant and caseload (gray) reported by the county during the same period. SARS-CoV-2 concentrations were normalized by PMMoV (pepper mild mottle virus) concentration to adjust for load changes. B. Lineage distribution in UCSD campus wastewater. C. Monthly lineage averages for wastewater collected at Point Loma Wastewater Treatment Plant during the Delta surge (N= 5, 20, 25, 7)

Extended Data Figure 6: Quantification of deconvolution uncertainty in first detection of VOCs. A-D. Bootstrap distributions of Freyja abundance estimates obtained by resampling read data from each sample corresponding to the first detection of that VOC in San Diego. Two samplers were found to contain Delta on the same day. First detections were also confirmed using a VOC qPCR panel, as shown in Figure 2 and Extended Data Table 3. 95% Confidence intervals for variant prevalence for each first detection event: A. Alpha: (0.232, 0.278), B. Delta: (0.336, 0.397), C. Delta: (0.676, 0.772), D. Omicron: (0.017, 0.021).

Extended Data Figure 7: Estimated proportion of Omicron sequences in clinical data.
Omicron estimates tracked via S-gene target failure, SGTF (characteristic of Omicron lineage
BA.1 and its descendants) qPCR assays for clinical samples in San Diego between November
27th, 2021-February 7th, 2022. First detection of Omicron through clinical genomic sequencing
in San Diego was December 8th. Dotted line shows a rolling average with a window size of
seven days.