Influenza A H1N1–mediated pre-existing immunity to SARS-CoV-2 predicts

COVID-19 outbreak dynamics

Nerea Martín Almazán[1,2,3]*, Afsar Rahbar [1,2]*†, Marcus Carlsson[4], Tove Hoffman[5], Linda Kolstad[5],

Bengt Rönnberg[5], Mattia Russel Pantalone[1,2], Ilona Lewensohn Fuchs[6,7], Anna Nauclér[1], Mats

Ohlin[8], Mariusz Sacharczuk[9,10], Piotr Religa[1,2,10], Stefan Amér[11], Christian Molnár[11,12], Åke

Lundkvist[5], Andres Susrud[13], Birger Sörensen[13], Cecilia Söderberg-Nauclér[1,2,†]

[1]Department of Medicine, Unit for Microbial Pathogenesis, Karolinska Institutet, Stockholm,

Sweden. [2]Department of Neurology, Karolinska University Hospital, Stockholm, Sweden.

[3]Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm,

Sweden. [4]Centre for the Mathematical Sciences, Lund University. [5]Zoonosis Science Center (ZSC),

Department of Medical Biochemistry and Microbiology (IMBIM), Uppsala University, Uppsala,

Sweden. [6]Division of Clinical Microbiology, Department of Labortory Medicine, Karolinska

Institutet, Stockholm, Sweden. [7]Department of Clinical Microbiology, Karolinska University

Hospital, Stockholm, Sweden. [8]Department of Immunotechnology and SciLifeLab Human Antibody

Therapeutics Infrastructure Unit, Lund University, Lund, Sweden. [9]Faculty of Pharmacy with the

Laboratory Medicine Division, Department of Pharmacodynamics, Medical University of Warsaw,

Centre for Preclinical Research and Technology, Banacha 1B, Warsaw, Poland. [10]Department of

Experimental Genomics, Institute of Genetics and Animal Biotechnology, Polish Academy of

Sciences, Postępu 36A, Jastrzebiec, Poland. [11]Familjeläkarna Saltsjöbaden, Saltsjöbaden, Sweden.

[12]Department of Neurobiology, Care Sciences and Society, NVS, Karolinska Institutet, Stockholm,

Sweden. [13]Immunor AS, Oslo, Norway.

*These authors contributed equally to this study.

* Correspondence:
Corresponding Authors
cecilia.naucler@ki.se and afsar.rahbar@ki.se

Keywords: SARS-CoV-2, influenza A H1N1, HLA types, pre-existing immunity, molecular
mimicry, mathematical models

Abstract

Background: Susceptibility to SARS-CoV-2 infections is highly variable, ranging from

asymptomatic and mild infections in most, to deadly outcome in few. This individual difference in

susceptibility and outcome could be mediated by a cross protective pre-immunity, but the nature of

this pre-immunity has remained elusive.

Methods: Antibody epitope sequence similarities and cross-reactive T cell peptides were searched

for between SARS-CoV-2 and other pathogens. We established an ELISA test, a Luminex Multiplex

bead array assay and a T cell assay to test for presence of identified peptide specific immunity in

blood from SARS-CoV-2 positive and negative individuals. Mathematical modelling tested if SARS-

CoV-2 outbreak dynamics could be predicted.

Findings: We found that peptide specific antibodies induced by influenza A H1N1 (flu) strains cross

react with the most critical receptor binding motif of the SARS-CoV-2 spike protein that interacts

with the ACE2 receptor. About 55–73% of COVID-19 negative blood donors in Stockholm had

detectable antibodies to this peptide, NGVEGF, in the early pre-vaccination phase of the pandemic,

and seasonal flu vaccination trended to enhance SARS-CoV-2 antibody and T cell immunity to this

peptide. Twelve identified flu/SARS-CoV-2 cross-reactive T cell peptides could mediate protection

against SARS-CoV-2 in 40–71% of individuals, depending on their HLA type. Mathematical

modelling taking pre-immunity into account could fully predict pre-omicron SARS-CoV-2

outbreaks.

Interpretation: The presence of a specific cross-immunity between Influenza A H1N1 strains and

SARS-CoV-2 provides mechanistic explanations to the epidemiological observations that influenza

vaccination protects people against SARS-CoV-2 infection.

Introduction

Although the COVID-19 epidemic was declared a pandemic in March 2020, it is still not fully under control, and emerging mutant strains continue to cause great concern. SARS-CoV-2 is considered a new virus to humans. Therefore, a major impact was expected. Mathematical modeling predicted an infection rate of at least 70% within a few months, suggesting catastrophic scenarios of collapsed health care systems and high death tolls if strict nonpharmacological mitigation strategies were not implemented (1). However, after the first wave, measured seroprevalence levels were less than 25% in a majority of hard-hit locations. In Stockholm, Sweden, a seroprevalence of 12% was reached in September 2020 after the first wave and under less strict nonpharmacological interventions than in most other western countries.

The SARS-CoV-2 infection affects people very differently. About 50% of infections caused by the original Wuhan strain were asymptomatic. Among people with symptomatic infections, 80% had mild symptoms; 20% developed severe disease and required hospital care, and 3–5% were admitted to the intensive care unit (2, 3). People over 70 years of age and those with obesity, type II diabetes, or hypertension are at higher risk of severe disease (4), and the impact on some patients, even young previously healthy people, can be disastrous. A possible explanation for differences in susceptibility is pre-existing protective immunity, suggested by the unexpected decline in infections during the first wave of the pandemic. In June 2020, 11% were estimated to be seropositive for SARS-CoV-2 in Stockholm, yet the decline started in early April, about 1 month after the onset of the first wave, despite rather limited mitigation strategies compared to other countries. From mid-March 2020, high schools and universities were on distance learning, people were expected to work from home and, if

possible, avoid public transportation. Frequent hand washing, social distancing, and staying at home

if feeling sick were the main recommendations from health authorities and the government. A lock-

down was never implemented, elementary and middle schools remained open, and until January

2021, face masks were not recommended, even in the care of vulnerable patients in hospitals.

The decline of cases in larger cities in Sweden continued from April, which implied some kind of

protective immunity among people. Infection rates on cruise ships during the first wave also peaked

at ~20%, even though many passengers were older than 70 (5, 6). By May 2020, 19.1% of 2149 staff

members at Danderyd´s Hospital in Stockholm tested positive for SARS-CoV-2 antibodies (7).

Furthermore, rarely more than 15–20% of household members became infected with pre-omicron

strains by a family member with confirmed COVID-19 (8). At several elder care homes in Stockholm

and Uppsala in Sweden, about 23% of personnel rapidly became antibody positive during the first

wave (9). In New York, seroprevalence was 23.6% after the spring of 2020 (10). We hypothesized

that this pattern of declining viral spread and the apparent protection of about 75-80% of the

population from severe COVID-19 disease before omicron evolved is best explained by a pre-

existing immunity, which would also contribute to herd immunity thresholds. To test this hypothesis,

we used mathematical models to study the effects of factors such as nonpharmacological

interventions, age, interactive patterns, mobility, and pre-immunity on viral outbreaks. It proved

impossible to match modeled and real data without incorporating a pre-existing immunity level of

50–60% (11, 12). We therefore set out to identify the source of the pre-immunity.

Pre-immunity to SARS-CoV-2 is most likely mediated by previous infections. Indeed, 40–60% of

healthy blood donors, including those giving blood before SARS-CoV-2 existed, respond to SARS-

CoV-2 peptides in vitro (13-17), and pre-existing polymerase-peptide specific T cells expand,

particularly in patients with abortive infections (16). Such T-cell pre-immunity was implied to be

caused by common cold coronaviruses (18-20) and could contribute to herd immunity levels (21-23).

Hence, in some people, T cells trained to recognize unrelated pathogen peptides may protect against

severe COVID-19 through cross-immunity (or molecular mimicry). However, as this virus can be

transmitted by aerosols (24), it is unlikely that T cells protect against infection with SARS-CoV-2 on

a population level. Instead, antibody protection is expected to be required.

Neutralizing antibodies to SARS-CoV-2 have been found in pre-pandemic sera (25) and must

therefore have been triggered by another pathogen, potentially by other common cold coronaviruses

(19). One report suggested that 44% of children and 5.7% of adults have antibodies to common

coronaviruses that can confer neutralizing activity against SARS-CoV-2 (20). However, although

coronavirus antibody titers are boosted by SARS-CoV-2 infection, these are not considered

protective (26). Furthermore, this level of pre-immunity would not explain the slow-down of SARS-

CoV-2 spread when about 20% of a population becomes infected. We therefore searched for potential

cross-reactivities between SARS-CoV-2 and other pathogens.

Results

Identification of potential cross-reactivities between SARS-CoV-2 and other pathogens

A BLAST search for cross-reactive protein sequences between SARS-CoV-2 and any other unrelated

pathogen identified SARS-CoV (2004) with 76% homology, MERS-CoV (2012) with 35.1%

homology and other pathogens such as non-human coronaviruses, human coronaviruses and of

interest the cysteine bond of the SARS-CoV-2 homology to the neuraminidase protein of H1N1

Nagasaki and Kyoto strains. This led us to further analyze any potential homologies to other

influenza strains which could possibly explain the SARS-CoV-2 pre-immunity. We next used a

method focusing on small 6-mer peptides aiming to search for cross-reactive epitopes that are

optimal for antibody binding, identified a peptide in SARS-CoV-2, NGVEGF (Fig. 1a), that is

identical to a peptide in the neuraminidase of two strains of influenza A H1N1 (swine flu):

Nagasaki/07N005/2008 and Kyoto/07K520/2008. NGVEGF was not found in any H1N1 strain

sequenced before 2008 (Table S1). A variant, NGVKGF, was present in 99.3% of swine flu strains (n

= 18,972) sequenced after 2008 and in 31.4% of strains (n = 1467) sequenced before 2008 (Table

S1). The NGVKGF peptide is present in SARS-CoV-2 variants from Brazil (Gamma, P1), South

Africa (Beta, B.1.351, V 501Y.V2), and New York (Iota, B.1.526), which carry an E484K mutation.

Remarkably and very interesting, the NGVEGF/NGVKGF peptide was present in the most

critical part of the receptor binding motif of the spike protein (amino acids (aa) N481 to F486, Fig.

1a) that interacts with the angiotensin converting enzyme 2 (ACE2) receptor (Fig. 1b) (37). In H1N1,

the NGVEGF/NGVKGF peptide is situated in a known immunodominant region of the

neuraminidase protein (Fig. 1c) and is hence expected to elicit an antibody response in a high

proportion of people infected with influenza A H1N1 strains after 2008 (38). Flu strains containing

NGVKGF are currently circulating around the globe and have been included in seasonal flu vaccines

during the last decade. It is therefore possible that antibodies to NGVEGF/NGVKGF developed

during an Influenza A H1N1 infection or after a seasonal flu vaccination could have protected some

people against pre-Omicron strains of SARS-CoV-2.

Receptor binding motif–specific antibodies are present in COVID-19-negative unvaccinated

individuals

To determine if NGVEGF peptide specific antibodies are present in healthy unvaccinated COVID-19

negative individuals, we collected plasma/serum samples from 328 healthy persons in September

2020 and analyzed the samples for the presence of IgG-specific antibodies to SARS-CoV-2 spike and

to the NGVEGF peptide, and developed a diagnostic ELISA method to detect NGVEGF peptide–

specific antibodies. Spike-specific antibodies were detected in 53 (16.2%) of 328 samples (Fig. 1d

and e) (30) with a multiplex assay and determined who among the donors that were positive or

negative for SARS-CoV-2.

Since most people have been exposed to influenza A H1N1 strains since 2008, and flu specific

antibodies can be passively transferred to infants via placenta before birth, we had no optimal

negative control samples to estimate a strict cut-off value for positivity in the ELISA test. Instead, we

relied on negative controls for peptide reactivity of irrelevant peptides, and also confirmed a lack of

NGVEGF specific antibodies in mice plasma (S Fig. 1). In mice, flu vaccination elicited

development of NGVEGF peptide specific IgM and IgG antibodies (S Fig. 1 a and b); these mice

serum samples also served as positive and negative controls for the ELISA test. For the human sera,

the prevalence of IgG positivity to NGVEGF was 73% at a threshold optical density (OD) value ≥0.2

(Fig. 1F and g), 68% at OD ≥0.3, and 55% at OD ≥0.4. Of note, only one of 53 COVID-19-positive

subjects had high titers of NGVEGF-specific antibodies (OD >2, Fig. 1f).

We next established a Luminex Multiplex bead array assay to compare antibody reactivity to 8

NGVEGF or NGVKGF spike peptides (11 or 17 aa in length, Table S2). The peptides were

synthesized with a cysteine bridge and biotinylated at the N or C terminus, respectively. Three

SARS-CoV-2 peptides, an adenovirus peptide, and an irrelevant peptide (Neglle1) served as controls.

Using this method, we confirmed a variable antibody reactivity to NGVEGF (peptide 3) and

NGVKGF (peptide 7) in human sera from different individuals (Fig. 2a–d). The highest reactivity

was observed to the longer NGVKGF peptide biotinylated at the N terminus containing the cysteine

bridge (peptide 7) and was found both in COVID-19-positive and -negative individuals (Fig. 2b and

d). Sera that contained NGVKGF- or NGVEGF-reactive antibodies however rarely recognized the

full-length recombinant spike protein. COVID-19-positive subjects generally had low median

absolute deviation (MAD) values to NGVKGF (peptide 7, Fig. 2 e and f), and some did not mount an

antibody response to NGVEGF (peptide 3, Fig. 2e). However, several subjects with anamnestic flu

and some with known family exposure to SARS-CoV-2 who did not become infected with SARS-

CoV-2 in spite of exposure, interestingly had antibodies with high MAD values to NGVKGF

(example in Fig. 2g). High levels of NGVKGF and NGVEGF specific antibodies were also prevalent

among COVID-19-negative blood donors with unknown SARS-CoV-2 exposure (example in Fig.

2h).

The NGVEGF peptide arose in swine flu in 2009 and the variant NGVKGF was subsequently

present in >99% of H1N1 strains. The NGVKGF peptide was also present in 31.4% of strains ($n =$

1467) sequenced before 2008. We hypothesized that these peptides could have generated cross-

protective immunity to SARS-CoV-2, and therefore examined the prevalence of NGVEGF-reactive

antibodies in patient sera collected before and after 2008 (52 samples from 2011 and 223 from 1996).

At an OD ≥0.2, 90% of sera from 2011 and 79% of sera from 1996 (Fig. 1h and i) contained

NGVEGF-reactive IgG antibodies (Fig. 1i). With a cut off of OD ≥0.3, 73% of sera from 2011 and

66% of sera from 1996 (Fig. 1h and i) contained NGVEGF-reactive IgG antibodies. Since antibody

prevalence to NGVEGF was higher than expected in serum from 1996, we searched for other

peptides that could have elicited an antibody response to the NGVEGF peptide that could cross-react

with the receptor binding domain of SARS-CoV-2. By structural analyses, we identified two

additional peptides—DGVKGF, and NGIKGF—that were similar to NGVEGF and hence could also

have contributed to a protective immune response to SARS-CoV-2. NGIKGF was only present in 1

(<0.01%) and DGVKGF in 990 (67.5%) of 1467 H1N1 strains before 2008 (Table S1). After 2008,

DGVKGF was present in 107 (0.56%) and NGIKGF in 12 (<0.01%) of 18,972 sequence H1N1

strains. Thus, the DGVKGF, NGVEGF, and NGVKGF peptides have been present in many influenza

A H1N1 strains over long periods of time and these peptides may have triggered antibody responses

mediating specific cross-protective immunity to the receptor binding motif of the SARS-CoV-2

Spike protein.

Inhibitory antibodies to SARS-CoV-2 and NGVEGF specific T cells can increase after flu

vaccination

To examine if NGVEGF specific antibody titers and NGVEGF specific T cells increase in flu and

COVID-19 vaccinated individuals and if these could neutralize SARS-CoV-2 infection, we collected

plasma and blood cells from 20 individuals before and after Flu and after COVID-19 vaccination.

Plasma samples from only 19 subjects were available for this test, as when ethical permission was

granted for the study in December of 2020, almost all flu vaccines had been administered in Sweden,

and COVID-19 vaccinations were ready to start. We first tested plasma samples from these

individuals collected before and after flu and COVID vaccination for the presence of NGVEGF

specific antibodies. We observed a trend for enhanced IgG titers in 7 healthy individuals, but not

among 12 elderly people who lived in an elderly care home and were considered "vulnerable due to

poor health" (Fig 3a and b). In all subjects, the NGVEGF specific antibodies decreased after COVID-

19 vaccination. We next tested the neutralizing capacity of antibodies in these plasmas in a virus

neutralization cell culture test, but the plasma samples were toxic to the cultured cells and resulted in

high cell death. This phenomenon was not observed when control sera were tested in the same assay.

As the plasmas were not possible to use in this cell culture assay aiming to test if antibodies present

in the plasma had an inhibitory effect on SARS-CoV-2 infection, we examined the ability of plasma

containing NGVEGF/NGVKGF specific antibodies to inhibit binding of the spike protein to the

ACE2 receptor in a SARS-CoV-2 surrogate virus neutralization test (sVNT). We observed some

enhanced potential protective SARS-CoV-2 immunity by flu vaccination (VaxigripTetra

Quadrivalent Flu vaccine, Sanofi Pasteur (39) as measured by enhanced NGVEGF specific

antibodies and the SARS-CoV-2 sVNT (Figure 3 c, d, e, and Table S5). Although the data obtained

from these analyses are limited and should only be considered descriptive in its nature, we made

some interesting observations. Three subjects with no recent anamnestic flu infection had low

binding inhibitory activity (mean 32.7% inhibition) that was enhanced to mean 55% by flu

vaccination) and this inhibitory capacity was further increased by COVID-19 vaccination (to mean

94%) (Fig. 3c), even though the NGVEGF specific antibodies were decreased (Figure 3a). Four

subjects with suspected flu infection within the past 2 years, had inhibitory antibody activity at levels

similar to those after flu vaccination (mean 47% neutralization), and this neutralizing capacity did not

increase more by flu vaccination (mean 47% inhibition), but was further enhanced by COVID-19

vaccination (to mean 72%) (Fig. 3d). Twelve of the subjects lived in an elderly care home and were

considered "vulnerable due to poor health" and only had a minor increase in the inhibitory effect of

antibodies after flu vaccination, from mean 34% before to a mean of 40% after flu vaccination and to

61% after COVID-19 vaccination (Fig. 3e and Table S5); for some of them the neutralizing capacity

to SARS-CoV2 spike was still concerningly low (Fig. 3e and Table S5). Nine (75%) had an adequate

response to the mRNA BNT162b2 COVID-19 vaccine from Pfizer-BioNTech (mean inhibition

68%), and three had an insufficient response (mean 39% inhibition) after two vaccine doses (Fig. 3f

and Table S5). This was also reflected in lower SARS-CoV-2 specific antibody levels. This

inhibitory effect was higher in the seven other subjects of various ages, and increased from 41%

before to 51% after flu vaccination and to 81% after COVID-19 vaccination (n=7, Fig. 3g), and they

all had adequate antibody levels to SARS-CoV-2 after vaccination.

Flu-mediated protection to SARS-CoV-2 may vary in different populations

While antibodies may protect people from becoming infected, cytotoxic T cells are crucial to resolve

life-threatening infections by killing virus-infected cells. In modeling analyses, we found that the

NGVEGF peptide in theory can be presented to CD8 T cells by some HLA class I molecules (HLA-

A*33:01, HLA-A*68:01, HLA-A*31:01, HLA-A*11:01); these are found in about 22.2% of

Scandinavians (Table S3). We confirmed that B and T cells from 20 healthy subjects recognized and

responded to NGVEGF peptides *in vitro* and that this reactivity was boosted in some people after

seasonal flu vaccination (Fig. 3h). This response was highly individual and differed for CD19, CD4,

and CD8 T cells. Flu-vaccinated subjects had significantly more IFN-γ producing CD8 T cells that

recognized the NGVEGF peptide (mean increase from 1.1% to 2.3%, $p = 0.009$) and B cells (mean

increase from 1.9% to 3.5%, $p = 0.003$); IFN-γ producing CD4 T cells reactive to the NGVEGF

peptide also trended higher after flu vaccination (mean increase from 1.6% to 2.0%, $p = 0.0567$, Fig.

3h). Interestingly, 7 of 20 (35%) individuals had a robust increase in the numbers of CD8 T cells

reactive to the NGVEGF peptide (mean increase of 4.3%); we predicted that about 22% of

Scandinavians have HLA types able to present this peptide efficiently to T cells. Thus, the number of

T cells reactive against the NGVEGF peptide increased prominently in some individuals after Flu

vaccination.

Further peptide screening identified 11 additional influenza H1N1 cross-reactive CD8 T-cell

peptides to SARS-CoV-2 (Table S4). Modeling implied that they could be presented by HLA types

found in about 71% of people in Scandinavia (mainly HLA-A*02:01 and HLA-A*01:01), but in only

40% of people worldwide (Table S3). These observations suggest that the strength of protective

immunity induced by influenza A H1N1 strains that could have mediated protection against SARS-

CoV-2 may vary around the globe.

Mathematical modeling supports that the existence of a pre-existing immunity to SARS-CoV-2

dampened pandemic spread of the virus before Omicron emerged

Evidently, a substantial proportion of the world´s population had antibodies to NGVEGF and T cells

reactive against flu peptides that could have provided protection from infection or severe COVID-19

disease during the first waves of the pandemic, and before vaccinations were introduced. To further

understand whether a pre-existing flu-mediated cross-immunity to SARS-CoV-2 could had

dampened the epidemic on a population level, we turned to mathematical models.

We implemented the SEIR-code by Britton et al. (40), of which there are two versions. The first

version takes interactive patterns between different age groups into account (Age SEIR), and the

second also considers variations in social activity (Age-Act SEIR) (Fig. 4a). We tested these models

on case data from Stockholm, using data from the second wave (Fig. 4a and b), as testing was not

reliable during the first wave. Data from the Swedish Public Health Agency allowed for separation of

cases by the original versus the alpha strain. The blue curve in Fig. 4 represents cases in the second

wave caused by the original strain only, readjusted to account for underreporting of cases. Based on

data from the Swedish Public Health Agency, seroprevalence was 10% in Stockholm at the start of

the second wave (early September 2020) and rose to 22.6% in mid-February 2021, between the

second and third waves (11, 12). These findings are consistent with the seroprevalence estimated

from our serology data: 16.2% ($n = 328$) in late September 2020 and 21.1% ($n = 450$) in late

February 2021. When we attempted to fit the curve of cases with either SEIR model using an

immunity level of 10%, the curves are nowhere near reality (Fig. 4a). However, an almost perfect fit

to measured data was observed when we used pre-immunity levels of 60% for Age-SEIR and 50%

for Age-Act SEIR, which corresponds to a pre-pandemic immunity protective level of 60–70% (Fig.

4b). This level correlates remarkably well with the measured seroprevalence of NGVEGF-specific

antibodies in people in Stockholm (55–73%, depending on OD cut-off for positives) and with the

modeled HLA class I–mediated protective T-cell immunity levels for a Scandinavian population

(estimated to be 71% according to expected HLA types in the population). No other parameter we

examined affected the model output in a similar manner. Thus, it was not possible to match modeled

data to actual case data without taking a substantial protective pre-existing immunity into

consideration (Fig. 4a and b), and the modelled data corresponded remarkably well with the

measured pre-immunity levels to the Influenza A H1N1.

We next ran the code on case data for India, which we expected had a lower pre-immunity

protection, as a result of their populations different HLA types compared to Scandinavians. A pre-

immunity protection in India is therefore expected to be effective at around 20–30% in the first

Wuhan strain wave and only be protective at a 10–20% level for India´s delta wave). We used

seroprevalence data released by the Indian Council of Medical Research; 7% in August/September

2020, 24% around January 1, 2021, and 67% in June/July 2021

(https://indianexpress.com/article/explained/explained-icmr-covid-fourth-serosurvey-findings-

7413949/). Using a 25% pre-immunity against the original strain and a 10% pre-immunity for the

delta variant, and assuming an antibody half-life of 16 months after natural infection (41), the model

fit the observed case data very well, implying that pre-immunity protection, as we hypothesized, was

lower for India than Stockholm. These observations give further validation to the real-life value of

our findings, i.e that a protection to SARS-CoV-2 could have been mediated by previous Influenza A

H1N1 infections and that this protection may be different in different populations who have different

HLA types and thereby variable capacity to present flu peptides to T cells.

Discussion

We discovered that the receptor-biding motif in the SARS-CoV-2 spike protein that interacts with the

ACE2 receptor is identical or very similar in structure to highly immunogenic peptides, NGVEGF,

NGVKGF and DGVKGF that are present in neuraminidase of many influenza A H1N1 strains that

have circulated around the globe during the last decades. We provide evidence that 55%–73% of

individuals have pre-pandemic existing antibodies to this peptide that could have mediated some

protection against SARS-CoV-2, and we provide evidence that flu vaccination can increase immune

protection to NGVEGF and hence potentially against SARS-CoV-2. Since a majority of people have

some NGVEGF-reactive antibodies, these new insights also affect the interpretation of the role of

NGVEGF-specific antibodies in SARS CoV-2 infected individuals, especially concerning their

protective effects against variant viral strains containing the E484K, E484Q or the new E484A

omicron mutation (42). Our modeling data give further support to the hypothesis that pre-existing

immunity to influenza A H1N1 strains indeed could have protected a large set of people from SARS-

CoV-2 infection or severe COVID-19 before omicron emerged in Scandinavia, whereas this

protection may have been less prominent in India with a population carrying other HLA types.

Indeed, India had a much worse second wave than Sweden. Importantly, this discovery provides

explanations to the epidemiological observations that seasonal flu vaccination appears to provide

significant protection against COVID-19 infections, hospitalizations, ICU admission, and death (43-

48); with estimated protection rates (40-80%) that are in range of what was estimated in the present

study (55-73%).

The identified cross-reactive pre-immunity is not expected to provide sterilizing immunity to

SARS-CoV-2. Rather, we mainly consider that it acted as a brake on the epidemic viral spread, as a

higher viral dose was needed to infect someone who had a substantial level of flu-mediated pre-

immunity under given non-pharmacological interventions. However, we also consider the possibility

that those individuals could have been better protected from severe COVID-19 disease. People with

high titers of antibodies and higher levels of T cells that cross-react with flu and SARS-CoV-2 were

likely better protected against SARS-CoV-2 before Omicron emerged, perhaps especially if they had

a recent H1N1 infection or seasonal flu vaccination. This hypothesis is supported by the observed

increased specific immune activity to NGVEGF in some flu vaccinated individuals in the present

study. Whether a person who had flu-mediated protective immunity became infected or not, and

whether such individual developed severe diseases or not may have depended on the infectious dose

and the level of immunological protection against flu at the time of exposure.

On a population level, this cross-protective immunity may explain the unexpectedly slow unfolding

of the pandemic in Sweden despite the absence of a lock-down, while India that we predicted had a

lower efficiency of the flu-mediated protection, was hit harder by the early SARS-CoV-2 strains. The

relevance and accuracy of this theory was strengthened by our modelling data. One can then

speculate over the following scenario; if this cross-protective immunity did not exist, could we have

experienced as rapid spread of the original strain as Omicron, which appears to avoid both natural

immunity, vaccination immunity and influenza pre-immunity. If the spread of the more pathogenic

Wuhan strain causing more severe disease would have been as fast as omicron, the pandemic could

have been even more devastating than so far observed.

Mutant strains have rapidly outcompeted the original strain and new "variants of concern" that are

more contagious have spread rapidly over the world, as natural immunity, vaccine immunity and

potentially flu mediate pre-immunity provide less protection against them (49, 50). Is it then possible

that flu-mediated immunity has selected for the early variants of concern? The Gamma (P1, Brazil),

Alpha (B1.1.7, Britain), and Beta (B.1.351, South Africa) strains contain an N501Y mutation that is

thought to enhance by 10-fold the binding affinity of the spike 1 protein for the ACE2 receptor (51-

53). The Delta strain (B.1.617) has a T478K mutation, and the NY Iota strain has a S477N mutation

in the receptor binding domain, which was predicted to reduce protection from vaccines (54, 55).

Interestingly, the N501Y, S477N, and T478K mutations flank the NGVEGF cysteine loop at aa 481

to 486, which interacts with the ACE2 receptor. As a result of these mutations, NGVKGF-specific

antibodies may have become less protective, enabling these viral strains to infect a higher proportion

of individuals at lower doses. It is then possible that the N501Y, S477N, and T487K mutations in

SARS-CoV-2 evolved through laws of Darwinian evolution to increase affinity for the ACE2

receptor and to evade NGVEGF-interacting antibodies, making these variants more contagious?

Under such scenario, the pre-existing flu-mediated immunity may have aided in the selection of the

first new viral variants of concern.

Our data also opens for speculations of the relevance of this finding for SARS-CoV-2

susceptibility especially among children. High numbers of hospitalized children infected with the

delta variant were reported in the US, the UK, and Israel, but not in Sweden. One could speculate

over the possibility that flu vaccine strategies could have affected SARS-CoV-2 severity among

children. Seasonal flu vaccinations are recommended for children in the US, the UK, and Israel, but

not in Sweden. In Sweden, flu vaccinations are only recommended to high-risk groups and for people

over 65 years of age. A flu-mediated pre-immunity to SARS-CoV-2 may be less efficient and less

sustainable after repeated flu vaccinations than after influenza A H1N1 infection that would elicit a

more robust B- and T-cell immunity. The flu vaccination recommendations and favorable HLA types

might explain Sweden´s lower incidence of severe COVID-19 disease in children under 10 years of

age of which many had swine flu as their first influenza A H1N1 infection, and hence developed a

robust immune response to NGVEGF/NGVKGF peptides. Before Omicron, Sweden did not have

high rates of severe infections among children requiring hospital care; however, multisystem

inflammatory syndrome in children was twice as prevalent in Sweden as in the US in November

2021. This syndrome develops 4–6 weeks after diagnosis of COVID-19 and does not seem to be

related to the severity of SARS-CoV-2 infection; its incidence may rather be related to the number of

infections. Sweden kept schools open during the pandemic and had high levels of transmission

among children, but few became severely ill in the acute phase of COVID-19. Flu vaccination

strategies together with unfavorable HLA types with lower capacity to present flu peptides to T cells

may hence explain differences in susceptibility to SARS-CoV-2 infection and the risk of developing

severe COVID-19 in different parts of the world. We observed that many children below 10 years of

age were hospitalized in Sweden during the Omicron wave, which was not observed during earlier

wave. This may be explained by their loss in previous influenza mediated protection due to the

E484A Omicron mutation.

Our experimental and modeling data have limitations. First, levels of antibodies against

NGVEGF, and the neutralization assay have no threshold to estimate protection on an individual

level and antibody protection should be considered to be more relevant on a population level, and we

had no suitable NGVEGF antibody negative human sera to use as a control. We have searched for

control sera and T cells from 6-9 months old children, as these should be predicted to have the lowest

prevalence of NGVEGV specific antibodies and T cell reactivities, but found no such samples among

contacted colleagues. Second, the mathematical model can only suggest a range of pre-immunity

levels that are likely to be true in reality. Both SEIR models we used in earlier work (11, 12) and the

method developed here are crude tools, and the results should be interpreted with caution. However,

SIR (susceptible, infective, recovered) and SEIR models that did not include pre-immunity

completely failed to predict the dynamics of SARS-CoV-2 spread (11, 12); pre-immunity levels that

we here found to be different in Stockholm versus in India. This is the main mathematical argument

for the existence of a pre-existing immunity, the exact level of which is hard to estimate with

certainty. On the other hand, the estimated pre-immunity levels from SEIR models and the

completely different mathematical tool we devised for this study yielded remarkably consistent

results, near the levels suggested by pre-immunity data observed here—supporting the existence of

flu-mediated antibodies and T cells that cross react with and protect against SARS-CoV-2 infection

or from severe COVID-19 disease in a high proportion of Swedish people. This scenario would also

explain why so many people in Sweden were not infected despite household exposure, had

asymptomatic infections, or experienced mild disease before Omicron emerged.

We conclude that the high prevalence of flu-mediated cross-protective immunity to SARS-CoV-2

is important for understanding SARS-CoV-2 susceptibility, vaccine responses, protection against

new variants, the natural course of COVID-19 in different individuals, as well as for the impact of

this virus and its mutants on people and our society. Learnings from such pre-immunity protection

studies are important to consider to better handle future pandemics.

Ethics Statement

The study was approved by the Ethics Committee at Karolinska Institutet, Stockholm, Sweden (Dnr

2020-06333: all vaccinated subjects gave written informed consent; blood donors from blood bank

were anonymous; Dnr 2020-07232: all subjects gave written informed consent; Dnr 06400: included

anonymous blood samples from healthy donors (HD) in 2011 (Dnr: 01-420) and in 1996 (Dnr: 95-

397 and 02-091). The study protocol for animal study was approved by II Local Ethics Committee

for Animal Experiments in Warsaw (Dnr: 16.07.2021).

Author Contributions

Conceptualization: CSN, MC, AS, BS. Methodology: CSN, AR, BR, PR, AS, SA, ÅL, TH, MO, BS,

CM, MC, LK, NMA. Software: MC. Visualization: AR, PR, AN, NMA. Validation: CSN, AR, BR,

PR, ÅL, TH, LK, NMA. Formal analysis: CSN, AR, BR, PR, AN, MC. Investigation: AR, BR, PR,

ILF, MRP, SA, ÅL, TH, MS, LK, NMA. Resources: PR, ILF, MRP, SA, ÅL, CM. Data Curation:

AR, TH, AN, LK, NMA. Funding acquisition: CSN, ÅL. Project administration: CSN, AR, PR.

Supervision: CSN, PR, ÅL. Writing – original draft: CSN. Writing – review & editing: CSN, AR,

BR, PR, ILF, MRP, AS, SA, AN, MS, ÅL, TH, MO, BS, CM, MC, LK, NMA

Declaration of Interests

BS, AS have submitted a patent application for a COVID-19 peptide vaccine. Other authors declare

that the research was conducted in the absence of any commercial or financial relationships that

could be construed as a potential conflict of interest.

Funding

This research was partially funded by the Swedish Research Council (VR: 2019-01736, 2017-05807,

2018-02569), The European Union's Horizon 2020 Research Innovation Program under grant

874735 (VEO), The Knut and Alice Wallenberg Foundation, and the Science for Life Laboratory

Uppsala (Projects: Nevermore Covid, SiCoV and Molres).

Acknowledgments

We thank the SciLifeLab Autoimmunity and Serology Profiling infrastructure unit for multiplex bead

array serology and Cecilia Hellström, Peter Nilsson and Sophia Hober for their valuable input. We

thank Benjamin Murrell and Daniel Scheward for performing an initial pseudo neutralization assay

for the project and for their valuable input in discussions of our data. We thank Jonna Hermansson

and Lisa Lindberg for their help with sample collection at the elderly care home, Koon Chu Yaiw for

sample preparations, Angela Silveira for providing the cohort of blood samples from 1996 and all

individuals who donated blood to this study. Finally, we thank Stephen Ordway for excellent help

with editing the manuscript.

Data Sharing Statement

The datasets generated in current study are available from the corresponding authors upon reasonable

request.

References

1.   N. M. Ferguson DL, G. Nedjati-Gilani, N. Imai, K. Ainslie, M. Baguelin, S. Bhatia, A.

Boonyasiri, Z. Cucunubá, G. Cuomo-Dannenburg, A. Dighe, I. Dorigatti, H. Fu, K. Gaythorpe, W.

Green, A. Hamlet, W. Hinsley, L. C. Okell, S. van Elsland, H. Thompson, R. Verity, E. Volz, H.

Wang, Y. Wang, P. G.T. Walker, C. Walters, P. Winskill, C. Whittaker, C. A Donnelly, S. Riley, A.

C. Ghani. Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19

mortality and healthcare demand. Imperial College London, . 2020;10(77482):491–7.

2.   Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected

with 2019 novel coronavirus in Wuhan, China. Lancet. 2020;395(10223):497-506.

3.   Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality

of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet.

2020;395(10229):1054-62.

4.   O'Driscoll M, Dos Santos GR, Wang L, Cummings DAT, Azman AS, Paireau J, et al. Age-

specific mortality and immunity patterns of SARS-CoV-2. Nature. 2020.

5.   Moriarty LF, Plucinski MM, Marston BJ, Kurbatova EV, Knust B, Murray EL, et al. Public

Health Responses to COVID-19 Outbreaks on Cruise Ships - Worldwide, February-March 2020.

MMWR Morb Mortal Wkly Rep. 2020;69(12):347-52.

6.   Expert Taskforce for the C-CSO. Epidemiology of COVID-19 Outbreak on Cruise Ship

Quarantined at Yokohama, Japan, February 2020. Emerg Infect Dis. 2020;26(11):2591-7.

7.   Rudberg AS, Havervall S, Manberg A, Jernbom Falk A, Aguilera K, Ng H, et al. SARS-CoV-

2 exposure, symptoms and seroprevalence in healthcare workers in Sweden. Nat Commun.

2020;11(1):5064.

8.   Madewell ZJ, Yang Y, Longini IM, Jr., Halloran ME, Dean NE. Household Transmission of

SARS-CoV-2: A Systematic Review and Meta-analysis. JAMA Netw Open. 2020;3(12):e2031756.

9.   Lindahl JF, Hoffman T, Esmaeilzadeh M, Olsen B, Winter R, Amer S, et al. High

seroprevalence of SARS-CoV-2 in elderly care employees in Sweden. Infect Ecol Epidemiol.

2020;10(1):1789036.

10.   Pathela P, Crawley A, Weiss D, Maldin B, Cornell J, Purdin J, et al. Seroprevalence of Severe

Acute Respiratory Syndrome Coronavirus 2 Following the Largest Initial Epidemic Wave in the

United States: Findings From New York City, 13 May to 21 July 2020. J Infect Dis.

2021;224(2):196-206.

11.   Carlsson M, Hatem G, Söderberg-Nauclér C. Mathematical modeling suggests pre-existing

immunity to SARS-CoV-2. MedRxiv. 2021.

12.   Carlsson M, Söderberg-Nauclér C. Pre-immunity to influenza A H1N1 affects COVID-19

outbreak dynamics, predicts herd immunity thresholds, and implies that Stockholm has reached herd

immunity twice. MedRxiv. 2021.

13.   Grifoni A, Weiskopf D, Ramirez SI, Mateus J, Dan JM, Moderbacher CR, et al. Targets of T

Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease and Unexposed

Individuals. Cell. 2020;181(7):1489-501 e15.

14.   Braun J, Loyal L, Frentsch M, Wendisch D, Georg P, Kurth F, et al. SARS-CoV-2-reactive T

cells in healthy donors and patients with COVID-19. Nature. 2020;587(7833):270-4.

15.   Le Bert N, Tan AT, Kunasegaran K, Tham CYL, Hafezi M, Chia A, et al. SARS-CoV-2-

specific T cell immunity in cases of COVID-19 and SARS, and uninfected controls. Nature.

2020;584(7821):457-62.

16.   Swadling L, Diniz MO, Schmidt NM, Amin OE, Chandran A, Shaw E, et al. Pre-existing

polymerase-specific T cells expand in abortive seronegative SARS-CoV-2. Nature. 2021.

17.   Sekine T, Perez-Potti A, Rivera-Ballesteros O, Stralin K, Gorin JB, Olsson A, et al. Robust T

Cell Immunity in Convalescent Individuals with Asymptomatic or Mild COVID-19. Cell.

2020;183(1):158-68 e14.

18.   Ma Z, Li P, Ikram A, Pan Q. Does Cross-neutralization of SARS-CoV-2 Only Relate to High

Pathogenic Coronaviruses? Trends Immunol. 2020;41(10):851-3.

19.   Jiang S, Du L. Effect of Low-Pathogenic Human Coronavirus-Specific Antibodies on SARS-

CoV-2. Trends Immunol. 2020;41(10):853-4.

20.   Ng KW, Faulkner N, Cornish GH, Rosa A, Harvey R, Hussain S, et al. Preexisting and de

novo humoral immunity to SARS-CoV-2 in humans. Science. 2020.

21.   Doshi P. Covid-19: Do many people have pre-existing immunity? BMJ. 2020;370:m3563.

22.   Sette A, Crotty S. Pre-existing immunity to SARS-CoV-2: the knowns and unknowns. Nat

Rev Immunol. 2020;20(8):457-8.

23.   Lipsitch M, Grad YH, Sette A, Crotty S. Cross-reactive memory T cells and herd immunity to

SARS-CoV-2. Nat Rev Immunol. 2020;20(11):709-13.

24.   Greenhalgh T, Jimenez JL, Prather KA, Tufekci Z, Fisman D, Schooley R. Ten scientific

reasons in support of airborne transmission of SARS-CoV-2. Lancet. 2021;397(10285):1603-5.

25.   Bertoglio F, Meier D, Langreder N, Steinke S, Rand U, Simonelli L, et al. SARS-CoV-2

neutralizing human recombinant antibodies selected from pre-pandemic healthy donors binding at

RBD-ACE2 interface. Nat Commun. 2021;12(1):1577.

26.   Anderson EM, Goodwin EC, Verma A, Arevalo CP, Bolton MJ, Weirick ME, et al. Seasonal

human coronavirus antibodies are boosted upon SARS-CoV-2 infection but not associated with

protection. Cell. 2021;184(7):1858-64 e10.

27.   Zhang Y, Aevermann BD, Anderson TK, Burke DF, Dauphin G, Gu Z, et al. Influenza

Research Database: An integrated bioinformatics resource for influenza virus research. Nucleic Acids

Res. 2017;45(D1):D466-D74.

28.   Reynisson B, Alvarez B, Paul S, Peters B, Nielsen M. NetMHCpan-4.1 and NetMHCIIpan-

4.0: improved predictions of MHC antigen presentation by concurrent motif deconvolution and

integration of MS MHC eluted ligand data. Nucleic Acids Res. 2020;48(W1):W449-W54.

29.   Ayoglu B, Mitsios N, Kockum I, Khademi M, Zandian A, Sjoberg R, et al. Anoctamin 2

identified as an autoimmune target in multiple sclerosis. Proc Natl Acad Sci U S A.

2016;113(8):2188-93.

30.   Hober S, Hellstrom C, Olofsson J, Andersson E, Bergstrom S, Jernbom Falk A, et al.

Systematic evaluation of SARS-CoV-2 antigens enables a highly specific and sensitive multiplex

serological COVID-19 assay. Clin Transl Immunology. 2021;10(7):e1312.

31.   Team, R.C.R. A language and environment for stastistical computing. R Foundation for

Stastistical Computing, Vienna, Austria URL https://wwwR-projectorg/. 2019.

32.   Team, Studio RR. Integrated Development for R. RStudio, Inc, Boston MA URL

http://wwwrstudiocom/. 2018.

33.   Rahbar A, Peredo I, Solberg NW, Taher C, Dzabic M, Xu X, et al. Discordant humoral and

cellular immune responses to (CMV) in glioblastoma patients whose tumors are positive for CMV.

Oncoimmunology. 2015;4(2):e982391.

34.   Hoffman T, Kolstad L, Lindahl JF, Albinsson B, Bergqvist A, Ronnberg B, et al. Diagnostic

Potential of a Luminex-Based Coronavirus Disease 2019 Suspension Immunoassay (COVID-19 SIA)

for the Detection of Antibodies against SARS-CoV-2. Viruses. 2021;13(6).

35.   Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate

time-varying reproduction numbers during epidemics. Am J Epidemiol. 2013;178(9):1505-12.

36.   Bi Q, Wu Y, Mei S, Ye C, Zou X, Zhang Z, et al. Epidemiology and transmission of COVID-

19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study.

Lancet Infect Dis. 2020;20(8):911-9.

37.   Yang J, Petitjean SJL, Koehler M, Zhang Q, Dumitru AC, Chen W, et al. Molecular

interaction and inhibition of SARS-CoV-2 binding to the ACE2 receptor. Nat Commun.

2020;11(1):4541.

38.   Liu WC, Lin CY, Tsou YT, Jan JT, Wu SC. Cross-Reactive Neuraminidase-Inhibiting

Antibodies Elicited by Immunization with Recombinant Neuraminidase Proteins of H5N1 and

Pandemic H1N1 Influenza A Viruses. J Virol. 2015;89(14):7224-34.

39.   Gresset-Bourgeois V, Leventhal PS, Pepin S, Hollingsworth R, Kazek-Duret MP, De Bruijn I,

et al. Quadrivalent inactivated influenza vaccine (VaxigripTetra). Expert Rev Vaccines.

2018;17(1):1-11.

40.   Britton T, Ball F, Trapman P. A mathematical model reveals the influence of population

heterogeneity on herd immunity to SARS-CoV-2. Science. 2020;369(6505):846-9.

41.   Townsend JP, Hassler HB, Wang Z, Miura S, Singh J, Kumar S, et al. The durability of

immunity against reinfection by SARS-CoV-2: a comparative evolutionary study. Lancet Microbe.

2021;2(12):e666-e75.

42.   Verghese M, Jiang B, Iwai N, Mar M, Sahoo MK, Yamamoto F, et al. A SARS-CoV-2

Variant with L452R and E484Q Neutralization Resistance Mutations. J Clin Microbiol.

2021;59(7):e0074121.

43.   Marin-Hernandez D, Schwartz RE, Nixon DF. Epidemiological evidence for association

between higher influenza vaccine uptake in the elderly and lower COVID-19 deaths in Italy. J Med

Virol. 2021;93(1):64-5.

44.   Fink G, Orlova-Fink N, Schindler T, Grisi S, Ferrer APS, Daubenberger C, et al. Inactivated

trivalent influenza vaccination is associated with lower mortality among patients with COVID-19 in

Brazil. BMJ Evid Based Med. 2020.

45.   Conlon A, Ashur C, Washer L, Eagle KA, Hofmann Bowman MA. Impact of the influenza

vaccine on COVID-19 infection rates and severity. Am J Infect Control. 2021;49(6):694-700.

46.   Debisarun PA, Gossling KL, Bulut O, Kilic G, Zoodsma M, Liu Z, et al. Induction of trained

immunity by influenza vaccination - impact on COVID-19. PLoS Pathog. 2021;17(10):e1009928.

47.   Jabr Alwazzeh M, Mohammed Telmesani L, Saud AlEnazi A, Abdulwahab Buohliqah L,

Talal Halawani R, Jatoi NA, et al. Seasonal influenza vaccination coverage and its association with

COVID-19 in Saudi Arabia. Inform Med Unlocked. 2021;27:100809.

48.   Tayar E, Abdeen S, Abed Alah M, Chemaitelly H, Bougmiza I, Ayoub HK, AH., et al.

ffectiveness of influenza vaccination against SARS-CoV-2 infection among healthcare workers in

Qatar. medRxiv. 2022.

49.   Rees-Spear C, Muir L, Griffith SA, Heaney J, Aldon Y, Snitselaar JL, et al. The effect of

spike mutations on SARS-CoV-2 neutralization. Cell Rep. 2021;34(12):108890.

50.   Gallagher KME, Leick MB, Larson RC, Berger TR, Katsis K, Yam JY, et al. SARS -CoV-2

T-cell immunity to variants of concern following vaccination. bioRxiv. 2021.

51.   Graham C, Seow J, Huettner I, Khan H, Kouphou N, Acors S, et al. Impact of the B.1.1.7

variant on neutralizing monoclonal antibodies recognizing diverse epitopes on SARS-CoV-2 Spike.

bioRxiv. 2021.

52.   Liu Y, Liu J, Plante KS, Plante JA, Xie X, Zhang X, et al. The N501Y spike substitution

enhances SARS-CoV-2 infection and transmission. Nature. 2021.

53.   Khan A, Wei DQ, Kousar K, Abubaker J, Ahmad S, Ali J, et al. Preliminary Structural Data

Revealed That the SARS-CoV-2 B.1.617 Variant's RBD Binds to ACE2 Receptor Stronger Than the

Wild Type to Enhance the Infectivity. Chembiochem. 2021;22(16):2641-9.

54.   Wang R, Chen J, Gao K, Wei GW. Vaccine-escape and fast-growing mutations in the United

Kingdom, the United States, Singapore, Spain, India, and other COVID-19-devastated countries.

Genomics. 2021;113(4):2158-70.

55.   Vasireddy D, Vanaparthy R, Mohan G, Malayala SV, Atluri P. Review of COVID-19

Variants and COVID-19 Vaccine Efficacy: What the Clinician Should Know? J Clin Med Res.

2021;13(6):317-25.

Figure legends

Figure. 1 Localization of peptide sequences within spike protein of SARS-CoV-2. (a)

Localization of the NGVEGF peptide in the spike protein of SARS-CoV-2. (b) NGVEGF is present

in the critical domain of SARS-CoV-2 that interacts with the ACE2 receptor. (c) NGVKGF is

situated in a highly immunodominant region of the neuraminidase enzyme of influenza A H1N1 and

is expected to elicit an antibody response in most people infected with influenza A H1N1.

Receptor binding motif–specific antibodies are present in COVID-19-negative unvaccinated

individuals. (d and e) Fifty-three of 328 subjects had IgG antibodies (mean MFI >6 SD) against

soluble pre-fusion stabilized trimeric spike glycoprotein (SPIKE-f (HEK). (f) NGVEGF-specific IgG

antibody levels in COVID-19-positive and COVID-19-negative subjects, measured by ELISA. (g)

Prevalence of NGVEGF-specific IgG antibody values (cut-off values are shown for OD >0.2, >0.3,

>0.35, and >0.4) in COVID-19-positive and COVID-19-negative subjects. (h) NGVEGF-specific

IgG antibody levels in serum samples from healthy donors in 2001 and 1996, measured by ELISA.

(i) Prevalence of NGVEGF-specific IgG antibody values (cut-off values are shown for OD >0.2,

>0.3, >0.35, and >0.4) in healthy donors (HD) from 2011 and 1996. OD: optical density,

Figure. 2 NGVEGF- and NGVKGF-specific IgG antibodies are present in both COVID-19-

positive and COVID-19-negative individuals. (a and c) Heat map of MAD values for peptides 3, 7,

and 11 in COVID-19 positive (A) and COVID-19-negative (c) cohorts. (b and d) The IgG antibody

levels (MAD values) to the NGVKGF (peptide 7) were higher than to the NGVEGF peptide (peptide

3) in both COVID-19-positive (b) and COVID-19-negative (d) cohorts. (e–h) Peptide-specific IgG

antibodies in two COVID-19-positive individuals (e and f), in a subject who was exposed to Flu and

SARS CoV-2 but did not get sick (g), and in an COVID-19-negative subject (h). One-way ANOVA

and Tukey's multiple comparison test were used for multiple comparisons. ∗∗∗∗$p < 0.00001$,

∗∗∗$p < 0.0001$, ∗∗$p < 0.001$, ∗$p < 0.01$, ns: no significant.

Figure. 3 Inhibitory antibodies to SARS-CoV-2 and NGVEGF specific T cells can increase after

flu vaccination. (a and b) Plasma from 19 individuals before and after Flu and after COVID-19

vaccination were examined for presence of NGVEGF specific antibodies using ELISA. A trend was

observed for enhanced IgG titers in 7 healthy individuals, but not among 12 elderly people who lived

in an elderly care home and were considered "vulnerable due to poor health". (a and b) In all

subjects, the NGVEGF specific antibodies decreased after COVID-19 vaccination. (c-e) Inhibitory

binding of Spike to the ACE2 receptor detected in plasma/serum with a surrogate virus neutralization

assay: Grey bars represent mean value of binding inhibition; individual signs represent MFI value for

Spike 1 reactive antibodies. (c) Individuals ($n = 3$) with no recent anamnestic flu infection had lower

binding inhibitory activity, which was boosted by flu vaccination. (d) Four subjects who had

evidence of anamnestic flu had binding inhibitory activity at levels similar to those after flu

vaccination; this immunity level was not further enhanced by flu vaccination. (e) In 12 subjects,

COVID-19 vaccination increased binding inhibitory activity against SARS-CoV-2 to a mean value of

61%. (f) Increased binding inhibitory activity after COVID-19 vaccination (mean values: 34.3%

before vaccination, 40.2% after flu vaccination, and 61% after SARS CoV-2 vaccination) in elderly

cohort in care home (g) and in healthy donors (mean values: 41.2% before vaccination, 50.6% after

flu vaccination, and 81.4% after COVID-19 vaccination). Expansion of IFN-γ producing NGVEGF

peptide specific T cells in Flu-vaccinated subjects. (h) Increased number of IFN-γ-producing B and T

cells stimulated with NGVEGF peptides and analyzed by flow cytometry in a group of 20 subjects

before and after seasonal flu vaccination (CD4, CD8, and CD19 cells). Nonparametric One-way

ANOVA test and Dunn`s multiple comparison test were used for multiple comparisons of percent of

antibody and antibody-binding inhibition before and after vaccinations. Paired *student t-test* was used

for comparison of IFN- γ producing cells before and after Flu vaccination.

Figure. 4 Mathematical modeling considering pre-existing immunity predicts COVID-19

outbreaks. Without taking pre-immunity into account, it was not possible to match the development

of the second wave in Stockholm County with two heterogeneous SEIR-models developed by Britton

et. al.(40): the Age-SEIR model, which takes variable social interactions between different age

groups into account, and the Age-Act-SEIR, which also takes variations in social activity within each

age group into account. Blue curves are for actual cases. (a) Attempts to fit actual cases in the

absence of pre-immunity, using different R_0 values. (b) Curves generated with Age-SEIR using 60%

pre-immunity and with Age-Act-SEIR using 50% pre-immunity. (c) The pandemic progression can

be accurately modeled for India, using a pre-immunity of 25% against the Wuhan strain and 10%

against the Delta variant.

Supplementary Figure. 1

NGVEGF peptide specific antibodies are induced in mice by Flu or SARS-CoV-2 peptide

vaccination. (a and b) Mice received one dose of VaxigripTetra Quadrivalent vaccine (Sanofi

Pasteur) followed 2 weeks later by a booster containing the same vaccine and 5 SARS-CoV-2

peptides (Table S2). Control mice were injected with saline. Sera were collected at 2 weeks (response

after first dose) and at 6 weeks (response after second dose) and were analyzed for IgM (a) and IgG

(b). OD values for IgM and IgG in control mice were <0.2. OD values for IgG after first dose

vaccination was <0.2 and increase after second dose vaccination to >0.2. OD values for IgM was

>0.2 at first dose vaccination and increased to >2.0. Nonparametric one-way ANOVA test and

Dunn`s multiple comparison test were used. OD: optical density, $****p<0.00001$, $***p<0.0001$,

$**p<0.001$, $*p<0.01$, ns: no significant.

Figure 1

a   b   c

Fig.1

Fig.1

Fig.1

Fig.2

Fig.2

c

MAD Values
SARS CoV-2 negative individuals (n=275)

Peptide 3   Peptide 7   Peptide 11

SPIKE RBD

d

MAD values

Peptide 3   Peptide 7   Peptide 11

SPIKE RBD

Fig.2

Fig. 3

Fig.3

Fig. 3

Fig. 3

Fig. 4

a

c