1 **Trends and country-level variation in age at first sex in sub-Saharan Africa among**

2 **birth cohorts entering adulthood between 1985 and 2020**

3 Van Kính Nguyen PhD[1,*], Jeffrey W Eaton PhD[1]

5 [1]MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial

6 College London, London, United Kingdom

* Corresponding author:

Van Kính Nguyen
MRC Centre for Global Infectious Disease Analysis, School of Public Health
Imperial College London
St. Mary's Hospital Campus
Norfolk Place
London
W2 1PG
k.nguyen@imperial.ac.uk

Abstract

Background: Debuting sexual intercourse marks exposure to pregnancy or fatherhood and

sexually transmitted infections (STIs), including HIV. In sub-Saharan Africa (SSA), sexual

debut varies according to cultural, religious, and economic factors, and encouraging delay

has been a longstanding component of behavioural HIV prevention strategies. Age at first

sex (AFS) is routinely collected in national household surveys, but data are affected by

reporting biases, limiting utility to monitor trends and guide sexual health interventions.

Methods: We collated individual-level data from nationally-representative household

surveys to analyse timing and national trends in AFS in 42 SSA countries. We used a log-

skew-logistic distribution to characterize the time to AFS in a Bayesian spatio-temporal

model, providing estimates of the sexual debut rate by sex, age, time, and country. We

statistically adjusted for reporting biases by comparing AFS reported by the same birth

cohorts in multiple survey rounds, allowing different reporting biases by sex and country.

Results: Median AFS in 2015 ranged from 15.8 among Angolan women to 25.3 among men

in Niger. AFS was younger for women than men in 37/40 countries. The gap was largest for

Sahel region countries and minimal in southern African countries. The distribution of female

AFS was asymmetric with half debuting sex in an age range of 3.9 years [IQR 3.4–5.0

across countries]. Median AFS increased slightly between 1985 and 2020, ranging 0.84

years [IQR 0.11–1.55] and 0.79 [IQR -0.23–1.98] for females and males, respectively. The

gender gap changed little over time in most countries. Female teens often reported higher

AFS compared to when asked in their late twenties while male teens reported lower AFS;

both sexes recalled a higher AFS in older ages compared to their thirties.

Conclusions: AFS increased slightly in most SSA countries, but changes were modest

relative to large and persistent variation between countries and sexes, indicating relatively

entrenched socio-cultural practices around sexual debut. Sexual health, family planning, and

HIV/STI prevention services should adapt to local practices rather than focusing

interventions to change AFS. These estimates for rates of sexual debut provide data to

guide programmatic prioritization and implementation of sexual health services.

Introduction

Debuting sexual intercourse is an important life course event for myriad developmental, social, and health reasons. Initiation of sexual activity marks the time when an individual becomes at risk for pregnancy or fatherhood and of acquiring and transmitting sexually transmitted infections (STIs) [1]. Younger age at first sex (AFS) has been associated with a higher rate of STIs [2–5] and increased risk of engaging risky sexual behaviours that facilitate STIs [6].

In sub-Saharan Africa (SSA), patterns and determinants of sexual debut have been of particular interest because of risk of exposure to HIV infection [7] and high rates of adolescent pregnancy [8], which is associated with increased risk of adverse maternal and child health outcomes. Age of first sex (AFS) varies between populations across SSA countries [9,10] according to cultural factors, religious practices, education, socio-economic status [11,12], and has changed over time along with these determinants. Secondary education and wealth quintile have been associated with later sexual debut among young women, though among men evidence is more mixed or in the opposite direction [13–15].

Delayed sexual debut has been associated with declines in HIV incidence [4,16], and encouraging young people to delay sexual debut was considered a priority for curtailing HIV epidemics in sub-Saharan Africa (SSA) during the 2000s as part of the "Abstinence, Be faithful, and use a Condom (ABC)" approach to HIV prevention [17]. Consequently, AFS in a population has become a key sexual risk indicator for determining and monitoring HIV and STI prevention activities [18,19].

Data on AFS are routinely collected in nationally-representative household surveys [20,21] conducted roughly every five years to monitor key health and development indicators. These data, reported retrospectively through face-to-face interviews, are susceptible to systematic response biases related to difficulty recalling dates of life events that happened long ago or desirable standards of society, culture, and health or political campaigns [22–25]. Studies in

which the same individuals have been surveyed longitudinally have estimated that the

percentage of inconsistent AFS reports ranged from 30% to 56% between survey rounds

[26,27] and documented a tendency to report older AFS as respondents age [10,27,28].

Discrepant reports were in both directions and mostly a few years in magnitude but up to 10

years [21,27]. As the cohort aged from teenager to adulthood, men tended to report higher

AFS while women reported lower AFS [10,29,30]. Including inconsistently reported individual

observations, however, did not substantially affect estimates of the population median AFS

[27,31] or predictors in multivariate modelling [21].

While many studies have documented evidence of reporting biases in AFS, few have

attempted to systematically correct for biases to reconstruct how AFS has changed within

and between populations in SSA. Moreover, analyses have tended to focus on the median

AFS in a population [9,32], but other features of the distribution of AFS are less well

characterised. The rate at which young people at each age initiate sexual activity determines

level of exposure to adolescent pregnancy, the range of ages at which sexual health

services need to be introduced, and sexual mixing dynamics that determine HIV and STI

transmission and need for prevention.

Guided by previous literature on systematic differences in reported AFS as respondents

aged [9,10,27], we previously developed and validated a survival analysis model to estimate

the rate of sexual debut from population survey data, adjusting for reporting biases [33]. In

this analysis, we extended and applied the model to incorporate all available AFS data in the

SSA region. Through this analysis, we aimed to (1) characterise variation in AFS across

countries and between sexes in SSA, (2) assess temporal changes in AFS over the last four

decades, including during the period of intensive behavioural HIV interventions in eastern

and southern Africa, and (3) provide data on the rate and full distribution of age at first sex

as a resource for programmatic prioritisation and implementation of sexual health services

and re-evaluation of strategies to identify the most at risk adolescent and young populations

across SSA.

Material and Methods

We used a skew logistic distribution to model the distribution of the log of reported age of

first sex in a time to event modelling framework (Supplemental Text S1). In methodological

comparisons, the log-skew-logistic distribution more accurately represented the sexual debut

hazard than other common survival distributions [33]. To characterise temporal trends and

geographic variation in the AFS distribution, we allowed the skew logistic distribution

parameters to vary by country and birth cohort (Supplemental Text S1). Changes in the AFS

over time for successive birth cohorts were represented by a second-order auto-regressive

(AR2) process separately for males and females in each country. Since members of each

birth cohort reported data about their AFS distribution in multiple household surveys spaced,

typically, approximately five years apart, differences in the AFS reported among the same

cohort in successive surveys quantified and permitted adjustment for systematic reporting

bias in the AFS according to the age of the cohort at the time of the survey as described

elsewhere [33]. The AFS reporting bias as function of age at report was smoothed with a

first-order random walk (RW1) term for each sex and country. To accommodate countries

with only one survey (making identification of country-specific bias from successive surveys

impossible), the effect was estimated borrowing from other countries with a hierarchical

model (Supplemental Text S1).

Household survey data

Nationally representative surveys which collected AFS were collated, including Demographic

and Health Surveys (DHS) [34], Multiple Indicators Cluster Surveys (MICS) [35], AIDS

Indicators Survey (AIS) [34], Performance Monitoring and Accountability (PMA) [36],

Population and Health Survey (PHS) [37], Population-based HIV Impact Assessment (PHIA)

[38], Sexual Behavior Survey (SBS) [39], and South Africa HIV Prevalence, HIV Incidence,

Behaviour and Communication Survey (SABSSM) [40]. Supplementary S1 Table lists the

analysed surveys. We used only surveys in which individual records dataset were available

for analysis. In these surveys, respondents were asked "How old were you (at which age)

when you had sex (sexual intercourse/vaginal sex) for the very first time (if ever)?" for which

valid responses were either never, integer age in years, or at first marriage/union which was

replaced by the reported age at first marriage. Respondents' age was calculated from the

century-month-code records of data of birth and date of interview; if those variables were not

presented, we calculated the century-month-code from the month and year of birth; if only

the year was available, we set month of birth to June; otherwise, the age of respondent

variable in the questionnaires were used.

Model for age at first sex

AFS was reported as integer ages. We modelled the integer AFS as an interval censored

event in which the exact event occurring between the reported AFS and AFS + 1. Individuals

who reported not yet sexually debuting were right censored at their age at the interview. The

likelihood for the reported AFS by each survey respondent was modelled using a time to

event, interval censored model. A weighted pseudo-likelihood approached was used to

account for unequal survey weights, which were normalized to the total sample size of each

survey [41]. The three-parameter skew log-logistic distribution was used to describe the AFS

distribution. This distribution permits a non-monotonic and asymmetric functional form of the

sexual debut hazard by age which other commonly used distributions, e.g., log-normal,

gamma, were not able to capture. It was also shown to have a better predictive value than

the simpler two-parameter log-logistic distribution, which has a symmetric hazard function

[42]. Illustrations of the distribution and its parameters are in Supplemental Text S1.

The model was fitted separately by sex but simultaneously for all countries, with

independent shape and skewness parameters for each country. The linear predictor was

modelled on the distribution's scale parameter as an addictive effect of the spatial

correlation, birth cohort, country-birth cohort interaction, age at report effect, and country-

age interaction. Birth cohort and age at report effects were structured as second-order

autoregressive (AR2) and first-order random walk (RW1) models, respectively; an intrinsic

conditional auto-regressive (ICAR) model [43] was used to model the potential correlation

between neighbouring countries. We assumed interactions that allowed the temporal trend

and age effect to vary independently between countries. Model parameters were estimated

with empirical Bayes using Template Model Builder [44]. Further technical specifications of

the model constraints, priors, and implementations are in Supplemental Text S1. We

computed posterior medians and 95% credible intervals of the outcomes of interests,

including median AFS and proportion ever had sex by specified age thresholds, using

posterior simulations of 3000 parameters samples.

Results

Survey data were collated for 42 SSA countries. Seven SSA countries were not included in

our analysis because they did not have AFS data (Cape Verde, Djibouti, Equatorial Guinea,

Mauritania, Mauritius, Seychelles, and Somalia). Among 253 surveys considered (consisting

of 251 female and 172 male datasets), 32 datasets had more than 10% missing the AFS

variables (AFS and ever had sex indicator variable), mainly from PHIA and MICS surveys; 6

datasets had more than 10% responded as has ever had sex but did not answer a specific

age (do not know or refused responses). These observations were removed from the

analyses. In addition, 136 datasets had reports of AFS under seven years, ranging 0.01%–

1.67% of all observations. We considered those observations irregularities and removed

from the analyses. We included only respondents aged 15–55 years at the time of interview

and born after 1950. The final dataset contained 3,088,315 individual survey respondents for

56 birth cohorts from 1950 through 2005 (entering adulthood at age 15 between 1965 and

2020). The sample size per dataset ranged from 1364 (male, Eswatini MICS 2014) to 41,821

(female, Nigeria DHS 2018). Further details are in Supplemental Table S1.

Spatial patterns of median AFS and proportion sexually active by age 18

For the birth cohort entering adulthood (turning age 15) in 2015, the median AFS ranged

from 15.8 [IQR: 15.5–16.2] in Angola to 25.3 [IQR: 24.5–26.1] in Niger (Figure 1). Females

had lower median AFS than males in most countries, but across countries the AFS between

the two sexes were correlated such that countries where females had low median AFS also

had males with low median AFS and vice versa. This was substantiated by a high correlation

between male and female's estimates of the skew-logistic distribution parameters

(Supplemental Figure S1). The median AFS for men was less than women only in Lesotho,

South Africa, and Comoros. Only females in Rwanda, Burundi, Comoros, and Eritrea had

median AFS above twenty.

The spatial variation in sexual debut is further evident in the percentage who ever had sex

before age 18 years (Figure 2). A very low percentage of men in the Sahel region of Africa

had debuted sex by age 18, with the lowest 4.4% [3.2–6.0%] percent in Niger. Among

women, the percentage sexually active by age 18 was lowest, below 45%, in the southern

countries, and as low as 18.4% [16.5–20.6%] in Burundi. In both sexes, Botswana, Burundi,

and Rwanda all had older AFS than the surrounding region whereas Mozambique had

notably younger AFS in the southeast Africa. At the other extreme, women in Liberia and

men in Angola were most likely to debut sex under eighteen with 86.1% [82.8–88.9%] and

77.5% [73.2–80.8%], respectively. AFS were spatially homogenous in the remained

locations. Similar patterns were observed for the percentage ever had sex under fifteen

(Supplemental Figure S2) where the largest estimate was 37.7% [32.9–42.7%] in Angola

male following by Chad female at 33.3% [31.5–35.1]. In nearly half of the cases (37/82) less

than 10% had initiated sexual activity by age 15. (Supplemental Figure S2, Supplemental

Table S2).

Time trends in AFS

The median AFS increased between birth cohorts entering adulthood in 1985 and 2020 in 60

of 82 time-series estimated by country and sex (Figure 3), and in 40 increased more than

one year. During this period, the average change in median AFS across all countries was

0.84 years [IQR: 0.11-1.55] for females and 0.79 [IQR: -0.23 -1.97] for males. The changes,

however, have not been monotonic but fluctuated over time (such as in Gambia, Guinea

Bissau, Uganda, and Rwanda) and varied between countries. The estimated trends were

similar for male and female in most countries with few exceptions of Comoros, South Africa,

and Lesotho in the most recent birth cohorts. Increases in AFS were especially large for men

in some West African countries (e.g., Senegal, Niger, Mali, Chad, Nigeria, Burkina Faso,

Benin). The largest estimated decreases were also among men (e.g., Tanzania, Sierra

Leon, Lesotho, Mozambique). Eastern and Southern Africa exhibited minimal changes in

AFS during the time frame with the average increase of 0.31 and 0.12 years, respectively.

In the same birth cohort, the gap in median AFS between women and men was under two

years in most countries and remained unchanged in the region. The gap was larger in the

West Africa countries, especially those in the Sahel and Nigeria, and in the Horn of Africa

(Ethiopia, Eritrea). Median AFS increased most in Western Africa (Figure 4). Middle Africa

had the lowest median AFS on average for both men and women. In Southern Africa, which

is dominated by the large population of South Africa, there was smaller sex difference in

median AFS and median AFS decreased since the 1985 adult cohort, in contrast to the

increasing trend in other regions.

Distribution of age at sexual debut by sex

The distribution of AFS was asymmetric in most countries with various degree of skewness

(Figure 1). The inter-quartile range (IQR) of the AFS distribution represent the age range

with the highest sexual debut activity and its magnitude reflects how fast the rate of most of

the population becomes sexually active. In most countries the IQR was under five years with

an average of 4.0 [2.8–8.2] years. The duration for 50% of a birth cohort to initiate sexual

activity was as short as 2.2 years (Liberia - female) or up to 9.6 years (Sudan - male).

Between sexes, in most countries (37/40, excluding two countries with only one sex data)

the IQR was shorter for females than for males; among these, 18 countries had a difference

of more than a year. The distribution of AFS spanned a longer range for countries of the

Sahel and Horn of Africa.

Biases in reported AFS by age at interview

The estimated pattern for the effect of age at the time of interview on the reported AFS for

each country are shown in Figure 5. Data in different countries yielded similar information on

the effect of age at report bias. Among females, there was a tendency to report higher AFS

in their teens compared to when they reached their late twenties, indicating under-reporting

of sexual activity among the youngest survey respondents. The opposite pattern was

observed among men—young men tended to report a younger AFS distribution. For both

sexes, the reported distribution of AFS increased as the cohort aged. The direction of both

bias's directions varied across countries. The coefficient estimates of the biases, however,

only show the relative difference between ages, but do not identify at which age people

would be both truthful and accurately recall their AFS. To arrive at a marginal estimate for

the AFS of each birth cohort, we chose age 23 as the reference age in both female and male

and report estimates using this reference age throughout the results presented.

Discussion

In this study, we systematically analysed trends and patterns in debuting sexual activity

across sub-Saharan Africa among cohorts entering adulthood from 1985 to 2020, a period

during which the continent combatted the HIV/AIDS pandemic and experienced rapid social

change, urbanisation, and economic development. Assessment of past sexual debut trends

and current status in each country are important to help evaluate the impact of the past

interventions, respond to the increasing sexual health needs for young people, and identify

the key ages at which to provide sexual health services. We applied a statistical model to

account for reporting biases that have previously hampered consistent interpretation of AFS

recorded in household surveys and used a parsimonious but flexible model to characterise

the distribution of AFS over ages.

The gap between AFS for men and women differed greatly between the southern and

northern countries with little evidence of changes over time in most countries. The gap in

AFS remained large and widened over time in the north while remained small and narrowed

over time in the south. In the Western Africa region, the gap of more than two years in

median AFS between male and female in many countries implies more disparity in sexual

partner mixing compared to countries in the south. Large gender gaps could make

adolescent girls more vulnerable to adverse sexual health outcomes [45].

The median AFS has increased in many parts of sub-Saharan Africa over the past three

decades, especially in the northern-most countries, but the timing and magnitude of changes

varied. In countries heavily affected by HIV/AIDS epidemic in southern and eastern Africa, in

which there has been a multitude of intervention programmes focused on reducing sexual

risk, median AFS only slightly increased or even decreased. Other analyses have reported

that trends towards earlier sexual debut are not pronounced [9] and median AFS in SSA

increased over time [32]. Previous findings from individual countries also found similar

results such as a small increase in men's AFS in Zambia [28] or both sexes in Uganda [46].

The only modest or decreasing AFS may indicate that efforts to encourage delayed sexual

initiation via expanding access to education [47] may not have had large effect on AFS,

though access to education may have improved knowledge and accessibility to safe sex

practices and sexual health services . In contrast to the Southern African countries, AFS has

increased more steeply in Western Africa where the HIV/AIDS epidemic is not as severe

[48]. Stricter cultural and religious norms about abstaining sex until marriage in this region

may have meant that gains in female education [49] leading to later marriage in for women

[50] resulted more directly in delayed sexual debut, while increasing economic uncertainty

has pressured men to delay marriage [51]. Taken together, health policies aimed at effecting

changes have had limited or inconsistent effects. Strategies adapt sexual health, family

planning, and HIV/STI prevention promotion to changes in AFS to appropriately target the

timing of the current and upcoming adult birth cohorts entering sexually active life are likely

to be more effective.

Reproductive health and HIV monitoring indicators have been focused on single measure

including median AFS in a population or the proportion initiating sexual activity by specified

ages, resulting a loss of age-specific sexual debut behaviour. Our analysis calculated the full

distribution of debuting sexual activity, reflecting the range of ages at which adolescents and

young adults may need to become engaged in sexual health, family planning, and HIV/STI

prevention services. The rate at which populations become sexually active varied across

countries, but typically was slightly more peaked for young women. For young women, the

percentage sexually experienced increased from 25% to 75% in around four years

compared to five years for men. This varied across countries, however. In some, sexual

debut of a population occurred in a very short span of age while in others in a gradual

manner (Figure 1). Two populations with the same median AFS, but different shapes

parameters may have different demand for reproductive health and STIs services where one

may need to cover a larger target age than the other. Programming prevention services

efforts should be directed accordingly to deliver services to the crucial period of a sexually

active life using our detailed estimate of the AFS distribution.

Previous research using longitudinal data or sexual activity biomarkers has identified

reporting biases in AFS that can obfuscate trends assessed from cross-sectional surveys

[27,32,45,52], yet it was difficult to address them in isolated analyses of specific surveys or

single countries considering wide varieties of pattern and magnitude in reporting biases. This

study addressed this problem by comparing reports among the same cohort in successive

surveys and learning from all the available data in SSA to extract the pattern of the bias.

Biases existed in most countries and we identified a few general patterns in the age at report

bias. Young men tended to report earlier AFS than when asked again in later surveys, while

young women tended to report older AFS. Above age 30, the reported AFS tended to

increase as the cohort aged, especially for men, indicative of recall biases for exact age for

events that occurred long ago. These results are consistent with previous findings

documenting biased self-reporting of sexual activity by norms that boys are expected to

engage in sexual activity and girls are expected to abstain until marriage [24][10,29,30].

Other country-specific analyses have also documented that respondents were less likely to

report a young AFS in later surveys [28]. While adjusting for these biases to reconstruct

consistent time trends is a strength of our model, the variation in magnitude and direction of

biases across countries implies caution about confidently distinguishing actual trends from

changing reporting biases among the youngest cohorts in new surveys. This will be

especially challenging in settings with fewer surveys from which to triangulate across

different time points in life of a birth cohort [24]. The results from our study still highlight that

age- and sex-specific reporting bias is prevalent and could lead to misjudgement of the

needs for sexual health services in adolescent and young population – especially for young

girls where more might be needed than the data suggest.

This study has several limitations. First, while we were able to adjust for relative reporting

differences by age at report, there was no 'gold standard' measure to identify at which age

reporting was most accurate. The chosen reference age of twenty-three is at most ten years

difference from the AFS and above the average age of marriage of women in SSA, thus it is

expected that reports at this age are less influenced by social desirability compared to a

younger age and recall more accurate compared to older reporting age. In addition, as digit

preference for age reporting ending in zero or five is a common issue [53], using ages

around 25 or 30 might introduce biases to the reported estimates. Second, country- and sex-

specific differences of potential confounders to median AFS such as religion [54] were not

modelled. Third, our model allowed the median AFS to change over time, and for the shape

of the distribution and reporting bias to vary by country but did not allow the shape or

reporting bias to change over time within each country. Fourth, survey series varied slightly

in the eligibility of respondents asked about AFS, the way AFS was recorded, and data

processing procedures. For example, respondents older than 24 were not asked AFS in

some in MICS or FHS. In DHS it is standard practice to recode internally inconsistent data

such as reported AFS that is older than the age at the time of survey, after the conception of

the first child, or after the first marriage (45), but this is not done in all surveys, as evidenced

in larger proportions of excluded observations (S1 Table). Fifth, we accounted for survey

design by incorporating survey weights into the model pseudo likelihood [41], but did not

model stratified clustered sampling designs. Finally, results in this paper are limited to SSA

at national level. Similar methods to ours could be applied for subnational estimates, which

may provide more granular and actionable characterisation of sexual risk dynamics.

Subnational analyses should consider urban-rural migration trajectories and implications of

respondents no longer residing in the location where retrospectively reported sexual debut

occurred.

Conclusions

AFS has increased slightly in most, but not all, SSA countries. However, in most cases

changes over-time within a country were small compared to large and relatively persistent

variation between countries and genders, reflecting deeply rooted cultural norms around

sexual debut. AFS trend remained relatively stagnant in the countries most affected by the

HIV epidemic suggests behavioural intervention programs encouraging delayed sexual

debut [55] have had relatively modest effects on AFS. Instead, adapting sexual health, sex

education, family planning, and STI and HIV prevention services to local norms around

sexual debut should be prioritised over seeking to intervene to change AFS. Our estimates

of age- and sex-specific sexual debut rates provides data to support locally-adapted health

programming and more detailed inputs to epidemic models of HIV and STIs that may also

help to shed light on national and regional variation in epidemic burdens and transmission

dynamics, for example the role of the larger age gaps in west and middle African countries

compared eastern and southern regions.

Declarations

Ethics approval and consent to participate

The study involved secondary analysis of anonymised publicly available data following

project approval from the Demographic and Health Surveys Program, the Performance

Monitoring for Action project, the Population-based HIV Impact Assessment project, and

South African National HIV Prevalence, Incidence, Behaviour and Communication Survey.

Primary survey protocols were reviewed and approved by the relevant national ethics

committee in each country.

Consent for publication

Not applicable

Availability of data and materials

The datasets analysed during the current study are available in the Demographic and Health

Surveys Program (https://dhsprogram.com/Data/), the Performance Monitoring for Action

project (https://www.pmadata.org/data), the Population-based HIV Impact Assessment

project (https://phia-data.icap.columbia.edu), and South African National HIV Prevalence,

Incidence, Behaviour and Communication Survey (http://www.hsrc.ac.za/).

Competing interests

The Authors declare that there is no conflict of interest.

Funding

This research was supported the Bill and Melinda Gates Foundation (OPP1190661),

National Institute of Allergy and Infectious Disease of the National Institutes of Health under

award numbers R01AI136664 and R01AI152721, and the MRC Centre for Global Infectious

Disease Analysis (reference MR/R015600/1), jointly funded by the UK Medical Research

Council (MRC) and the UK Foreign, Commonwealth & Development Office (FCDO), under

the MRC/FCDO Concordat agreement and is also part of the EDCTP2 programme

supported by the European Union. The funders have no roles in the study design or

manuscript preparation.

Authors' contributions

VKN and JWE conceived the work. VKN and JWE designed the work. VKN implemented the

models. Both authors critically reviewed model results throughout the model development

process. VKN wrote the first draft of the manuscript. Both authors critically edited the

manuscript for intellectual content.

Acknowledgements

This research was supported the Bill and Melinda Gates Foundation (OPP1190661). We

acknowledge joint MRC Centre funding from the UK Medical Research Council and

Department for International Development via MRC MR/R015600/1.

 Figures

Figure 1. Distribution of the age at first sex by region, country, and sex. The median of each

distribution is shown with the vertical lines. Estimates are for the birth cohort turning fifteen in

2015 and assumed age 23 years at survey report. Within each region, countries sequenced

by increase median female AFS. Numerical values and 95% uncertainty intervals for

estimates are reported in Supplemental Table S3. Country names and region allocations

corresponding to the three-letter are ISO 3166-1 alpha-3 code labels were as follows:

Western* Africa:*** BEN – Benin; BFA – Burkina Faso; CIV – Cote d'Ivoire; GHA – Ghana;

GIN – Guinea; GMB – Gambia; GNB – Guinea-Bissau; LBR – Liberia; MLI – Mali; NER –

Niger; NGA – Nigeria; SEN – Senegal; SLE – Sierra Leone; TGO – Togo; ***Middle* Africa***:***

AGO - Angola; CAF - Cent. Afr. Republic; CMR - Cameroon; COD - Dem. Rep. Congo; COG

- Congo; GAB - Gabon; STP - Sao Tome and Principe; TCD - Chad; ***Eastern* Africa***:*** BDI -

Burundi; COM - Comoros; ERI - Eritrea; ETH - Ethiopia; KEN - Kenya; MDG - Madagascar;

MOZ - Mozambique; MWI - Malawi; RWA - Rwanda; SDN - Sudan; SSD - South Sudan;

TZA - Tanzania; UGA - Uganda; ZMB - Zambia; ZWE - Zimbabwe; ***Southern* Africa***:*** BWA -

Botswana; LSO - Lesotho; NAM - Namibia; SWZ - Eswatini; ZAF - South Africa *(also in*

Supplemental Table S4).

Figure 2. Percentage ever had sex by age 18 among the birth cohort turning 15 in the year

2015. The island countries Comoros and São Tomé & Príncipe are represented in the

bottom left. Countries without data are coloured the same as the background. Numerical

estimates and 95% uncertainty interval are in Supplemental Table S2. Country names

corresponding to the three-letter ISO 3166-1 alpha-3 code labels are in Figure 1 caption and

Supplemental Table S4.

Figure 3. Trend in median age at sexual debut for men and women in Sub Saharan-Africa

countries by year turned age 15 (x-axis). For countries where the most recent survey is

before 2015, trends are extrapolated for recent birth cohorts based on the overall trend

across the region and its neighbours. Countries without AFS data are excluded from this

figure. Country names corresponding to the three-letter ISO 3166-1 alpha-3 code labels are

in Figure 1 caption and Supplemental Table S4.

Figure 4. Aggregated trend of median age at first sex by region for men and women. Sudan

was included in Eastern Africa. The regional trend is a weighted average of the countries

included in the analyses by their respective population size in 2020 sourced from the UN

World Population Prospects 2019. Country classifications in each region are defined in the

Figure 1 *caption and Supplemental Table S4.*

Figure 5. Direction of the effect of age at report bias on the median age at first sex estimate

relative to other ages. A positive coefficient in the figure corresponds to an increase in

median AFS estimate reported by the respective age compared to a median AFS reported

by the age with that coefficient equals zero.

Supplemental Materials

Text S1. Skew log-logistic distribution & specifications of the survival model

Figure S1. Parameter distribution correlation

Figure S2. Spatial distribution of the percentage ever had sex under fifteen.

Figure S3. Proportion ever had sex and fitted value by country and sex

Table S1. List of surveys used in the analyses

Table S2. Parameter estimates by country and sex

Table S3. Proportion ever had sex under 15 &18 by country and sex

References

[1] WHO. Report on global sexually transmitted infection surveillance 2018. 2018.

[2] Pettifor AE, van der Straten A, Dunbar MS, Shiboski SC, Padian NS. Early age of first
sex. AIDS 2004. https://doi.org/10.1097/01.aids.0000131338.61042.b8.

[3] Kaestle CE, Halpern CT, Miller WC, Ford CA. Young age at first sexual intercourse and
sexually transmitted infections in adolescents and young adults. Am J Epidemiol
2005;161:774–80. https://doi.org/10.1093/aje/kwi095.

[4] Hallett TB, Gregson S, Lewis JJC, Lopman BA, Garnett GP. Behaviour change in
generalised HIV epidemics: Impact of reducing cross-generational sex and delaying
age at sexual debut. Sex Transm Infect 2007;83:i50-54.
https://doi.org/10.1136/sti.2006.023606.

[5] Wand H, Ramjee G. The relationship between age of coital debut and HIV
seroprevalence among women in Durban, South Africa: a cohort study. BMJ Open
2012;2. https://doi.org/10.1136/bmjopen-2011-000285.

[6] Kaplan DL, Jones EJ, Olson EC, Yunzal-Butler CB. Early Age of First Sex and Health
Risk in an Urban Adolescent Population. J Sch Health 2013.
https://doi.org/10.1111/josh.12038.

[7] Howard AL, Pals S, Walker B, Benevides R, Massetti GM, Oluoch RP, et al. Forced
Sexual Initiation and Early Sexual Debut and Associated Risk Factors and Health
Problems Among Adolescent Girls and Young Women — Violence Against Children
and Youth Surveys, Nine PEPFAR Countries, 2007–2018. Morb Mortal Wkly Rep
2021;70:1629–34. https://doi.org/10.15585/mmwr.mm7047a2.

[8] Mbabazi C, Kintu A, Asiimwe JB, Ssekamatte JS, Shah I, Canning D. Proximate and
distal factors associated with the stall in the decline of adolescent pregnancy in
Uganda. BMC Public Health 2021;21:1875. https://doi.org/10.1186/s12889-021-11403-
6.

[9] Wellings K, Collumbien M, Slaymaker E, Singh S, Hodges Z, Patel D, et al. Sexual
behaviour in context: a global perspective. Lancet Lond Engl 2006;368:1706–28.
https://doi.org/10.1016/s0140-6736(06)69479-8.

[10]   Zaba B, Pisani E, Slaymaker E, Boerma JT. Age at first sex: Understanding recent
trends in African demographic surveys. Sex Transm Infect 2004;80:28–35.
https://doi.org/10.1136/sti.2004.012674.

[11]   Upchurch DM, Levy-Storms L, Sucoff CA, Aneshensel CS. Gender and ethnic
differences in the timing of first sexual intercourse. Fam Plann Perspect 1998;30:121–
7.

[12]   Zaba B, Pisani E, Slaymaker E, Boerma JT. Age at first sex: Understanding recent
trends in African demographic surveys. Sex Transm Infect 2004;80:28–35.
https://doi.org/10.1136/sti.2004.012674.

[13]   Gupta N, Mahy M. Sexual initiation among adolescent girls and boys: trends and
differentials in sub-Saharan Africa. Arch Sex Behav 2003;32:41–53.
https://doi.org/10.1023/a:1021841312539.

[14]   McGrath N, Nyirenda M, Hosegood V, Newell M-L. Age at first sex in rural South Africa.
Sex Transm Infect 2009;85:i49–55. https://doi.org/10.1136/sti.2008.033324.

[15]   Madise N, Zulu E, Ciera J. Is poverty a driver for risky sexual behaviour? Evidence
from national surveys of adolescents in four African countries. Afr J Reprod Health
2007;11:83–98.

[16] Gregson S, Garnett GP, Nyamukapa CA, Hallett TB, Lewis JJC, Mason PR, et al. HIV
decline associated with behavior change in eastern Zimbabwe. Science 2006;311:664–
6. https://doi.org/10.1126/science.1121054.

[17] Halperin DT, Steiner MJ, Cassell MM, Green EC, Hearst N, Kirby D, et al. The time has
come for common ground on preventing sexual transmission of HIV. The Lancet
2004;364:1913–5. https://doi.org/10.1016/S0140-6736(04)17487-4.

[18] UNAIDS. Monitoring the Declaration of Commitment on HIV/AIDS: guidelines on
construction of core indicators n.d.

[19] Dehne, K. L., Gabriele, Riedner. Sexually transmitted infections among adolescents :
the need for adequate health services. n.d.

[20] Aisha NZD, Philipp U, Vladimíra K. Sexual Activity by Marital Status and Age: A
Comparative Perspective, Technical Paper No. 11 2017.

[21] Assche SB Van. Are we measuring what we want to measure? Individual consistency
in survey response in rural Malawi. Demogr Res 2003;9:77–108.
https://doi.org/10.4054/DemRes.2003.S1.3.

[22] Catania JA. A framework for conceptualizing reporting bias and its antecedents in
interviews assessing human sexuality. J Sex Res 1999;36:25–38.
https://doi.org/10.1080/00224499909551964.

[23] Wellings K, Collumbien M, Slaymaker E, Singh S, Hodges Z, Patel D, et al. Sexual
behaviour in context: a global perspective. Lancet Lond Engl 2006;368:1706–28.
https://doi.org/10.1016/S0140-6736(06)69479-8.

[24] Gersovitz M. HIV, ABC and DHS: age at first sex in Uganda. Sex Transm Infect
2007;83:165–8. https://doi.org/10.1136/sti.2006.021576.

[25] United Nations. World population monitoring: adolescents and youth. U N 2012:1.

[26] Eggleston E, Leitch J, Jackson J. Consistency of Self-Reports of Sexual Activity among
Young Adolescents in Jamaica. Int Fam Plan Perspect 2000;26:79–83.
https://doi.org/10.2307/2648271.

[27] Wringe A, Cremin I, Todd J, McGrath N, Kasamba I, Herbst K, et al. Comparative
assessment of the quality of age-atevent reporting in three HIV cohort studies in sub-
Saharan Africa. Sex Transm Infect 2009;85. https://doi.org/10.1136/sti.2008.033423.

[28] Slaymaker E. Monitoring trends in sexual behaviour in Zambia, 1996-2003. Sex
Transm Infect 2004;80:ii85–90. https://doi.org/10.1136/sti.2004.012054.

[29] Mensch BS, Hewett PC, Erulkar AS. The reporting of sensitive behavior by
adolescents: A methodological experiment in Kenya. Demography 2003;40:247–68.
https://doi.org/10.2307/3180800.

[30] Luke N, Clark S, Zulu EM. The relationship history calendar: improving the scope and
quality of data on youth sexual behavior. Demography 2011;48:1151–76.
https://doi.org/10.1007/s13524-011-0051-2.

[31] Cremin I, Mushati P, Hallett T, Mupambireyi Z, Nyamukapa C, Garnett GP, et al.
Measuring trends in age at first sex and age at marriage in Manicaland, Zimbabwe. Sex
Transm Infect 2009;85:i34–40. https://doi.org/10.1136/sti.2008.033431.

[32] Slaymaker E, Scott RH, Palmer MJ, Palla L, Marston M, Gonsalves L, et al. Trends in
sexual activity and demand for and use of modern contraceptive methods in 74
countries: a retrospective analysis of nationally representative surveys. Lancet Glob
Health 2020;8:e567–79. https://doi.org/10.1016/S2214-109X(20)30060-7.

[33] Nguyen VK, Eaton JW. A model for reconstructing trends and distribution in age at first
sex from multiple household surveys with reporting biases. 2021.
https://doi.org/10.1101/2021.12.14.21267770.

[34] ICF. Demographic and Health Surveys (various). Funded USAID Rockv Md ICF Distrib
2020.

[35] UNICEF. Multiple Indicator Cluster Surveys (MICS). 2020.

[36] Elizabeth Heger Boyle, Devon Kristianse, Matthew Sobek. IPUMS PMA: Version 4.1
[dataset]. Minneapolis, MN: IPUMS. 2020. https://doi.org/10.18128/D081.V4.1.

[37] UNICEF. Eritrea Population and Health Survey. 2020.

[38] PHIA Collaborating Institutions. Population-based HIV Impact Assessment (PHIA)
2020. https://doi.org/10.18128/D081.V4.1.

[39] MEASURE Evaluation Project. Zambia sexual behaviour surveys. 2020.
https://doi.org/10.18128/D081.V4.1.

[40] The Human Sciences Research Council. South African National HIV Prevalence,
Incidence and Behaviour Survey. 2020. https://doi.org/10.18128/D081.V4.1.

[41] Asparouhov T. General Multi-Level Modeling with Sampling Weights. Commun Stat -
Theory Methods 2006;35:439–60. https://doi.org/10.1080/03610920500476598.

[42] Van Kinh Nguyen, Jeffrey W Eaton. Model the age at first sex distribution and age at
reporting bias in household survey data. MedRxiv 2021.

[43] Besag J, Kooperberg C. On Conditional and Intrinsic Autoregression. Biometrika
1995;82:733–46. https://doi.org/10.2307/2337341.

[44] Kristensen K, Nielsen A, Berg CW, Skaug H, Bell BM. TMB: Automatic differentiation
and laplace approximation. J Stat Softw 2016;70. https://doi.org/10.18637/jss.v070.i05.

[45] Melesse DY, Mutua MK, Choudhury A, Wado YD, Faye CM, Neal S, et al. Adolescent
sexual and reproductive health in sub-Saharan Africa: who is left behind? BMJ Glob
Health 2020;5:e002231. https://doi.org/10.1136/bmjgh-2019-002231.

[46] Slaymaker E, Bwanika JB, Kasamba I, Lutalo T, Maher D, Todd J. Trends in age at first
sex in Uganda: evidence from Demographic and Health Survey data and longitudinal
cohorts in Masaka and Rakai. Sex Transm Infect 2009;85:i12–9.
https://doi.org/10.1136/sti.2008.034009.

[47] Shapiro D, Gebreselassie T. Marriage in Sub-Saharan Africa: Trends, Determinants,
and Consequences. Popul Res Policy Rev 2014;33:229–55.
https://doi.org/10.1007/s11113-013-9287-4.

[48] Dwyer-Lindgren L, Cork MA, Sligar A, Steuben KM, Wilson KF, Provost NR, et al.
Mapping HIV prevalence in sub-Saharan Africa between 2000 and 2017. Nature
2019;570:189–93. https://doi.org/10.1038/s41586-019-1200-9.

[49] Barro Robert, Jong-Wha Lee. A New Data Set of Educational Attainment in the World,
1950-2010. J Dev Econ 2013;104:184–98.
https://doi.org/10.1016/j.jdeveco.2012.10.001.

[50] Bongaarts J, Mensch BS, Blanc AK. Trends in the age at reproductive transitions in the
developing world: The role of education. Popul Stud 2017;71:139–54.
https://doi.org/10.1080/00324728.2017.1291986.

[51] Smith DJ. Masculinity, Money, and the Postponement of Parenthood in Nigeria. Popul
Dev Rev 2020;46:101–20. https://doi.org/10.1111/padr.12310.

[52] Plummer ML, Ross DA, Wight D, Changalucha J, Mshana G, Wamoyi J, et al. "A bit
more truthful": The validity of adolescent sexual behaviour data collected in rural
northern Tanzania using five methods. Sex. Transm. Infect., vol. 80, 2004, p. ii49–56.
https://doi.org/10.1136/sti.2004.011924.

[53] Amos M, Stones T. Trends in Demographic and Health Survey data quality: An
analysis of age heaping over time in 34 countries in Sub Saharan Africa between 1987
and 2015. BMC Res Notes 2017;10. https://doi.org/10.1186/s13104-017-3091-x.

[54] Westoff, Charles F. K Bietsch. Religion and Reproductive Behavior in Sub Saharan
Africa. DHS Anal Stud No 48 2015.

[55] Halperin DT, Steiner MJ, Cassell MM, Green EC, Hearst N, Kirby D, et al. The time has
come for common ground on preventing sexual transmission of HIV. The Lancet
2004;364:1913–5. https://doi.org/10.1016/s0140-6736(04)17487-4.

Female

Male

Percentage ever had sex by age 18

20% 40% 60% 80%

Median AFS
by year turned age 15