medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

The Canadian COVID-19 Experiences Survey: Study Protocol

Peter A. Hall*[1,2], Geoffrey T. Fong*[1,2,3], Sara C. Hitchman[4], Anne C.K. Quah[2], Thomas Agar[2], Gang Meng[2], Mohammad N. Sakib[1], Anna Hudson[1] and Christian Boudreau[5]

Author affiliations:

1. School of Public Health Sciences, University of Waterloo, Waterloo, Canada
2. Department of Psychology, University of Waterloo, Waterloo, Canada
3. Ontario Institute for Cancer Research, Toronto, Ontario, Canada
4. Department of Communication and Media Research, University of Zurich, Zurich, Switzerland
5. Department of Statistics and Actuarial Science, University of Waterloo, Waterloo, Ontario, Canada

*Corresponding authors.
Peter A. Hall, School of Public Health Sciences, University of Waterloo, 200 University Avenue West, Waterloo, ON, Canada; pahall@uwaterloo.ca; Phone: 519.888.4567 x48110.
Geoffrey T. Fong, Department of Psychology, University of Waterloo, 200 University Avenue West, Waterloo, ON, Canada; gfong@uwaterloo.ca; Phone: 519.888.4567 x45811.

Abstract

Introduction: Vaccine hesitancy remains a significant challenge even through the second year of the COVID-19 pandemic, even in the most well-supplied countries. In Canada, despite high vaccine uptake of the first wave of vaccinations, there are signs of declining uptake of vaccine boosters. Likewise, recommended COVID-19 mitigation behaviors such as mask wearing, distancing, and hand hygiene have been challenging to sustain over the pandemic. The aim of the Canadian COVID-19 Experiences Survey (CCES) is to use a variety of social-cognitive and neurocognitive variables to predict the uptake and sustaining of vaccinations, boosters, and COVID-19 mitigation behaviors across the Canadian population through the second and third years of the pandemic.

Methods and analyses: Vaccine hesitant and fully vaccinated adults (*N*=1,958) were recruited to the 35-minute CCES web survey, using quota sampling, with sampling weights being applied to ensure representativeness of the Canadian population. In Wave 1, conducted between 28 September and 21 October, 2021, 43.3% (*n*=848) reported not having received any vaccinations, 50.2% (*n*=983) reported being fully vaccinated, and 6.5% (*n*=127) reported having had one vaccination with no intention of having a second dose. Social cognitive variables being measured include intentions, beliefs, reasons, and attitudes. Neurocognitive variables being measured include executive dysfunction, delay discounting, and temporal orientation. Other key variables being measured include trust in science, political orientation, mood, and long-COVID symptoms.

Ethics and dissemination: The CCES has received ethical review and approval by the University of Waterloo Office of Research Ethics. Findings will be disseminated through peer-reviewed publications and presentations at scientific meetings.

Keywords:

COVID-19, SARS-CoV-2, behavior, prevention, vaccine, mitigation.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Background

The emergence of the COVID-19 pandemic has brought to the forefront the importance of population-wide readiness for pandemic response. For governments, this means maintaining availability of personal protective equipment, medical supplies, treatment equipment and facilities, as well as maintaining alert and communication networks to respond to the risk in a coordinated manner. Policies and procedures that facilitate intra- and international cooperation are also critical. The manner in which governments of all levels communicate with population members is critical, and determining the most effective communication strategies is particularly important with a large amount of competition from other messaging sources.

On the level of population members, pandemic response inevitably involves absorbing information about the threat, engaging in mitigation behaviors as well as taking vaccines or other medical measures when they become available. Critical determinants of vaccine uptake and mitigation behaviors include social norms, perceived risk, and beliefs about each of these and intention strength. In a post hoc manner, reasons for taking or not taking these precautions are also critical, and the array of SARS-CoV-2 vaccines available opens the possibilities for reasons to take or not take vaccines, but also reasons for preference of one over another.

Finally, and less readily apparent, some aspects of the brain and its information processing modes can have influence on both the receipt of information communicated about personal and population level risk from an infectious agent, and behavioral responses to it. Together, the nature of population wide health communications, psychological factors influence behavioral response of population members, and neural systems coordinating information received and behavioral response are ultimately all important for pandemic response.

The Canadian COVID-19 Experiences Survey and its laboratory counterparts were devised as a way of testing multiple levels of pandemic response, with an eye toward growing the critical knowledge base of how health communications, psychological processes, and levels of influence can be all harnessed for the purpose of optimally effective response to the threat posed by SARS-CoV-2 to the Canadian population, and indeed the world.

Methods

Participants

The CCES consists of bi-annual survey waves with replenishment targeting 2,000 participants 18 to 54 years of age per survey wave of 35 minutes. A critical feature of the CCES is that participants are recruited using a quota system within six geographic regions (British Columbia, Alberta, Prairies (Saskatchewan and Manitoba), Ontario, Quebec, and Atlantic (New Brunswick, Nova Scotia, Prince Edward Island, and Newfoundland and Labrador)), three age groups (18-24, 25-39, and 40-54) ensuring a 50/50 balance of fully vaccinated and vaccine hesitant individuals. Eligible participants were recruited from Leger Opinion, the largest proprietary nationally representative probability-based panel in Canada. Three eligible criteria were those who have yet to receive any dose of vaccine approved by Health Canada, received only one dose of a two-dose vaccine and have not yet decided to have a second dose, or have received both doses of a two-dose vaccine at Wave 1. In subsequent waves, the fully

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

vaccinated eligibility criterion will also include the one-dose vaccine that has been approved by Health Canada.

In Wave 1 of the survey, after removing speeders from the sample, the final total sample size is 1958 (Figure 1). 50% (*N*=983) of the sample was fully vaccinated, approximately 45% (*N*=848) unvaccinated, and 5% (n=127) had a single dose without an intention to seek a second dose. The survey was administered online in English and French. Data collection for Wave 1 of CCES (*N*=1958) took place between 28 September and 21 October, 2021, with additional waves scheduled for every 6-month period, e.g., Wave 2 of CCES is planned for February 2022 (Figure 2).

Sampling

A typical objective of the survey is to estimate descriptive and analytical parameters of the population. Weighting was used to represent the Canadian population based on information gathered from the sample and population figures from the Canadian census. Samples were first post stratified by geographic/language regions: Alberta, British Columbia, Manitoba + Saskatchewan, Ontario, Quebec English, Quebec French, and the Atlantic provinces (Nova Scotia, New Brunswick, Prince Edward Island, Newfoundland and Labrador). For each of the vaccinated and vaccine hesitant group separately, sampling weights were then computed using a raking procedure and calibrated to target marginal joint population distributions of the geographic/language regions, and the gender and age group combinations, based on population benchmarks in the 2016 Canadian census data and the disposition code in the sample, thus allowing generalization to the Canadian population. Two key survey statistics were computed for Wave 1 data: the response rate, defined the number of eligible respondents who completed the survey divided by the estimated number of eligible respondents that were selected/contacted (the American Association for Public Opinion Research (AAPOR) Response Rate 4 (RR4)), and the cooperation rate, defined as the proportion of eligible respondents (i.e., those who have completed all eligibility questions and have been found to be eligible) who completed the survey (AAPOR Cooperation Rate 4 (COOP4).[1] The cooperation rate is 94.8% and the response rate is 13.0%.

Measures

All measures were completed online, and questions were delivered in the native language of the respondent (English or French). Question categories are described in the sections that follow:

COVID-19 Infection History and Symptom Severity

SARS-CoV-2 infection history. All participants were asked, “What best describes YOUR experience with COVID-19 infection?” Those who reply “I have NOT been infected” are then asked, “How do you know that you HAVE NOT BEEN infected with COVID-19?” followed by a number of options to indicate lack of symptoms, a negative test or other reasons. Those who respond “I have been infected” to the original question are instead asked, “How do you know that you HAVE BEEN infected with COVID-19?”, followed by a number of reasons, including symptoms and a positive test result.

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

COVID-19 Disease Severity. Among those who reported being infected, respondents are then asked to describe the severity level of the symptoms using the following item stem: "How severe was your COVID-19 illness?", with the following response options: "Not at all severe", "Slightly severe", "Moderately severe", "Very severe", "Extremely severe". Care-level indicators of COVID-19 severity were assessed via two items: 1) hospitalization history for COVID-19, and 2) whether or not they were fitted with a ventilator during a hospitalization.

COVID-19 Mitigation Behaviors

Participants were asked to indicate their level of consistency with respect to implementation of COVID-19 mitigation behaviours using a general measure ("How consistently do you follow the recommendations by your local or provincial public health officials about social distancing?") as well as behavior-specific measures ("When outside your home, how consistently do you currently maintain a distance from others of at least 2 meters?"). Specific behaviors assessed include distancing (maintaining 2-meter separation), mask wearing, and hand hygiene.

Cognitive Variables

Executive dysfunction. Executive function is a set of cognitive processes that constitute the ability to inhibit impulses, the ability to work with information in an "online" state, and switch flexibly back and forth between modes of responses (Barkley & Fischer, 2010; Miyake et al., 2000). Executive dysfunction was assessed using four items from the "self-restraint" subscale of the Barkley Deficits in Executive Function Scale (BDEFS; Barkley et al., 2011). Respondents were asked, "How often have you experienced each of the problems listed below during the past 6 months?" Following this, respondents were required to indicate frequency ranging from "Never or rarely" to "Very often". Items included the following: "I am unable to inhibit my reactions or responses to events or to other people", and "I act without thinking", "I make impulsive comments to others", and "I am likely to do things without considering the consequences for doing them".

Delay discounting. Delay discounting is the tendency to devalue rewards disproportionately to their absolute value, as a function of delay time to receipt (Bickel & Marsch, 2001). Those who have strong delay discounting have a tendency to impulsively reach for a lesser but immediately available reward over a much later but delayed alternative. Delay discounting was assessed using a dynamic 5-question version previously validated by Koffarnus and colleagues (Koffarnus et al., 2014). The module introduction started with the following instructions: "For the next 5 questions, you will be asked to choose between receiving different amounts of money at different points in time. You will see two options. You are requested to choose one of them. There are no right or wrong answers, but please take your time, answer thoughtfully, and pay careful attention to each of the options." Participants were then given a series of monetary offers for the participant to respond to, starting with "Would you rather have $ 500 now or $ 1000 in 3 weeks?" These questions continued with the same fixed monetary amounts, but with a varied time delay for the larger amount (e.g., 1 day, 2 years). The item responses are converted to a "k" value, where higher values of k reflect higher levels of discounting of rewards as a function of time delay to receipt.

Time perspective. Time perspective was assessed with four survey items derived from the Time Perspective Questionnaire (TPQ; Hall et al., 2015), two pairs of items: one pair reflecting a present orientation and a second pair reflecting a future orientation. The item pairs

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

items may be used to assess separate values for each orientation in an orthogonal manner, or subtracted (future item pair score – present item pair score) such that higher scores on the metric reflect relatively stronger future than present orientation. Instructions for the module were as follows: "For each of the statements below, indicate your level of agreement or disagreement by using the following scale". For each statement, participants were required to rate their level of agreement from "Strongly agree" to "Strongly disagree". Present oriented items were as follows: "Living for the moment is more important than planning for the future." and "I spend a lot more time thinking about today than thinking about the future." Future oriented items were as follows: "I spend a lot of time thinking about how my present actions will have an impact on my life later on." and "I consider the long-term consequences of an action before I do it."

Mood symptoms (starting in Wave 2)

Anxiety. Anxiety symptoms are measured using the Generalized Anxiety Disorder 7-item scale (GAD-7; Spitzer et al., 2006). This scale consists of 7 items assessing a variety of symptoms of anxiety, such as: "feeling nervous, anxious, or on edge", and "Feeling afraid, as if something awful might happen". Responses are given on a 4-point scale, where 1 = "Not at all", 2 = "Several days", 3 = "More than half of the days", 4 = "Nearly every day". The time window for responses was 2 weeks following recovery from COVID-19.

Depression. Symptoms of depressions are assessed using the Centres of Epidemiological Scale – 10 item Scale (CESD-10; Andresen et al., 1994). This scale consists of 10 items assessing symptoms of low mood, lack of energy, and negative thinking styles. Sample items include "I was bothered by things that usually don't bother me.", and "I felt lonely.". Responses are given on a 4-point scale, where 1 = "Rarely or none of the time", 2 = "Some or a little of the time", 3 = "Occasionally or a moderate amount of the time", 4 = "All of the time". The time window for responses was 1 week following recovery from COVID-19.

Agitation. Agitation is assessed using three items custom designed for the survey. Respondents are asked to indicate how often they have experienced the following symptoms during the week following recovery from COVID-19: ""I felt agitated.", "I lashed out at other people in a way that was not like me." "I was much more irritable than usual." Responses are given on a 4-point scale, where 1 = "Rarely or none of the time", 2 = "Some or a little of the time", 3 = "Occasionally or a moderate amount of the time", 4 = "All of the time". The time window for responses was 1 week following recovery from COVID-19.

Social Cognitive Variables

Social Norms

Descriptive Norms: A descriptive norm is the perceived number of people that someone believes engage in a behaviour. Respondents were asked about their perceived descriptive norms of vaccination, mask wearing, social distancing, and handwashing among: 1) their five most important people, 2) their province, and 3) in Canada. Respondents were first asked to list the five most important people in their life. They were then asked to report the vaccination status of their five most important people (fully vaccinated, one vaccine, no vaccines). Finally, respondents were asked to estimate what % of people in their province and what % of people in Canada are fully vaccinated or will be fully vaccinated in the next four months. For social

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

distancing, mask wearing, and hand washing, respondents were asked to report how many of their five most important people consistently engage in the behaviour.

Injunctive Norms: Injunctive norms are people's perspectives about what others believe. Respondents were asked whether they "Strongly agree" to "Strongly disagree" on a 5-point scale that 1) their five most important people believe that it is important to be vaccinated, 2) people in their province believe that it is important to be vaccinated, and 3) that people in Canada believe that it is important to be vaccinated. They were then asked the same sequence of questions about social distancing, wearing masks, and washing hands.

Dynamic Injunctive Norms: Dynamic injunctive norms are people's beliefs about how they think the beliefs of others are changing over time. Respondents were asked which statement best describes how the beliefs about vaccines of their five most important people, people in their province, and people in Canada have changed in the last four months, with the following response options: 1) more of them believe that it is important to be fully vaccinated, 2) there is no change in the number that believe it is important to be fully vaccinated, and 3) fewer of them believe that it is important to be fully vaccinated. They were then asked questions about how the beliefs about social distancing, wearing masks, and hand hygiene of their five most important people, people in their province, and people in Canada had changed in the last four months using similar response options.

Reasons

Reasons for vaccination outcomes are assessed using a three-point response scale including the following options: "No, not at all a reason for me", "Yes, this is somewhat of a reason for me", and "Yes, this is an important reason for me." The source list of reasons is presented in Table 1.

Reasons for not getting vaccinated and not getting fully vaccinated: Respondents who were not vaccinated or had only received one dose with no plans for a second were asked reasons for not getting vaccinated or fully vaccinated.

Perceived reasons for not getting vaccinated: Fully vaccinated people were then presented with a list of reasons for not getting vaccinated and then asked to indicate what % of people who are not vaccinated would give those reasons.

Reasons for getting fully vaccinated: Among fully vaccinated people who were initially hesitant about getting vaccinated, reasons for getting fully vaccinated were asked.

COVID-19 Severity Beliefs and Worry

Severity Beliefs. Beliefs about COVID-19 severity were assessed using the following stem: "Below are a number of statements about COVID-19. For each, indicate your level of agreement." Responses were given on a 5-point scale with response options ranging from "Strongly agree" to "Strongly disagree." Items are presented in Table 2.

Worry. A number of worry items were added, pertaining to the severity of COVID-19 impact on the self (e.g., "How worried are you that you will get very sick from COVID-19?"), as well as the impact on others (e.g., "How worried are you that a family member or a close friend will get very sick from COVID-19?"). Responses were given on a 5-point scale from "Not at all worried" to "Extremely worried."

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Perceived Effectiveness and Beliefs About Mitigation Behaviors

Participants were asked about perceived levels of effectiveness for recommended COVID-19 mitigation behaviors using the following stem: "Below are a number of statements about masks. For each, indicate your level of agreement." Responses were given on a 5-point scale from "Strongly agree" to "Strongly disagree." Examples of items include, "If worn properly, masks can protect the wearer from getting infected by COVID-19.", and "Masks make it hard to breathe properly." Behavioral targets included mask wearing, social distancing, hand hygiene, and vaccinations.

Vaccination Intentions

Vaccination intentions were assessed with two items, one focussed on the original vaccination and the other focussed on a hypothetical booster shot recommended in the future. The items were as follows: "What best describes your intention to get fully vaccinated in the next 4 months?" and "What best describes your intention to get an additional vaccine shot in the future (e.g., a booster shot) if the government says that it is necessary?" Response options were as follows: [I have] "No intention to get fully vaccinated", "A very low intention", "A low intention", "A moderate intention", "A strong intention", and "A very strong intention."

Political orientation, Trust in Science, Trust in Information Sources

Political orientation. Political orientation measures included political orientation at the provincial and national level, and where people place themselves on the political spectrum on a 7-point scale ranging from "Extremely liberal" to "Extremely conservative", which internationally equates to "Extremely left" to "Extremely right" due to differences in the political meaning of liberal. Trust in COVID-19 information from political leaders and COVID-19 measures was also measured at the provincial and national level.

Trust in science. Trust in science was assessed using level of agreement for a variety of statements, listed in Table 3. Responses were on a 5-point scale from "Strongly agree" to "Strongly disagree." Items were obtained from Nadelson and colleagues (Nadelson et al., 2021); these items have been previously validated with respect to the COVID-19 context (Travis et al., 2021).

Information sources. Sources for obtaining COVID-19 information were assessed with the following item: "How much, if at all, do you currently get information about COVID-19 from each of the following sources?" Respondents were asked to rate on a 4-point scale from "Not at all" to "Very much" each of the following options: Church/religious group, doctor, newspapers/magazines, television, social media, or other sources.

PSA Evaluations

A collection of public service announcements (PSAs) encouraging vaccination and COVID-19 mitigation behaviors and vaccination were professionally developed for the purpose of the CCES. Videos are 25 seconds in duration, and were produced in both official languages (i.e., English and French). Following each video clip, respondents rated the video on a number of evaluative dimensions, which are listed in Table 4. Most items assess the extent to which the PSAs were effective at capturing attention, motivating behavior, and encouraging reflection.

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Several items assessed the extent to which each PSA adhered to its intended thematic content (e.g., "How effective was this ad in getting people to think about the future benefits vs. the short-term costs of [getting vaccinated/performing mitigation behaviors]"). Rating responses were provided on a 5-point scale from "Not at all" to "Extremely."

Demographics

Age in years, gender, marital status, income, education, ethnicity were assessed via self-report. Financial strain was assessed using two items: "In the last 30 days, because of a shortage of money, were you unable to pay any important bills on time, such as electricity, telephone or rent bills?" and "Are you currently receiving assistance or income from any federal, provincial or local programs, such as food bank use, welfare/social assistance, employment insurance, etc.?" Additional items were included to assess the extent to which individuals lost a job and/or received government benefits of various types. Demographic features of the Wave 1 CCES sample are presented in Table 5.

Discussion

The CCES is one of many studies examining predictors of vaccination and adherence in the wake of the COVID-19 crisis. For this reason, findings from this study may be considered collectively with other studies to draw the most reliable conclusions possible about the predictors of vaccine hesitancy and mitigation behavior implementation. There are, however, several features that distinguish the CCES from other studies, including the integration of quota sampling and the inclusion of neurocognitive measures. The former ensures that the CCES will have maximum statistical power to identify reliable predictors of vaccine hesitancy, perhaps more so than much larger studies that do not employ quota sampling. The equal numbers of vaccinated and vaccine hesitant individuals in the sample mean that limited power will be an unlikely reason for any null predictors; conversely, we are unlikely to falsely identify significant predictors, for the same reasons.

The inclusion of neurocognitive predictors of mitigation behaviors is also potentially novel, though not entirely unique to our study. We will however be able to examine interactions between neurocognitive and social cognitive predictors, in part because of the wide breadth of the latter measures included. Finally, the timing of the first wave ensured wide availability of vaccines in every province in Canada, ensuring that limited access is not likely to be a confound for any effects that we might observe.

Finally, a primary purpose for the conduct of this survey was to evaluate a large pool of custom developed PSAs for promoting vaccination and COVID-19 mitigation behaviors among members of the general public. The CCES will enable identification of effective communication approaches for use in future public-facing media campaigns for infectious disease mitigation. Combining this component with the aforementioned array of social cognitive and neurocognitive assessments, it will be possible to identify which messaging strategies work for which members of the population, as a function of social cognitive (and even neurocognitive) characteristics. Among the most important objectives of the CCES will be to identify thematic and perceptual features of PSAs that appeal differentially to those who are vaccine hesitant versus those who are already fully compliant.

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Some limitations and logistical challenges remain however. First, as the proportion of vaccinated individuals continues to rise, this subgroup will be increasingly difficult to identify and reach. In the current study context, the proportion of fully vaccinated individuals was well above 70% already at Wave 1. However, it is anticipated that booster shots coinciding with future waves will provide an opportunity to shift focus from two vaccines to three. Finally, as the media becomes increasingly saturated with communications about COVID-19, and as the threat posed by the pandemic waxes and wanes, modelling temporal factors over time will become increasingly important.

Conclusions

In conclusion, the CCES provides an opportunity to learn more about what drives vaccination and mitigation behavior during an active pandemic situation. What we learn in the context of COVID-19 will continue to provide important clues as to how to manage future infectious disease threats on the population level, and what factors within the social cognitive domain and neurocognitive domain might predict uptake of recommendations. Beyond this, the CCES will provide highly novel information about what dimensions of PSA content are most appealing to the general public, and how this may vary systematically by vaccine hesitancy status.

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

References

American Association for Public Opinion Research. Standard definitions: Final dispositions of case codes and outcome rates for surveys. Retrieved from. 2016 Jan 28.

Andresen EM, Malmgren JA, Carter WB, Patrick DL. Screening for depression in well older adults: Evaluation of a short form of the CES-D. American journal of preventive medicine. 1994 Mar 1;10(2):77-84.

Barkley RA. *Barkley deficits in executive functioning scale (BDEFS)*. Guilford Press; 2011 Feb 1.

Barkley RA, Fischer M. Predicting impairment in major life activities and occupational functioning in hyperactive children as adults: Self-reported executive function (EF) deficits versus EF tests. Developmental neuropsychology. 2011 Jan 31;36(2):137-61.

Bickel WK, Marsch LA. Toward a behavioral economic understanding of drug dependence: delay discounting processes. Addiction. 2001 Jan;96(1):73-86.

Hall PA, Fong GT, Sansone G. Time perspective as a predictor of healthy behaviors and disease-mediating states. In Time perspective theory; review, research and application 2015 (pp. 339-352). Springer, Cham.

Koffarnus MN, Bickel WK. A 5-trial adjusting delay discounting task: accurate discount rates in less than one minute. Experimental and clinical psychopharmacology. 2014 Jun;22(3):222.

Miyake A, Friedman NP, Emerson MJ, Witzki AH, Howerter A, Wager TD. The unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks: A latent variable analysis. Cognitive psychology. 2000 Aug 1;41(1):49-100.

Nadelson L, Jorcyk C, Yang D, Jarratt Smith M, Matson S, Cornell K, et al. I Just Don’t Trust Them: The Development and Validation of an Assessment Instrument to Measure Trust in Science and Scientists. School Science and Mathematics. 2014;114[2]:76–86.

Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. Archives of internal medicine. 2006 May 22;166(10):1092-7.

Travis J, Harris S, Fadel T, Webb G. Identifying the determinants of COVID-19 preventative behaviors and vaccine intentions among South Carolina residents. Plos one. 2021 Aug 25;16(8):e0256178.

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Table 1

Reasons items

1. The chances of me becoming really sick from COVID-19 are low.
2. I already had COVID-19 so I don't need the vaccine.
3. I do not think that the vaccine is effective against COVID-19.
4. I am healthy and at low risk for COVID-19.
5. I practice enough other behaviours (e.g., mask wearing, social distancing) to avoid the COVID-19 virus so I don't need to get fully vaccinated.
6. Getting vaccinated is not necessary because COVID-19 is not a real threat.
7. I would rather build my immunity by being infected rather than by getting a vaccine.
8. I am against vaccination in general.
9. I have concerns about the safety of the COVID-19 vaccine.
10. I have concerns about side effects of the COVID-19 vaccine.
11. The COVID-19 vaccine could cause autism and other serious diseases
12. The COVID-19 vaccine could cause COVID illness.
13. The COVID-19 vaccine could alter my DNA.
14. I am worried about unknown future effects of the COVID-19 vaccine.
15. I do not like needles and injections.
16. I need more information about the safety and effectiveness of the COVID-19 vaccine.
17. I do not have enough time to get vaccinated.
18. I don't need to be vaccinated because there will be enough others who will get vaccinated.
19. I am concerned that vaccines are being used for some other purpose.
20. I am not ready to decide whether to get fully vaccinated.
21. I do not trust the authorities who are recommending the vaccine.
22. The scientists who developed the vaccine are not sure that it is safe and effective.
23. The government has no right to tell me what I have to do with my body—that's my choice.
24. Some of my family and/or friends have concerns about the safety of the vaccine.
25. Some of my family and/or friends have concerns about the effectiveness of the vaccine.
26. None or very few of my family and/or friends have gotten fully vaccinated.
27. Information about the vaccine from TV makes me concerned about the safety and/or effectiveness of the vaccine.
28. Information about the vaccine from social media (e.g., Facebook, Twitter, Instagram, YouTube, chat rooms) makes me concerned about the safety and/or effectiveness of the vaccine.
29. I know people who have had a bad reaction to a COVID vaccine.
30. I know people (adults or children) who have had bad reactions to other vaccines.
31. I don't know anyone who has gotten seriously sick from COVID-19.

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Table 2

Beliefs about COVID-19 severity, origins, and mitigation measures

1. COVID-19 is an extremely serious threat to public health.
2. COVID-19 was manufactured by humans.
3. The number of people dying from COVID-19 is being exaggerated by the authorities.
4. Public health officials are doing the best they can to fight COVID-19.
5. COVID-19 is being made into a bigger deal than it actually is.
6. Certain groups want COVID-19 to continue because they profit from it.
7. The shutdown of the economy due to COVID-19 has caused more damage than the COVID-19 illness itself.
8. COVID-19 is no worse than the flu.
9. COVID-19 started when the virus escaped from a lab in China.
10. COVID-19 started when the virus was transmitted from animals in a Chinese market to humans.
11. Closing schools to prevent COVID-19 transmission has caused more harm than good in the long-term.
12. Lockdowns to prevent COVID-19 transmission hurt the economy in the long-term.
13. I have made efforts to improve air quality at home (e.g., by using a better ventilation filter or a HEPA air purifier)

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Table 3

Trust in science items

1. We can trust science to find the answers that explain the natural world.
2. There are other perspectives besides science to uncover the truth.
3. We can trust scientists to find solutions to major problems.
4. Scientists work to help people.
5. Scientific findings often contradict each other so it's hard to figure out what is true.
6. Scientists are honest and ethical in their work
7. We cannot trust scientists because their findings are often driven by their desire to advance their careers.
8. Scientists are arrogant.

From: Nadelson L, Jorcyk C, Yang D, Jarratt Smith M, Matson S, Cornell K, et al. I Just Don't Trust Them: The Development and Validation of an Assessment Instrument to Measure Trust in Science and Scientists. School Science and Mathematics. 2014;114[2]:76–86.

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Table 4

PSA Evaluation questions

1. How much did you like this ad?
2. How engaging was this ad?
3. How much did this ad make you stop and think about [how consistently you (wear masks where and when advised/ follow advice about social distancing/ wash your hands) / your vaccination status]?
4. How attention-grabbing was this ad?
5. How relevant was this ad to your life?
6. How eye-catching was this ad?
7. How much did this ad make you self-reflect?
8. How effective would this ad be in motivating people to [get fully vaccinated/wear masks/socially distance/handwash frequently and thoroughly]?
9. How effective would this ad be in motivating YOU to [get fully vaccinated (or to get future shots) if recommended/wear masks (or to continue wearing masks/socially distance (or to continue to socially distance)/wash your hands frequently and thoroughly]?
10. How effective was this ad in describing the benefits of [getting fully vaccinated/wearing masks/socially distancing/handwashing frequently and thoroughly]?
11. How effective was this ad in describing the costs of NOT [getting fully vaccinated/wearing masks/socially distancing/handwashing frequently and thoroughly]?
12. How effective was this ad in getting people to think about the future benefits vs. the short-term costs of [getting fully vaccinated/wearing masks/socially distancing/handwashing frequently and thoroughly]?
13. How effective was this ad in highlighting the increase in Canadians who are [getting fully vaccinated/wearing masks/socially distancing/handwashing frequently and thoroughly]?
14. How worried did this ad makes you feel?

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Table 5

Demographic features of the CCES Wave 1 sample.

variable	value	Frequency	%
Gender	Male	768	39.22
	Female	1190	60.78
Age group	18-24	322	16.45
	25-39	789	40.3
	40-54	847	43.26
Income	Low	305	15.58
	Moderate	436	22.27
	High	1031	52.66
	No answer	186	9.5
Education	Low	408	20.84
	Moderate	710	36.26
	High	810	41.37
	No answer	30	1.53
Geographic regions	AB	245	12.51
	BC	244	12.46
	MB+SK	119	6.08
	Maritimes	111	5.67
	ON	737	37.64
	QC-EN	134	6.84
	QC-FR	368	18.79
Vaccination status	I have NOT received any vaccine shot	848	43.31
	Received ONE vaccine shot	127	6.49
	Received TWO vaccine shots	983	50.2
COVID19 infection history	Not infected	1640	83.76
	infected: Not at all severe	57	2.91
	infected: Slightly severe	46	2.35
	infected: Moderately severe	52	2.66
	infected: Very severe	18	0.92
	infected: Extremely severe	4	0.2
	not stated	134	6.84
	infected: severity not stated	7	0.36
Hospitalization history	Not infected	1640	83.76
	not stated	134	6.84
	infected not Hospitalized	174	8.88
	infected hospitalized: no ventilator	5	0.26
	infected hospitalized: ventilator	5	0.26

Note: *N*=1,968.

medRxiv preprint doi: https://doi.org/10.1101/2021.12.24.21268387; this version posted December 27, 2021. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

Informed Consent Statement:

Informed consent was obtained on the web survey from all study participants prior to completing the survey.

Ethics:

The survey protocols and all materials, including the survey questionnaire, received ethical approval from the Office of Research Ethics, University of Waterloo, Canada (ORE# 43569).

Conflict of interest statement:

The authors have no conflicts.

Acknowledgments:

The authors would like to acknowledge and thank all those that contributed to The Canadian COVID-19 Experiences Survey: all study investigators and collaborators, and the project staff at their respective institutions, namely Jamie Wheeler, Kai Jiang, Shelly Jordan, Simon Thompson, Lindsey Webster of the University of Waterloo, Andrew Mattern and Samantha Rochon of Leger Opinion.

Author contributions:

PH, GF and SH conceived the study, obtained the research funding and led the writing of the manuscript; AQ and TA participated in the design and implementation of the survey, as well as the monitoring of the data collection process; GM and CB devised and implemented the sample weighting procedures; MS and AH participated in the writing of the manuscript.

Funding Statement

This research was supported by an operating grant to P. Hall, G. Fong and S. Hitchman by the Canadian Institutes for Health Research (CIHR), Institute for Population and Public Health (GA3-177733).



Figure 1. Study recruitment diagram for Wave 1 of the CCES. "1 dose"=1 vaccine dose with no intention to get a second dose (applies to 2 dose regimens only). Target recruitment for Wave 1 and future waves is set provisionally at 2,000.

*Sampling frame defined as those Canadians with fixed broadband internet access (estimated as 94% of the Canadian population as of 2020; StatsCan).



Figure 2. Prospective survey design for the first three waves of CCES data collection.