Can individuals with suboptimal antibody responses to conventional

antiviral vaccines acquire adequate antibodies from SARS-CoV-2 mRNA

vaccination?

Wataru Ogura[1], Kouki Ohtsuka[2*], Sachiko Matsuura[1], Takahiro Okuyama[1], Satsuki

Matsushima[2], Satoko Ymasaki[2], Hiroyuki Miyagi[1], Kumiko Sekiguchi[1], Hiroaki

Ohnishi[2], Takashi Watanabe[2]

[1] Department of Clinical Laboratory, Kyorin University Hospital, Mitaka, Tokyo, Japan

[2] Department of Laboratory Medicine, Kyorin University School of Medicine, Mitaka,

Tokyo, Japan

* Corresponding author

E-mail: kouki7@ks.kyorin-u.ac.jp

Funding Statement

This study did not receive any funding

Reprint billing address

Kouki Ohtsuka

Department of Laboratory Medicine

Kyorin University School of Medicine

6-20-2, Shinkawa, Mitaka, Tokyo, 181-8611, Japan

E-mail: kouki7@ks.kyorin-u.ac.jp

Abstract:

Objective

In Japan, healthcare workers (HCWs) are vaccinated against contagious viruses (measles, rubella, chickenpox, mumps, and hepatitis B) to prevent nosocomial infection; however, some do not produce sufficient antibodies (suboptimal responders). Whether suboptimal responders to live attenuated viruses or inactivated viruses vaccines can produce adequate antibodies to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) mRNA vaccines remains to be elucidated.

Methods

In this prospective cohort study, SARS-CoV-2 anti-spike antibodies were measured 11 times, from before the first BNT162b2 vaccination to 5 months after the second vaccination. Antibody titers of suboptimal and normal responders were compared. SARS-CoV-2 neutralizing antibody activity was measured twice in suboptimal responders, 1 week to 1 month, and 5 months after the second vaccination.

Patients

This study included 50 HCWs who received two doses of mRNA BNT162b2 vaccine 3 weeks apart.

Results

After vaccination, the SARS-CoV-2 anti-spike antibody was detectable in the samples from suboptimal and normal responders at each timepoint. The median SARS-CoV-2 anti-spike antibody titer was higher in suboptimal responders than in normal responders 1 week after receiving the second dose of BNT162b2 vaccine (3721.0 vs. 2251.5, P=0.029). Suboptimal responders had SARS-CoV-2 neutralizing antibody

activity 1 week to 1 month, and 5 months after the second vaccination, which exceeded

the positive threshold 5 months after the second vaccination.

Conclusion

After BNT162b2 vaccination, suboptimal responders acquired adequate SARS-

CoV-2 anti-spike and SARS-CoV-2 neutralizing antibodies to prevent SARS-CoV-2.

These results suggest that vaccination with mRNA vaccine against SARS-CoV-2 should

also be recommended for suboptimal responders to conventional vaccines.

Keywords:

coronavirus disease 2019 (COVID-19), severe acute respiratory syndrome coronavirus 2

(SARS-CoV-2), mRNA vaccine, anti-spike antibody, neutralizing antibody

Introduction

Coronavirus disease (COVID-19), which is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has been declared a pandemic (1-4). In view of the high number of cases of severe illness and death due to COVID-19, vaccination against SARS-CoV-2 infection is crucial.

SARS-CoV-2 has an enveloped, single, positive-stranded RNA genome that encodes four major viral structural proteins, namely, the nucleocapsid, spike, envelope, and membrane proteins; the latter three proteins are found in its membrane. The spike protein guides viral entry into host cells by binding to angiotensin-converting enzyme 2 (ACE2) receptors, which are widely expressed on epithelial cells and macrophages (5-7) and is thus an ideal target for messenger RNA (mRNA) vaccine development.

The BNT162b2 mRNA vaccine (Pfizer–BioNTech) is highly effective at preventing clinically significant COVID-19 (8-10), reducing the incidence of asymptomatic infection and associated infectivity (11,12), and preventing the spread of COVID-19 (9). The BNT162b2 vaccine was the first SARS-CoV-2 vaccine to be approved in Japan, and priority vaccination of healthcare workers (HCWs) began in February 2021 in order to safeguard the medical delivery system (13).

HCWs also have a high risk of infection with other contagious viruses such as measles, rubella, chickenpox, mumps, and hepatitis B. Therefore, in Japan, HCWs are vaccinated against these viruses to prevent nosocomial infections and viral antibody titers are measured to ensure the presence of protective antibody levels (14). However, some individuals do not develop adequate antibody titers against contagious viruses even after vaccination. Such individuals are called suboptimal responders. It is unclear

whether suboptimal responders to conventional vaccines can produce adequate SARS-

CoV-2 antibodies, including neutralizing antibodies, after vaccination with the

BNT162b2 vaccine to protect them against SARS-CoV-2 infection. Therefore, in this

study, we investigated whether suboptimal responders acquired SARS-CoV-2 anti-spike

and SARS-CoV-2 neutralizing antibodies after vaccination with the BNT162b2 vaccine.

Materials and methods

Participants

The study was conducted at Kyorin University Hospital in Tokyo, Japan.

HCWs at the hospital received two doses of the BNT162b2 vaccine 3 weeks apart. The

participants were HCWs at the hospital who agreed to participate in the study and to

periodically donate blood samples before and after vaccination. Those whose antibody

levels were below the standard values for preventing infection after receiving two or

more doses of vaccine against measles, rubella, chickenpox, mumps, or hepatitis B

viruses were classified as suboptimal responders.

SARS-CoV-2 anti-spike antibody assay

SARS-CoV-2 anti-spike antibody measurements were performed on serum

obtained from centrifuged peripheral blood using the Elecsys Anti-SARS-CoV-2 S RUO

(Roche Diagnostics International Ltd, Rotkreuz, Switzerland) on a Cobas 6000 (Roche

Diagnostics) analyzer via the double-antigen sandwich enzyme-linked immunoassay

method (15). Samples were collected before the first dose of BNT162b2 vaccine was

administered; 3 days after the first vaccination; 1, 2, and 3 weeks after the first

vaccination; 1 week after the second vaccination; and 1, 2, 3, 4, and 5 months after the

second vaccination. The measurement range of the antibody assay was 0.4–250, and

samples with an antibody titer of ≥250 U/mL were assayed again after dilution.

SARS-CoV-2 neutralizing antibody assay

Using serum obtained from the peripheral blood of suboptimal responders after

vaccination, SARS-CoV-2 neutralizing antibody activity was measured at Osaka

University Microbial Disease Research Association Laboratory. A 50% inhibitory

dilution factor (ID_{50}) using pseudovirus (16) was measured, and ID_{50} values of ≥50 was

considered positive (17). For the neutralization assay, 60 μL of pseudovirus, which is

equivalent to 2.5×10^6 relative light units (RLU)/mL, was incubated with 60 μL of serial

dilutions of serum samples for 1 h at 37□. After incubation, 100 μL of the mixture was

added to the Vero cells in a 96-well plate. The cells were lysed 24 h post-infection,

luciferase was activated using the Luciferase Assay System (E1501, Promega,

Wisconsin, USA), and RLU were measured using a Synergy LX analyzer (BioTek,

USA). Percent neutralization was calculated using GraphPad Prism 8.0.2 (GraphPad

Software, San Diego, California, USA). Percentages of RLU reduction (inhibition rate)

were calculated as follows: 1 − (RLU of sample sera − pseudovirus only wells) / (RLU

from medium only wells − pseudovirus only wells) × 100%. The titers of neutralizing

sera were calculated as ID_{50}. The transition of neutralizing antibody activity values after

vaccination in suboptimal responders was examined 1 week to 1 month and 5 months

after receiving the second dose of the BNT162b2 vaccine.

Statistical analysis

Nonparametric continuous data are reported as median and interquartile range

(IQR). The Mann–Whitney U test was performed to assess whether differences between

cohorts were statistically significant. Two-sided *P*-values of <0.05 were considered

statistically significant. The Wilcoxon matched-pairs signed rank test was performed to

assess whether differences in antibody levels within an individual at different time

points were statistically significant. To compare the age (ratio of the number of

individuals aged 20–40 years and those aged 50–60 years) and sex compositions of the

suboptimal responders and normal responders, Fisher's exact test was performed to test

the statistical significance. The Mann–Whitney U test was performed to test for

statistically significant differences in antibody levels between suboptimal responders

and normal responders 1 week after the second vaccination. All statistical analyses were

performed with EZR (Saitama Medical Center, Jichi Medical University, Saitama,

Japan), which is a graphical user interface for R (The R Foundation for Statistical

Computing, Vienna, Austria) (18). Specifically, EZR is a modified version of R

commander (version 1.54) that has been designed to add statistical functions frequently

used in biostatistics.

Results

The suboptimal responders comprised three men (L1, L2, L4) with negative

(titer <16.0) anti-measles immunoglobulin G (IgG) antibody titer after measles virus

vaccination and one woman (L3) with a negative (titer <4.0) anti-mumps IgG antibody

titer after mumps virus vaccination. Forty-six participants, comprising 12 men and 34

women, had antiviral antibody titers after vaccination against other viruses above the

standard value (normal responders). The four suboptimal responders were aged 20–40

years. Normal responders were older than suboptimal responders (the ratios of the

number of those aged 50–60 years were 57% vs. 0%, respectively), but no significant

difference was observed (P >0.99). In addition, the male/female ratio was higher in

suboptimal responders than in normal responders (74% vs. 15%), but no significant

difference was observed (P >0.99). All suboptimal responders were non-smokers and

non-obese, and none of them had a history of immunosuppressive drug use or having a

medical condition associated with immunodeficiency.

SARS-CoV-2 anti-spike antibody levels

The median (IQR) SARS-CoV-2 anti-spike antibody titer was 2251.5 (1639.0–

3276.3) in the normal responders and 3721.0 (3255.0–4983.3) in the suboptimal

responders 1 week after receiving the second dose of BNT162b2 vaccine. This

difference was statistically significant (P=0.029) (Fig. 1A). For the total antibody level

1 week after the second vaccination, there was no significant difference in the median

antibody levels between those aged 50–60 years and those aged 20–40 years(P=0.66).

The median (IQR) SARS-CoV-2 anti-spike antibody in normal responders was

2251.5 (1639.0–3276.3) and 779.3 (534.6–1024.3) measured 1 week and 5 months,

respectively, after receiving the second dose of BNT162b2 vaccine (Fig. 1B). In

suboptimal responders, the median (IQR) SARS-CoV-2 anti-spike antibody titters were

3721.0 (3255.0–4983.3) and 666.7 (578.9–880.6) measured 1 week and 5 months,

respectively, after receiving the second dose of BNT162b2 vaccine (Fig. 1C).

SARS-CoV-2 anti-spike antibody was produced by both suboptimal and normal

responders, but the level was lower at 5 months than at 1 week after the second dose of

vaccine (Fig. 1).

(A) SARS-CoV-2 anti-spike antibody responses over the whole study period from

before the first dose to 5 months after the second dose of vaccine.

(B) SARS-CoV-2 anti-spike antibody titers in normal responders 1 week and 5 months

after receiving the second dose of the BNT162b2 vaccine. The antibody titers were

significantly lower at 5 months than at 1 week after the second dose ($P<0.001$).

(C) SARS-CoV-2 anti-spike antibody titers in suboptimal responders 1 week and 5

months after receiving the second dose of the BNT162b2 vaccine. The antibody titers

decreased, but the decrease was not statistically significant ($P=0.068$).

The box plot comparisons were performed using the Mann–Whitney U test. *P*-values

<0.05 were considered statistically significant. Suboptimal responders had low-level

antibody responses to previous measles or mumps vaccines, and normal responders had

normal antibody responses to previous antiviral vaccines.

NaN; not a number

SARS-CoV-2 neutralizing antibody activity in participants with

suboptimal antibody responses to previous antiviral vaccines

SARS-CoV-2 neutralizing antibody titers were 1154.0 in L1 at 1 week after

receiving the second dose of BNT162b2 vaccine and 774.2, 638.3, and 654.6 in L2, L3,

and L4, respectively, at 1 month after receiving the second dose of BNT162b2 vaccine

(Fig. 2). In the 5 months after the second vaccination, SARS-CoV-2 neutralizing

antibody titers were 105.5, 147.7, 485.2, and 894.4 in L1, L2, L3, and L4, respectively

(Fig. 2). All four suboptimal responders had a neutralizing antibody activity within the

protective range (≥50), from the 1 week to 1 month and 5 months after the second

vaccination (Fig. 2). The SARS-CoV-2 neutralizing antibody level in suboptimal

responders was lower at 5 months after receiving the second dose of BNT162b2 vaccine

than at 1 week to 1 month after receiving the second dose, but this difference was not

statistically significant (P=0.38) (Fig. 2).

Data are presented as 50% inhibitory dilutions (ID_{50}). The dashed line indicates the

cutoff for ID_{50} (17).

Discussion

The BNT162b2 vaccine has been highly effective in preventing clinically

significant COVID-19 (8-10). However, limited information is available about the

effectiveness of the BNT162b2 vaccine in suboptimal responders to conventional

vaccines. In this study, suboptimal responders produced adequate SARS-CoV-2 anti-

spike antibodies and neutralizing antibodies.

Muik et al. (19) measured the neutralization efficacy of sera from 40 healthy

German individuals immunized with the BNT162b2 vaccine and a 50% inhibitory

dilution factor using pseudovirus according to the same method as was used in this

study. The SARS-CoV-2 neutralizing activity of the suboptimal responders in this study

1 week to 1 month after receiving the second dose of BNT162b2 vaccine was

comparable to that of the participants in the German study. All four suboptimal

responders developed protective levels of neutralizing antibodies. The SARS-CoV-2

neutralizing antibody activity value in the fifth month after the second dose of

BNT162b2 vaccine did not show a significant decrease in activity compared with that 1

week to 1 month after receiving the second dose of BNT162b2 vaccine. The SARS-

CoV-2 anti-spike antibody titer was lower at 5 months after the second dose of

BNT162b2 vaccine than at 1 week after the second dose, but the decrease was not

statistically significant in suboptimal responders. These results suggest that mRNA

vaccines can invoke a protective immune response even in suboptimal responders who

do not develop adequate antibody titers against viruses in response to conventional

vaccines against viral diseases such as mumps and measles.

Measles, rubella, chickenpox, and mumps vaccines are live attenuated vaccines

that weaken the toxicity of pathogenic microorganisms, and hepatitis B vaccines are

inactivated vaccines that cause pathogenic microorganisms to lose their ability to infect

(20). In previous studies, approximately 2–10% of healthy individuals do not produce

protective antibody levels in response to viral vaccines (21). The risk factors for low

response to these conventional viral vaccines include male sex, age of >40 years,

smoking habit, obesity, chronic diseases, and genetic factors. However, the factors

associated with a low antibody response to SARS-CoV-2 mRNA vaccines are not well

understood. Smokers, patients with cancer and hematological malignancies, and organ

transplant recipients have been reported to have a lower SARS-CoV-2 anti-spike

antibody response than individuals without these characteristics after receiving SARS-

CoV-2 vaccines (22-26). It is intriguing that the suboptimal responders in this study who

did not produce adequate antibodies against measles or mumps produced adequate

SARS-CoV-2 anti-spike antibodies after receiving the BNT162b2 vaccine. They were

healthy and did not have risk factors such as an organ transplant and

immunosuppression. One possible reason for this discrepancy is the different

mechanism of antibody production between conventional vaccines and SARS-CoV-2

mRNA vaccines.

The BNT162b2 vaccine is an mRNA vaccine that incorporates mRNA

encoding spike protein (antigen) on the surface of the SARS-CoV-2 into human cells.

The produced antigen directly functions as a template to induce the acquired immune

system (27). On the other hand, with live attenuated vaccines or inactivated vaccines, it

is necessary for recipients to process the vaccine and recognize it as an antigen. The

difference in the mechanism of action of vaccines could be a factor contributing to the

discrepancy in response to the BNT162b2 vaccine from the response to conventional

live attenuated vaccines among suboptimal responders. Such a hypothesis should be

tested in future studies. It is also unclear why suboptimal responders had higher

antibody titers than normal responders in the first week after the second dose of vaccine.

Of note, the SARS-CoV-2 anti-spike antibody level is known to peak and then

decrease within months after the second vaccination, which is a much shorter period

than that of antibodies against live attenuated vaccines (28,29). In order to achieve

adequate COVID-19 disease control, a third dose of SARS-CoV-2 mRNA vaccine is

needed, which is already being administered in many countries. In view of the results of

the present study, a third dose of the BNT162b2 vaccine is recommended for both

suboptimal and normal responders to conventional vaccines. (28)

This study has some limitations. The number of participants was limited, and

all participants were HCWs vaccinated at a single hospital. In addition, there were no

suboptimal responders to inactivated vaccines (such as hepatitis B) included in this

study. Multicenter studies with a larger sample size are needed to confirm the results of

the present study. In addition, further immunological studies are necessary to clarify the

mechanism by which humoral immunity is acquired after mRNA vaccination in

suboptimal responders to conventional vaccines.

In conclusion, after two doses of the BNT162b2 vaccine, suboptimal

responders to conventional antiviral vaccines developed protective levels of SARS-

CoV-2 anti-spike and SARS-CoV-2 neutralizing antibodies. These results suggest that

vaccination with mRNA vaccine against SARS-CoV-2 should also be recommended to

suboptimal responders against conventional vaccines.

Ethics consideration

282 　　　We carried out a prospective cohort study of 50 HCWs. This study was

283 conducted with the approval of the Ethics Committee of Kyorin University School of

284 Medicine (Number R02-041), and was in accordance with the Declaration of Helsinki.

285 Informed consent was obtained from all study participants.

Acknowledgments

288 　　　We thank the HCWs of the Department of Clinical Laboratory, Kyorin

289 University Hospital involved in the SARS-CoV-2 antibody monitoring for their valuable

290 contribution.

References

293 1. Coronaviridae Study Group of the International Committee on Taxonomy of

294   Viruses, Baker SC, Baric RS, et al. The species severe acute respiratory

295   syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-

296   2. Nat Microbiol 5:536-544, 2020.

297 2. Wu F, Zhao S, Yu B, et al. A new coronavirus associated with human respiratory

298   disease in China. Nature 579:265-269, 2020.

299 3. Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new

300   coronavirus of probable bat origin. Nature 579:270-273, 2020.

301 4. Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with

302   pneumonia in China, 2019. N Engl J Med 382:727-733, 2020.

303 5. Song Z, Xu Y, Bao L, et al. From SARS to MERS, Thrusting coronaviruses into

304   the spotlight. Viruses 11:59, 2019.

305 6. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 cell entry

306   depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease

307   inhibitor. Cell 181:271-280.e8, 2020.

308 7. Jiang S, Lu L, Liu Q, Xu W, Du L. Receptor-binding domains of spike proteins of

309   emerging or re-emerging viruses as targets for development of antiviral vaccines.

310   Emerg Microbes Infect 1:e13, 2012.

311 8. Haas EJ, Angulo FJ, McLaughlin JM, et al. Impact and effectiveness of mRNA

312   BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases,

313   hospitalisations, and deaths following a nationwide vaccination campaign in

314   Israel: an observational study using national surveillance data. Lancet 397:1819-

315   1829, 2021.

9.   Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2

mRNA Covid-19 vaccine. N Engl J Med 383:2603-2615, 2020.

10.   Dagan N, Barda N, Kepten E, et al. BNT162b2 mRNA Covid-19 vaccine in a

nationwide mass vaccination setting. N Engl J Med 384:1412-1423, 2021.

11.   Regev-Yochay G, Amit S, Bergwerk M, et al. Decreased infectivity following

BNT162b2 vaccination: a prospective cohort study in Israel. Lancet Reg Health

Eur 7:100150, 2021.

12.   Petter E, Mor O, Zuckerman N, et al. Initial real world evidence for lower viral

load of individuals who have been vaccinated by BNT162b2. Preprint; 2021.

13.   Pharmaceuticals and Medical Devices Agency. Special Approval for Emergency

on first COVID-19 vaccine in Japan [media release]. 2021. Available from:

pmda.go.jp. [Cited 2022 February 20] Available from:

https://www.pmda.go.jp/english/about-pmda/0003.pdf

14.   Kumakura S, Shibata H, Onoda K, Nishimura N, Matsuda C, Hirose M.

Seroprevalence survey on measles, mumps, rubella and varicella antibodies in

healthcare workers in Japan: sex, age, occupational-related differences and

vaccine efficacy. Epidemiol Infect. Cambridge University Press 142:12-19, 2014.

15.   Riester E, Findeisen P, Hegel JK, et al. Performance evaluation of the Roche

Elecsys anti-SARS-CoV-2 S immunoassay. J Virol Methods 297:114271, 2021.

16.   Nie J, Li Q, Wu J, et al. Quantification of SARS-CoV-2 neutralizing antibody by a

pseudotyped virus-based assay. Nat Protoc 15:3699-3715, 2020.

17.   McMahan K, Yu J, Mercado NB, et al. Correlates of protection against SARS-

CoV-2 in rhesus macaques. Nature 590:630-634, 2021.

18.   Kanda Y. Investigation of the freely available easy-to-use software 'EZR' for

medical statistics. Bone Marrow Transplant 48:452-458, 2013.

19.   Muik A, Wallisch AK, Sänger B, et al. Neutralization of SARS-CoV-2 lineage

B.1.1.7 Pseudovirus by BNT162b2 vaccine-elicited human sera. Science

371:1152-1153, 2021.

20.   Krammer F. SARS-CoV-2 vaccines in development. Nature 586:516-527, 2020.

21.   Wiedermann U, Garner-Spitzer E, Wagner A. Primary vaccine failure to routine

vaccines: why and what to do? Hum Vaccin Immunother 12:239-243, 2016.

22.   Shem-Tov N, Yerushalmi R, Danylesko I, et al. Immunogenicity and safety of the

BNT162b2 mRNA COVID-19 vaccine in haematopoietic stem cell transplantation

recipients. Br J Haematol 196:884-891, 2022.

23.   Nomura Y, Sawahata M, Nakamura Y, et al. Age and smoking predict antibody

titres at 3 months after the second dose of the BNT162b2 COVID-19 vaccine.

Vaccines (Basel) 9:1042, 2021.

24.   Herzog Tzarfati K, Gutwein O, Apel A, et al. BNT162b2 COVID-19 vaccine is

significantly less effective in patients with hematologic malignancies. Am J

Hematol 96:1195-1203, 2021.

25.   Monin L, Laing AG, Muñoz-Ruiz M, et al. Safety and immunogenicity of one

versus two doses of the COVID-19 vaccine BNT162b2 for patients with cancer:

interim analysis of a prospective observational study. Lancet Oncol 22:765-778,

2021.

26.   Shroff RT, Chalasani P, Wei R, et al. Immune responses to two and three doses of

the BNT162b2 mRNA vaccine in adults with solid tumors. Nat Med 27:2002-

2011, 2021.

27.   Mascellino MT, Di Timoteo F, De Angelis M, Oliva A. Overview of the main anti-

364   SARS-CoV-2 vaccines: mechanism of action, efficacy and safety. Infect Drug

365   Resist 14:3459-3476, 2021.

366   28.   Israel A, Merzon E, Schäffer AA, et al. Elapsed time since BNT162b2 vaccine and

367   risk of SARS-CoV-2 infection: test negative design study. BMJ 375 :e067873,

368   2021.

369   29.   Levin EG, Lustig Y, Cohen C, et al. Waning immune humoral response to

370   BNT162b2 Covid-19 vaccine over 6 months. N Engl J Med 385:e84-e84, 2021.

Figure legends

Fig 1. SARS-CoV-2 anti-spike antibody responses in suboptimal/normal

responders before and after vaccination with the BNT162b2 vaccine

Fig 2. SARS-CoV-2 neutralizing antibody activity responses after the second dose

of BNT162b2 vaccine in suboptimal responders

Fig. 1A

Fig. 1B

Fig. 1C

Fig. 2