1

Effective connectivity of LSD-induced ego dissolution

Devon Stoliker[1*], Leonardo Novelli[1], Franz X. Vollenweider[3], Gary F. Egan[1,2], Katrin H. Preller[3 +], and Adeel Razi[1,2,4,5 +]

[1]Turner Institute for Brain and Mental Health, Monash University, Clayton, VIC

[2] Monash Biomedical Imaging, Monash University, Clayton, VIC

[3] Department of Psychiatry, Psychotherapy & Psychosomatics, University Hospital for Psychiatry, Zurich, Switzerland

[4] Wellcome Centre for Human Neuroimaging, UCL, London, United Kingdom

[5]CIFAR Azrieli Global Scholars Program, CIFAR, Toronto, Canada

+ Joint Senior Authors

* Corresponding author

Name: Devon Stoliker

Address: Monash Biomedical Imaging,

762-772 Blackburn Rd,

Clayton VIC 3168. Australia.

Email: devon.stoliker@monash.edu

Keywords: Ego dissolution; LSD; resting state functional MRI; dynamic causal modelling; predictive coding

Abstract:

Classic psychedelic-induced ego dissolution involves a shift in the sense of self and blurring of boundary

between the self and the world. A similar phenomenon is identified in psychopathology and is associated

to the balance of anticorrelated activity between the default mode network (DMN) –

which directs attention inwards – and the salience network (SN) – which recruits the dorsal attention

network (DAN) to direct attention outward. To test whether change in anticorrelated networks underlie

the peak effects of LSD, we applied dynamic causal modeling to infer effective connectivity of resting

state functional MRI scans from a study of 25 healthy adults who were administered 100mg of LSD, or

placebo. We found that change in inhibitory effective connectivity from the SN to DMN became

excitatory, and inhibitory effective connectivity from DMN to DAN decreased under the peak effect of

LSD. These changes in connectivity reflect diminution of the anticorrelation between resting state

networks that may be a key neural mechanism of LSD-induced ego dissolution. Our findings suggest the

hierarchically organised balance of resting state networks is a central feature in the construct of self.

Significance:

The findings can inform the parallel between the maintenance of subject-object boundary and changes to

anticorrelated canonical resting state brain networks. Effective connectivity informs the hierarchical

organisation of brain networks underlying modes of perception. Moreover, the anticorrelation of brain

networks is an important measure of mental function. Understanding the neural mechanisms of

anticorrelation change under psychedelics help identify its relationship to psychosis and its association to

psychedelic assisted therapeutic outcomes.

Keywords – fMRI, psychedelics, resting state connectivity, Dynamic causal modelling, LSD, ego dissolution

Introduction:

Classic psychedelics are powerful substances with low toxicity that can temporarily alter brain activity

and produce profound changes to consciousness (Carhart-Harris, 2018; Johnson, Richards, & Griffiths,

2008; Preller & Vollenweider, 2018; F. X. Vollenweider, 1998; F. X. Vollenweider et al., 1997).

Historical use of their mind altering effects that are undergoing translation into modern clinical therapies

may constitute a crucial component of their therapeutic efficacy (Roseman, Nutt, & Carhart-Harris, 2018;

Yaden & Griffiths, 2020). The subjective effects of classic psychedelics are characterised by *ego*

dissolution (Preller & Vollenweider, 2018; Stoliker, Egan, Friston, & Razi, 2021), described as the shift

in the sense of self and a loss of boundary between the subjective and the objective world (Dittrich, 1998;

Studerus, Gamma, & Vollenweider, 2010; Vollenweider and Smallridge, 2021; Grof, 1980; Lebedev et

al., 2015). Ego dissolution is a validated construct (Dittrich, 1998; Matthew M. Nour, Evans, Nutt, &

Carhart-Harris, 2016; Studerus et al., 2010) and is thought to involve changes to resting state network

(RSN) activity (Müller, Dolder, Schmidt, Liechti, & Borgwardt, 2018; Preller et al., 2018; Stoliker et al.,

2021).

Functional magnetic resonance imaging (fMRI) investigations indicate activity across the brain is

functionally integrated and forms multiple RSNs (Raichle, 2015). RSNs are associated to mental activity

and the balance of connectivity between them is associated to the direction of conscious attention

(Michael D. Fox et al., 2005; K. J. Friston, 2011; Greicius & Menon, 2004). The default mode network

(DMN) composed of the medial prefrontal cortex (mPFC), posterior cingulate cortex (PCC), and

(bilateral) angular gyrus (AG) is a RSN that activates primarily in the absence of immediate external goal-

directed attention (Raichle et al., 2001). Its function in self-focused thinking and attention suggests its

close relationship to the ego (Andrews-Hanna, Smallwood, & Spreng, 2014). In contrast, the dorsal

attention network (DAN) is a RSN activated during external-focused task-driven attention (Corbetta &

Shulman, 2002; Michael D. Fox & Raichle, 2007) and is usually considered to be composed of the frontal

eye field (FEF) and intraparietal sulcus (IPS) bilaterally. The activity of the DMN and DAN are identified

as anticorrelated and predictably alternate with the inward or outward switching of attention (Michael D.

Fox et al., 2005). The DMN-DAN anticorrelation can be hypothesised to be a mechanism maintaining the

boundary between the subject (observer) and object (observation) that is altered during experiences of

psychedelic ego dissolution.

A third resting state network, the salience network (SN), acts as the switching mechanism coordinating

the direction of attention between internal and external stimuli (Liang et al., 2015; Seeley et al., 2007).

The SN's cardinal regions are the dorsal anterior cingulate cortex (dACC) and anterior insula (AI). Both

the dACC and AI are consistently coactivated across cognitive tasks (Swick, Ashley, & Turken, 2011),

however the dACC is more involved in response selection and conflict monitoring (Ide, Shenoy, Yu, &

Li, 2013; Menon, 2011) while the AI receives greater multimodal sensory input (Averbeck & Seo, 2008)

(Vogt & Pandya, 1987), detects behaviourally relevant stimuli (Menon, 2015) and coordinates the

dynamic interactions of anticorrelated networks (Menon & Uddin, 2010; Sridharan, Levitin, & Menon,

2008).

Coordinated interactions between these networks produce important biopsychological functions. SN

coactivation with the DAN detects bottom-up features in the visual environment that are infrequent or

biologically significant (Egner et al., 2008; Fecteau & Munoz, 2006; Szczepanski, Pinsk, Douglas,

Kastner, & Saalmann, 2013) and also enables the detection of resources relevant to higher-order goals

(Menon, 2015). Furthermore, anticorrelated function between the SN and DMN is a biomarker of

efficient cognition (Chand, Wu, Hajjar, & Qiu, 2017; Putcha, Ross, Cronin-Golomb, Janes, & Stern,

2016). Trauma to the white matter tracts within the SN that connect the rAI and dACC predicts

dysregulated DMN function (Bonnelle et al., 2012). Importantly, abnormality of SN connectivity and its

anticorrelated interactions is indicative of schizophrenia (Manoliu et al., 2014), psychosis (Palaniyappan

& Liddle, 2012; Wotruba et al., 2013) and internalizing disorders (Peterson, Thome, Frewen, & Lanius,

2014) (See Menon 2015 for review of SN associations to psychopathology) (Menon, 2011).

The DMN and SN have been previously investigated in relation to unique senses of self. The SN has been

suggested to be involved in an aspect of the self defined as the basic sense of being rooted within a body,

termed the *minimal* or *embodied* self (Blanke & Metzinger, 2009; Lebedev et al., 2015; Legrand & Ruby,

2009). This association is supported by changes to the SN documented in psychopathology (Liu et al.,

2018), meditation (Doll, Hölzel, Boucard, Wohlschläger, & Sorg, 2015; Ramirez Barrantes et al., 2019)

and psychedelic-induced ego dissolution (Lebedev et al., 2015) (see Supplementary Table 1 for a subset

of psychedelic findings related to networks and regions of interest). The pre-reflective qualities that

define the minimal self have also been suggested as antecedents of the *narrative* aspect of self (Ho,

Preller, & Lenggenhager, 2020; M. M. Nour & Carhart-Harris, 2017). The narrative aspect of self is

believed to be under the control of the DMN and describes self-related mental activity and personal

identity (Metzinger, 2003; Millière, 2017) that strongly parallels the classic Freudian construct of ego

(Carhart-Harris & Friston, 2010; Cieri & Esposito, 2019). These parallels have led to exploratory

investigations of the DMN under psychedelic-induced ego dissolution that indicate a general pattern of

reduced connectivity (Carhart-Harris & Friston, 2010; Cieri & Esposito, 2019; Ruban & Kolodziej, 2018;

Franz X. Vollenweider & Preller, 2020).

Reduced DMN activity has also been identified in meditation and is a feature of improved mental health

(Batchelor, 2008; Judson A. Brewer et al., 2011; Hasenkamp, Wilson-Mendenhall, Duncan, & Barsalou,

2012; Millière, Carhart-Harris, Roseman, Trautwein, & Berkovich-Ohana, 2018). Psychedelics have been

reported to reduce symptoms of patients experiencing internalising mental health disorders (Carhart-

Harris et al., 2018; Nutt & Carhart-Harris, 2020; Ruban & Kolodziej, 2018). Altered self-boundaries may

be important to these therapeutic outcomes (Griffiths et al., 2016; Roseman et al., 2018; Ross et al.,

2016). Nascent alignment between the free energy principle and psychoanalytic processes proposes that

high-order, secondary processes of the ego function to minimise free energy of lower-orders (Cieri &

Esposito, 2019; Karl Friston, 2009; K. Friston, Kilner, & Harrison, 2006; Herzog et al., 2020).

Psychedelic-induced free energy increase in networks such as the DMN is suggested to disinhibit high

order inhibitory selection of bottom-up channels and enable consideration of alternate hypotheses

underlying making sense of the world (Carhart-Harris & Friston, 2019; Safron, 2021; Stoliker et al.,

2021). These mechanisms may broadly account for experiences of ego dissolution (Stoliker, Egan, &

Razi, 2021). An early interpretation of psychedelics noted in *filtration theory* similarly suggests

improvements follow psychedelic reduction of ego-protective defenses (Connolly, 2018; Freud, 1894;

Stoliker et al., 2021; Swanson, 2018). Understanding network changes in ego dissolution may inform how

network interactions relate to clinical outcomes. Meditation is also recognised for its benefits to wellbeing

and resembles a similar trajectory of ego disarmament by seeking cessation of the self (Batchelor, 2008;

Millière et al., 2018). The practice of meditation has demonstrated clinical utility (Goyal et al., 2014) and

can enhance wellbeing by altering the relationship between self and other (Dambrun, 2017). Pertinently, a

form of meditation termed *nondual awareness* meditation reduces the anticorrelation of extrinsic and

intrinsic activated brain regions and produces a subjective experience of dissolving subject-object

boundaries (Josipovic, Dinstein, Weber, & Heeger, 2012). These findings indicate anticorrelation under

control of the SN as a neural mechanism that underlies ego dissolution.

Anticorrelation between the DMN and task positive networks under psilocybin has previously been

investigated with the findings demonstrating reduced anticorrelation when participants experienced ego

dissolution, under psilocybin (intravenous infusion, 2mg dissolved in 10ml) (Carhart-Harris et al., 2012).

However, a similar investigation, under ayahuasca (oral brew, 2.2. mL/kg body weight, containing 0.8

mg/ml DMT and 0.21 mg/ml harmine) failed to identify anticorrelation changes (Palhano-Fontes et al.,

2015). The inability of functional connectivity analyses to determine the direction of connectivity

between networks in these studies suggests the value of adopting mechanistic approaches to determine

changes in effective connectivity of networks under psychedelics. Dynamic Causal Modeling (DCM) is a

Bayesian method of inference based on task based or resting state fMRI time series activity of brain

regions (K. J. Friston, Harrison, & Penny, 2003; Karl J. Friston, Kahan, Biswal, & Razi, 2014). DCM can

disentangle hierarchical RSN and regional interactions by determination of the directionality of

connectivity. DCM has been previously applied to investigate thalamic connectivity to the cortex under

LSD (Preller et al., 2019) that indicated the SN is at the apex of the DMN and DAN triple network

hierarchy (Zhou et al., 2018). The SN's position in this hierarchy and its mediating role to control the

switching of DAN and DMN activity suggests that change to the SN by psychedelics may influence their

patterns of anticorrelated activity. The SN and DMN share associations to aspects of self and the

importance of their connectivity in psychopathology suggests change in their connectivity may be a

mechanism of ego dissolution that underlies a shift in the sense of self (Northoff et al., 2006; Scalabrini et

al., 2020).

Therefore, to understand the neural mechanisms of ego dissolution and inform the biological basis of the

subject-object relationship, the directed changes to these networks under the classic psychedelic lysergic

acid diethylamide (LSD) was investigated. LSD effects were examined across placebo (two weeks apart

from LSD administration), peak effects at 75 minutes and later effects at 300 minutes post LSD

administration using DCM analysis to reveal regional and network connectivity changes. Ego dissolution

(quantified as oceanic boundlessness (OBN)) was measured on the five dimensions of altered states of

consciousness scale (5D-ASC) (Studerus et al., 2010). Based on previous research that associated the

networks under investigation to self and subject-object boundaries, we hypothesised the association

between effective connectivity would change during the peak effects of LSD and ego dissolution. DMN-

DAN was tested for change to subject-object boundaries and SN-DMN was tested for change to the self.

However, the direction of excitatory-inhibitory connectivity change remained exploratory. Additionally,

we measured connectivity of regions composing the networks of interest and the (hierarchical)

connectivity strength between networks.

Results:

Acute Drug Effects

Functional connectivity changes

Functional connectivity across groups demonstrated a fading of the pattern of anticorrelation between

DMN and DAN and SN, from placebo to peak effects of LSD before showing evidence of restoration in

the later effects at 300 minutes. Group level regional functional connectivity matrix across each condition

(placebo, 75 minutes after LSD administration and 300 minutes post LSD administration) is provided in

Supplementary Fig. 1.

Between networks changes in effective connectivity

The first resting-state fMRI scan was done 75 minutes post administration of LSD which is during the

peak effects of LSD. The connectivity strength between networks was computed as the difference

between averaged and unsigned efferent and afferent connection parameters between networks (see

methods for further details) (Zhou et al., 2018). As shown in Fig. 1a), at this time, group level, between

networks, effective connectivity increased from the SN to DMN causing the directed connection to

become excitatory. A similar change was observed in the excitatory connectivity from the DMN to the

DAN resulting in reduced inhibitory connectivity (see Fig. 1c). SN to DMN and DMN to DAN changes

from placebo are greatest during the peak effects at 75 minutes and reduce in the later effects at 300

minutes (see Fig. 1a) and 1b). These changes show reduced hierarchical connectivity strength (i.e.,

increased afferent connections) of the SN and increased hierarchical connectivity strength (i.e., increased

efferent connections) of the DMN and DAN from placebo (SN= -1.16 Hz, DMN = + 0.32 Hz, DAN =

+0.41 Hz, see Supplementary S4 for explanation and quantitative analysis).

Fig 1. Network effective connectivity change under peak effects of LSD. a) Highlighted connections show

changes in effective connectivity compared to placebo signifying peak effect. b) Network effective

connectivity change graphed across placebo, peak effects and later effects. Same data as a) and b) but

plotted as a line graph for better visualization. Values display effect sizes (posterior expectations) of

connections in Hz. All results are for posterior probability > 0.99.

Between regions changes in effective connectivity.

Between regions effective connectivity that underly the SN to DMN changes include increased inhibitory

self-connectivity of the dACC and lFEF and effective connectivity from the lAG to rAG, lFEF to the lIPS,

lIPS to rAI. These regions are highlighted because the change in their effective connectivity show large

effect sizes during the peak effects than in later effects of LSD. Moreover, lAI to dACC and rAG to rAI

effective connectivity show large effect size change from inhibitory in the peak effects of LSD to excitatory

in the later effects (see Fig. 2, panel a) below and Supplementary Fig. 3 for untranslated effective

connectivity matrices across conditions).

Fig 2. Region effective connectivity signifying a) peak and b) lasting effects of LSD. Red lines and lettering

indicate excitatory change; blue lines and lettering indicate inhibitory change. Only connections with

change from placebo to 75 minutes post LSD > 0.1 Hz are included. Panel a) illustrates effective

connectivity with change > 0.1 Hz from peak effects at 75 minutes to the later effects at 300 minutes,

signifying connectivity unique to the peak effects of LSD. See Supplementary Table 2 for posterior

expectations (effect sizes) and credible intervals. Panel b) illustrates effective connectivity with change <

0.1 Hz from peak effects at 75 minutes to the later effects at 300 minutes, signifying connectivity

changes that last over the effects of LSD. See Supplementary Table 3 for posterior expectations (effect

size) and credible intervals. All results are for posterior probability > 0.99.

Lasting effects

Between networks changes in effective connectivity.

LSD effects are also distinguished by changes from placebo that last across time under LSD. Increased
DAN to DMN and decreased DMN to SN effective connectivity at 75 minutes remains evident 300 minutes
after LSD (see Fig. 1, panel a) above, and see Supplementary Fig. 3 for effect size and posterior
probabilities).

238 *Between regions changes in effective connectivity.*

239 Effects lasting across time under LSD are identified by change in lAG and rIPS self-inhibition, lFEF to

240 mPFC inhibition and lIPS to PCC excitation. These connections show large effect sizes from placebo to

241 peak effect that remain to the later effects of LSD (see Fig. 2, panel b) above and Supplementary Fig. 3 for

242 untranslated effective connectivity matrices across conditions).

243
The results demonstrate increased effective connectivity of the DMN to DAN and SN to the DMN during

the peak effects of LSD. These changes correspond with reduced SN hierarchical connectivity strength (i.e.,

increased afferent connections) and coincide with a fading of the functional anticorrelation (see

Supplementary Fig. 1 for functional connectivity results).

Discussion:

This investigation seeks to understand how effective connectivity between anticorrelated large-scale brain

networks is related to ego dissolution. Our analysis reveals between network effective connectivity changes

that occur with a diminution of the pattern of anticorrelation under the peak effects of LSD. Bidirectional

changes in effective connectivity between the DMN and DAN were investigated for their relationship to

subject-object boundaries. We identified reduced inhibition of the DMN to the DAN under the peak effects

of LSD. The reduced inhibition is largely lost in the later effects, when ego dissolution dissipates. This

indicates reduced inhibition of the DMN to DAN as a feature of peak LSD effects that may relate to the

fading of the functional anticorrelation between the two networks. Reduced DMN to DAN inhibition may

also represent an increased transmission and connection of the narrative self to the sense of object.

Moreover, hierarchical organisation and strength of networks were calculated using efferent vs afferent

connections (Zhou et al., 2018). The DMN and DAN showed increased hierarchical connectivity strength

during the peak effects of LSD and segregate from SN, under the peak effects of LSD (See Supplementary

S4). The increase in hierarchical connectivity strength of the DMN and DAN reinforces evidence of their

fading anticorrelation that may contribute to the dissolution of the boundary between the subject and the

object. The opposite effective connectivity from DAN to DMN also displays reduced inhibition under LSD.

However, this change remains over the course of time suggesting it is not a primary mechanism of the

reduced functional anticorrelation between them or ego dissolution.

Network level DMN-DAN changes are supported by the lAG to rAG connectivity. The AG serves

functions in autobiographical memory and bodily awareness (Bréchet, Grivaz, Gauthier, & Blanke, 2018).

AG also operates complex language functions and makes meaning out of visually perceived self-related

words (Yaoi, Osaka, & Osaka, 2015). Excitatory change in the connection from lAG to rAG in placebo to

the peak effects of LSD is greater than in the later effects suggesting the importance of this connection in

the peak effects.

Inclusion of the SN in this analysis enabled measurement of its effective connectivity to the DMN and

DAN under LSD. The change to the coordinated balance of networks, under the control of SN, by LSD

may be an important but overlooked neural mechanism of ego dissolution suggested by the superiority of

SN in this hierarchy of triple networks (Zhou et al., 2018). Previous reports of reduced anticorrelation

(Carhart-Harris et al., 2012), reduced SN integrity (Lebedev et al., 2015) in occurrences of ego

dissolution, and its hypothesised function in basic conscious-awareness also indicate the value of

measuring SN connectivity in the anticorrelation between DMN and DAN. Moreover, ego dissolution has

previously been suggested to involve the breakdown of sub-personal processes underlying the minimal

self, a suggestion that is consistent with Bayesian models of phenomenal selfhood in which the subjective

structure of conscious experience is determined from the optimisation of prediction in perception and

action (Clark, 2013; Karl Friston, 2010; Millière, 2017). The change in the SN effective connectivity

under the peak effects of LSD is our most pronounced finding. SN connectivity to the DMN changes

from inhibitory to excitatory before returning to inhibitory in the later effects. This flip of valence

suggests SN connectivity change to the DMN is mechanistic in the peak effects of LSD and may

indirectly influence the DMN-DAN interactions. The opposite connection, the DMN to SN, shows an

inverse pattern of change from placebo, suggesting reduced DMN influence over the SN which lasts over

time. The SN to DMN connectivity change may therefore be a more likely mechanism of ego dissolution

representing a quietening of narrative self in the peak effects of LSD that reduces in the later effects. This

change in DMN function to minimise free energy and suppress prediction errors that in Freudian terms

may liberate the ego from the reality principle (Carhart-Harris & Friston, 2010; Cieri & Esposito, 2019). SN to

DMN change is accompanied by reduced hierarchical connectivity strength of the SN that is characterised

by increased SN afferent connections and increased DMN efferent connections. The SN has previously

been identified to be hierarchically superior to the DMN and DAN (Zhou et al., 2018). We demonstrate

the divergence of SN and DMN hierarchical strength under the peak effects of LSD shift the hierarchical

order of the SN beneath the DMN.

The changes to region level effective connectivity underlying the SN are demonstrated by dACC self-

connectivity under the peak effects of LSD. The dACC is a central brain hub involved in cognitive

functions, social emotions (Andrews-Hanna et al., 2014; Etkin, Egner, & Kalisch, 2011; Preller et al., 2015)

and performance monitoring (Ham et al., 2014) and has previously been noted in social cognition under

psilocybin (Preller et al., 2016). Enlargement of the ACC has also been reported after regular use of

ayahuasca (Carlos Bouso et al., 2015). We find dACC self-connectivity inhibition and inhibitory

connectivity from the lAI to the dACC under the peak effects of LSD, compared to placebo. In the later

effects dACC inhibition become less inhibitory while the lAI changes to excitatory. The AI is involved in

interoceptive awareness (Chong, Ng, Lee, & Zhou, 2017) and the shift of internal and external perception

(Menon & Uddin, 2010; Sridharan et al., 2008). Damage to SN connectivity predicts dysregulated DMN

function (Menon, 2011) and alteration of SN anticorrelated interactions share association with psychosis

and internalising disorders (Bonnelle et al., 2012). dACC and lAI inhibition at the peak effects of LSD may

indicate the role of SN inhibition that is relevant to ego dissolution and psychedelic assisted therapy.

Connectivity between the DMN and SN is also noted in region changes from the rAG to rAI. This

connection follows the lAI to dACC pattern of inhibition in the peak effects and excitation in the later

effects, suggesting a reduced influence of the DMN to the SN during peak effects, when ego dissolution

occurs that may suggest a shifting sense of self.

The PCC and dACC represent cardinal regions of the DMN and SN respectively and are related to the

narrative and minimal aspects of self (Lebedev et al., 2015; Millière, 2017). The connectivity changes

between them are reported in association to meditation and the mindfulness capacities of meditators (Judson

A. Brewer et al., 2011; Millière et al., 2018; Palhano-Fontes et al., 2015). We provided evidence of the

excitation from the dACC to PCC in our effective connectivity analysis under peak effects (see

Supplementary Fig. 3), which may suggest dACC cognitive control functions influence on self-related

functions of PCC. Interestingly, self-connectivity of the PCC, a hub region of DMN, increases excitation

during the peak effects of LSD. PCC involvement in self-awareness is well documented (J. A. Brewer,

Garrison, & Whitfield-Gabrieli, 2013; Northoff et al., 2006) and activity of this region is commonly

observed to decrease under psychedelics (see Supplementary Table 1).

Other notable regions in our analysis include the FEF and IPS. The FEF operates visual attention

(Thompson, Biscoe, & Sato, 2005) and the IPS serves functions directing attention and memory (Corbetta

& Shulman, 2002; Michael D. Fox, Corbetta, Snyder, Vincent, & Raichle, 2006). The nearby right inferior

parietal lobule (rIPL) is also associated with the sense of agency and self-other discrimination (Chaminade

& Decety, 2002; Lucina Q. Uddin, Molnar-Szakacs, Zaidel, & Iacoboni, 2006). Smaller effect sizes

involving these regions were identified in our analysis (see Supplementary Fig. 3) suggesting more subtle

widespread brain changes occur under the effects of LSD.

Taken together, under the peak effects of LSD, a strong increase in change of effective connectivity, and a

widening of gap between the hierarchical connectivity strength of the SN vs the DMN and DAN shifts in

a direction antithetical to normal hierarchical organisation. This may be said to resemble a collapse – or

flattening – of the hierarchy, during peak effects of LSD, when ego dissolution occurs (See Fig. 1 and

Supplementary S4). Effective connectivity explains this effect as increased excitatory connectivity from

the SN to the DMN. These directed connection changes and the hierarchical strength changes may

function to alter the relationship between the minimal and narrative senses of self. The direction of this

change suggests the influence of the minimal-self traverses over the narrative-self and may relate to the

shift in sense of self described under ego dissolution.

Modelling the connectivity of ego dissolution can provide a means to determine the neural mechanisms that

underlie the perception of inner and outer reality. The present research establishes important steps to

identify the change in network interactions associated with the dichotomy of the subject-object relationship

under LSD. The networks involved in this interaction are important in cognitive function and mental

wellbeing, suggesting that understanding their change due to psychedelics may elucidate mechanisms of

psychedelic clinical therapy. Consideration of practical and theoretical challenges may aid future research

directed to study ego dissolution and the anticorrelation between brain networks.

Our relatively small sample size (n=20) is clearly a limitation. Sample size and processing pipeline strongly

impact the reliability of results. Small sample size may also account for unexpected placebo effective

connectivity in our group of subjects. A second practical limitation is the large variance of participant

subjective responses to a standard dose of LSD (100mg). Averaging the connectivity of participants

experiencing highly variable subjective shifts in consciousness may dilute the effective connectivity

representing LSD's subjective effects. An alternative to increased sample size may be predetermining

participant dose-response and including only participants with high subjective responses in the analysis.

Replicating the investigation of anticorrelated networks with alternate classic psychedelic substances

including psilocybin, ayahuasca and mescaline in addition to LSD will be a worthy direction to validate our

results. Furthermore, comparisons with similar altered forms of consciousness reporting ego change such

as psychosis and meditation may also help discern the function of components in the anticorrelated

networks.

The selection of regions and region coordinates composing the networks of interest require careful attention.

For example, some studies define the DAN as composed by the posterior prefrontal cortex, the inferior

precentral sulcus, the superior occipital gyrus, the middle temporal motion complex, and the superior

parietal lobule (Andrews-Hanna et al., 2014). Similarly, the DMN may be composed using the inferior

parietal lobule (Di & Biswal, 2014) closely situated to the IPS used in the DAN in this analysis. The

composition of regions composing networks can strongly impact results. Furthermore, selection of region

centroids can also affect the results. Previous work using this LSD data set used different coordinates for

the PCC and identified its increased self-inhibition (Preller et al., 2019), although our analysis identified

decreased PCC inhibition. Avoiding this problem is challenging due to anatomical variability between

participants, variability in the method of determining coordinates and the selection of regions composing

networks. However, the standardisation of these considerations across studies is important to improve the

accuracy and reliability of findings across studies.

Future work to extend the current scope of analysis to include connectivity dynamics of additional task

positive networks anticorrelated to the DMN and under control of the SN is required. For example, non-

psychedelic research involving the CEN (central executive network, also known as the frontoparietal central

executive network (FPCEN)) has been conducted to investigate schizophrenia (Leptourgos et al., 2020;

Menon, 2011), meditation (Doll et al., 2015) and control of attention (Corbetta & Shulman, 2002). Its

importance is further signified by investigations of large-scale network interactions and anticorrelations

with the DMN (Andrews-Hanna et al., 2014; Bolton et al., 2020; Menon, 2018). Inclusion of the CEN in

anticorrelation investigations may provide a more complete account of anticorrelated network changes

associated to ego dissolution.

Theoretical challenges include consensus on the nature and situation of ego dissolution within the taxonomy

of psychedelic-induced effects. Psychometric investigations of psychoanalytic based concepts of ego-

functions and measurement of their brain connectivity are recommended future directions. Psychedelics

may contribute to these investigations by altering and allowing measurement of brain mechanisms

hypothesised to underlie psychoanalytic constructs. Integration of psychoanalytic theory and brain

connectivity processes may be important to identifying processes involved in enabling the sense of self and

experiences of ego dissolution.

Conclusions:

The between network balance of anticorrelated activity of specific RSNs depends on subtle adjustments to

network activation. Mental wellbeing and efficient cognition rely on the balance of these interactions. LSD

appears to shift the balance of network activation and diminish the anticorrelation between brain networks

responsible for internal and external modes of perception. Observed increases in effective connectivity from

the DMN to DAN and increased hierarchal strength of these networks under peak effects may account for

the blurring of boundary between subject and object experienced in ego dissolution. Ego dissolution also

involves a shift in the sense of self that may be explained by changes to the interactions between SN and

DMN. These networks are related to distinct aspects of self. Increased effective connectivity from the SN

to the DMN and decreased SN hierarchical connectivity strength emphasise hierarchical flattening and

increased salience functions reaching the DMN under the peak effects of LSD. Future research could model

how changes to these networks effective connectivity may support psychedelic therapeutic outcomes. The

neuroscientific study of psychedelic-induced ego dissolution reminds us that constructs and representations

of self and internal and external reality exist in connectivity dynamics. This intriguing understanding may

inspire future investigations of sentience and consciousness to learn how normal brain function mechanisms

contribute to the subject-object relationship and frame our perspective of reality.

Methods:

Participants

The data analysed in this paper were collected as part of a larger study (registered at ClinicalTrials.gov (NCT02451072)), which is reported in (Preller et al., 2017) and (Preller et al., 2018), and was approved by the Cantonal Ethics Committee of Zurich. 25 subjects (19 males and 6 females; mean age = 25.24 y; SD = 3.72 y; range = 20–34 y) were recruited through advertisements at universities in Zurich, Switzerland. All participants were deemed healthy after screening for medical history, physical examination, blood analysis, and electrocardiography. The Mini-International Neuropsychiatric Interview (MINI-SCID), the Diagnostic and Statistical Manual of Mental Disorders, fourth edition self-rating questionnaire for Axis-II personality disorders (SCID-II), and the Hopkins Symptom Checklist (SCL-90-R) were used to exclude subjects with present or previous psychiatric disorders or a history of major psychiatric disorders in first-degree relatives. Participants were asked to abstain from prescription and illicit drug use two weeks prior to first testing and throughout the duration of the study, and abstain from alcohol use 24 hours prior to testing days. Urine tests were also used to exclude pregnancy. Further exclusion criteria included left-handedness, poor knowledge of the German language, cardiovascular disease, history of head injury or neurological disorder, history of alcohol or illicit drug dependence, MRI exclusion criteria, including claustrophobia, and previous significant adverse reactions to a hallucinogenic drug. All participants provided written informed consent statements in accordance with the declaration of Helsinki before participation in the study. Subjects received written and oral descriptions of the study procedures, as well as details regarding the effects and possible risks of drug treatment.

Design

A double blind, randomized, placebo-controlled, cross-over study was performed. Testing days occurred two weeks apart and participants were orally administered either LSD after pretreatment with 179 mg

Mannitol and 1 mg Aerosil (LSD condition) or 179 mg Mannitol and 1 mg Aerosil after pre-treatment with

179 mg Mannitol and 1 mg Aerosil (placebo condition). Resting state scans (10 minutes each) were taken

75 and 300 minutes following administration. Subjective experience was assessed using the retrospective

5-Dimensions Altered States of Consciousness (5D-ASC) questionnaire. Participants completed the 5D-

ASC 720 min following treatment administration.

MRI Data Acquisition and Preprocessing

MRI data were acquired on a Philips Achieva 3.0T whole-body scanner. A 32-channel receive head coil

and MultiTransmit parallel radio frequency transmission was used. Images were acquired using a whole-

brain gradient-echo planar imaging (EPI) sequence (repetition time, 2,500 ms; echo time, 27 ms; slice

thickness, 3 mm; 45 axial slices; no slice gap; field of view, 240×240 mm^2; in-plane resolution, 3×3 mm;

sensitivity-encoding reduction factor, 2.0). Additionally, high-resolution anatomical images (voxel size, 0.7

$\times 0.7 \times 0.7$ mm^3) were acquired using a standard T1-weighted 3D magnetization prepared rapid-acquisition

gradient echo sequence. The acquired images were analysed using SPM12 (https://www.fil.ion.ucl.ac.uk).

The pre-processing steps of the images consisted of slice-timing correction, realignment, spatial

normalization to the standard EPI template of the Montreal Neurological Institute (MNI), and spatial

smoothing using a Gaussian kernel of 6-mm full-width at half maximum. Head motion was investigated

for any excessive movement, but movement did not exceed 3 mm in any participant.

Extraction of region coordinates across subjects

Group ICA for fMRI Toolbox (GIFT, http://mialab.mrn.org/ software/gift) (Calhoun et al. 2001) was used

to identify the three resting state networks of interest from placebo scans. Pre-processed resting state fMRI

data were spatially sorted into 20 components (Biswal et al. 2010; Shirer et al. 2012; Tsvetanov et al. 2016)

and spatially matched with pre-existing network templates (Shirer et al. 2012).

Networks were composed of cardinal regions constituting a core part of DMN (Andrews-Hanna, Reidler,

Sepulcre, Poulin, & Buckner, 2010; Dixon et al., 2017) which reliably show anticorrelation with the DAN

and SN (Chen, Glover, Greicius, & Chang, 2017; Dixon et al., 2017; M. D. Fox, Zhang, Snyder, &

Raichle, 2009; Fransson, 2005; L. Q. Uddin, Kelly, Biswal, Castellanos, & Milham, 2009) and followed

the selection of regions in a related investigation by Zhou and colleagues (Zhou et al., 2018).

Identification of cardinal nodes within each intrinsic network – averaged across our subjects – was

located using peak RSN activity of clusters within networks (p=.05) visualized using xjView toolbox

(https://www.alivelearn.net/xjview). Associations between peak coordinates and cardinal nodes of

network regions of interest (ROI) were determined by expert visual inspection. The MNI coordinates of

the selected ROIs are listed in Table 4.

A generalized linear model (GLM) was used to regress 6 head motion parameters (3 translation and 3

rotational), white matter and cerebrospinal fluid signals from preprocessed data. One subject was excluded

as no activation was found in one or more regions of interest. We also used global signal regression in our

pre-processing pipeline. The time series for each ROI was computed as the first principal component of the

voxel activity within a 6 mm sphere centred on the ROI coordinates (as listed in Table 4).

Region	MNI coordinates			Network
	x	y	z	
PCC	0	-73	38	DMN
mPFC	0	53	32	DMN
lAG	-54	-61	29	DMN
rAG	57	-58	20	DMN
dACC	0	17	41	SN
lAI	-48	14	-7	SN
rAI	48	14	-7	SN
lFEF	-30	-10	68	DAN
rFEF	30	-7	68	DAN
lIPS/SPL	-42	-43	62	DAN
rIPS/SPL	39	-40	65	DAN

Table 4 Coordinates of regions of interest. The DMN comprised of the posterior cingulate cortex (PCC),

medial prefrontal cortex (mPFC), and left and right angular gyrus (lAG/rAG); the SN comprised of the dorsal

anterior cingulate cortex (dACC), left and right anterior insula (lAI/rAI); and the DAN comprised the left

and right frontal eye field (lFEF/rFEF) and the left and right inferior parietal sulcus/superior parietal lobule

(lIPS(SPL)/rIPS(SPL).

Anticorrelation Functional Connectivity Validation

The functional connectivity matrix between regions of the DMN, SN, and DAN in the placebo condition

was computed for all subjects. Since our arguments rest on the premise of functional anticorrelation

between the resting-state networks (and similar second-order features are used for model estimation by the

following spectral DCM analysis), 4 subjects that did not show evidence of the anticorrelation in the placebo

condition were excluded from further analysis. As such, 20 subjects were left available for the following

DCM analysis.

Specification and Inversion of DCM

A fully-connected DCM was specified using the 11 ROIs defined in Table 4, without any exogenous inputs.

The DCM for each subject was then inverted using spectral DCM (Karl J. Friston et al., 2014; Razi, Kahan,

Rees, & Friston, 2015) to infer the effective connectivity that best explains the observed cross-spectral

density for each subject. This procedure was repeated for each of the three testing conditions. The DCM

fitted the data very well and explained variance was over 85% across all subjects, and averaged 91% (see

Supplementary Fig. 5).

Second Level Analysis Using Parametric Empirical Bayes

The effective connectivity inferred by spectral DCM for each subject are taken to the second (group) level

to test hypotheses about between-subject effects. A General Linear Model (GLM) is employed to

decompose individual differences in effective connectivity into hypothesised group-average connection

strengths plus unexplained noise. Hypotheses on the group-level parameters are tested within the Parametric

Empirical Bayes (PEB) framework (K. J. Friston et al., 2016), where both the expected values and the

covariance of the parameters are taken into account. That is, precise parameter estimates influence the

group-level result more strongly than uncertain estimates, which are down-weighted. Bayesian model

reduction (BMR) is used as an efficient form of Bayesian model selection (K. J. Friston et al., 2016).

Network Level Effective Connectivity and Hierarchical Organization

The expected network-level connectivity was computed as the sum of the expected effective connectivity

values between the corresponding ROIs. Then, following Zhou et. al., 2018 (Zhou et al., 2018) the

hierarchical connectivity strength of each network was obtained by computing the difference between its

averaged efferent and afferent connections (i.e., absolute values, see Supplementary S4). A similar

approach was used for analysing hierarchical projections in the monkey brain (Goulas, Uylings, & Stiers,

2012) and prefrontal cortex hierarchical organization in humans (Nee & D'Esposito, 2016; Zhou et al.,

2018).

References

Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R., & Buckner, R. L. (2010). Functional-
anatomic fractionation of the brain's default network. *Neuron, 65*(4), 550-562.
doi:10.1016/j.neuron.2010.02.005
Andrews-Hanna, J. R., Smallwood, J., & Spreng, R. N. (2014). The default network and self-
generated thought: component processes, dynamic control, and clinical relevance.
Annals of the New York Academy of Sciences, 1316(1), 29-52. doi:10.1111/nyas.12360
Averbeck, B. B., & Seo, M. (2008). The Statistical Neuroanatomy of Frontal Networks in the
Macaque. *PLOS Computational Biology, 4*(4), e1000050.
doi:10.1371/journal.pcbi.1000050
Batchelor, S. (2008). *Buddhism without beliefs : a contemporary guide to awakening*. London:
Bloomsbury.
Blanke, O., & Metzinger, T. (2009). Full-body illusions and minimal phenomenal selfhood.
Trends Cogn Sci, 13(1), 7-13. doi:10.1016/j.tics.2008.10.003
Bolton, T., Wotruba, D., Buechler, R., Theodoridou, A., Michels, L., Kollias, S., . . . Van De Ville, D.
(2020). Triple Network Model Dynamically Revisited: Lower Salience Network State
Switching in Pre-psychosis. *Frontiers in Physiology, 11*. doi:10.3389/fphys.2020.00066
Bonnelle, V., Ham, T. E., Leech, R., Kinnunen, K. M., Mehta, M. A., Greenwood, R. J., & Sharp, D.
J. (2012). Salience network integrity predicts default mode network function after

traumatic brain injury. *Proc Natl Acad Sci U S A, 109*(12), 4690-4695.
doi:10.1073/pnas.1113455109
Bréchet, L., Grivaz, P., Gauthier, B., & Blanke, O. (2018). Common Recruitment of Angular Gyrus
in Episodic Autobiographical Memory and Bodily Self-Consciousness. *Frontiers in*
Behavioral Neuroscience, 12(270). doi:10.3389/fnbeh.2018.00270
Brewer, J. A., Garrison, K. A., & Whitfield-Gabrieli, S. (2013). What about the "Self" is Processed
in the Posterior Cingulate Cortex? *Front Hum Neurosci, 7*, 647.
doi:10.3389/fnhum.2013.00647
Brewer, J. A., Worhunsky, P. D., Gray, J. R., Tang, Y.-Y., Weber, J., & Kober, H. (2011). Meditation
experience is associated with differences in default mode network activity and
connectivity. *Proceedings of the National Academy of Sciences, 108*(50), 20254-20259.
doi:10.1073/pnas.1112029108
Carhart-Harris, R. L. (2018). How do psychedelics work? *Current Opinion in Psychiatry*, 1.
doi:10.1097/YCO.0000000000000467
Carhart-Harris, R. L., Bolstridge, M., Day, C. M. J., Rucker, J., Watts, R., Erritzoe, D. E., . . . Nutt,
D. J. (2018). Psilocybin with psychological support for treatment-resistant depression:
six-month follow-up. *Psychopharmacology, 235*(2), 399-408. doi:10.1007/s00213-017-
4771-x
Carhart-Harris, R. L., & Friston, K. J. (2010). The default-mode, ego-functions and free-energy: a
neurobiological account of Freudian ideas. *Brain, 133*(Pt 4), 1265-1283.
doi:10.1093/brain/awq010
Carhart-Harris, R. L., & Friston, K. J. (2019). REBUS and the Anarchic Brain: Toward a Unified
Model of the Brain Action of Psychedelics. *Pharmacological Reviews, 71*(3), 316-344.
doi:10.1124/pr.118.017160
Carhart-Harris, R. L., Leech, R., Erritzoe, D., Williams, T. M., Stone, J. M., Evans, J., . . . Nutt, D. J.
(2012). Functional Connectivity Measures After Psilocybin Inform a Novel Hypothesis of
Early Psychosis. *Schizophrenia Bulletin, 39*(6), 1343-1351. doi:10.1093/schbul/sbs117
Carlos Bouso, J., Palhano-Fontes, F., Rodriguez-Fornells, A., Ribeiro, S., Sanches, R., Crippa, J. A.
S., . . . Riba, J. (2015). Long-term use of psychedelic drugs is associated with differences
in brain structure and personality in humans. *European Neuropsychopharmacology,*
25(4), 483-492. doi:10.1016/j.euroneuro.2015.01.008
Chaminade, T., & Decety, J. (2002). Leader or follower? Involvement of the inferior parietal
lobule in agency. *Neuroreport, 13*(15), 1975-1978. doi:10.1097/00001756-200210280-
00029
Chand, G. B., Wu, J., Hajjar, I., & Qiu, D. (2017). Interactions of the Salience Network and Its
Subsystems with the Default-Mode and the Central-Executive Networks in Normal Aging
and Mild Cognitive Impairment. *Brain Connectivity, 7*(7), 401-412.
doi:10.1089/brain.2017.0509
Chen, J. E., Glover, G. H., Greicius, M. D., & Chang, C. (2017). Dissociated patterns of anti-
correlations with dorsal and ventral default-mode networks at rest. *Human Brain*
Mapping, 38(5), 2454-2465. doi:https://doi.org/10.1002/hbm.23532
Chong, J. S. X., Ng, G. J. P., Lee, S. C., & Zhou, J. (2017). Salience network connectivity in the
insula is associated with individual differences in interoceptive accuracy. *Brain Structure*
and Function, 222(4), 1635-1644. doi:10.1007/s00429-016-1297-7

Cieri, F., & Esposito, R. (2019). Psychoanalysis and Neuroscience: The Bridge Between Mind and
Brain. *Frontiers in Psychology, 10*(1983). doi:10.3389/fpsyg.2019.01983
Clark, A. (2013). Whatever next? Predictive brains, situated agents, and the future of cognitive
science. *Behav Brain Sci, 36*(3), 181-204. doi:10.1017/s0140525x12000477
Connolly, P. (2018). Expected Free Energy Formalizes Conflict Underlying Defense in Freudian
Psychoanalysis. *Front Psychol, 9*, 1264. doi:10.3389/fpsyg.2018.01264
Corbetta, M., & Shulman, G. (2002). Control of Goal-Directed and Stimulus-Driven Attention in
the Brain. *Nature reviews. Neuroscience, 3*, 201-215. doi:10.1038/nrn755
Dambrun, M. (2017). Self-centeredness and selflessness: happiness correlates and mediating
psychological processes. *PeerJ, 5*, e3306. doi:10.7717/peerj.3306
Di, X., & Biswal, B. B. (2014). Identifying the default mode network structure using dynamic
causal modeling on resting-state functional magnetic resonance imaging. *NeuroImage,*
86, 53-59. doi:https://doi.org/10.1016/j.neuroimage.2013.07.071
Dittrich, A. (1998). The standardized psychometric assessment of altered states of
consciousness (ASCs) in humans. *Pharmacopsychiatry, 31 Suppl 2*, 80-84. doi:10.1055/s-
2007-979351
Dixon, M. L., Andrews-Hanna, J. R., Spreng, R. N., Irving, Z. C., Mills, C., Girn, M., & Christoff, K.
(2017). Interactions between the default network and dorsal attention network vary
across default subsystems, time, and cognitive states. *NeuroImage, 147*, 632-649.
doi:10.1016/j.neuroimage.2016.12.073
Doll, A., Hölzel, B. K., Boucard, C. C., Wohlschläger, A. M., & Sorg, C. (2015). Mindfulness is
associated with intrinsic functional connectivity between default mode and salience
networks. *Frontiers in Human Neuroscience, 9*, 461. doi:10.3389/fnhum.2015.00461
Egner, T., Monti, J. M. P., Trittschuh, E. H., Wieneke, C. A., Hirsch, J., & Mesulam, M. M. (2008).
Neural Integration of Top-Down Spatial and Feature-Based Information in Visual Search.
The Journal of Neuroscience, 28(24), 6141. doi:10.1523/JNEUROSCI.1262-08.2008
Etkin, A., Egner, T., & Kalisch, R. (2011). Emotional processing in anterior cingulate and medial
prefrontal cortex. *Trends in Cognitive Sciences, 15*(2), 85-93.
doi:10.1016/j.tics.2010.11.004
Fecteau, J. H., & Munoz, D. P. (2006). Salience, relevance, and firing: a priority map for target
selection. *Trends Cogn Sci, 10*(8), 382-390. doi:10.1016/j.tics.2006.06.011
Fox, M. D., Corbetta, M., Snyder, A. Z., Vincent, J. L., & Raichle, M. E. (2006). Spontaneous
neuronal activity distinguishes human dorsal and ventral attention systems. *Proceedings*
of the National Academy of Sciences, 103(26), 10046-10051.
doi:10.1073/pnas.0604187103
Fox, M. D., & Raichle, M. E. (2007). Spontaneous fluctuations in brain activity observed with
functional magnetic resonance imaging. *Nature Reviews Neuroscience, 8*(9), 700-711.
doi:10.1038/nrn2201
Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005).
The human brain is intrinsically organized into dynamic, anticorrelated functional
networks. *Proceedings of the National Academy of Sciences of the United States of*
America, 102(27), 9673. doi:10.1073/pnas.0504136102

Fox, M. D., Zhang, D., Snyder, A. Z., & Raichle, M. E. (2009). The global signal and observed
anticorrelated resting state brain networks. *J Neurophysiol, 101*(6), 3270-3283.
doi:10.1152/jn.90777.2008
Fransson, P. (2005). Spontaneous low-frequency BOLD signal fluctuations: an fMRI investigation
of the resting-state default mode of brain function hypothesis. *Hum Brain Mapp, 26*(1),
15-29. doi:10.1002/hbm.20113
Freud, S. (1894). *The neuro-psychoses of defence* (Vol. 1). London: Hogarth Press.
Friston, K. (2009). The free-energy principle: a rough guide to the brain? *Trends in Cognitive
Sciences, 13*(7), 293-301. doi:https://doi.org/10.1016/j.tics.2009.04.005
Friston, K. (2010). The free-energy principle: a unified brain theory? *Nature Reviews
Neuroscience, 11*, 127. doi:10.1038/nrn2787
https://www.nature.com/articles/nrn2787#supplementary-information
Friston, K., Kilner, J., & Harrison, L. (2006). A free energy principle for the brain. *J Physiol Paris,
100*(1-3), 70-87. doi:10.1016/j.jphysparis.2006.10.001
Friston, K. J. (2011). Functional and effective connectivity: a review. *Brain Connect, 1*(1), 13-36.
doi:10.1089/brain.2011.0008
Friston, K. J., Harrison, L., & Penny, W. (2003). Dynamic causal modelling. *NeuroImage, 19*(4),
1273-1302. doi:10.1016/s1053-8119(03)00202-7
Friston, K. J., Kahan, J., Biswal, B., & Razi, A. (2014). A DCM for resting state fMRI. *NeuroImage,
94*, 396-407. doi:https://doi.org/10.1016/j.neuroimage.2013.12.009
Friston, K. J., Litvak, V., Oswal, A., Razi, A., Stephan, K. E., van Wijk, B. C. M., . . . Zeidman, P.
(2016). Bayesian model reduction and empirical Bayes for group (DCM) studies.
NeuroImage, 128, 413-431. doi:10.1016/j.neuroimage.2015.11.015
Goulas, A., Uylings, H. B. M., & Stiers, P. (2012). Mapping the Hierarchical Layout of the
Structural Network of the Macaque Prefrontal Cortex. *Cerebral Cortex, 24*(5), 1178-
1194. doi:10.1093/cercor/bhs399
Goyal, M., Singh, S., Sibinga, E. M., Gould, N. F., Rowland-Seymour, A., Sharma, R., . . .
Haythornthwaite, J. A. (2014). Meditation programs for psychological stress and well-
being: a systematic review and meta-analysis. *JAMA Intern Med, 174*(3), 357-368.
doi:10.1001/jamainternmed.2013.13018
Greicius, M. D., & Menon, V. (2004). Default-Mode Activity during a Passive Sensory Task:
Uncoupled from Deactivation but Impacting Activation. *J Cogn Neurosci, 16*(9), 1484-
1492. doi:10.1162/0898929042568532
Griffiths, R. R., Johnson, M. W., Carducci, M. A., Umbricht, A., Richards, W. A., Richards, B. D., . .
. Klinedinst, M. A. (2016). Psilocybin produces substantial and sustained decreases in
depression and anxiety in patients with life-threatening cancer: A randomized double-
blind trial. *Journal of Psychopharmacology, 30*(12), 1181-1197.
doi:10.1177/0269881116675513
Grof, S. (1980). *LSD Psychotherapy*. Alameda, CA: Hunter House Publishers.
Ham, T. E., Bonnelle, V., Hellyer, P., Jilka, S., Robertson, I. H., Leech, R., & Sharp, D. J. (2014).
The neural basis of impaired self-awareness after traumatic brain injury. *Brain : a journal
of neurology, 137*(Pt 2), 586-597. doi:10.1093/brain/awt350
Hasenkamp, W., Wilson-Mendenhall, C. D., Duncan, E., & Barsalou, L. W. (2012). Mind
wandering and attention during focused meditation: a fine-grained temporal analysis of

fluctuating cognitive states. *NeuroImage, 59*(1), 750-760.
doi:10.1016/j.neuroimage.2011.07.008

Herzog, R., Mediano, P. A. M., Rosas, F. E., Carhart-Harris, R., Perl, Y. S., Tagliazucchi, E., &
Cofre, R. (2020). A mechanistic model of the neural entropy increase elicited by
psychedelic drugs. *Scientific Reports, 10*(1), 17725. doi:10.1038/s41598-020-74060-6

Ho, J. T., Preller, K. H., & Lenggenhager, B. (2020). Neuropharmacological modulation of the
aberrant bodily self through psychedelics. *Neurosci Biobehav Rev, 108*, 526-541.
doi:10.1016/j.neubiorev.2019.12.006

Ide, J. S., Shenoy, P., Yu, A. J., & Li, C. S. (2013). Bayesian prediction and evaluation in the
anterior cingulate cortex. *The Journal of Neuroscience, 33*(5), 2039-2047.
doi:10.1523/jneurosci.2201-12.2013

Johnson, M., Richards, W., & Griffiths, R. (2008). Human hallucinogen research: guidelines for
safety. In (Vol. 22, pp. 603-620). London, England.

Josipovic, Z., Dinstein, I., Weber, J., & Heeger, D. (2012). Influence of meditation on anti-
correlated networks in the brain. *Frontiers in Human Neuroscience, 5*, 183. Retrieved
from https://www.frontiersin.org/article/10.3389/fnhum.2011.00183
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3250078/pdf/fnhum-05-00183.pdf

Lebedev, A. V., Lövdén, M., Rosenthal, G., Feilding, A., Nutt, D. J., & Carhart-Harris, R. L. (2015).
Finding the self by losing the self: Neural correlates of ego-dissolution under psilocybin.
Human Brain Mapping, 36(8), 3137-3153. doi:10.1002/hbm.22833

Legrand, D., & Ruby, P. (2009). What is self-specific? Theoretical investigation and critical
review of neuroimaging results [American Psychological Association
doi:10.1037/a0014172]. Retrieved

Leptourgos, P., Fortier-Davy, M., Carhart-Harris, R., Corlett, P., Dupuis, D., Halberstadt, A., . . .
Jardri, R. (2020). Hallucinations Under Psychedelics and in the Schizophrenia Spectrum:
An Interdisciplinary and Multiscale Comparison. *Schizophrenia Bulletin*.
doi:10.1093/schbul/sbaa117

Liang, X., He, Y., Salmeron, B. J., Gu, H., Stein, E. A., & Yang, Y. (2015). Interactions between the
salience and default-mode networks are disrupted in cocaine addiction. *The Journal of
neuroscience : the official journal of the Society for Neuroscience, 35*(21), 8081-8090.
doi:10.1523/JNEUROSCI.3188-14.2015

Manoliu, A., Riedl, V., Zherdin, A., Mühlau, M., Schwerthöffer, D., Scherr, M., . . . Sorg, C. (2014).
Aberrant dependence of default mode/central executive network interactions on
anterior insular salience network activity in schizophrenia. *Schizophr Bull, 40*(2), 428-
437. doi:10.1093/schbul/sbt037

Menon, V. (2011). Large-scale brain networks and psychopathology: a unifying triple network
model. *Trends Cogn Sci, 15*(10), 483-506. doi:10.1016/j.tics.2011.08.003

Menon, V. (2015). *Salience Network* (Vol. 2): Elsevier.

Menon, V. (2018). The Triple Network Model, Insight, and Large-Scale Brain Organization in
Autism. *Biological Psychiatry, 84*(4), 236-238. doi:10.1016/j.biopsych.2018.06.012

Menon, V., & Uddin, L. Q. (2010). Saliency, switching, attention and control: a network model of
insula function. *Brain structure & function, 214*(5-6), 655-667. doi:10.1007/s00429-010-
0262-0

Metzinger, T. (2003). *Being no one: The self-model theory of subjectivity*. Cambridge, MA, US: MIT Press.

Millière, R. (2017). Looking for the Self: Phenomenology, Neurophysiology and Philosophical Significance of Drug-induced Ego Dissolution. *Frontiers in Human Neuroscience, 11*(245). doi:10.3389/fnhum.2017.00245

Millière, R., Carhart-Harris, R. L., Roseman, L., Trautwein, F.-M., & Berkovich-Ohana, A. (2018). Psychedelics, Meditation, and Self-Consciousness. *Frontiers in Psychology, 9*(1475). doi:10.3389/fpsyg.2018.01475

Müller, F., Dolder, P. C., Schmidt, A., Liechti, M. E., & Borgwardt, S. (2018). Altered network hub connectivity after acute LSD administration. *NeuroImage. Clinical, 18*, 694-701. doi:10.1016/j.nicl.2018.03.005

Nee, D. E., & D'Esposito, M. (2016). The hierarchical organization of the lateral prefrontal cortex. *eLife, 5*. doi:10.7554/eLife.12112

Northoff, G., Heinzel, A., de Greck, M., Bermpohl, F., Dobrowolny, H., & Panksepp, J. (2006). Self-referential processing in our brain--a meta-analysis of imaging studies on the self. *NeuroImage, 31*(1), 440-457. doi:10.1016/j.neuroimage.2005.12.002

Nour, M. M., & Carhart-Harris, R. L. (2017). Psychedelics and the science of self-experience. *Br J Psychiatry, 210*(3), 177-179. doi:10.1192/bjp.bp.116.194738

Nour, M. M., Evans, L., Nutt, D., & Carhart-Harris, R. L. (2016). Ego-Dissolution and Psychedelics: Validation of the Ego-Dissolution Inventory (EDI). *Frontiers in Human Neuroscience, 10*. doi:10.3389/fnhum.2016.00269

Nutt, D., & Carhart-Harris, R. (2020). The Current Status of Psychedelics in Psychiatry. *JAMA Psychiatry*. doi:10.1001/jamapsychiatry.2020.2171

Palaniyappan, L., & Liddle, P. F. (2012). Does the salience network play a cardinal role in psychosis? An emerging hypothesis of insular dysfunction. *J Psychiatry Neurosci, 37*(1), 17-27. doi:10.1503/jpn.100176

Palhano-Fontes, F., Andrade, K. C., Tofoli, L. F., Santos, A. C., Crippa, J. A., Hallak, J. E., . . . de Araujo, D. B. (2015). The psychedelic state induced by ayahuasca modulates the activity and connectivity of the default mode network. *Plos One, 10*(2), e0118143. doi:10.1371/journal.pone.0118143

Peterson, A., Thome, J., Frewen, P., & Lanius, R. A. (2014). Resting-State Neuroimaging Studies: A New Way of Identifying Differences and Similarities among the Anxiety Disorders? *The Canadian Journal of Psychiatry, 59*(6), 294-300. doi:10.1177/070674371405900602

Preller, K. H., Burt, J. B., Ji, J. L., Schleifer, C. H., Adkinson, B. D., Stämpfli, P., . . . Anticevic, A. (2018). Changes in global and thalamic brain connectivity in LSD-induced altered states of consciousness are attributable to the 5-HT2A receptor. *eLife, 7*, e35082. doi:10.7554/eLife.35082

Preller, K. H., Herdener, M., Pokorny, T., Planzer, A., Kraehenmann, R., Stampfli, P., . . . Vollenweider, F. X. (2017). The Fabric of Meaning and Subjective Effects in LSD-Induced States Depend on Serotonin 2A Receptor Activation. *Curr Biol, 27*(3), 451-457. doi:10.1016/j.cub.2016.12.030

Preller, K. H., Pokorny, T., Hock, A., Kraehenmann, R., Stampfli, P., Seifritz, E., . . . Vollenweider, F. X. (2016). Effects of serotonin 2A/1A receptor stimulation on social exclusion processing. *Proc Natl Acad Sci U S A, 113*(18), 5119-5124. doi:10.1073/pnas.1524187113

Preller, K. H., Pokorny, T., Kraehenmann, R., Scheidegger, M., Dziobek, I., Staempfli, P., & Vollenweider, F. X. (2015). The 5-HT2A/1A agonist psilocybin reduces social pain and enhances empathy in healthy volunteers. *European Neuropsychopharmacology, 25*, S301-S301. doi:10.1016/s0924-977x(15)30364-3

Preller, K. H., Razi, A., Zeidman, P., Stämpfli, P., Friston, K. J., & Vollenweider, F. X. (2019). Effective connectivity changes in LSD-induced altered states of consciousness in humans. *Proceedings of the National Academy of Sciences, 116*(7), 2743-2748. doi:10.1073/pnas.1815129116

Preller, K. H., & Vollenweider, F. X. (2018). Phenomenology, Structure, and Dynamic of Psychedelic States. *Curr Top Behav Neurosci, 36*, 221-256. doi:10.1007/7854_2016_459

Putcha, D., Ross, R. S., Cronin-Golomb, A., Janes, A. C., & Stern, C. E. (2016). Salience and Default Mode Network Coupling Predicts Cognition in Aging and Parkinson's Disease. *Journal of the International Neuropsychological Society : JINS, 22*(2), 205-215. doi:10.1017/S1355617715000892

Raichle, M. E. (2015). The brain's default mode network. *Annu Rev Neurosci, 38*, 433-447. doi:10.1146/annurev-neuro-071013-014030

Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L. (2001). A default mode of brain function. *Proc Natl Acad Sci U S A, 98*(2), 676-682. doi:10.1073/pnas.98.2.676

Ramirez Barrantes, R., Arancibia, M., Stojanova, J., Aspé-Sánchez, M., Córdova, C., & Henríquez, R. A. (2019). Default Mode Network, Meditation, and Age-Associated Brain Changes: What Can We Learn from the Impact of Mental Training on Well-Being as a Psychotherapeutic Approach? *Neural Plasticity, 2019*, 1-15. doi:10.1155/2019/7067592

Razi, A., Kahan, J., Rees, G., & Friston, K. J. (2015). Construct validation of a DCM for resting state fMRI. *NeuroImage, 106*, 1-14. doi:https://doi.org/10.1016/j.neuroimage.2014.11.027

Roseman, L., Nutt, D. J., & Carhart-Harris, R. L. (2018). Quality of Acute Psychedelic Experience Predicts Therapeutic Efficacy of Psilocybin for Treatment-Resistant Depression. *Frontiers in Pharmacology, 8*, 974. doi:10.3389/fphar.2017.00974

Ross, S., Bossis, A., Guss, J., Agin-Liebes, G., Malone, T., Cohen, B., . . . Schmidt, B. L. (2016). Rapid and sustained symptom reduction following psilocybin treatment for anxiety and depression in patients with life-threatening cancer: a randomized controlled trial. *Journal of Psychopharmacology, 30*(12), 1165-1180. doi:10.1177/0269881116675512

Ruban, A., & Kolodziej, A. A. (2018). Changes in default-mode network activity and functional connectivity as an indicator of psychedelic-assisted psychotherapy effectiveness. *Neuropsychiatria I Neuropsychologia, 13*(3), 91-97. doi:10.5114/nan.2018.81249

Safron, A. (2020, November 30). On the Varieties of Conscious Experiences: Altered Beliefs Under Psychedelics (ALBUS). https://doi.org/10.31234/osf.io/zqh4b

Scalabrini, A., Vai, B., Poletti, S., Damiani, S., Mucci, C., Colombo, C., . . . Northoff, G. (2020). All roads lead to the default-mode network-global source of DMN abnormalities in major depressive disorder. *Neuropsychopharmacology, 45*(12), 2058-2069. doi:10.1038/s41386-020-0785-x

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., . . . Greicius, M. D. (2007). Dissociable intrinsic connectivity networks for salience processing and

executive control. *The Journal of Neuroscience, 27*(9), 2349-2356.
doi:10.1523/jneurosci.5587-06.2007

Sridharan, D., Levitin, D. J., & Menon, V. (2008). A critical role for the right fronto-insular cortex
in switching between central-executive and default-mode networks. *Proceedings of the*
National Academy of Sciences, 105(34), 12569. doi:10.1073/pnas.0800005105

Stoliker, D., Egan, G., Friston, K., & Razi, A. (2021). Neural Mechanisms and Psychology of
Psychedelic Ego Dissolution. In: PsyArXiv.

Stoliker, D., Egan, G., & Razi, A. (2021). Reduced precision underwrites ego dissolution and
therapeutic outcomes under psychedelics. In: PsyArXiv.

Studerus, E., Gamma, A., & Vollenweider, F. X. (2010). Psychometric evaluation of the altered
states of consciousness rating scale (OAV). *Plos One, 5*(8), e12412-e12412.
doi:10.1371/journal.pone.0012412

Swanson, L. R. (2018). Unifying Theories of Psychedelic Drug Effects. *Frontiers in Pharmacology,*
9(172). doi:10.3389/fphar.2018.00172

Swick, D., Ashley, V., & Turken, U. (2011). Are the neural correlates of stopping and not going
identical? Quantitative meta-analysis of two response inhibition tasks. *NeuroImage,*
56(3), 1655-1665. doi:10.1016/j.neuroimage.2011.02.070

Szczepanski, S. M., Pinsk, M. A., Douglas, M. M., Kastner, S., & Saalmann, Y. B. (2013).
Functional and structural architecture of the human dorsal frontoparietal attention
network. *Proceedings of the National Academy of Sciences, 110*(39), 15806.
doi:10.1073/pnas.1313903110

Thompson, K. G., Biscoe, K. L., & Sato, T. R. (2005). Neuronal basis of covert spatial attention in
the frontal eye field. *The Journal of Neuroscience, 25*(41), 9479-9487.
doi:10.1523/jneurosci.0741-05.2005

Uddin, L. Q., Kelly, A. M., Biswal, B. B., Castellanos, F. X., & Milham, M. P. (2009). Functional
connectivity of default mode network components: correlation, anticorrelation, and
causality. *Hum Brain Mapp, 30*(2), 625-637. doi:10.1002/hbm.20531

Uddin, L. Q., Molnar-Szakacs, I., Zaidel, E., & Iacoboni, M. (2006). rTMS to the right inferior
parietal lobule disrupts self–other discrimination. *Social cognitive and affective*
neuroscience, 1(1), 65-71. doi:10.1093/scan/nsl003

Vogt, B. A., & Pandya, D. N. (1987). Cingulate cortex of the rhesus monkey: II. Cortical afferents.
Journal of Comparative Neurology, 262(2), 271-289.
doi:https://doi.org/10.1002/cne.902620208

Vollenweider, F. X. (1998). Advances and pathophysiological models of hallucinogenic drug
actions in humans: a preamble to schizophrenia research. *Pharmacopsychiatry, 31 Suppl*
2, 92-103. doi:10.1055/s-2007-979353

Vollenweider, F. X., Leenders, K. L., Scharfetter, C., Maguire, P., Stadelmann, O., & Angst, J.
(1997). Positron emission tomography and fluorodeoxyglucose studies of metabolic
hyperfrontality and psychopathology in the psilocybin model of psychosis.
Neuropsychopharmacology, 16(5), 357-372. doi:10.1016/s0893-133x(96)00246-1

Vollenweider, F. X., & Preller, K. H. (2020). Psychedelic drugs: neurobiology and potential for
treatment of psychiatric disorders. *Nature Reviews Neuroscience, 21*(11), 611-624.
doi:10.1038/s41583-020-0367-2

Wotruba, D., Michels, L., Buechler, R., Metzler, S., Theodoridou, A., Gerstenberg, M., . . .
Heekeren, K. (2013). Aberrant Coupling Within and Across the Default Mode, Task-
Positive, and Salience Network in Subjects at Risk for Psychosis. *Schizophrenia Bulletin,*
40. doi:10.1093/schbul/sbt161
Yaden, D. B., & Griffiths, R. R. (2020). The Subjective Effects of Psychedelics Are Necessary for
Their Enduring Therapeutic Effects. *ACS Pharmacology & Translational Science*.
doi:10.1021/acsptsci.0c00194
Yaoi, K., Osaka, M., & Osaka, N. (2015). Neural correlates of the self-reference effect: evidence
from evaluation and recognition processes. *Frontiers in Human Neuroscience, 9*(383).
doi:10.3389/fnhum.2015.00383
Zhou, Y., Friston, K. J., Zeidman, P., Chen, J., Li, S., & Razi, A. (2018). The Hierarchical
Organization of the Default, Dorsal Attention and Salience Networks in Adolescents and
Young Adults. *Cereb Cortex, 28*(2), 726-737. doi:10.1093/cercor/bhx307

861