Gene-Environment Pathways to Cognitive Intelligence and Psychotic-Like Experiences in Children

Junghoon Park[1]*, Eunji Lee[2]*, Gyeongcheol Cho[3], Heungsun Hwang[3], Bogyeom Kim[2], Gakyung Kim[4], Yoonjung Yoonie Joo[2,5,6†], Jiook Cha[1,2,4†]

1. Interdisciplinary Program in Artificial Intelligence, College of Engineering, Seoul National University, Seoul, South Korea

2. Department of Psychology, College of Social Sciences, Seoul National University, Seoul, South Korea

3. Department of Psychology, McGill University, Montréal, Canada

4. Department of Brain and Cognitive Sciences, College of Natural Sciences, Seoul National University, Seoul, South Korea

5. Department of Digital Health, Samsung Advanced Institute for Health Sciences & Technology (SAIHST), Sungkyunkwan University, Seoul, South Korea

6. Samsung Medical Center, Seoul, South Korea

* Shared co-first authorship; † shared corresponding authorship

Correspondence to:

Yoonjung Yoonie Joo, PhD

    Irwon-ro 115, Gangnamgu, Seoul, 06335, South Korea

    yoonjungjoo@skku.edu

Jiook Cha, PhD

    Gwanak-ro 1, Building 16, Suite M512, Gwanakgu, Seoul, 08826, South Korea

    connectome@snu.ac.kr

Abstract

Background

In children, psychotic-like experiences (PLEs) are related to risk of psychosis, schizophrenia, and other mental disorders. Maladaptive cognitive functioning is a well-known risk factor and early marker for psychosis, schizophrenia, and other mental disorders. Since cognitive functioning is linked to various genetic and environmental factors during development, we hypothesize that it mediates the effects of those factors on childhood PLEs. Using large longitudinal data, we tested the relationships of genetic and environmental factors (such as familial and neighborhood environment) with cognitive intelligence and their relationships with current and future PLEs in children.

Methods

To estimate associations against potential confounding bias, we leveraged large-scale multimodal data of 6,602 children (aged 9-10 years old; 47.15% females; 5,211 European-ancestry) from the Adolescent Brain and Cognitive Development Study. Linear mixed model and a novel structural equation modeling (SEM) method that allows estimation of both components and factors were used to estimate the joint effects of cognitive phenotypes polygenic scores (PGSs), familial and neighborhood socioeconomic status (SES), and supportive environment on NIH Toolbox cognitive intelligence and PLEs. We adjusted for ethnicity (genetically defined), schizophrenia PGS, and additionally unobserved confounders (using computational confound modeling).

Results

We identified that lower cognitive intelligence and higher PLEs correlated significantly with several genetic and environmental variables: i.e., lower PGSs for cognitive phenotypes, lower familial SES, lower neighborhood SES, lower supportive parenting behavior, and lower positive school environment. In SEM, lower cognitive intelligence significantly mediated the genetic and environmental influences on higher PLEs (Indirect effects of PGS: β range=-0.0355~ -0.0274; Family SES: β range=-0.0429~ -0.0331; Neighborhood SES: β range=0.0126~ 0.0164; Positive Environment: β range=-0.0039~ -

0.003). Supportive parenting and a positive school environment had the largest total impact on PLEs (β range=-0.152~ -0.1316) than any other genetic or environmental factors.

Conclusions

Our results reveal the role of genetic and environmental factors on children's PLEs via its negative impact on cognitive intelligence. Our findings have policy implications in that improving the school and family environment and promoting local economic development might be a way to enhance cognitive and mental health in children.

Keywords

Cognitive intelligence; psychotic-like experiences; genetic-environmental pathway; structural equation modeling; confounding adjustments

Key Points

- While existing research shows the association between cognitive decline and PLEs, the genetic and environmental pathways to cognitive intelligence and psychotic risk in children remain unclear.

- We identified the significant role of genetic and environmental factors (family, neighborhood, and school) on children's PLEs via a negative impact on cognitive intelligence.

- Leveraging large samples with multimodal longitudinal data and advanced computational modeling for adjustment of observed/unobserved confounding bias, our results underscore the importance of incorporating socioeconomic policies into children's cognitive and mental health programs.

Introduction

Childhood is the critical developmental period in human life. Cognitive intelligence and mental health in this period significantly impact key life outcomes at later ages, including academic performance, economic productivity, physical health, intelligence, and psychopathology (Shonkoff, 2012; Walker et al., 2021). Literature shows the significant impact of social adversities on cognitive ability and mental health in early childhood. Lower family socioeconomic status (SES), particularly household income, is linked to lower neurocognitive ability and higher risk of psychopathology in childhood (Hair et al., 2015; Noble et al., 2015; Peverill et al., 2021; Tomasi & Volkow, 2021; Weissman et al., 2018).

Additional to family SES, the importance of neighborhood social environment on children's neurocognitive ability has been also emphasized (Gard et al., 2021; Tooley et al., 2020). Adverse neighborhood environment, such as the percent of families below poverty line, low education levels, and exposure to violence, is associated with lower cognitive performance and a greater risk for psychosis in children (Butler et al., 2018; Karcher, Schiffman, et al., 2021; Rakesh et al., 2021; Taylor et al., 2020). Conversely, as protective factors against familial and neighborhood socioeconomic challenges, supportive parenting (Brody et al., 2017; Brody et al., 2019; Holmes et al., 2018; Luby et al., 2012; Luby et al., 2016) and positive school environment (Gard et al., 2021; Piccolo et al., 2019; Rakesh et al., 2021) have been highlighted to improve child cognition and mental health.

Psychotic-like experiences (PLEs), which are prevalent in childhood, indicate the risk of psychosis (van der Steen et al., 2019; Van Os & Reininghaus, 2016). Although they are not a direct precursor of schizophrenia, children reporting PLEs in ages of 9-11 years are at higher risk of psychotic disorders in adulthood (Kelleher & Cannon, 2011; Poulton et al., 2000). PLEs also point towards the potential for other psychopathologies including mood, anxiety, and substance disorders (van der Steen et al., 2019), are linked to deficits in cognitive intelligence (Cannon et al., 2002; Kelleher & Cannon, 2011) and show a stronger association with environmental risk factors during childhood than other internalizing/externalizing symptoms (Karcher, Schiffman, et al., 2021).

Maladaptive cognitive intelligence may act as a mediator for the effects of genetic and environmental risks on the manifestation of psychotic symptoms (Cannon et al., 2000; Keefe et al., 2006; Reichenberg et al., 2005). Abnormal neurodevelopment, influenced by genetic factors, combined with disrupted cognitive processes resulting from socioenvironmental adversity, may eventually give rise to the positive symptoms of

schizophrenia, relevant to PLEs (Garety et al., 2001; Howes & Murray, 2014). Family studies show a decline in cognitive intelligence preceding psychotic symptoms is related to genetic risk (Cosway et al., 2000; Curtis et al., 2001). In more recent studies these associations have led to the model positing that cognitive intelligence mediates the genetic risk for psychopathology and PLEs (Karcher, Paul, et al., 2021; Pat et al., 2021).

To minimize potential bias in the estimates of environmental effects, it is crucial to adjust for genetic confounding (Sariaslan et al., 2016), given the substantial genetic influence on intelligence (Bouchard & McGue, 1981; Deary et al., 2006; Plomin & Spinath, 2004) and PLEs (Bentall & Fernyhough, 2008; Maxwell et al., 2023). Recent advances in genetics have led to the development of the polygenic score (PGS) approach: a computational method to estimate the genetic loading for a complex trait using statistical associations of each single-nucleotide polymorphism (SNP) identified by genome-wide association studies (GWAS) (Choi et al., 2020). Particularly, PGS for two related but distinct phenotypes—cognitive performance and educational attainment—holds significant importance. As the two most frequently used proxies of cognitive intelligence in genetic studies, PGSs for cognitive performance and educational attainment are positively correlated with intelligence, educational attainment, income, self-rated health, and height (Judd et al., 2020; Lee et al., 2018; Okbay et al., 2022; Selzam et al., 2019). Furthermore, the PGS for educational attainment is associated with a wide range of biological and social outcomes, including brain morphometry (Judd et al., 2020; Karcher, Paul, et al., 2021), psychopathologies such as PLEs, autism, depression, Alzheimer's disease, neuroticism (Karcher, Paul, et al., 2021; Okbay et al., 2022), cognitive decline (Joo et al., 2022; Karcher, Paul, et al., 2021; Ritchie et al., 2020), BMI, time spent watching television, geographic residence (Abdellaoui et al., 2022), and wealth inequality (Barth et al., 2020). Similar to the terms used in prior research, we will collectively refer to these two PGSs of focus as 'cognitive phenotypes PGSs' throughout this paper (Joo et al., 2022; Okbay et al., 2022; Selzam et al., 2019). An important gap in the literature is the lack of integrated assessment of the effects of genetic and environmental factors at multiple levels (e.g., familial vs neighborhood) to dissect the genetic and environmental effects underlying abnormal cognitive intelligence and the PLEs. Addressing this with large multimodal data will allow for a more complete understanding of the factors related to the development of PLEs.

In this study, we systematically explore the longitudinal trajectories of genetic and environmental influences on PLEs, mediated through cognitive intelligence. Toward this goal, we firstly assess the associations of cognitive phenotype PGSs, family and

neighborhood SES, and supportive environment with children's cognitive intelligence and longitudinally measured PLEs. To maintain robustness of our assessment, we employed statistical and computational approaches to carefully consider potential confounding. We then test the mediating effect of cognitive intelligence on the relationship between genetic and environmental factors and PLEs. Our investigation traces these effects from the baseline and through the 1-year and 2-year follow-ups, providing a nuanced understanding on the role of cognitive phenotype polygenic scores, family socioeconomic status, neighborhood socioeconomic status, and positive family and school environments in shaping PLEs in children aged 9-10 years.

Materials and Methods

Study Participants

We used the multi-modal genetic and environmental data of 11,878 preadolescent children aged 9-10 years old collected from 21 research sites of the Adolescent Brain Cognitive Development (ABCD) Study, one of the largest longitudinal studies for children's neurodevelopment in the United States. We analyzed the baseline, first year, and second year follow-up datasets included in ABCD Release 4.0, downloaded on January 25, 2022. After k-nearest neighbor imputation of missing values of covariates (categorical variables: sex, genetic ancestry, marital status of the caregiver, ABCD research sites; continuous variables: age, BMI, family history of psychiatric disorders; 4.67% of total observations imputed) using the R package VIM (Kowarik & Templ, 2016), we removed participants with missing data on study variables (missing genotype: N=3,260; follow-up observations: N=1,180; neighborhood information: N=694; cognitive intelligence tests: N=126; PLEs: N=5; positive environment: N=11). The final samples included 6,602 multiethnic children, which comprised 890 of African ancestry (13.48%), 91 of East Asian ancestry (1.38%), 5,211 of European ancestry (78.93%), 229 of Native American ancestry (3.47%), and 181 not specified (2.74%).

Data

NIH Toolbox Cognitive Performance

Children's neurocognitive abilities were assessed using the NIH Toolbox Cognitive Battery, which has seven cognitive instruments for examining executive function, episodic memory, language abilities, processing speed, working memory, and attention (Thompson et al., 2019). We utilized baseline observations of uncorrected composite scores of fluid intelligence (Dimensional Change Card Sort Task, Flanker Test, Picture Sequence Memory Test, List Sorting Working Memory Test), crystallized intelligence (Picture Vocabulary Task and Oral Reading Recognition Test), and total intelligence (all seven instruments) provided in the ABCD Study dataset.

Psychotic-Like Experiences (PLEs)

Baseline and 1- and 2-year follow-up of PLEs were measured using the children's responses to the Prodromal Questionnaire-Brief Child Version. In line with previous research (Karcher et al., 2018; Karcher et al., 2020; Karcher, Schiffman, et al., 2021), we computed *Total Score* and *Distress Score*, each indicating the number of psychotic symptoms and levels of total distress. Considering self-reports and parent-reports of psychopathology may differ (Achenbach, 2006), we additionally used parent-rated PLEs derived from four items of the Child Behavior Checklist according to previous studies (Karcher et al., 2018; Karcher et al., 2020; Karcher, Schiffman, et al., 2021). Self-reported PLEs and parent-reported PLEs had significant positive correlation (Pearson's correlation of baseline year: $r = 0.095$~0.0989, p<0.0001; 1-year follow-up: $r = 0.1322$~0.1327, p<0.0001; 2-year follow-up: $r = 0.1569$~0.1632, p<0.0001).

Polygenic Scores (PGS)

To investigate the aggregated effect of genetic components, we estimated PGS of two representative cognitive phenotypes for each participant: educational attainment (EA) and cognitive performance (CP) (Choi et al., 2020). We used the summary statistics released from a GWAS (Lee et al., 2018) of European-descent individuals for EA (n=1,131,881) and CP (n=257,841). EA was measured as the years of schooling; CP, measured as the respondent's score on cognitive ability assessments of general cognitive function and verbal-numerical reasoning, was assessed in participants from the COGENT consortium and the UK Biobank. To construct PGS of schizophrenia for sensitivity analyses, we used the summary statistics from the multiple GWAS of European sample (n=65,967; Ruderfer et al. (2018)) and East Asian sample (n=58,140; Lam et al. (2019)). We applied PRS-CSx, a high-dimensional Bayesian regression framework that improves cross-population prediction via continuous shrinkage prior to SNP effect sizes (Ge et al., 2019) (For details, see **Appendix S1**). The two PGSs for cognitive phenotypes had a positive significant correlation (Pearson's correlation: r =0.4331, p<0.0001).

Family-, Neighborhood-, and School-level Environment

We assessed children's family-level SES with family income, parental education, and family's financial adversity based on parent self-reporting (Karcher et al., 2020; Taylor et al.,

2020; Tomasi & Volkow, 2021). Higher family income and parental education and lower

family's financial adversity denote a higher family SES.

Neighborhood-level SES was assessed using the *Area Deprivation Index* (ADI), the

percentage of individuals below -125% of the poverty level (henceforth "*poverty*"), and *years*

of residence, which were associated with PLEs in prior research (Karcher, Schiffman, et al.,

2021). Higher values of ADI and *poverty* and fewer *years of residence* indicate a lower

neighborhood SES.

Based on existing literature (Karcher, Schiffman, et al., 2021; Rakesh et al., 2021),

we measured the level of positive parenting behavior and positive school environment to

assess the effect of positive family and school environment on each individual.

Statistical Modeling

In this study, we employ linear mixed models and a novel structural equation

modelling (SEM) method to examine the longitudinal trajectories of genetic and

environmental influences on PLEs mediated by cognitive intelligence. We specifically

investigate the mediating role of cognitive intelligence within the impacts of cognitive

phenotype polygenic scores (PGSs), high family socioeconomic status (SES), low

neighborhood SES, and positive family and school environments on PLEs. These influences

are examined across three periods of PLEs observations: baseline, 1-year follow-up, and 2-

year follow-up (**Figure 1**).

Linear Mixed Models

Using linear mixed models to minimize potential population stratification from

different household environments and geographical locations (Choi et al., 2020), we analyzed

the genetic and environmental effects on cognitive intelligence and PLEs. Key variables of

interest were PGS, family SES, neighborhood SES, and positive environment. To avoid

multicollinearity, CP and EA PGSs were used separately in two other models. As a random

factor, the variable indicating ABCD research sites was used. Variables within each model

had no signs of multicollinearity (Fox & Monette, 1992) (generalized variance inflation

factor < 2.0 for every variable in all models). The child's sex, age, genetic ancestry, BMI,

marital status of the caregiver, and family history of psychiatric disorders were included as

covariates in the model (Karcher, Schiffman, et al., 2021). All continuous variables were

standardized (z-scaled) beforehand to get standardized estimates, and the analyses were

conducted with the *lme4* package in R version 4.1.2. Throughout this paper, threshold for

statistical significance was set at $p<0.05$, with correction for multiple comparisons using the

false discovery rate. 95% bootstrapped confidence intervals were obtained with 5,000

iterations.

Path Modeling

To test the plausibility of whether cognitive intelligence may mediate the association

between genetic and environmental factors and PLEs, we used an up-to-date SEM method,

integrated generalized structured component analysis (IGSCA) (Hwang, Cho, Jung, et al.,

2021). This approach is suited to our study using the multi-modal genetic and environmental

variables in that it estimates models with both factors and components as statistical proxies

for the constructs.

Standard SEM using latent factors (i.e., indirectly measured indicators that explain

the covariance among observed variables) to represent indicators such as PGS or family SES

relies on the assumption that observed variables within each construct share a common

underlying factor. If this assumption is violated, standard SEM cannot effectively control for

estimation biases. The IGSCA method addresses this limitation by allowing for the use of

composite indicators (i.e., components)—defined as a weighted sum of observed variables—

as constructs in the model, more effectively controlling bias in estimation compared to the

standard SEM. During estimation, the IGSCA determines weights of each observed variable

in such a way as to maximize the variances of all endogenous indicators and components.

We assessed path-analytic relationships among the six key constructs: cognitive

phenotypes PGSs, family SES, neighborhood SES, positive family and school environment,

general intelligence, and PLEs. Considering that the observed variables of the PGSs, family

SES, neighborhood SES, positive family and school environment, and PLEs are evaluated as

a composite index by prior research, the IGSCA method can mitigate bias more effectively by

representing these constructs as components. Notably, investigations carried out by Judd et al.

(2020) and Martin et al. (2015) utilized composite indicators to examine the genetic influence

on educational attainment and ADHD. Moreover, socioenvironmental influences are often

treated as composite indicators as highlighted in Judd et al. (2020). When considering the

psychosis continuum, studies like that of van Os et al. (2009) postulate that psychotic

disorders are likely underpinned by a multiplicity of background factors rather than a single

common factor. This perspective is substantiated by a multitude of prior research that deploys

composite indices for the measurement of psychotic symptoms. For these reasons, we

statistically represented these constructs as the weighted sums of their observed variables or

components (Cho et al., 2021). On the other hand, we represented general intelligence as a

common factor that determines the underlying covariance pattern of fluid and crystallized

intelligence, based on the classical g theory of intelligence (Jensen, 1998; Spearman, 1904).

The IGSCA model included the same covariates used in the linear mixed model as

well as the ABCD research site as an additional covariate. We applied GSCA Pro 1.1

(Hwang, Cho, & Choo, 2021) to fit the IGSCA model to the data and checked the model's

goodness-of-fit index (GFI) (Jöreskog & Sörbom, 1986), standardized root mean square

residual (SRMR), and total variance of all indicators and components explained (FIT) to

assess its overall goodness-of-fit. Ranging from 0 to 1, a larger FIT value indicates more

variance of all variables is explained by the specified model (e.g., FIT=0.50 denotes that the

model explains 50% of the total variance of all variables) (Hwang, Cho, & Choo, 2021). The

rules-of-thumb cutoff criteria in IGSCA is GFI≥0.93 and SRMR≤0.08 for an acceptable fit

(Cho et al., 2020). Finally, we conducted conditional process analyses to investigate further

the indirect and total effects of the constructs in the model. As a trade-off for obtaining robust

nonparametric estimates without distributional assumptions for normality, the IGSCA method

does not return exact p-values (Hwang, Cho, Jung, et al., 2021). As a reasonable alternative,

we obtained 95% confidence intervals based on 5,000 bootstrap samples to test the statistical

significance of parameter estimates.

Sensitivity Analyses

To ensure robustness of the main analyses results, we conducted multiple sensitivity

analyses. As the European-descent-based GWAS was used for constructing PGS, we reran the

main analyses using participants of European ancestry (n=5,211) to adjust for ethnic

confounding. Next, we tested effects of gene x environment interactions on cognitive

intelligence and PLEs, respectively. We also tested the effects of cognitive phenotypes PGS

adjusting for schizophrenia PGS, given the association of schizophrenia PGS and cognitive

deficit in psychosis patients (Shafee et al., 2018) and individuals at-risk of psychosis (He et

al., 2021). Lastly, we adjusted for unobserved confounding bias in the linear mixed model, using a recently developed framework for causal inference based on null treatments approach (Miao et al., 2022). Designed to discern causal effects from multiple treatment variables within non-randomized, observational data, the null treatments approach hinges on the assumption that no fewer than half of the confounded treatments exert no causal influence on the outcome. It circumvents the need for prior knowledge regarding which treatments are null and eliminates the necessity for independence among treatments. Given our model's inclusion of numerous treatment variables with shared variances due to the presence of unobserved confounders (Abdellaoui et al., 2022; Okbay et al., 2022)—including cognitive phenotypes PGS, family and neighborhood SES, positive family and school environments— we opted to employ this method.

Results

Demographics

Table 1 presents the demographic characteristics of the final samples. For multiethnic subjects (main analyses, n=6,602), 47.15% were female, and the parents of 70.21% were married. In European ancestry samples (sensitivity analyses, n=5,211), 46.71% were female, and the parents of 77.47% were married. Children of European ancestry showed significantly different marital status (p<0.0001), lower BMI (p<0.0001) and family history of psychiatric disorders (p<0.0001) compared to children of other genetic ancestries. Our linear mixed model and IGSCA analyses were adjusted using sex, age, marital status, BMI, family history of psychiatric disorders, and ABCD research sites as covariates.

Genetic influence on cognitive phenotypes correlates positively with intelligence and negatively with PLEs

As shown in **Figure 2**, higher PGSs of cognitive capacity phenotypes correlated significantly with higher intelligence (CP PGS: β=0.1113~0.1793; EA PGS: β=0.0699~0.1567). While CP PGS was associated only with lower baseline year *Distress Score* PLEs (β=-0.0323), EA PGS was associated with lower baseline year and follow-up

PLEs of all measures (baseline: β=-0.0518~-0.0519; 1-year: β=-0.0423~-0.043; 2-year: β=-0.036~ -0.0463). No significant correlations were found between CP PGS and follow-up PLEs (p>0.05). The effects of EA PGS were larger on baseline year PLEs than follow-up PLEs. The effect sizes of EA PGS on children's PLEs were larger than those of CP PGS (EA PGS: β=-0.036~-0.0519; CP PGS: β=-0.0323). (**Table S1**).

Family and neighborhood SES correlates positively with intelligence and negatively with PLEs

Parental education associated positively with all types of intelligence (β=0.0699~0.1745) and negatively with baseline year *Total* and *Distress Score* PLEs (β=-0.0528~-0.043), 1-year follow-up PLEs (β=-0.0538~-0.0449), and 2-year follow-up PLEs (β=-0.0459~-0.0389). Family income correlated positively with intelligence (β=0.0723~0.1365) and negatively with 2-year follow-up parent-rated PLEs (β=-0.0503~ -0.0502). Family's financial disadvantage correlated negatively with baseline year parent-rated PLEs (β=0.0726~0.0728), 1-year follow-up PLEs of all types (β=0.0307~0.0577), and 2-year follow-up PLEs of all types (β=0.0461~0.0581).

The ADI correlated significantly negatively with all types of intelligence (β=-0.0684~-0.054). Additionally, a higher ADI correlated significantly with higher baseline year PLEs (β=0.0587~0.0914), 1-year follow-up PLEs (β=0.0523~0.0613), and 2-year follow-up PLEs (β=0.0397~0.0449).

We found no significant associations of *poverty* with any of the target variables (p> 0.05). *Years of residence* correlated significantly with crystallized intelligence (β=0.035~0.0372) and baseline year *Total Score* PLEs (β=-0.029~-0.0273). (**Table S1**).

Positive family and school environments correlate positively with intelligence and negatively with the influence of PLEs

Positive parenting behaviors showed significant negative correlations with baseline year PLEs (β=-0.0702~-0.0419), 1-year follow-up PLEs (β=-0.0588~-0.0397), and 2-year follow-up PLEs (β=-0.0623~-0.0356) (**Figure 2**). Positive school environment was associated

positively with total intelligence (β=0.0353~0.0397) and fluid intelligence
(β=0.0514~0.0545) and negatively with all three measures of baseline year PLEs (β=-
0.1193~-0.0468), 1-year follow-up PLEs (β=-0.1078~-0.04), and 2-year follow-up PLEs (β=-
0.1068~-0.0586) (**Table S1**).

Structural Equation Modeling-Integrated Generalized Structured Component Analysis

The IGSCA model showed that intelligence mediated the effects of genes and
environments on the risk for psychosis (PLEs) (**Figure 3** and **Table 2**). Estimated factor
loadings of latent factor variable and weights of component variables are presented in **Table
S2**. The model showed a good model fit with a GFI of 0.9735, SRMR of 0.0359, and FIT
value of 0.4912 (Cho et al., 2020). Intelligence was under significant direct influences of the
cognitive phenotypes PGS (β=0.2427), family SES (β=0.2932), neighborhood SES (β=-
0.1121), and positive environment (β=0.0268). Family SES and positive environment had
significant negative direct effects on PLEs of all years. Cognitive phenotypes PGS and
neighborhood SES showed no significant direct effects on any of the PLEs (p>0.05).
Intelligence significantly mediated the effects of the PGS, family and neighborhood SES, and
positive environment on PLEs of all years: the PGS (baseline year: β=-0.035; 1-year: β=-
0.0355; 2-year: β=-0.0274), family SES (baseline year: β=-0.0423; 1-year: β=--0.0429; 2-
year: β=-0.0331), neighborhood SES (baseline year: β=0.0162; 1-year: β=0.0164; 2-year:
β=0.0126), and positive environment (baseline year: β=-0.0039; 1-year: β=-0.0039; 2-year:
β=-0.003).

For all observed years, positive environment had largest total effects on PLEs
(baseline year: β=-0.152; 1-year: β=-0.1316; 2-year: β=-0.1364), followed by family SES
(baseline year: β=-0.1216; 1-year: β=-0.1119; 2-year: β=-0.1164), neighborhood SES
(baseline year: β=0.0504; 1-year: β=0.0329; 2-year: β=0.0192), and PGS (baseline year: β=-
0.0498; 1-year: β=-0.036; 2-year: β=-0.0365). The total effects of each indicator on PLEs
were significant except for those of neighborhood SES (p>0.05).

Sensitivity Analyses

As sensitivity analyses, we reran our main analyses with adjustment for ethnic

confounding, schizophrenia PGS, and unobserved confounding, respectively. Results of linear mixed model and IGSCA analyses were consistent (**Table S3-S7**). See **Appendix S2** for detailed results.

Discussion

This study investigated the relationships of the genetic and environmental influences on the development of intelligence and the PLEs in youth, leveraging genetic data from the large epidemiological samples and a multi-level environmental (socioeconomic) data. Our results support the model that genetic factors (PGS for cognitive phenotypes), socioeconomic conditions, and family and school environments may influence cognitive intelligence in children, and this impact may lead to the individual variability of the current and future PLEs in children. Our analysis with integrated data shows the contributions of genetic and environmental factors, respectively, to cognitive and mental wellness in children, and further provides policy implications to improve them.

Our SEM shows that cognitive intelligence mediates the environmental and genetic influence on the current and future PLEs. The environmental factors (family SES, neighborhood SES, and positive parenting and schooling) and PGS of cognitive phenotypes exhibit significant indirect effects via cognitive intelligence on PLEs. Prior research identifying the mediation of cognitive intelligence focused on either genetic (Karcher, Paul, et al., 2021) or environmental factors (Lewis et al., 2020) alone. Studies with older clinical samples have shown that cognitive deficit may be a precursor for the onset of psychotic disorders (Eastvold et al., 2007; Fett et al., 2020; Vorstman et al., 2015). Our study advances this by demonstrating the integrated effects of genetic and environmental factors on PLEs through the cognitive intelligence in 9-11 years old children. Such comprehensive analysis contributes to assessing the relative importance of various factors influencing children's cognition and mental health, and it can aid future studies designed for identifying health policy implications. Considering the directions and magnitudes of the effects, though the effects of PGS remain significant, aggregated effects of environmental factors account for much greater degrees on PLEs.

Our results of cognitive intelligence mediating the genetic and environmental effects on PLEs may be related to several potential mechanisms. Children raised in higher family SES may have sufficient nutrition and cognitive stimulants, whereas children living in

deprived neighborhoods may be exposed to higher rates of crime, air pollution, and substance abuse (Lewis et al., 2020; Marshall et al., 2020; Taylor et al., 2020; Tomasi & Volkow, 2021). Environmental enrichment may be associated with longer periods of neural plasticity (e.g. myelination, maturation of brain circuitry), leading to higher cognitive ability and lower risk of mental disorders like PLEs (Tooley et al., 2021). This may be further linked to the cognitive reserve theory. The theory suggests that genetic influence for cognitive phenotypes and environmental enrichment promotes more efficient, flexible brain networks, which may lead to greater resilience against psychopathology (Stern, 2009). Indeed, prior clinical studies show the linkage between cognitive reserve and psychosis (Amoretti et al., 2018; Leeson et al., 2011).

Our results indicate that genetic influences on cognitive phenotypes are significantly linked to PLEs. PGSs for CP and EA were strongly correlated with PLEs (baseline year, 1-year follow-up, and 2-year follow-up). These associations were robust after adjustment for schizophrenia PGS, ethnic confounding and unobserved confounders. Cognitive phenotypes PGS generally show higher predictive performance than PGS of any other traits (Lee et al., 2018; Okbay et al., 2022; Plomin & von Stumm, 2018). Genetic variants associated with CP and EA are related to complex traits across the life span, including neuroticism, depressive symptoms, smoking in adulthood, cognitive decline at a later age (Joo et al., 2022), risk for Alzheimer's disease (Lee et al., 2018; Okbay et al., 2022), brain volume, area, and thickness, as well as psychotic disorders (Karcher, Paul, et al., 2021). Prior gene expression studies suggest that polygenic signals for schizophrenia, bipolar disorder, and EA are significantly enriched in the central nervous system, particularly the cerebellum (Finucane et al., 2018). Our findings emphasize the importance of cognitive phenotypes PGS as a biomarker which not only implicates cognitive traits but also exhibits genetic overlap with the PLEs.

The differing magnitudes of the PGS impact between EA and CP warrant attention. The effects of the EA PGS on the PLEs of all years were 160.68%~371.67% larger than those of CP PGS. This discrepancy may result from that the larger sample size of EA GWAS than that of CP GWAS. Alternatively, the discrepancies in effect sizes may suggest different genetic compositions between EA and CP. Recent literature documents that more than half of the polygenic signal for EA is related to noncognitive and social skills required for successful educational attainment (Demange et al., 2021), whereas CP may rather be linked to cognitive skills. This observation also supports the well-established relationships of the EA PGS with socioeconomic and life-course outcomes (e.g., social mobility (Belsky et al., 2018), voter turnout (Aarøe et al., 2020), BMI, income, time spent watching television, geographic

residence (Abdellaoui et al., 2022), and wealth inequality (Barth et al., 2020), which may be

influenced by unobserved environmental factors (Young et al., 2019). In our analysis, the

utilization of two PGSs a more comprehensive evaluation, contributing to an estimation of

the genetic and environmental factors that attempted to minimize confounding bias.

Furthermore, the significant effects of cognitive phenotypes PGS on cognitive

intelligence (β=0.0699~0.1793) remained robust and similar in magnitude after adjusting for

genetic ancestry (β=0.0754~0.1866) and other (unobserved) confounding (β=0.0546~0.1776).

As we controlled for family-, neighborhood-, and school-level environmental factors and

unobserved confounders, our results may be interpreted as significant genetic influences on

individual's cognitive intelligence. This interpretation is supported by a recent study (Isungset

et al., 2022): Despite of the socioeconomic differences in Norway (a typical social democratic

welfare state) and the US (a typical liberal welfare state), the magnitudes of genetic influence

on cognitive intelligence were similar (Norway: β=0.18; US: β=0.17). Cognitive phenotypes

PGS is an important genetic factor across the nations and societies. Therefore, analyses

omitting the genetic influence may be subject to overestimation of the socioeconomic impact

(Plomin & von Stumm, 2018; Sariaslan et al., 2016).

This study shows that a high SES and positive environment, particularly positive

parenting behavior and school environment, is associated with higher intelligence and a lower

risk for PLEs in children. While prior research has emphasized the dominant role of family

SES (e.g. family income) (Tomasi & Volkow, 2021), our SEM analyses (IGSCA) showed

that positive environmental factors such as supportive parenting and schooling have a greater

impact on children's PLEs. Specifically, the effect sizes were the highest in supportive family

and school environment, followed by family and neighborhood SES. Even after adjusting for

genetic ancestry and unobserved confounders, the strong associations of positive parenting

and schooling with higher intelligence and fewer PLEs remained significant. These findings

suggest that interventions that target positive family and school environments may be

particularly effective. Recent research supports this notion, showing that interventions that

promote supportive parenting and inclusive school environments can improve neurocognitive

ability, academic performance, and decrease risk behaviors such as drinking and emotional

eating (Brody et al., 2017; Brody et al., 2019; Holmes et al., 2018).

Moreover, our results showed that positive parenting and schooling in baseline year

were associated not only with baseline year PLEs but also with PLEs 1-2 years later. This is

in line with prior research showing that intervention focused on parenting behavior and

school environment have long-lasting positive effects that extend into adulthood and even

across generations (Cunha & Heckman, 2007; Hill et al., 2020).

While policy implications in observational studies like ours might be limited, our

findings show the importance of comprehensive approaches considering the entire ecosystem

of children's lives—including residential, family, and school environment—for future

research aimed to enhance children's intelligence and mental health. When we combine the

total effect sizes of neighborhood and family SES, as well as positive school environment and

parenting behavior ($\sum |\beta| = 0.2718 \sim 0.3242$), they considerably surpass the total effect

sizes of cognitive phenotypes PGSs ($|\beta| = 0.0359 \sim 0.0502$). It has been suggested that a

holistic and quantitative approach that takes into account the comprehensive ecosystem of

family, school, and residential environments may ensure policy effects and efficient use of

resources (Cree et al., 2018; Garner & Yogman, 2021; Shonkoff, 2012). For example, the

Health Impact in 5 Years Initiative of the US Centers for Disease Control and Prevention

(CDC, 2018) includes 14 evidence-based interventions, such as providing school-based

prevention programs, public transportation, home improvement loans, and earned income tax

credits, to tackle the social determinants of public health. Our study strengthens the idea that

an interdisciplinary science-driven, coordinated approach to intervening in the select

environmental factors may allow practical improvements in child development, particularly

in those who are at a disadvantage.

Our study has some limitations. First, due to data availability constraints in the

ABCD study, we only utilized baseline observations for NIH Toolbox cognitive

intelligence, and we could not test whether PLEs might be a mediator of intelligence.

Second, the generalizability of our findings may be limited since most of the participants

included in our analysis are from European ancestry. Although the ABCD Study aimed to

achieve its representativeness by recruiting from an array of school systems located around

each of the 21 research sites, chosen for their diversity in geography, demographics, and

socioeconomic status, it is not fully representative of the US population (Compton et al.,

2019). Third, the duration of the follow-up period utilized in this study is relatively short

(1-year and 2-year follow-up), which may limit the interpretability of our findings for

understanding cognitive and psychiatric development during later childhood. Future

research could potentially benefit from employing longer follow-up periods, as more

follow-up observations are being collected in the ABCD Study. Fourth, while we used a

wide range of statistical methods to adjust for confounding bias from observed and

unobserved variables (e.g., genetic ancestry), we did not account for other types of

potential bias such as sample selection bias. Fifth, despite a number of causal inference methods used in this study, the ABCD Study is a non-randomized dataset. Given the observational nature of the ABCD Study, interpreting our results as actual causality requires more caution. Finally, we did not include all important environmental variables, such as air pollution (Marshall et al., 2020) and social capital (Krabbendam, 2005), which are not collected in the ABCD study.

In conclusion, our study provides potential pathways of genetic factors of cognitive phenotypes and environmental factors of family, school, and neighborhood to cognitive and mental wellness in children. Our findings underscore the importance of a comprehensive approach that considers both biological and socioeconomic features in promoting young children's cognitive ability and mental health. Given the importance of child development, it requires joint efforts from multiple disciplines.

Funding and Disclosure

This work was supported by the New Faculty Startup Fund from Seoul National University, a Research grant from Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, South Korea (No. 2021R1I1A1A01054995, 2021R1C1C1006503, 2021K1A3A1A2103751212, 2021M3E5D2A01022515, RS-2023-00266787, RS-2023-00265406, RS-2023-00250759), Creative-Pioneering Researchers Program through Seoul National University (No. 200-20230058), Semi-Supervised Learning Research Grant by SAMSUNG(No.A0426-20220118), Identify the network of brain preparation steps for concentration Research Grant by LooxidLabs (No.339-20230001), and Institute of Information & communications Technology Planning & Evaluation (IITP) grant funded by the Korea government (MSIT) [NO.2021-0-01343, Artificial Intelligence Graduate School Program (Seoul National University)]. The authors have nothing to disclose.

Author Contributions

JP, EL, and GC conducted the main analyses. HH, YYJ, and JC supervised the study. JP, EL preprocessed data. JP, EL, GC, HH performed statistical analyses. EL, BK, GK, and

565 YYJ constructed PGS. JP, EL, GC, YYJ, and JC drafted and critically reviewed the

566 manuscript. All authors read and approved the final draft.

Data Availability

All codes used in this study can be found at

https://github.com/Transconnectome/Intell_PLE_Pathway. Due to ABCD Study's policy in

data sharing, we provide a synthetic dataset instead of real observations. This synthetic

dataset was generated with conditional GAN (Xu et al., 2019) to imitate the data structure of

our final study samples. After automatic hyperparameter optimization with Optuna (Akiba et

al., 2019), the synthetic dataset showed overall quality score of 84.15%. Note that analyses

results from the synthetic data may not be 100% identical to the results presented in this

paper, due to the differences between synthetic vs original dataset.

References

Aarøe, L., Appadurai, V., Hansen, K. M., Schork, A. J., Werge, T., Mors, O., Børglum, A. D., Hougaard, D. M., Nordentoft, M., & Mortensen, P. B. (2020). Genetic predictors of educational attainment and intelligence test performance predict voter turnout. *Nature Human Behaviour*, 1-11. DOI: https://doi.org/10.1038/s41562-020-00952-2

Abdellaoui, A., Dolan, C. V., Verweij, K. J. H., & Nivard, M. G. (2022). Gene–environment correlations across geographic regions affect genome-wide association studies. *Nature Genetics*. DOI: 10.1038/s41588-022-01158-0

Achenbach, T. M. (2006). As Others See Us. *Current Directions in Psychological Science*, *15*(2), 94-98. DOI: 10.1111/j.0963-7214.2006.00414.x

Akiba, T., Sano, S., Yanase, T., Ohta, T., & Koyama, M. (2019, 25 July). Optuna: A Next-Generation Hyperparameter Optimization Framework.*KDD '19* Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining, Anchorage, AK, USA.

Akimova, E. T., Breen, R., Brazel, D. M., & Mills, M. C. (2021). Gene-environment dependencies lead to collider bias in models with polygenic scores. *Scientific Reports*, *11*(1). DOI: 10.1038/s41598-021-89020-x

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, *19*(9), 1655-1664. DOI: 10.1101/gr.094052.109

Amoretti, S., Cabrera, B., Torrent, C., Mezquida, G., Lobo, A., González-Pinto, A., Parellada, M., Corripio, I., Vieta, E., De La Serna, E., Butjosa, A., Contreras, F., Sarró, S., Penadés, R., Sánchez-Torres, A. M., Cuesta, M., & Bernardo, M. (2018). Cognitive reserve as an outcome predictor: first-episode affective versus non-affective psychosis. *Acta Psychiatrica Scandinavica*, *138*(5), 441-455. DOI: 10.1111/acps.12949

Barth, D., Papageorge, N. W., & Thom, K. (2020). Genetic endowments and wealth inequality. *Journal of Political Economy*, *128*(4), 1474-1522. DOI: https://doi.org/10.1086/705415

Belsky, D. W., Domingue, B. W., Wedow, R., Arseneault, L., Boardman, J. D., Caspi, A., Conley, D., Fletcher, J. M., Freese, J., & Herd, P. (2018). Genetic analysis of social-class mobility in five longitudinal studies. *Proceedings of the National Academy of Sciences*, *115*(31), E7275-E7284. DOI: 10.1073/pnas.1801238115

Bentall, R. P., & Fernyhough, C. (2008). Social Predictors of Psychotic Experiences: Specificity and Psychological Mechanisms. *Schizophrenia Bulletin*, *34*(6), 1012-1020. DOI: 10.1093/schbul/sbn103

Bouchard, T. J., Jr., & McGue, M. (1981). Familial studies of intelligence: a review. *Science*, *212*(4498), 1055-1059. DOI: 10.1126/science.7195071

Brody, G. H., Gray, J. C., Yu, T., Barton, A. W., Beach, S. R. H., Galván, A., Mackillop, J., Windle, M., Chen, E., Miller, G. E., & Sweet, L. H. (2017). Protective Prevention Effects on the Association of Poverty With Brain Development. *JAMA Pediatrics*, *171*(1), 46. DOI: 10.1001/jamapediatrics.2016.2988

Brody, G. H., Yu, T., Nusslock, R., Barton, A. W., Miller, G. E., Chen, E., Holmes, C., McCormick, M., & Sweet, L. H. (2019). The Protective Effects of Supportive Parenting on the Relationship Between Adolescent Poverty and Resting-State Functional Brain Connectivity During Adulthood. *Psychological Science*, *30*(7), 1040-1049. DOI: 10.1177/0956797619847989

Butler, O., Yang, X. F., Laube, C., Kühn, S., & Immordino-Yang, M. H. (2018). Community violence exposure correlates with smaller gray matter volume and lower IQ in urban adolescents. *Human Brain Mapping*, *39*(5), 2088-2097. DOI: 10.1002/hbm.23988

Cannon, M., Caspi, A., Moffitt, T. E., Harrington, H., Taylor, A., Murray, R. M., & Poulton, R. (2002). Evidence for Early-Childhood, Pan-Developmental Impairment Specific to Schizophreniform Disorder. *Archives of General Psychiatry*, *59*(5), 449. DOI: 10.1001/archpsyc.59.5.449

Cannon, T. D., Bearden, C. E., Hollister, J. M., Rosso, I. M., Sanchez, L. E., & Hadley, T. (2000). Childhood Cognitive Functioning in Schizophrenia Patients and Their Unaffected Siblings: A Prospective Cohort Study. *Schizophrenia Bulletin*, *26*(2), 379-393. DOI: 10.1093/oxfordjournals.schbul.a033460

CDC. (2018). *Health Impact in 5 years*. Retrieved October 23, 2021 from

https://www.cdc.gov/policy/hst/hi5/index.html

Cho, G., Hwang, H., Sarstedt, M., & Ringle, C. M. (2020). Cutoff criteria for overall model fit indexes in generalized structured component analysis. *Journal of Marketing Analytics*, *8*(4), 189-202. DOI: 10.1057/s41270-020-00089-1

Cho, G., Sarstedt, M., & Hwang, H. (2021). A comparative evaluation of factor- and component-based structural equation modeling approaches under (in)correct construct representations. *British Journal of Mathematical and Statistical Psychology*. DOI:

Choi, S. W., Mak, T. S., & O'Reilly, P. F. (2020). Tutorial: a guide to performing polygenic risk score analyses. *Nature Protocols*, *15*(9), 2759-2772. DOI: 10.1038/s41596-020-0353-1

Compton, W. M., Dowling, G. J., & Garavan, H. (2019). Ensuring the Best Use of Data: The Adolescent Brain Cognitive Development Study. *JAMA Pediatrics*, *173*(9), 809-810. DOI: 10.1001/jamapediatrics.2019.2081

Cosway, R., Byrne, M., Clafferty, R., Hodges, A., Grant, E., Abukmeil, S. S., Lawrie, S. M., Miller, P., & Johnstone, E. C. (2000). Neuropsychological change in young people at high risk for schizophrenia: results from the first two neuropsychological assessments of the Edinburgh High Risk Study. *Psychological Medicine*, *30*(5), 1111-1121. DOI: 10.1017/s0033291799002585

Cree, R. A., Bitsko, R. H., Robinson, L. R., Holbrook, J. R., Danielson, M. L., Smith, C., Kaminski, J. W., Kenney, M. K., & Peacock, G. (2018). Health Care, Family, and Community Factors Associated with Mental, Behavioral, and Developmental Disorders and Poverty Among Children Aged 2–8 Years — United States, 2016. *MMWR. Morbidity and Mortality Weekly Report*, *67*(50), 1377-1383. DOI: 10.15585/mmwr.mm6750a1

Cunha, F., & Heckman, J. (2007). The technology of skill formation. *American Economic Review*, *97*(2), 31-47. DOI: 10.1257/aer.97.2.31

Curtis, C. E., Calkins, M. E., Grove, W. M., Feil, K. J., & Iacono, W. G. (2001). Saccadic Disinhibition in Patients With Acute and Remitted Schizophrenia and Their First-Degree Biological Relatives. *American Journal of Psychiatry*, *158*(1), 100-106. DOI: 10.1176/appi.ajp.158.1.100

Deary, I. J., Spinath, F. M., & Bates, T. C. (2006). Genetics of intelligence. *European Journal of Human Genetics*, *14*(6), 690-700. DOI: 10.1038/sj.ejhg.5201588

Demange, P. A., Malanchini, M., Mallard, T. T., Biroli, P., Cox, S. R., Grotzinger, A. D., Tucker-Drob, E. M., Abdellaoui, A., Arseneault, L., Van Bergen, E., Boomsma, D. I., Caspi, A., Corcoran, D. L., Domingue, B. W., Harris, K. M., Ip, H. F., Mitchell, C., Moffitt, T. E., Poulton, R., . . . Nivard, M. G. (2021). Investigating the genetic architecture of noncognitive skills using GWAS-by-subtraction. *Nature Genetics*, *53*(1), 35-44. DOI: 10.1038/s41588-020-00754-2

Eastvold, A. D., Heaton, R. K., & Cadenhead, K. S. (2007). Neurocognitive deficits in the (putative) prodrome and first episode of psychosis. *Schizophrenia Research*, *93*(1-3), 266-277. DOI: 10.1016/j.schres.2007.03.013

Fett, A.-K. J., Velthorst, E., Reichenberg, A., Ruggero, C. J., Callahan, J. L., Fochtmann, L. J., Carlson, G. A., Perlman, G., Bromet, E. J., & Kotov, R. (2020). Long-term Changes in Cognitive Functioning in Individuals With Psychotic Disorders. *JAMA Psychiatry*, *77*(4), 387. DOI: 10.1001/jamapsychiatry.2019.3993

Finucane, H. K., Reshef, Y. A., Anttila, V., Slowikowski, K., Gusev, A., Byrnes, A., Gazal, S., Loh, P.-R., Lareau, C., Shoresh, N., Genovese, G., Saunders, A., Macosko, E., Pollack, S., Perry, J. R. B., Buenrostro, J. D., Bernstein, B. E., Raychaudhuri, S., McCarroll, S., . . . Price, A. L. (2018). Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell types. *Nature Genetics*, *50*(4), 621-629. DOI: 10.1038/s41588-018-0081-4

Fox, J., & Monette, G. (1992). Generalized Collinearity Diagnostics. *Journal of the American Statistical Association*, *87*(417), 178-183. DOI: 10.1080/01621459.1992.10475190

Gard, A. M., Maxwell, A. M., Shaw, D. S., Mitchell, C., Brooks-Gunn, J., McLanahan, S. S., Forbes, E. E., Monk, C. S., & Hyde, L. W. (2021). Beyond family-level adversities: Exploring the developmental timing of neighborhood disadvantage effects on the brain. *Developmental Science*, *24*(1). DOI: 10.1111/desc.12985

Garety, P. A., Kuipers, E., Fowler, D., Freeman, D., & Bebbington, P. E. (2001). A cognitive model of the positive symptoms of psychosis. *Psychological Medicine*, *31*(2), 189-195. DOI: 10.1017/S0033291701003312

Garner, A., & Yogman, M. (2021). Preventing Childhood Toxic Stress: Partnering With Families and Communities to Promote Relational Health. *Pediatrics*, *148*(2), e2021052582. DOI: 10.1542/peds.2021-052582

Ge, T., Chen, C.-Y., Ni, Y., Feng, Y.-C. A., & Smoller, J. W. (2019). Polygenic prediction via Bayesian regression and continuous shrinkage priors. *Nature Communications*, *10*(1). DOI: 10.1038/s41467-019-09718-5

Hair, N. L., Hanson, J. L., Wolfe, B. L., & Pollak, S. D. (2015). Association of Child Poverty, Brain Development, and Academic Achievement. *JAMA Pediatrics*, *169*(9), 822-829. DOI: 10.1001/jamapediatrics.2015.1475

He, Q., Jantac Mam-Lam-Fook, C., Chaignaud, J., Danset-Alexandre, C., Iftimovici, A., Gradels Hauguel, J., Houle, G., Liao, C., Amado, I., Bourgin, J., Daban-Huard, C., Magaud, E., Plaze, M., Rivollier, F., Dion, P. A., Rouleau, G. A., Kebir, O., Krebs, M.-O., & Chaumette, B. (2021). Influence of polygenic risk scores for schizophrenia and resilience on the cognition of individuals at-risk for psychosis. *Translational Psychiatry*, *11*(1). DOI: 10.1038/s41398-021-01624-z

Hill, K. G., Bailey, J. A., Steeger, C. M., Hawkins, J. D., Catalano, R. F., Kosterman, R., Epstein, M., & Abbott, R. D. (2020). Outcomes of Childhood Preventive Intervention Across 2 Generations. *JAMA Pediatrics*, *174*(8), 764. DOI: 10.1001/jamapediatrics.2020.1310

Holmes, C. J., Barton, A. W., Mackillop, J., Galván, A., Owens, M. M., McCormick, M. J., Yu, T., Beach, S. R. H., Brody, G. H., & Sweet, L. H. (2018). Parenting and Salience Network Connectivity Among African Americans: A Protective Pathway for Health-Risk Behaviors. *Biological Psychiatry*, *84*(5), 365-371. DOI: 10.1016/j.biopsych.2018.03.003

Howes, O. D., & Murray, R. M. (2014). Schizophrenia: an integrated sociodevelopmental-cognitive model. *The Lancet*, *383*(9929), 1677-1687. DOI: https://doi.org/10.1016/S0140-6736(13)62036-X

Hwang, H., Cho, G., & Choo, H. (2021). *GSCA Pro 1.1 User's Manual*. https://doi.org/10.13140/RG.2.2.28162.61127/1

Hwang, H., Cho, G., Jung, K., Falk, C. F., Flake, J. K., Jin, M. J., & Lee, S. H. (2021). An approach to structural equation modeling with both factors and components: Integrated generalized structured component analysis. *Psychological Methods*, *26*(3), 273-294. DOI: 10.1037/met0000336

Isungset, M. A., Conley, D., Zachrisson, H. D., Ystrom, E., Havdahl, A., Njølstad, P. R., & Lyngstad, T. H. (2022). Social and genetic associations with educational performance in a Scandinavian welfare state. *Proceedings of the National Academy of Sciences*, *119*(25). DOI: 10.1073/pnas.2201869119

Jensen, A. R. (1998). *The g Factor: The Science of Mental Ability*. Praeger Publishers/Greenwood Publishing Group.

Joo, Y. Y., Cha, J., Freese, J., & Hayes, M. G. (2022). Cognitive Capacity Genome-Wide Polygenic Scores Identify Individuals with Slower Cognitive Decline in Aging. *Genes*, *13*(8), 1320. DOI: 10.3390/genes13081320

Jöreskog, K. G., & Sörbom, D. (1986). *PRELIS: a program for multivariate data screening and data summarization*. Scientific Software, Mooresville.

Judd, N., Sauce, B., Wiedenhoeft, J., Tromp, J., Chaarani, B., Schliep, A., Van Noort, B., Penttilä, J., Grimmer, Y., Insensee, C., Becker, A., Banaschewski, T., Bokde, A. L. W., Quinlan, E. B., Desrivières, S., Flor, H., Grigis, A., Gowland, P., Heinz, A., . . . Klingberg, T. (2020). Cognitive and brain development is independently influenced by socioeconomic status and polygenic scores for educational attainment. *Proceedings of the National Academy of Sciences*, *117*(22), 12411-12418. DOI: 10.1073/pnas.2001228117

Karcher, N. R., Barch, D. M., Avenevoli, S., Savill, M., Huber, R. S., Simon, T. J., Leckliter, I. N., Sher, K. J., & Loewy, R. L. (2018). Assessment of the Prodromal Questionnaire–Brief Child Version for Measurement of Self-reported Psychotic-like Experiences in Childhood. *JAMA Psychiatry*, *75*(8), 853. DOI: 10.1001/jamapsychiatry.2018.1334

Karcher, N. R., Paul, S. E., Johnson, E. C., Hatoum, A. S., Baranger, D. A. A., Agrawal, A., Thompson, W. K., Barch, D. M., & Bogdan, R. (2021). Psychotic-like Experiences and Polygenic Liability in the Adolescent Brain Cognitive Development Study. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*. DOI:

https://doi.org/10.1016/j.bpsc.2021.06.012
Karcher, N. R., Perino, M. T., & Barch, D. M. (2020). An item response theory analysis of the
Prodromal Questionnaire-Brief Child Version: Developing a screening form that informs
understanding of self-reported psychotic-like experiences in childhood. *Journal of Abnormal*
Psychology, *129*(3), 293-304. DOI: 10.1037/abn0000502
Karcher, N. R., Schiffman, J., & Barch, D. M. (2021). Environmental Risk Factors and Psychotic-like
Experiences in Children Aged 9–10. *Journal of the American Academy of Child & Adolescent*
Psychiatry, *60*(4), 490-500. DOI: 10.1016/j.jaac.2020.07.003
Keefe, R. S., Perkins, D. O., Gu, H., Zipursky, R. B., Christensen, B. K., & Lieberman, J. A. (2006). A
longitudinal study of neurocognitive function in individuals at-risk for psychosis. *Schizophr*
Res, *88*(1-3), 26-35. DOI: 10.1016/j.schres.2006.06.041
Kelleher, I., & Cannon, M. (2011). Psychotic-like experiences in the general population:
characterizing a high-risk group for psychosis. *Psychological Medicine*, *41*(1), 1-6. DOI:
10.1017/s0033291710001005
Kowarik, A., & Templ, M. (2016). Imputation with the R Package VIM. *Journal of Statistical*
Software, *74*(7), 1 - 16. DOI: 10.18637/jss.v074.i07
Krabbendam, L. (2005). Schizophrenia and Urbanicity: A Major Environmental Influence--
Conditional on Genetic Risk. *Schizophrenia Bulletin*, *31*(4), 795-799. DOI:
10.1093/schbul/sbi060
Lam, M., Chen, C.-Y., Li, Z., Martin, A. R., Bryois, J., Ma, X., Gaspar, H., Ikeda, M., Benyamin, B.,
Brown, B. C., Liu, R., Zhou, W., Guan, L., Kamatani, Y., Kim, S.-W., Kubo, M.,
Kusumawardhani, A. A. A. A., Liu, C.-M., Ma, H., . . . the, N. (2019). Comparative genetic
architectures of schizophrenia in East Asian and European populations. *Nature Genetics*,
51(12), 1670-1678. DOI: 10.1038/s41588-019-0512-x
Lee, J. J., Wedow, R., Okbay, A., Kong, E., Maghzian, O., Zacher, M., Nguyen-Viet, T. A., Bowers, P.,
Sidorenko, J., Karlsson Linnér, R., Fontana, M. A., Kundu, T., Lee, C., Li, H., Li, R., Royer,
R., Timshel, P. N., Walters, R. K., Willoughby, E. A., . . . Cesarini, D. (2018). Gene discovery
and polygenic prediction from a genome-wide association study of educational attainment in
1.1 million individuals. *Nature Genetics*, *50*(8), 1112-1121. DOI: 10.1038/s41588-018-0147-3
Leeson, V. C., Sharma, P., Harrison, M., Ron, M. A., Barnes, T. R. E., & Joyce, E. M. (2011). IQ
Trajectory, Cognitive Reserve, and Clinical Outcome Following a First Episode of Psychosis:
A 3-Year Longitudinal Study. *Schizophrenia Bulletin*, *37*(4), 768-777. DOI:
10.1093/schbul/sbp143
Lewis, G., Dykxhoorn, J., Karlsson, H., Khandaker, G. M., Lewis, G., Dalman, C., & Kirkbride, J. B.
(2020). Assessment of the Role of IQ in Associations Between Population Density and
Deprivation and Nonaffective Psychosis. *JAMA Psychiatry*, *77*(7), 729. DOI:
10.1001/jamapsychiatry.2020.0103
Luby, J. L., Barch, D. M., Belden, A., Gaffrey, M. S., Tillman, R., Babb, C., Nishino, T., Suzuki, H.,
& Botteron, K. N. (2012). Maternal support in early childhood predicts larger hippocampal
volumes at school age. *Proceedings of the National Academy of Sciences*, *109*(8), 2854-2859.
DOI: 10.1073/pnas.1118003109
Luby, J. L., Belden, A., Harms, M. P., Tillman, R., & Barch, D. M. (2016). Preschool is a sensitive
period for the influence of maternal support on the trajectory of hippocampal development.
Proceedings of the National Academy of Sciences, *113*(20), 5742-5747. DOI:
10.1073/pnas.1601443113
Marshall, A. T., Betts, S., Kan, E. C., McConnell, R., Lanphear, B. P., & Sowell, E. R. (2020).
Association of lead-exposure risk and family income with childhood brain outcomes. *Nature*
Medicine, *26*(1), 91-97. DOI: 10.1038/s41591-019-0713-y
Martin, J., Hamshere, M. L., Stergiakouli, E., O'Donovan, M. C., & Thapar, A. (2015).
Neurocognitive abilities in the general population and composite genetic risk scores for
attention-deficit hyperactivity disorder. *Journal of Child Psychology and Psychiatry*, *56*(6),
648-656. DOI: https://doi.org/10.1111/jcpp.12336
Maxwell, J., Ronald, A., Cardno, A. G., Breen, G., Rimfeld, K., & Vassos, E. (2023). Genetic and
Geographical Associations With Six Dimensions of Psychotic Experiences in Adolesence.
Schizophrenia Bulletin, *49*(2), 319-328. DOI: 10.1093/schbul/sbac149
Miao, W., Hu, W., Ogburn, E. L., & Zhou, X.-H. (2022). Identifying Effects of Multiple Treatments in

the Presence of Unmeasured Confounding. *Journal of the American Statistical Association*, 1-15. DOI: 10.1080/01621459.2021.2023551

Noble, K. G., Houston, S. M., Brito, N. H., Bartsch, H., Kan, E., Kuperman, J. M., Akshoomoff, N., Amaral, D. G., Bloss, C. S., Libiger, O., Schork, N. J., Murray, S. S., Casey, B. J., Chang, L., Ernst, T. M., Frazier, J. A., Gruen, J. R., Kennedy, D. N., Van Zijl, P., . . . Sowell, E. R. (2015). Family income, parental education and brain structure in children and adolescents. *Nature Neuroscience*, *18*(5), 773-778. DOI: 10.1038/nn.3983

Okbay, A., Wu, Y., Wang, N., Jayashankar, H., Bennett, M., Nehzati, S. M., Sidorenko, J., Kweon, H., Goldman, G., Gjorgjieva, T., Jiang, Y., Hicks, B., Tian, C., Hinds, D. A., Ahlskog, R., Magnusson, P. K. E., Oskarsson, S., Hayward, C., Campbell, A., . . . Young, A. I. (2022). Polygenic prediction of educational attainment within and between families from genome-wide association analyses in 3 million individuals. *Nature Genetics*, *54*(4), 437-449. DOI: 10.1038/s41588-022-01016-z

Pat, N., Riglin, L., Anney, R., Wang, Y., Barch, D. M., Thapar, A., & Stringaris, A. (2021). Motivation and Cognitive Abilities as Mediators Between Polygenic Scores and Psychopathology in Children. *Journal of the American Academy of Child & Adolescent Psychiatry*. DOI: https://doi.org/10.1016/j.jaac.2021.08.019

Peverill, M., Dirks, M. A., Narvaja, T., Herts, K. L., Comer, J. S., & McLaughlin, K. A. (2021). Socioeconomic status and child psychopathology in the United States: A meta-analysis of population-based studies. *Clinical Psychology Review*, *83*, 101933. DOI: https://doi.org/10.1016/j.cpr.2020.101933

Piccolo, L. R., Merz, E. C., & Noble, K. G. (2019). School climate is associated with cortical thickness and executive function in children and adolescents. *Developmental Science*, *22*(1), e12719. DOI: 10.1111/desc.12719

Plomin, R., & Spinath, F. M. (2004). Intelligence: Genetics, Genes, and Genomics. *Journal of Personality and Social Psychology*, *86*, 112-129. DOI: 10.1037/0022-3514.86.1.112

Plomin, R., & von Stumm, S. (2018). The new genetics of intelligence. *Nature Reviews Genetics*, *19*(3), 148-159. DOI: 10.1038/nrg.2017.104

Poulton, R., Caspi, A., Moffitt, T. E., Cannon, M., Murray, R., & Harrington, H. (2000). Children's Self-Reported Psychotic Symptoms and Adult Schizophreniform Disorder. *Archives of General Psychiatry*, *57*(11), 1053. DOI: 10.1001/archpsyc.57.11.1053

Rakesh, D., Seguin, C., Zalesky, A., Cropley, V., & Whittle, S. (2021). Associations between neighborhood disadvantage, resting-state functional connectivity, and behavior in the Adolescent Brain Cognitive Development (ABCD) Study®: Moderating role of positive family and school environments. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*. DOI: https://doi.org/10.1016/j.bpsc.2021.03.008

Reichenberg, A., Weiser, M., Rapp, M. A., Rabinowitz, J., Caspi, A., Schmeidler, J., Knobler, H. Y., Lubin, G., Nahon, D., Harvey, P. D., & Davidson, M. (2005). Elaboration on Premorbid Intellectual Performance in Schizophrenia. *Archives of General Psychiatry*, *62*(12), 1297. DOI: 10.1001/archpsyc.62.12.1297

Ritchie, S. J., Hill, W. D., Marioni, R. E., Davies, G., Hagenaars, S. P., Harris, S. E., Cox, S. R., Taylor, A. M., Corley, J., Pattie, A., Redmond, P., Starr, J. M., & Deary, I. J. (2020). Polygenic predictors of age-related decline in cognitive ability. *Molecular Psychiatry*, *25*(10), 2584-2598. DOI: 10.1038/s41380-019-0372-x

Ruderfer, D. M., Ripke, S., McQuillin, A., Boocock, J., Stahl, E. A., Pavlides, J. M. W., Mullins, N., Charney, A. W., Ori, A. P. S., Loohuis, L. M. O., Domenici, E., Di Florio, A., Papiol, S., Kalman, J. L., Trubetskoy, V., Adolfsson, R., Agartz, I., Agerbo, E., Akil, H., . . . Kendler, K. S. (2018). Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell*, *173*(7), 1705-1715.e1716. DOI: https://doi.org/10.1016/j.cell.2018.05.046

Sariaslan, A., Fazel, S., D'Onofrio, B. M., Långström, N., Larsson, H., Bergen, S. E., Kuja-Halkola, R., & Lichtenstein, P. (2016). Schizophrenia and subsequent neighborhood deprivation: revisiting the social drift hypothesis using population, twin and molecular genetic data. *Translational Psychiatry*, *6*(5), e796-e796. DOI: 10.1038/tp.2016.62

Selzam, S., Ritchie, S. J., Pingault, J.-B., Reynolds, C. A., O'Reilly, P. F., & Plomin, R. (2019). Comparing Within- and Between-Family Polygenic Score Prediction. *The American Journal*

of Human Genetics, *105*(2), 351-363. DOI: https://doi.org/10.1016/j.ajhg.2019.06.006

Shafee, R., Nanda, P., Padmanabhan, J. L., Tandon, N., Alliey-Rodriguez, N., Kalapurakkel, S., Weiner, D. J., Gur, R. E., Keefe, R. S. E., Hill, S. K., Bishop, J. R., Clementz, B. A., Tamminga, C. A., Gershon, E. S., Pearlson, G. D., Keshavan, M. S., Sweeney, J. A., McCarroll, S. A., & Robinson, E. B. (2018). Polygenic risk for schizophrenia and measured domains of cognition in individuals with psychosis and controls. *Translational Psychiatry*, *8*(1). DOI: 10.1038/s41398-018-0124-8

Shonkoff, J. P. (2012). Leveraging the biology of adversity to address the roots of disparities in health and development. *Proceedings of the National Academy of Sciences*, *109*(Supplement_2), 17302-17307. DOI: 10.1073/pnas.1121259109

Spearman, C. (1904). "General Intelligence," Objectively Determined and Measured. *The American Journal of Psychology*, *15*(2), 201-292. DOI: 10.2307/1412107

Stern, Y. (2009). Cognitive reserve. *Neuropsychologia*, *47*(10), 2015-2028. DOI: https://doi.org/10.1016/j.neuropsychologia.2009.03.004

Taylor, R. L., Cooper, S. R., Jackson, J. J., & Barch, D. M. (2020). Assessment of Neighborhood Poverty, Cognitive Function, and Prefrontal and Hippocampal Volumes in Children. *JAMA Network Open*, *3*(11), e2023774. DOI: 10.1001/jamanetworkopen.2020.23774

Thompson, W. K., Barch, D. M., Bjork, J. M., Gonzalez, R., Nagel, B. J., Nixon, S. J., & Luciana, M. (2019). The structure of cognition in 9 and 10 year-old children and associations with problem behaviors: Findings from the ABCD study's baseline neurocognitive battery. *Developmental Cognitive Neuroscience*, *36*, 100606. DOI: 10.1016/j.dcn.2018.12.004

Tomasi, D., & Volkow, N. D. (2021). Associations of family income with cognition and brain structure in USA children: prevention implications. *Molecular Psychiatry*. DOI: 10.1038/s41380-021-01130-0

Tooley, U. A., Bassett, D. S., & Mackey, A. P. (2021). Environmental influences on the pace of brain development. *Nature Reviews Neuroscience*, *22*(6), 372-384. DOI: 10.1038/s41583-021-00457-5

Tooley, U. A., Mackey, A. P., Ciric, R., Ruparel, K., Moore, T. M., Gur, R. C., Gur, R. E., Satterthwaite, T. D., & Bassett, D. S. (2020). Associations between Neighborhood SES and Functional Brain Network Development. *Cerebral Cortex*, *30*(1), 1-19. DOI: 10.1093/cercor/bhz066

van der Steen, Y., Myin-Germeys, I., Van Nierop, M., Ten Have, M., De Graaf, R., Van Dorsselaer, S., Van Os, J., & Van Winkel, R. (2019). 'False-positive' self-reported psychotic experiences in the general population: an investigation of outcome, predictive factors and clinical relevance. *Epidemiology and Psychiatric Sciences*, *28*(5), 532-543. DOI: 10.1017/s2045796018000197

van Os, J., Linscott, R. J., Myin-Germeys, I., Delespaul, P., & Krabbendam, L. (2009). A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness–persistence–impairment model of psychotic disorder. *Psychological Medicine*, *39*(2), 179-195. DOI: 10.1017/S0033291708003814

Van Os, J., & Reininghaus, U. (2016). Psychosis as a transdiagnostic and extended phenotype in the general population. *World Psychiatry*, *15*(2), 118-124. DOI: 10.1002/wps.20310

Vorstman, J. A. S., Breetvelt, E. J., Duijff, S. N., Eliez, S., Schneider, M., Jalbrzikowski, M., Armando, M., Vicari, S., Shashi, V., Hooper, S. R., Chow, E. W. C., Fung, W. L. A., Butcher, N. J., Young, D. A., McDonald-Mcginn, D. M., Vogels, A., Van Amelsvoort, T., Gothelf, D., Weinberger, R., . . . Bassett, A. S. (2015). Cognitive Decline Preceding the Onset of Psychosis in Patients With 22q11.2 Deletion Syndrome. *JAMA Psychiatry*, *72*(4), 377. DOI: 10.1001/jamapsychiatry.2014.2671

Walker, S. P., Chang, S. M., Wright, A. S., Pinto, R., Heckman, J. J., & Grantham-Mcgregor, S. M. (2021). Cognitive, psychosocial, and behaviour gains at age 31 years from the Jamaica early childhood stimulation trial. *Journal of Child Psychology and Psychiatry*. DOI: 10.1111/jcpp.13499

Weissman, D. G., Conger, R. D., Robins, R. W., Hastings, P. D., & Guyer, A. E. (2018). Income change alters default mode network connectivity for adolescents in poverty. *Developmental Cognitive Neuroscience*, *30*, 93-99. DOI: 10.1016/j.dcn.2018.01.008

Xu, L., Skoularidou, M., Cuesta-Infante, A., & Veeramachaneni, K. (2019). Modeling Tabular Data Using Conditional GAN.*Proceedings of the 33rd International Conference on Neural*

913   *Information Processing Systems* 33rd Conference on Neural Information Processing Systems
914   (NeurIPS 2019), Vancouver, Canada.
915 Young, A. I., Benonisdottir, S., Przeworski, M., & Kong, A. (2019). Deconstructing the sources of
916   genotype-phenotype associations in humans. *Science*, *365*(6460), 1396-1400. DOI:
917   10.1126/science.aax3710

Figures

Figure 1: Study diagram for longitudinal trajectories of genetic and environmental influences on PLEs through cognitive intelligence. This study examines the mediating role of cognitive intelligence within the effects of cognitive phenotype PGSs, high family SES, low neighborhood SES, and positive family and school environments on PLEs observed at baseline, 1-year follow-up, and 2-year follow-up in children aged 9-10 years. Both direct and indirect effects, as well as total effects, were evaluated for statistical significance.

Figure 2: Linear models testing genetic, socioeconomic, and environmental factors
associated with cognitive intelligence and PLEs. Standardized coefficients of a linear
mixed model with CP PGS (**A**, **B**) and EA PGS (**C**, **D**). The analyses included 6,602 samples
of multiethnicity. Cognitive intelligence and PLEs correlated with the PGSs, residential
disadvantage, positive environment, and family SES in the opposite directions. Error bars
indicate 95% bootstrapped confidence intervals with 5,000 iterations. CP and EA denote
cognitive performance and education attainment, respectively; PGS, polygenic scores; SES,
socioeconomic status; PLEs, psychotic-like experiences; ADI, Area Deprivation Index;
Poverty, percentage of individuals below -125% of the poverty level; Years, years of
residence. (*p-FDR<0.05)

D

Psychotic-Like Experiences (1-year, 2-year follow-up)

Figure 3: Direct/Indirect effects of gene-environment factors to cognitive and PLEs outcomes

A. Direct pathways from PGS, high family SES, low neighborhood SES, and positive environment to cognitive intelligence and PLEs. Standardized path coefficients are indicated on each path as direct effect estimates (significance level *p<0.05). **B**. Indirect pathways to PLEs via intelligence were significant for polygenic scores, high family SES, low neighborhood SES, and positive environment, indicating the significant mediating role of intelligence. **C**. Relative effect sizes of direct and indirect pathways within the total effects on PLEs. The standardized effect sizes of direct pathways are colored within each bar. In **A-B**, child sex, genetic ancestry, BMI, marital status, family history of psychiatric disorders, and ABCD research sites were included as covariates. CP PGS and EA PGS denote polygenic scores of cognitive performance and education attainment, respectively; SES, socioeconomic status; PLEs, psychotic-like experiences; Crystallized and Fluid, crystallized and fluid intelligence; ADI, Area Deprivation Index; Poverty, percentage of individuals below -125% of the poverty level; Years, years of residence.

Note: * indicates a statistically significant parameter estimate at α=0.05.

A) Direct Effects

B) Indirect Effects

C) Relative Effect Sizes

Tables

Table 1. **Demographic characteristics of the study participants.**

Of the initial 11,878 ABCD samples, we obtained data for the variables of interest for 6,602 multiethnic children. For multiethnic subjects (main analyses, n=6,602), 47.15% were female, and the parents of 70.21% were married. In European ancestry samples (sensitivity analyses, n=5,211), 46.71% were female, and the parents of 77.47% were married. Children of European ancestry had significantly different marital status (p<0.0001), lower BMI (p<0.0001) and higher family history of psychiatric disorders (p<0.0001) than children of multiethnic ancestries. There were no significant differences in other characteristics between the two ancestry groups.

The 6,602 multiethnic participants consisted of 890 African-ancestry (13.48%), 229 Native American ancestry (3.47%), 91 East Asian ancestry (1.38%), 181 not specified (2.74%), and 5,211 European ancestry (78.93%) children. Differences between genetic ancestry groups were calculated using χ^2 tests for categorical variables and t tests for continuous variables.

Demographic Characteristics		European Ancestry (n=5,211)			Multiethnic (n=6,602)			Test Statistics	
		N	Ratio (%)	Mean (SD)	N	Ratio (%)	Mean (SD)	$t(df)/\chi^2(df)$	p value
Sex	Male	2,777	53.29%		3,489	52.85%		-0.4795 (11811)	0.6316
	Female	2,434	46.71%		3,113	47.15%			
Marital Status of the Caregiver	Married	4,037	77.47%		4,635	70.21%		-10.2326 (11811)	<0.0001
	Widowed	38	0.73%		50	0.76%			
	Divorced	485	9.31%		610	9.24%			
	Separated	155	2.97%		232	3.51%			
	Never Married	275	5.28%		718	10.88%			
	Living with Partner	221	4.24%		357	5.41%			
Age (rounded to chronological month)		5,211		118.99 (7.46)	6,602		118.94 (7.41)	0.3652 (11811)	0.715
BMI				18.29 (3.67)			18.72 (4.12)	-5.8889 (11811)	<0.0001
Family History of Psychiatric Disorders (proportion of first-				0.10 (0.11)			0.09 (0.11)	4.4296 (11811)	<0.0001

degree relatives who experienced mental illness)									
Genetic Ancestry	**African**	-			890	13.48%			
	Native American	-			229	3.47%			
	East Asian	-			91	1.38%			
	European	5,211	100%		5,211	78.93%			
	Not Specified	-			181	2.74%			

Table 2. **IGSCA analysis of multiethnic samples.** Sex, age, genetic ancestry, BMI, parental education, marital status of the caregiver, household income, and family's financial adversity based on parents' self-report, family history of psychiatric disorders, and ABCD research sites were included as covariates. Family socioeconomic status was included to confirm that the aassociations of PGS, neighborhood disadvantage, and positive environment are meaningful. SE and CI represent standard error and confidence intervals, respectively. Significant effects are marked with a star (*).

Analysis of Total/Direct/Indirect Effects					
Effect Type	**Paths**	**Estimate**	**SE**	**95% CI**	**Significance**
1. Effects from PGS to Intelligence (baseline year)					

Direct Effect	PGS→Intelligence	0.242736	0.01277	0.218202	0.267954	∗
2. Effects from High Family SES to Intelligence (baseline year)						
Direct Effect	High Family SES →Intelligence	0.293171	0.016737	0.260337	0.326413	∗
3. Effects from Low Neighborhood SES to Intelligence (baseline year)						
Direct Effect	Low Neighborhood SES →Intelligence	-0.1121	0.016768	-0.14568	-0.08118	∗
4. Effects from Positive Environment to Intelligence (baseline year)						
Direct Effect	Positive Environment→Intelligence	0.026793	0.012552	0.003984	0.052633	∗
5. Effects from Intelligence to Psychotic-like Experiences (all years)						
Direct Effect	Intelligence→ Psychotic-like Experiences (baseline year)	-0.14421	0.027683	-0.20344	-0.09516	∗
Direct Effect	Intelligence→ Psychotic-like Experiences (1-year follow-up)	-0.14638	0.027507	-0.20834	-0.09983	∗
Direct Effect	Intelligence→ Psychotic-like Experiences (2-year follow-up)	-0.11276	0.028708	-0.17428	-0.063	∗
6. Effects from PGS to Psychotic-like Experiences (baseline year)						
Total Effect	PGS→ Psychotic-like Experiences	-0.05017	0.011354	-0.07292	-0.02853	∗
Indirect Effect	PGS→ Intelligence→ Psychotic-like Experiences	-0.035	0.007126	-0.0508	-0.02273	∗
Direct Effect	PGS→ Psychotic-like Experiences	-0.01516	0.01347	-0.04085	0.012389	

7 Effects from High Family SES to Psychotic-like Experiences (baseline year)						
Total Effect	High Family SES→ Psychotic-like Experiences	-0.12126	0.019087	-0.15851	-0.08313	*
Indirect Effect	High Family SES→ Intelligence→ Psychotic-like Experiences	-0.04228	0.008652	-0.06139	-0.02707	*
Direct Effect	High Family SES→ Psychotic-like Experiences	-0.07898	0.020747	-0.11856	-0.03698	*
8. Effects from Low Neighborhood SES to Psychotic-like Experiences (baseline year)						
Total Effect	Low Neighborhood SES→ Psychotic-like Experiences	0.050374	0.018277	0.013545	0.085934	*
Indirect Effect	Low Neighborhood SES→ Intelligence→ Psychotic-like Experiences	0.016166	0.003944	0.009843	0.025298	*
Direct Effect	Low Neighborhood SES→ Psychotic-like Experiences	0.034209	0.0184	-0.00268	0.069813	
9. Effects from Positive Environment to Psychotic-like Experiences (baseline year)						
Total Effect	Positive Environment→ Psychotic-like Experiences	-0.15256	0.013871	-0.17965	-0.1252	*
Indirect Effect	Positive Environment→ Intelligence→ Psychotic-like Experiences	-0.00386	0.002065	-0.00859	-0.00058	*
Direct Effect	Positive Environment→ Psychotic-like Experiences	-0.14869	0.014025	-0.17573	-0.12073	*
10. Effects from PGS to Psychotic-like Experiences (1-year follow up)						
Total Effect	PGS→ Psychotic-like Experiences	-0.035895	0.011646	-0.058499	-0.013458	*

Indirect Effect	PGS→ Intelligence→ Psychotic-like Experiences	-0.03553	0.007062	-0.05176	-0.02376	*
Direct Effect	PGS→ Psychotic-like Experiences	-0.00036	0.013579	-0.02566	0.028107	
11. Effects from High Family SES to Psychotic-like Experiences (1-year follow up)						
Total Effect	High Family SES→ Psychotic-like Experiences	-0.11184	0.018291	-0.1478	-0.07584	*
Indirect Effect	High Family SES→ Intelligence→ Psychotic-like Experiences	-0.04291	0.008569	-0.06242	-0.0288	*
Direct Effect	High Family SES→ Psychotic-like Experiences	-0.06892	0.019586	-0.10522	-0.02866	*
12. Effects from Low Neighborhood SES to Psychotic-like Experiences (1-year follow up)						
Total Effect	Low Neighborhood SES→ Psychotic-like Experiences	0.032947	0.018055	-0.00264	0.068773	
Indirect Effect	Low Neighborhood SES→ Intelligence→ Psychotic-like Experiences	0.016409	0.004003	0.010133	0.025893	*
Direct Effect	Low Neighborhood SES→ Psychotic-like Experiences	0.016538	0.018503	-0.02066	0.051855	
13. Effects from Positive Environment to Psychotic-like Experiences (1-year follow up)						
Total Effect	Positive Environment→ Psychotic-like Experiences	-0.13149	0.013154	-0.15756	-0.10589	*
Indirect Effect	Positive Environment→ Intelligence→ Psychotic-like Experiences	-0.00392	0.00208	-0.0087	-0.00059	*
Direct Effect	Positive Environment→ Psychotic-like Experiences	-0.12757	0.013237	-0.15343	-0.10137	*

14. Effects from PGS to Psychotic-like Experiences (2-year follow up)						
Total Effect	PGS→ Psychotic-like Experiences	-0.03643	0.012196	-0.06027	-0.01272	∗
Indirect Effect	PGS→ Intelligence→ Psychotic-like Experiences	-0.02737	0.007142	-0.04307	-0.01508	∗
Direct Effect	PGS→ Psychotic-like Experiences	-0.00906	0.014737	-0.03696	0.021152	
15. Effects from High Family SES to Psychotic-like Experiences (2-year follow up)						
Total Effect	High Family SES→ Psychotic-like Experiences	-0.11632	0.018067	-0.15258	-0.08174	∗
Indirect Effect	High Family SES→ Intelligence→ Psychotic-like Experiences	-0.03306	0.008796	-0.05228	-0.01807	∗
Direct Effect	High Family SES→ Psychotic-like Experiences	-0.08326	0.019462	-0.12066	-0.04392	∗
16. Effects from Low Neighborhood SES to Psychotic-like Experiences (2-year follow up)						
Total Effect	Low Neighborhood SES→ Psychotic-like Experiences	0.01921	0.018684	-0.01767	0.055261	
Indirect Effect	Low Neighborhood SES→ Intelligence→ Psychotic-like Experiences	0.012641	0.003814	0.006533	0.0215	∗
Direct Effect	Low Neighborhood SES→ Psychotic-like Experiences	0.006569	0.019176	-0.03173	0.042823	
17. Effects from Positive Environment to Psychotic-like Experiences (2-year follow up)						
Total Effect	Positive Environment→ Psychotic-like Experiences	-0.13627	0.013881	-0.1635	-0.10926	∗

Indirect Effect	Positive Environment→ Intelligence→ Psychotic-like Experiences	-0.00302	0.001703	-0.0069	-0.00043	*
Direct Effect	Positive Environment→ Psychotic-like Experiences	-0.13325	0.014009	-0.16069	-0.10565	*

Supporting Information

Figure S1. Component correlation matrix of IGSCA analysis. A. Correlation between all component / factor variables of the IGSCA model. **B.** Correlation between all observed variables used to construct the relevant component / factor variables in the IGSCA model. CP PGS and EA PGS denote polygenic scores of cognitive performance and education attainment, respectively; PLEs, psychotic-like experiences; Crystallized and Fluid, crystallized and fluid intelligence; ADI, Area Deprivation Index; Poverty, percentage of individuals below -125% of the poverty level; Years, years of residence.

A

B

Appendix S1. Supplementary Methods

Genotype data

The ABCD study collected saliva samples of study participants at the baseline visit and shipped the samples from the collection site to the Rutgers University Cell and DNA Repository. Genomic DNA was extracted and genotyped using the Affymetrix NIDA Smokescreen array (733,293 SNPs). We removed any inferior SNPs with (i) genotype call rate<95%, (ii) sample call rate<95%, and (iii) rare variants with minor allele frequency (MAF)<0.01. Variant imputation was performed with the Michigan Imputation Server (Das et al., 2016) using the 1000 Genome phase 3 version 5 multiethnic Grch37/hg19 reference panel with Eagle ver2.4 phased output (Loh et al., 2016). For the imputed 12,046,090 SNPs, we only retained data from any individuals with <5% missing genotypes; without extreme heterozygosity (F coefficient < 3 standard deviations from the population mean); and SNPs with >0.4 imputation quality INFO score, <5% missingness rate, >0.01 MAF and Hardy-Weinberg equilibrium ($p>10^{-6}$). The genetic ancestry of each participant was determined with the fastSTRUCTURE algorithm (Raj, Stephens, & Pritchard, 2014), available from ABCD Release 4.0. Considering the diverse ethnic and racial backgrounds of the study participants, we estimated both kinship coefficients (K.C.s) and ancestrally informative principal components (P.C.s) to additionally control familial relatedness and ancestry admixture using PC-Air (M. P. Conomos, Miller, & Thornton, 2015) and PC-Relate (Matthew P. Conomos, Reiner, Weir, & Thornton, 2016). We selected unrelated participants that were inferred to be more distant than 4[th]-degree relatives (K.C.>0.022) and removed any outliers that fell significantly outside (>6 S.D. limits) the center in P.C. space. In the rest of this paper, we used final genotype data (11,301,999 variants) of 10,199 unrelated multiethnic samples, including 7,893 European-ancestry participants, after Q.C.

Polygenic Scores (PGS)

Hyperparameters of PGSs were optimized in the held-out validation set of 1,579 unrelated participants, consisting of 88 of African ancestry (5.57%), 25 of East Asian ancestry (1.58%), 1,365 of European ancestry (86.45%), 88 of Native American ancestry (2.91%), and 55 not specified (3.48%). The

validation set was created during the quality control process of the study genotype data when we selected unrelated study participants of the original ABCD samples with pairwise kinship coefficients of less than 0.022 among them. To select the optimal hyperparameter, we fitted linear regression models for intelligence composite scores within the validation set and evaluated the model performance in terms of the highest R^2 and effect size (beta). The models were adjusted for age, sex, and the first ten principal components of genotype data. For the PGS of multiethnic participants, genetic ancestry determined by ADMIXTURE (Alexander et al., 2009) was additionally included as a covariate. The validation samples were only used for hyperparameter tuning for PGS optimization and were excluded from any further analyses. The PGSs were residualized against the first ten ancestrally informative P.C.s to adjust for population stratification.

Parent-rated Psychotic-like Experiences

As self-reports and parent-reports of psychopathology often differ, parent-rated PLEs derived from four items of the Child Behavior Checklist:

1) "Hears sounds or voices that other people think aren't there."

2) "Sees things that other people think aren't there."

3) "Does things that other people think are strange."

4) "Has thoughts that other people would think are strange."

Each question was scored from 0=not true, 1=somewhat or sometimes true, and 2=very true or often true.

Family and Neighborhood Socioeconomic Status (SES)

We assessed children's family SES based on family Income, parental Education, and family's financial adversity. All three variables were based on self-reported responses of children's primary caregiver. For family income, the caregivers were asked "*What is your TOTAL COMBINED FAMILY INCOME for the past 12 months? This should include income (before taxes and deductions) from all sources, wages, rent from properties, social security, disability and/or veteran's benefits, unemployment*

benefits, workman's compensation, help from relative (include child payments and alimony), and so on. If Separated/Divorced, please average the two household incomes." Answer choices are shown below:

1) Less than $5,000

2) $5,000 through $11,999

3) $12,000 through $15,999

4) $16,000 through $24,999

5) $25,000 through $34,999

6) $35,000 through $49,999

7) $50,000 through $74,999

8) $75,000 through $99,999

9) $100,000 through $199,999

10) $200,000 and greater

Parental education was measured as the highest grade or level of school completed or highest degree received by the primary caregiver. ("What is the highest grade or level of school you have completed or the highest degree you have received?"). Answer choices were:

0) Never attended/Kindergarten only

1) 1st grade

2) 2nd grade

3) 3rd grade

4) 4th grade

5) 5th grade

6) 6th grade

7) 7th grade

8) 8th grade

9) 9th grade

10) 10th grade

11) 11th grade

12) 12th grade

13) High school graduate

14) GED or equivalent Diploma

15) Some college

16) Associate degree: Occupational

17) Associate degree: Academic Program

18) Bachelor's degree (ex. BA)

19) Master's degree (ex. MA)

20) Professional School degree (ex. MD)

21) Doctoral degree (ex. PhD)

Family's financial adversity is measured with Parent-Reported Financial Adversity Questionnaire, reflecting family's financial ability to pay for basic life expenses (Diemer et al., 2013). The questionnaire asked whether the child's caregiver and family experienced any of the following difficulties within the past 12 months:

1) "Needed food but couldn't afford to buy it or couldn't afford to go out to get it?"

2) "Were without telephone service because you could not afford it?"

3) "Didn't pay the full amount of the rent or mortgage because you could not afford it?"

4) "Were evicted from your home for not paying the rent or mortgage?"

5) "Had services turned off by the gas or electric company, or the oil company wouldn't deliver oil because payments were not made?"

6) "Had someone who needed to see a doctor or go to the hospital but didn't go because you could not afford it?"

7) "Had someone who needed a dentist but couldn't go because you could not afford it?"

Each of the seven items was scored with 0=no or 1=yes.

Neighborhood SES was measured by Residential History Derived Scores based on the census tracts of each respondent's primary address. The national percentile score of Area Deprivation Index (ADI) (Karcher, Schiffman, et al., 2021; Rakesh et al., 2021) was calculated from the 2011–2015 American Community Survey 5-year summary. Components used for deriving ADI includes:

1) Percentage of population aged >=25 y with <9 y of education

2) Percentage of population aged >=25 y with at least a high school diploma

3) Percentage of employed persons aged >=16 y in white collar occupations

4) Median family income

5) Income disparity defined by Singh as the log of 100 x ratio of the number of households with <10000 annual income to the number of households with >50000 annual income.

6) Median home value

7) Median gross rent

8) Median monthly mortgage

9) Percentage of owner

10) Percentage of occupied housing units with >1 person per room (crowding)

11) Percentage of civilian labor force population aged >=16 y unemployed (unemployment rate)

12) Percentage of families below the poverty level

13) Percentage of population below 138% of the poverty threshold

14) Percentage of single

15) Percentage of occupied housing units without a motor vehicle

16) Percentage of occupied housing units without a telephone

17) Percentage of occupied housing units without complete plumbing (log)

Positive Family and School Environment

Positive parenting behavior was assessed using the ABCD Children's Report of Parental Behavioral Inventory. We used the average values of mean summary scores of five questionnaire items about first and second caregivers. Single values were used for respondents lacking a second caregiver. Positive school environment was assessed as the sum of children's responses to twelve items in the ABCD School Risk and Protective Factors Survey.

Prior work used response items about the first caregiver for positive parenting behavior and the first six items about the school environment for positive school environment (Rakesh et al., 2021). However, to obtain a comprehensive and accurate assessment, we also included items about the second caregiver and six additional questions about the school environment.

Children were asked to choose between 1=Not like him/her, 2=Somewhat like him/her, or 3=A lot like him/her about the five questions regarding the first and second caregiver:

1) Smiles at me very often.

2) Is able to make me feel better when I am upset.

3) Believes in showing his/her love for me.

4) Is easy to talk to.

5) Makes me feel better after talking over my worries with him/her

The following questions were asked to assess positive school environment:

1) In my school, students have lots of chances to help decide things like class activities and rules.

2) I get along with my teachers.

3) My teacher(s) notices when I am doing a good job and lets me know about it.

4) There are lots of chances for students in my school to get involved in sports, clubs, or other school activities outside of class.

5) I feel safe at my school.

6) The school lets my parents know when I have done something well.

7) I like school because I do well in class.

8) I feel I'm just as smart as other kids my age.

9) There are lots of chances to be part of class discussions or activities.

10) In general, I like school a lot.

11) Usually, school bores me.

12) Getting good grades is not so important to me.

Each of the 12 items was scored with 1=NO!, 2=no, 3=yes, 4=YES!. Item 11 and 12 were reverse coded when obtaining summary scores.

Appendix S2. Supplementary Results

Results of linear mixed models with European samples

As the European-descent-based GWAS was used for constructing PGS, we reran the main analyses using participants of European ancestry (n=5,211) to adjust for ethnic confounding (females 46.71%, mean age 118.99 [SD 7.46]).

The results of linear mixed models were similar to those of main analyses (**Table S3**). Higher intelligence was significantly associated with higher CP and EA PGS (βs> 0.0754, p<0.0001), lower ADI (βs<-0.0503, p=0.0253), more *years of residence* (βs> 0.0268, p=0.0311), positive school environment (βs> 0.0365, p=0.004), higher parental education (βs>0.1067, p<0.0001), more family income (βs>0.0561, p=0.0166), and less family's financial disadvantages (βs<-0.0427, p=0.011). No significant association of *poverty* and supportive parenting behavior with intelligence was found (p>0.05).

More Total and Distress Score PLEs were significantly associated with lower CP and EA PGS (baseline: βs<-0.0316, p=0.0212; 1-year: βs<-0.0422, p=0.0016; 2-year: βs <-0.0489, p=0.0002), higher ADI (baseline: βs>0.0554, p=0.0421), less supportive parenting (baseline: βs<-0.0722, p<0.0001; 1-year: βs<-0.0473, p=0.0013; 2-year: βs<-0.0693, p<0.0001), less positive school environment (baseline: βs<-

0.1076, p<0.0001; 1-year: βs<-0.0896, p=0.0013; 2-year: βs<-0.0847, p<0.0001), lower parental education (baseline: βs<-0.0632, p=0.0011; 1-year: βs<-0.0567, p=0.004; 2-year: βs<-0.0463, p=0.0198), more family's financial disadvantage (1-year: βs>0.0605, p=0.005; 2-year: βs>0.0646, p<0.0001). No significant association of *years of residence*, *poverty*, and family income with Total and Distress Score PLEs was found (p>0.05). Parent-rated PLEs was positively associated with ADI (baseline: β=0.0486, p=0.0378; 1-year: βs>0.0527, p=0.0225) and negatively with positive school environment (baseline: βs<-0.0522, p=0.0018; 2-year: βs<-0.0601, p=0.0009) and family's financial disadvantage (2-year: βs>0.0501, p=0.0216).

Results of IGSCA with European samples

The results of IGSCA in European ancestry samples were similar to those in multiethnic participants (**Table S4**). It showed a good model fit with a GFI of 0.9695, SRMR of 0.0397, and FIT value of 0.4854.

Intelligence was under a significant direct influence of the cognitive phenotypes PGS (β=0.2987 [95% CI=0.2673~0.3281]), neighborhood SES (β=-0.0931 [95% CI=-0.1303~-0.0586]), family SES (β=0.3034 [95% CI=0.2683~0.3413]), and positive environment (β=0.0396 [95% CI=0.0104~0.0698]). Neighborhood SES (β=0.0411 [95% CI=0.0037~0.0784]), family SES (β=-0.0531 [95% CI=-0.0956~-0.0101]), and positive environment (β=-0.1473 [95% CI=-0.1779~-0.1163]) showed significant direct influence on baseline year PLEs, but cognitive phenotypes PGS did not. Constructs that had significant direct effects on PLEs of 1-year and 2-year follow-up were family SES (βs<-0.0850) and positive environment (βs<-0.1222).

Intelligence had a significant mediating effect on the pathways of the PGS (β=-0.0555 [95% CI=-0.0754~-0.0392]), family SES (β=-0.0563 [95% CI=-0.07749~-0.0392]), neighborhood SES (β=0.0173 [95% CI=0.0098~0.0271]), and positive environment (β=-0.0074 [95% CI=-0.0142~-0.0019]) to baseline year PLEs. For PLEs of 1-year and 2-year follow-up, the mediation effects of intelligence were also significant with all four constructs: PGS (1-year: β=-0.0397 [95% CI=-0.0586~-0.0235]; 2-year: β=-0.0204 [95% CI=-0.0389~-0.0036]), family SES (1-year: β=-0.0404 [95% CI=-0.0601~-0.0238]; 2-year: β=-0.0207 [95% CI=-0.0399~-0.0036]), neighborhood SES (1-year: β=0.0124 [95% CI=0.0064~0.0208]; 2-year: β=0.0064 [95% CI=0.0011~0.0132]), and positive environment (1-year: β=-0.0053 [95% CI=-

0.0108~-0.0012]; 2-year: β=-0.0027 [95% CI=-0.0067~-0.0002]). Positive environment had the largest total effects on PLEs among the constructs (baseline year: β=-0.1547; 1-year: β =-0.1274; 2-year: β=-0.1386).

Results of linear mixed models adjusted for schizophrenia PGS with multiethnic samples

We also assessed whether the effects of cognitive phenotypes PGS in the linear mixed model are significant including schizophrenia PGS. The results showed that inclusion of schizophrenia PGS in the models did not change much of the main results (Table S5). Schizophrenia PGS was negatively associated with total and fluid intelligence in EA PGS model (Total intelligence β=-0.0293, 95% CI=-0.0488~-0.01, p=0.0048; Fluid intelligence β=-0.0317, 95% CI=-0.0522~-0.0097, p=0.0062). No significant association was found between schizophrenia PGS and PLEs of all three time points (p>0.05).

Results of linear mixed models adjusted for unobserved confounders with multiethnic samples

Unobserved confounding variables may bias linear regression estimates. In particular, linear estimates may be subject to collider bias when those unobserved confounders and genetic factors are jointly associated with environmental factors and target traits (Akimova et al., 2021). Thus, bias from unobserved confounding variables were adjusted using the null treatments approach (Miao et al., 2022). This approach can identify the causal effects of multiple treatment variables in presence of unobserved confounders. Assuming that fewer than half of the treatments have causal effects on the outcome, it first estimates the joint distribution of treatments-confounders to obtain asymptotic bias using a factor model. Second, it estimates the relationship between treatments, confounders, and outcome using standard density estimation methods (e.g., least squares regression). Finally, by eliminating the asymptotic bias from the estimated treatments-confounders-outcome relationship, it can adjust for unobserved confounding even without specifying which treatments are null.

The significance of effects of PGSs was mostly preserved after adjusting for unobserved confounders (**Table S6**). Higher cognitive phenotypes PGSs correlated significantly with higher intelligence (CP PGS: βs>0.1111, p<0.0001; EA PGS: βs>0.0546, p=0.0002]). CP PGS was associated

with lower baseline year Distress Score PLEs (β= -0.0348, p=0.021) and EA PGS was associated with lower Total and Distress Score PLEs of baseline year and follow-up years (baseline: βs<-0.0534, p=0.0002; 1-year: β=-0.0375, p=0.0149; 2-year: βs<-0.0391, p=0.017). We found no significant associations of neighborhood SES variables with any of the target variables (p>0.05).

While positive parenting behaviors did not have significant association with intelligence, positive school environment was significantly associated with higher total and fluid intelligence (βs>0.0328, p=0.0242). Positive parenting behaviors had significant negative correlations with all types of PLEs of baseline year and follow-up years (baseline: βs<-0.0412, p=0.036; 1-year: βs<-0.0435, p=0.0316; 2-year: βs<-0.0558, p=0.0017). Positive school environment was also negatively associated with all types of PLEs of baseline year and follow-up years (baseline: βs<-0.0464, p=0.036; 1-year: βs<-0.0952, p<0.0001; 2-year: βs<-0.0583, p=0.0095).

Parental education and family income were not significantly associated with all types of intelligence and PLEs of all three time points. Family's financial disadvantage did not show significant association with intelligence and baseline year PLEs, but it had positive associations with 1-year follow-up Total Score (β=0.0604, p=0.0113) and Distress Score PLEs (β=0.0532, p=0.0197) and 2-year follow up Total Score (β>0.0544, p=0.0284) and Distress Score PLEs (β=0.0584, p=0.0224).

Supplementary References

Akimova, E. T., Breen, R., Brazel, D. M., & Mills, M. C. (2021). Gene-environment dependencies lead to collider bias in models with polygenic scores. *Scientific Reports*, *11*(1). doi:10.1038/s41598-021-89020-x

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research, 19*(9), 1655-1664.

Conomos, M. P., Miller, M. B., & Thornton, T. A. (2015). Robust Inference of Population Structure for Ancestry Prediction and Correction of Stratification in the Presence of Relatedness. *Genetic Epidemiology, 39*(4), 276-293.

Conomos, Matthew P., Reiner, Alexander P., Weir, Bruce S., & Thornton, Timothy A. (2016). Model-free Estimation of Recent Genetic Relatedness. *The American Journal of Human Genetics, 98*(1), 127-148.

Das, S., Forer, L., Schönherr, S., Sidore, C., Locke, A. E., Kwong, A., et al. (2016). Next-generation genotype imputation service and methods. *Nature Genetics, 48*(10), 1284-1287.

Diemer, M. A., Mistry, R. S., Wadsworth, M. E., López, I., & Reimers, F. (2013). Best Practices in

Conceptualizing and Measuring Social Class in Psychological Research. *Analyses of Social Issues and Public Policy*, *13*(1), 77-113. doi:10.1111/asap.12001

Karcher, N. R., Barch, D. M., Avenevoli, S., Savill, M., Huber, R. S., Simon, T. J., . . . Loewy, R. L. (2018). Assessment of the Prodromal Questionnaire–Brief Child Version for Measurement of Self-reported Psychotic-like Experiences in Childhood. *JAMA Psychiatry*, *75*(8), 853. doi:10.1001/jamapsychiatry.2018.1334

Karcher, N. R., Schiffman, J., & Barch, D. M. (2021). Environmental Risk Factors and Psychotic-like Experiences in Children Aged 9–10. *Journal of the American Academy of Child & Adolescent Psychiatry*, 60(4), 490-500.

Loh, P.-R., Danecek, P., Palamara, P. F., Fuchsberger, C., A Reshef, Y., K Finucane, H., et al. (2016). Reference-based phasing using the Haplotype Reference Consortium panel. *Nature Genetics, 48*(11), 1443-1448.

Miao, W., Hu, W., Ogburn, E. L., & Zhou, X.-H. (2022). Identifying Effects of Multiple Treatments in the Presence of Unmeasured Confounding. *Journal of the American Statistical Association*, 1-15. doi:10.1080/01621459.2021.2023551

Raj, A., Stephens, M., & Pritchard, J. K. (2014). fastSTRUCTURE: Variational Inference of Population Structure in Large SNP Data Sets. *Genetics, 197*(2), 573-589.

Rakesh, D., Seguin, C., Zalesky, A., Cropley, V., & Whittle, S. (2021). Associations between neighborhood disadvantage, resting-state functional connectivity, and behavior in the Adolescent Brain Cognitive Development (ABCD) Study®: Moderating role of positive family and school environments. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*.