Running Title: Shared brain and genetic architectures between mental health and physical

activity

Wei Zhang[1], Sarah E. Paul[2], Anderson Winkler[3], Ryan Bogdan[2], Janine D. Bijsterbosch[1]

[1]Radiology Department, Washington University School of Medicine, St. Louis
[2]Department of Psychological & Brain Sciences, Washington University in St. Louis
[3]National Institute of Mental Health / National Institutes of Health

8
9
10

Abstract

Physical activity is correlated with, and effectively treats various forms of psychopathology.
However, whether biological correlates of physical activity and psychopathology are shared
remains unclear. Here, we examined the extent to which the neural and genetic architecture of
physical activity and mental health are shared. Using data from the UK Biobank (N=6,389),
canonical correlation analysis was applied to estimate associations between the amplitude and
connectivity strength of sub-networks of three major neurocognitive networks (default mode,
DMN; salience, SN; central executive networks, CEN) with accelerometer-derived measures of
physical activity and self-reported mental health measures (primarily of depression, anxiety
disorders, neuroticism, subjective well-being, and risk-taking behaviors). We estimated the
genetic correlation between mental health and physical activity measures, as well as putative
causal relationships by applying linkage disequilibrium score regression, genomic structural
equational modeling, and latent causal variable analysis to genome-wide association summary
statistics (GWAS N=91,105-500,199). Physical activity and mental health were associated with
connectivity strength and amplitude of the DMN, SN, and CEN (r's≥0.12, p's<0.048). These
neural correlates exhibited highly similar loading patterns across mental health and physical
activity models even when accounting for their shared variance. This suggests a largely shared
brain network architecture between mental health and physical activity. Mental health and
physical activity (including sleep) were also genetically correlated (|rg|=0.085-0.121), but we
found no evidence for causal relationships between them. Collectively, our findings provide
empirical evidence that mental health and physical activity have shared brain and genetic
architectures and suggest potential candidate sub-networks for future studies on brain
mechanisms underlying beneficial effects of physical activity on mental health.

Keywords: mental health, physical exercise, sleep, canonical correlation, multivariate analysis,

genetic correlation

1. Introduction

Mental health and physical activity have both been linked to emotion, cognition, and brain

correlates[1–4]. Furthermore, physical activity is known to improve various psychiatric conditions[5–

7]. Yet, it remains elusive whether mental health and physical activity overlap in neural and

genetic architectures. Here, in a large population sample (n=6,389 from the UK Biobank [8]), we

estimate the extent to which physical activity and general mental health share patterns of resting-

state functional MRI network measures and genetic architectures.

1.1 Potential overlap between mental health and physical activity in neural networks

Functional alterations in large-scale brain networks have been consistently implicated in a wide

range of psychiatric disorders[9]. Dysfunctional configuration of neurocognitive networks such as

the default mode, salience, and central executive networks has been hypothesized to characterize

major psychiatric disorders including depression and anxiety (DMN, SN, CEN)[10]. In line with

this hypothesis, findings from meta-analyses have shown that core cognitive and affective

abnormalities in major depression can be accounted for by hypo-connectivity within the CEN

and hyper-connectivity within the DMN[11], together with hypo-connectivity between the control

systems (i.e., CEN) and salience, emotion processing systems (i.e., SN)[12]. Similarly, a recent

meta-analysis suggests that anxiety disorders are characterized by hypo-connectivity between

subcortical limbic circuits that partially overlap with the SN, CEN, and DMN, as well as

decoupling between the CEN and DMN[13]. Furthermore, the personality trait neuroticism, which

is considered a risk marker for psychopathology, has also been linked to alterations in functional

brain networks[14]. These same networks have also been linked to physical activity. Evidence from

fMRI studies on physical activity demonstrated the changes in activity of and functional

connectivity between these network hub regions including the hippocampus, parahippocampus,

dorsal anterior cingulate cortex, and ventromedial prefrontal cortex that primarily subserve

executive functions such as working memory, attention, and inhibition[15–19]. At the more system

level, a 12-month aerobic walking intervention was found to increase resting-state functional

connectivity between subnetworks of DMN, and between subnetworks of SN[20], whereas

connectivity in the CEN was found to increase after multiple sessions of high intensity interval

training[21]. Interestingly, the intensity of physical exercise appeared to modulate functional

connectivity changes in the hub regions of the CEN[22], as well as in the DMN subsystems acutely

and after 3 months of training[23]. Additionally, although sleep might not be considered a type of

physical activity for its inactive nature, it is closely connected to mental health such that sleep

problems have been considered as a risk factor for subsequent development of depressive

symptoms[24,25], and disrupted sleep is often seen in patients with various types of mood and

anxiety disorders[26–28].

Additionally, sleep disturbance can reduce physical activity levels and increase the risk of

exercise-related injuries[29]. Although sleep has been largely overlooked in the literature for

physical activity in relation to mental health, we included it in the current study for its close

relationships with both mental health and physical activity. Sleep duration can be accurately

measured with ecological sampling approaches such as wrist-worn accelerometers, which have

been adopted in medical and mental health research[30,31]. At the neural network level, sleep

deprivation has been linked to reduction in functional connectivity within the DMN [32–34],

whereas increased sleep duration has been linked to strengthened connectivity within the DMN

but reduced connectivity between DMN and SN[35]. Fluctuations in arousal (i.e., indication of

drowsiness) during resting-state scan acquisitions have also been linked to the fMRI signal

amplitude in sensorimotor networks[36]. These studies together highlight the engagement of the

DMN, SN and CEN in both mental health and physical activity and demonstrate that the

interplays between these large-scale intrinsic networks and subnetworks are coupled with

changes in mental health symptoms and physical exercises. Yet, it remains unclear whether

similar connectivity patterns or signal changes of these networks are shared by mental health and

physical activity.

1.2 Potential overlap between mental health and physical activity in genetic variance

Physical activity is known to promote resilience to various psychiatric conditions, alleviating

symptoms of depression, anxiety, and negative mood[37,38]. This buffering effect may be rooted in

the shared genetic variance between mental health and physical activity. For instance, recent

research employing summary statistics from independent genome-wide associate studies (GWAS)

showed that higher polygenic risk scores for depression are associated with increased odds of

incident depression, whereas self-reported physical activities such as walking, jogging, running,

dancing and yoga appeared to reduce the odds with similar magnitude [39]. This effect has also

been observed using more objective measurement of physical activity such that reduced activity

levels measured by accelerometer were found to associate with diagnoses of schizophrenia,
bipolar disorder, depression, or autism spectrum disorders (ASD) and that healthy participants
without disorder diagnoses were observed to perform less physical activity if they had a higher
polygenic risk score f or schizophrenia, depression, and ASD [40].
Additionally, overlapping genetic architectures may exist across various psychiatric disorders
including anxiety and depression[41–43] as psychiatric phenotypes are highly polygenic [44]. Thus,
latent genetic factors capturing shared variance across clusters of psychiatric symptoms may
improve identification of associations between mental health and physical activity. For instance,
genomic contributions to disorders such as depression and anxiety are captured by a genetic
factor for internalizing disorders that are primarily characterized by these two disorders as
indicated by a confirmatory factor analysis, and this genetic factor is positively genetically
associated with various adverse health outcomes and negatively genetically associated with
physical movement patterns[45]. These findings point to the possibility that the genetic
architectures of mental health and physical activity may overlap. The degree and phenotypic
specificity of overlap remains to be tested.
119
1.3 Shared brain and genetic architectures?
Together, the studies reviewed above show that mental health and physical activity both involve
large-scale brain networks such as the DMN, SN and CEN. Additionally, mental health and
physical activity may have partially overlapping genetic architectures, with evidence showing
associations between genetic liabilities of psychiatric disorders and physical activity, as well as
genetic associations between latent factors of psychopathology and physical activity. As mental
health and physical activity are also tightly related at the behavioral level (e.g., emotion,
cognition), it is reasonable to speculate that these two constructs may partially overlap in the
underlying neurobiological mechanisms. In this study, we aim to determine whether mental
health and physical activity have shared variance in brain and genetic architectures, using brain
network measures and genomic summary statistics.
131
132

2. Methods and Materials

2.1 Participants

The UK Biobank is an openly accessible population dataset with neuroimaging data collection, in addition to extensive demographic, behavioral, lifestyle, and cognitive measures[8,46]. An initial sample of N=8,378 participants from the UK Biobank (UKB) was considered for this study. These participants had participated in accelerometer-based physical activity evaluations and visited the assessment center, where the resting-state fMRI and mental health data were acquired. Data quality assurance resulted in exclusion of N=64 participants for insufficient accelerometer data (see details below in section 2.2.2), and N=1,925 participants with considerable missing data in the mental health questionnaire (see details below in section 2.2.1). The final sample had N=6,389 participants with 2,994 (46.9%) females (sample mean age=63.74 ± SD 7.53). All participants provided informed consent. UK Biobank has ethical approval from the North West Multi-Centre Research Ethics Committee (MREC). Data access was obtained under UK Biobank application ID 47267.

2.2 Data Acquisition and Preprocessing

2.2.1   General mental health measures

The UKB general mental health questionnaire consisting of 41 items was conducted on the same day as fMRI data acquisition (https://biobank.ndph.ox.ac.uk/showcase/label.cgi?id=100060). This self-reported questionnaire primarily measured depression, anxiety, and neuroticism, as well as subjective well-being. These variables had varying degrees of missing data points partly due to responses such as "do not know", "prefer not to answer", or "none of above," and partly due to question dependencies. To ensure robust model estimation while maximizing statistical power, individual variables or questions that had more than 30% missing values were excluded (N=10; see full descriptions for each included individual question in *Table S1*). A multivariate imputation procedure was then leveraged to handle the missingness in the remaining data. By default, this procedure implements multiple imputations with separate imputation models for each incomplete variable[47]. Predictive mean matching (PMM) approach was employed for imputing continuous variables, which first estimates a linear regression model for the target variable (e.g., Y) from all other variables in the data (e.g., non-Y variables) with complete observations. New coefficients are then drawn from the posterior predictive distribution of the

estimated regression coefficients and used to calculate the predicted values for the missing

entries in Y. The predictive values for the observed Y are also calculated using the estimated

regression coefficients. Finally, a small set of candidate donors is formed from the observed Y

(i.e., usually 3 or 5 donors) that have the closest predicted values to the missing Y, and the

observed value from one donor will be randomly selected to replace the missing value[47]. Using

PMM, 20 iterations was performed for each incomplete variable of mental health, and the final

imputed value for any given missing entry was averaged across all iterations.

As the current mental health questionnaire covers a broad range of multiple constructs, including

depression and anxiety symptoms, neuroticism, and subjective well-being, we performed data

decomposition using principal component analysis (PCA) on the imputed data to extract the most

relevant information about general psychopathology, using a R package (see details below in

section 2.4). We retained the top principal components that collectively explained more than 50%

variance of the data in the subsequent statistical analyses. PCA loadings of each individual

question per component can be found in *Table 1*.

178
179

2.2.2   Physical activity measures

Accelerometer data were acquired for a subset of UKB participants during a seven-day

monitoring period (https://biobank.ndph.ox.ac.uk/ukb/label.cgi?id=1008). This enabled real-time

measuring of physical activity for the participants throughout the entire week. Following a

recommendation for quality control [48], data from participants who had less than 72h device

wearing time or had no data in each one-hour period of the 24h cycle were excluded (N=64).

Using a publicly available machine learning algorithm, we extracted measures of five types of

physical activity including sleep, sedentary, walking, light task, and moderate activities[48]. This

algorithm applied random forest and hidden Markov models to a 126-dimental vector that

represented a range of time and frequency domain features for every non-overlapping 30-sec

epoch. The resulting probability of each physical activity was then defined as the count of

predicted activity type per 30-sec epoch divvied by the number of epochs[48]. In addition to these

probability measures, the average acceleration magnitude, and metabolic equivalents of task

(MET) were included to indicate overall activity intensity. The mean values of these features

were calculated across weekdays and weekends, respectively, as well as across the entire

monitoring period (i.e., average over weekdays and weekends). To account for variation in each

individual physical activity measure at different time points, all mean values were standardized

by standard deviations for each participant. In total, 21 standardized physical activity measures

were included in our analyses.

2.2.3   Resting-state fMRI data and brain network measure

Resting-state fMRI was acquired using a multiband sequence with an acceleration factor of 8

(TR=0.735; voxel size=2.4x2.4x2.4mm3). Preprocessing steps included motion correction,

grand-mean intensity normalization, high-pass temporal filtering, unwarping and ICA-FIX

denoising (Alfaro-Almagro et al., 2018). Full details can be found in UK Biobank Brain Imaging

Documentation (https://biobank.ctsu.ox.ac.uk/crystal/crystal/docs/brain_mri.pdf).

For this study, we used IDPs (imaging-derived phenotypes) that were generated and released by

the UKB [46]. Specifically, partial connectivity matrices and network amplitudes from ICA with

dimensionalities of 100 (ICA100) were considered (see *Supplementary Results* for the

comparison with ICA25). ICA is a data-driven approach that can estimate the resting-state brain

networks reliably and reproducibly, including the three networks of interest. It further helps

eliminate noise in the data by separating noise and signal components [49]. These advantages of

ICA make it an unbiased and powerful technique to study the resting-state networks. The

mapping of the DMN, SN and CEN onto the ICA components and the calculation of partial

connectivity matrices and network amplitudes are described below.

2.2.3.1 Mapping the networks of interest

We used the Stanford FIND atlas [50] to construct canonical spatial maps as the reference to

identify the three brain networks of interest, namely the DMN, SN and CEN in our sample.

Notably, the FIND atlas is a functional connectivity-based parcellation atlas, from which we

selected seven functional parcels that are all part of these three intrinsic, including dorsal (i.e.

posterior cingulate cortex and medial prefrontal cortex) and ventral default mode networks (i.e.,

retrosplenial cortex and medial temporal lobe), precuneus network, anterior (i.e., anterior insula

and dorsal anterior cingulate cortex) and posterior salience networks (i.e., posterior insula), as

well as left and right executive control networks (i.e., dorsolateral prefrontal cortex and parietal

cortex in the left and right hemispheres respectively). These selected parcels well represent the

subsystems of the three intrinsic networks that have been associated with physical exercises (e.g.,

via cognitive and interoceptive processing) [5,6] and various psychiatric disorders including

depression and anxiety [7–9]. We therefore selected them as our reference networks. To identify

ICA components that can be mapped onto these networks of interest, we examined fifty-five

signal components that were generated from ICA100 and detected the best matching components

corresponding to each of seven selected network parcels, based on spatial correlations. Seven

ICA components showing the highest spatial correlations with the FIND atlas were identified

(mean r = 0.37). We refer to these ICA components as 'subnetworks' from the large-scale DMN,

SN and CEN as they represented the subsystems of the three intrinsic networks.

2.2.3.2 Extracting subnetwork edges

Dual regression was performed to obtain timeseries for each ICA component or subnetwork at

the individual level [56]. These extracted subnetwork timeseries were then used to calculate the

partial connectivity matrix and amplitude measures of interest. As described in Miller et al.[46],

pairwise partial correlation coefficient were estimated using L2-regularized partial correlations

between all 55 signal components from ICA100, which are deemed to be non-artifactual. This

analysis resulted in a 55x55 partial correlation matrix, from which we selected the correlation

coefficients corresponding to our seven components of interest (i.e., subnetwork edges). In total,

21 subnetwork edges were extracted and included in the subsequent analyses.

2.2.3.3 Calculating subnetwork amplitudes

In additional to pairwise partial correlations between these 7 subnetworks, we further considered

the signal amplitude of each subnetwork as the brain network variables. The amplitude is defined

as the standard deviation of the ICA component timeseries [36]. Previous work has shown that

amplitudes capture the overall signal fluctuations within each subnetwork and can offer

complementary information in relation to behavioral measures. For example, a recent study using

the UK Biobank dataset demonstrated independent associations between network amplitudes and

behavioral measures in addition to connectivity strength [46]. Thus, we also included seven

subnetwork amplitudes in the subsequent analyses.

2.2.3.4 Overview of resting state measures

In summary, subnetwork edges from the partial connectivity matrix indicate the connection
strength between each pair of 7 subnetworks while controlling for all other ICA components,
whereas subnetwork amplitudes capture the variance of signal changes within each subnetwork.
In total, 28 brain measures including 21 partial connectivity strength measures (i.e., subnetwork
edges) and 7 amplitudes were included (see the overlayed subnetworks in *Figure 1;* also see
subnetwork selection in *Figure S1*). All of these resting-state imaging measures are available
from the UK Biobank showcase (bulk field IDs 25753 and 25755).

265
2.3   Statistical Analysis
Separate statistical analyses were performed to examine the shared brain network architecture
and genetic architecture between mental health and physical activity (see *Figure 1* for an
overview for the relevant variables and analyses).

2.3.1   Shared variance between mental health and physical activity
Pearson's correlation was used to identify the shared variance between phenotypes of mental
health and physical activity. As we decomposed the data of mental health into principal
components, all correlations were performed using the individual-specific component scores. We
further calculated the false discovery rate (FDR) to account for multiple testing on all pairwise
correlations between mental health and physical activity phenotypes.

2.3.2   Brain Associations with mental health and physical activity
Canonical Correlation Analysis (CCA) has been recognized as a key tool for population
neuroimaging that allows for investigating associations between imaging and non-imaging
variables[57]. Here in this study, CCA was performed to investigate the associations of brain
measures with physical activity and with mental health separately (i.e., simple CCA models).
Specifically, CCA finds a linear combination of brain measures that is maximally correlated with
a linear combination of mental health or physical activity variables respectively, as defined in $Y *$
$A = U \sim V = X * B$[58] where Y is the set of brain measures, X the set of mental health or physical
activity measures, A and B are the linear weights, and U and V the canonical variables or
canonical variate pair. The canonical correlation for each pair of canonical variates is defined as

288 the correlation between U and V. Canonical loadings that indicate the shared variance between

289 the original observations and canonical variables are calculated as the correlations between Y and

290 U, or between X and V.

291 To further identify unique brain associations with physical activity and with mental health

292 respectively, variance in brain network data explained by one set of variables was partialled out

293 in the CCA model for the other set of variables (i.e., the unique CCA model included physical

294 activity measures as covariates in the model for assessing brain-mental health associations and

295 vice versa). Confounding variables were included in all four CCA models and statistical

296 inference for CCA results was made via 1,000 permutations (i.e., breaking correspondence of

297 participant identity with brain measures and mental health/physical activity measures), as

298 implemented in the permCCA package[57]. Notably, the CCA model with the largest number of

299 non-confounding variables included 28 brain measures and 21 physical activity measures,

300 resulting in a ratio of approximately 130 observations (i.e., individuals) per feature. This is

301 expected to ensure sufficient stability for our study[59].

302 To further investigate whether the patterns of brain measures in relation to mental health and

303 physical activity overlap, post-hoc analyses were carried out to test the significance of canonical

304 loadings for each individual brain variable. Specifically, we aimed to determine whether the

305 same brain measures contributed significantly to the canonical associations both with mental

306 health and physical activity, and thus could indicate a shared brain basis. These analyses were

307 conducted only for significant canonical variates within each individual CCA model using

308 permutation testing, where correspondence between brain measures and mental health/physical

309 activity measures for each individual participant is shuffled. Canonical loading for each brain

310 variable was recorded per permutation, which resulted in separate null distributions of loadings

311 for each brain variable. The loadings from the true (unpermuted) CCA were then compared

312 against the matching null distributions for each individual brain variable. Statistical significance

313 was determined as the proportion of permuted loadings equal or higher than the observed

314 loadings from the unpermuted analysis, divided by the total number of 1,000 permutations.

315 These permutation-derived p values were further corrected for the number of significant

316 canonical variates within each model (i.e., record the permuted loadings across canonical

317 variates). To compare brain variable patterns across different CCA models, we matched the first

318 significant canonical variates from each model based on the correlations between the canonical

variate for the brain measures (i.e., correlating the vector *U* obtained from the mental health

CCA models with the vector *U* obtained from the physical activity CCA models).

To characterize the individual mental health questions and physical activity types in relation to

the tested brain associations, we further examined the loading patterns of each individual

question and physical activity type for the first canonical variate from all models, without testing

for statistical significance.

2.3.3   Shared genetic architecture between mental health and physical activity

Genetic correlations between mental health and physical activity were examined by leveraging

GWAS summary statistics for the relevant phenotypes.

2.3.3.1 Summary statistics

The mental health questionnaire used in this study includes items that measure neuroticism,

anxiety, subjective well-being, depression, and risk taking. We therefore sought to obtain

summary statistics for these psychopathological phenotypes. First, we extracted summary

statistics for *Neuroticism* from a GWAS meta-analysis of self-reported neuroticism in the UKB

(using the same questions as in our study) and Psychiatric Genetics Consortium (using the NEO-

FFI personality inventory)[60]. For *Generalized Anxiety Disorder,* we leveraged summary statistics

from a GWAS of self-reported Generalized Anxiety Disorder 2-item scale scores in the Million

Veteran Program[61]. We further obtained summary statistics for *Subjective Well-Being* from a

GWAS meta-analysis of life satisfaction, positive affect, or both life satisfaction and positive

affect across 59 cohorts[62]. For *Major Depressive Disorder*, we meta-analyzed summary statistics

from case-control GWAS in the UK Biobank and Psychiatric Genomics Consortium[63] and the

Million Veteran Program[64] (see *Supplementary Results* for further details). Lastly, we obtained

summary statistics for *Risk Taking* from a GWAS study using the UKB data, which included the

same question of risk taking as in our study[65]. Although these psychopathological phenotypes

obtained from the independent GWAS studies were not directly equivalent to the phenotypes

derived from mental health questionnaire in our brain association analyses due to different

measurements, the underlying constructs of depression, anxiety disorders, and neuroticism,

subjective well-being and risk-taking are identical. The phenotypes obtained to conduct genetic

association analyses are therefore similar to those in the brain association analyses.

Summary statistics for accelerometer data-derived physical activity phenotypes, including
moderate activity, overall activity, sedentary activity, walking, and sleep duration were derived
from a GWAS of N=91,105 participants of European ancestry in the UK Biobank[48]. The exact
phenotypes were also included in our brain association analyses.

Please refer to *Table 2* for an overview for all the summary statistics used in this study, including
sample size and SNP heritability.

2.3.3.2 Genetic correlations

We used linkage disequilibrium score regression (LDSR) and genomic structural equation
modeling (gSEM) to test whether the genomic architecture associated with general mental
health is shared with physical activity. LDSR leverages GWAS summary statistics to estimate
genetic correlations by regressing the SNP statistics on the SNP linkage disequilibrium (LD)
scores, or correlations between nearby genomic loci due to population stratification (i.e.,
systematic differences in allele frequencies due to differences in ancestry). gSEM characterizes
the latent genetic architecture across phenotypes based on the LDSR-derived genetic correlation
matrices [66]. To this end, we first applied LDSR to existing GWAS summary statistics of
psychopathological phenotypes (i.e., neuroticism, generalized anxiety disorder, subjective well-
being, major depressive disorder, and risk taking) and physical activity phenotypes (i.e., overall
activity, moderate activity, sedentary activity, sleep duration, and walking), respectively, to
estimate pairwise genetic correlations within each construct (i.e., within mental health and within
physical activity respectively). We also examined genetic correlations between mental health
phenotypes and physical activity phenotypes adjusted for sex and BMI. We then applied gSEM
to the covariance matrix of psychopathology and that of physical activity separately, allowing
one single latent factor to load freely within each model. Metrics indicating model fit (i.e., CFI,
comparative fit index; SRMR, standardized root mean squared residual) and factor loadings from
each of these models were used to determine whether one common genetic factor fit the physical
activity and mental health data well, respectively.

Because our results indicated poor model fit for some gSEM analyses (see details below in
Results section 3.3), we focused on the model of mental health, where "risk-taking" was
excluded to generate a latent factor of "negative affect" across other phenotypes. Specifically, we
explored genetic correlations between the latent factor of "negative affect" (i.e., without risk-

taking) and each of the five physical activity phenotypes. In addition, we examined genetic

correlations between "risk-taking" alone and each individual physical activity phenotype, using

LDSR. False discovery rate correction was used to correct for multiple testing (N=10 tests). We

also repeated analyses that returned significant results, with adjustment for sex and BMI.

Adjusted summary statistics from Doherty and colleagues (2018) were used in these analyses.

2.3.3.3 Causal relationships

To examine plausible causal associations between physical activity phenotypes and negative

affect, we conducted Latent Causal Variable Analysis (LCV). This approach finds a latent

variable that mediates the genetic correlation between two traits, such as negative affect and

sleep duration. Generally, if the latent variable has a stronger genetic correlation with trait 1 (e.g.,

sleep duration) than with trait 2 (e.g., negative affect), part of the genetic component of trait 1 is

thought to be causal for trait 2. This partial causality can be quantified using the genetic causality

proportion (GCP) of trait 1 (sleep duration) on trait 2 (negative affect), which can range between

0 (no partial genetic causality) and 1 (full genetic causality) [67]. The advantages of using LCV

over the more traditional Mendelian Randomization for causal inference include increased power

by leveraging SNPs across the genome and less susceptibility to confounding by horizontal

pleiotropy [67].

2.4   Confounding variables

Based on the literature, BMI, smoking and drinking status were included as confounding factors

in all statistical analyses for brain associations[68,69]. Age, sex, head motion during rs-fMRI

acquisition (i.e., mean frame-wise displacement), time difference in days between accelerometer

recording (i.e., start date) and assessment center visit date (i.e., acquisition date for both mental

health questionnaire and rs-fMRI), as well as the scanning site were further included. Due to

varying degrees of missingness in the confound variables (i.e., up to 22%), the same imputation

procedure, as described in *Section 2.2.2*, was performed except that additional usage of

multinomial logistic regression was employed to impute categorical data with more than two

levels (i.e., smoking and drinking status). Complete observations from all variables including the

confounding variables were used in the imputation procedure. When calculating the final

predicted values to replace missing data, the predicted values for continuous variables were

averaged across 20 iterations and the level with the highest count across iterations was selected.

All categorical variables after imputation were dummy coded for subsequent statistical analyses.

In the genetic correlation analyses, to ensure robust test effects, we repeated models with

significant results including sex and BMI as covariates.

416
417
2.5   Data and Code Availability

The UK Biobank data used in this study can be accessed by researchers upon application

(https://www.ukbiobank.ac.uk/register-apply). A machine learning algorithm shared on github

was   used   to   extract   physical   activity   measures

(https://github.com/activityMonitoring/biobankAccelerometerAnalysis). The Matlab code for

running permutation inference for CCA are also available on github, including the new extension

for testing canonical variable loadings (https://github.com/andersonwinkler/PermCCA). We

performed Pearson's correlations and imputations in R (version 4.1.0) [70], using basic *stat*

function and *MICE* package [47], respectively. Genetic analyses including LDSR and gSEM were

performed using the GenomicSEM package also in R [66], with code available on github

(https://github.com/GenomicSEM/GenomicSEM). LCV was performed using the open-source

LCV software (https://github.com/lukejoconnor/LCV). We also shared the derived data from

statistical analyses and code for producing the figures on the Open Science Framework

(https://osf.io/p3fzv/). To note, the GWAS summary statistics used in the current study are hosted

elsewhere, for which we only shared the downloading links.

433
434

3. Results (1,054 words)

3.1   Correlations between mental health and physical activity measures

In this sample, phenotypic measures of mental health and physical activity demonstrated overall

small but significant correlations, with coefficients ranging between –0.08 and 0.11 (FDR-

corrected p-values $\leq$ 0.045). As shown in *Table 1*, the mental health measures (i.e., principal

component scores of the seven principal components) were broadly associated with all physical

activity measures. In particular, sleep showed the strongest association with the first principal

component (PC1) of mental health such that greater sleep duration was associated with poorer

overall mental health (i.e., loadings of PC1 were mostly negative as shown in Table S1) and

walking had the greatest association with the fifth principal component (PC5) of mental health

such that higher depression scores was linked to reduced walking time. The overall coefficient

patterns were largely consistent for correlations between physical activity and mental health

measures despite of minor differences across different time windows (i.e., overall, weekdays,

weekend; see full correlations in *Table 1*).

3.2   Shared neural correlates of mental health and physical activity

3.2.1   Brain associations with mental health and physical activity

In simple CCA models (i.e., without accounting for the potentially shared variance in brain

measures between physical activity and mental health), 2 and 3 significant canonical variate pairs

were observed respectively for brain-mental health (r_1=0.16, p_1=0.001; r_2=0.12, p_2=0.001) and

brain-physical activity associations (r_1=0.23, p_1=0.001; r_2=0.15, p_2=0.001; r_3=0.13, p_3=0.047).

When controlling for the shared variance in brain measures between mental health and physical

activity (i.e., unique models), we found 2 significant canonical variates for both brain-mental

health (r_1=0.14, p_1=0.001; r_2=0.12, p_2=0.002) and brain-physical activity associations (r_1=0.21,

p_1=0.001; r_2=0.15, p_2=0.001), with slightly decreased canonical correlation coefficients

comparing to those from the simple models. The canonical variates for brain measures after

accounting for the shared variance (i.e., from the unique models) mapped well with those from

the simple models, as indicated by correlation coefficients between brain canonical variables (i.e.,

$\square$; all r's > 0.96; see full results in *Table S2*). These results are in line with our expectations that

both mental health and physical activity are closely associated with the functional networks

under investigation and that mental health and physical activity have shared variance in the

functioning of these networks, as indicated by the reduced canonical correlation coefficients

from the unique models.

3.2.2   Canonical loadings of brain measures

Findings from the post-hoc analyses on the significant canonical variates showed considerable

overlap between the brain measures with significant loadings associated with mental health and

with physical activity. Overall, the amplitude of subnetworks that indicates the overall signal

fluctuations in each subnetwork over time loaded higher than the edges, with the highest

loadings on the amplitude of dorsal DMN or left CEN for the first canonical variates in both the

mental health and physical activity models (*permuted p-values* $\leq$ 0.001; *Figure 2*). This

suggested that the magnitude of fluctuations of intra-network signal (i.e., amplitude) had higher

contributions to the observed canonical associations with both mental health and physical

activity than the inter-network connectivity strength. The similar loading pattern of all brain

measures were largely retained even when the shared variance was partialled out for mental

health and for physical activity, respectively, in the unique models (*permuted* p-values $\leq$ 0.035).

In addition to amplitude, connectivity between the dorsal DMN and left CEN (i.e., subnetwork

edge) also exhibited statistically significant loadings in the first canonical variates of both the

mental health and the physical activity models (simple and unique models; *permuted* p-values $\leq$

0.001; *Figure 2*). The brain variable loadings for the second canonical variate were highly

similar between simple and unique models for either the mental health or physical activity

models, with the amplitude and subnetwork edges showing evenly important involvement. Yet,

the patterns of these loadings differed between the mental health and physical activity model

(*Figure S2*).

3.2.3   Canonical loadings of mental health questions and physical activities

Canonical loadings of both individual mental health questions and physical activity types for the

first canonical variate also exhibited similar patterns between the simple and unique models

(*Figure S3*). In the models for brain-mental health associations, "risk taking" and "ever irritable/

argumentative for 2 days" had the highest loadings in both simple and unique models, whereas in

the models for brain-physical activity associations, "walking" in all time windows (i.e., overall,

weekdays, weekend) showed the greatest importance across models. Interestingly, among all

physical activity types, only "sleep" showed the opposite direction in canonical loadings (*Figure*

S3).

3.3   Genetic correlations

3.3.1   Genetic correlations for individual phenotypes

Pairwise LDSR was performed separately for psychopathological phenotypes and physical

activity phenotypes. All results were significant after controlling for multiple comparisons,

except for the genetic correlations between neuroticism and risk-taking ($rg = 0.039$, $p = 0.075$),

and between subjective well-being and risk-taking ($rg = 0.054$, $p = 0.146$; *Figure 3*). Specifically,

among psychopathological phenotypes, the smallest significant genetic correlation was observed

for generalized anxiety disorder and risk-taking ($rg = 0.151$, $p = 2.69\text{e-}05$, *FDR-corrected p-*

value $= 3.36\text{e-}05$), and the largest effect was observed for MDD and generalized anxiety disorder

($rg = 0.768$, $p = 2.62\text{e-}90$, *FDR-corrected p-value* $= 1.31\text{e-}89$). For physical activity, genetic

correlations ranged from -0.217 ($p = 1.03\text{e-}04$, *FDR-corrected p-value* $= 1.03\text{e-}04$) for sleep

duration and walking to 0.796 ($p = 1.77\text{e-}31$, *FDR-corrected p-value* $= 8.85\text{e-}31$) for moderate

and overall activity (*Figure 3*). Physical activity phenotypes adjusted for sex and BMI largely

recapitulated these results, with genetic correlations ranging from -0.209 ($p = 2.49\text{e-}04$, *FDR-*

corrected p-value $= 2.49\text{e-}04$) for sleep duration and walking, to 0.780 ($p = 2.12\text{e-}23$, *FDR-*

corrected p-value $= 1.06\text{e-}22$) for moderate and overall activities.

3.3.2   Genetic correlations for latent factors

Analyses using gSEM across all mental health phenotypes returned results indicating suboptimal

estimations with a low factor loading on risk taking from the model for a general

psychopathology latent factor (standardized loading $= 0.065$), which likely reflected its

conceptually distinct construct from all other phenotypes (i.e., depression, anxiety, neuroticism,

and subjective well-being). Additionally, initial gSEM results for physical activity phenotypes

indicated overall poor model fit (CFI $= 0.569$, SRMR $= 0.160$; *Figure S4*). Thus, we performed

gSEM for psychopathological phenotypes grouped as one "negative affect" latent factor after

excluding "risk-taking", with an effective sample size of 571,170 and good model fit (CFI $=$

0.987, SRMR $= 0.0535$; *Figure S5*). After correction for multiple testing, genetic variance in

negative affect was significantly and positively correlated with sleep duration ($rg = 0.121$, $p =$

1.25e-05, *FDR-corrected p* = 1.25e-04), and negatively correlated with moderate (*rg* = -0.117, *p*

= 1.33e-03, *FDR-corrected p* = 4.43e-03) and overall activity (*rg* = -0.085, *p* = 7.70e-04, *FDR-*

corrected p = 3.85e-03; *Figure 4;* also see the results for a latent negative affect factor including

"risk-taking" in *the Supplementary Results*). No significant genetic correlation was observed

between negative affect and sedentary activity or walking after multiple comparison corrections

(|rgs| ≤ 0.062, p-values ≥ 0.036, FDR-corrected p-values ≥ 0.060; *Table 3*). Post-hoc analyses

revealed that after adjustment for sex and BMI, only the genetic correlation between negative

affect and sleep duration remained significant (*rg* = 0.122, *p* = 9.38e-06). Using LDSR, risk-

taking was genetically correlated with both overall activity and walking (|rgs| ≥ 0.074, *p*s ≤ 0.020,

*ps*FDR ≤ 0.040) but not moderate or sedentary activity or sleep duration (|rgs| ≤ 0.081, *p*s ≥

0.092).

3.3.3 Latent Causal Variable Analysis.

Results from the LCV analyses did not indicate causal relationships between any physical

activity phenotypes and negative affect, in either direction (|GCPs| < 0.425; ps > 0.163).

4. Discussion (1,107 words)

In this study, we investigated whether mental health and physical activity have shared brain and

genetic architectures using the UK Biobank cohort. Our findings showed significant associations

of mental health and physical activity separately with a set of brain measures that represent the

connectivity strength and amplitude of subnetworks from the DMN, SN and CEN. Critically,

these significant associations exhibited highly similar patterns of brain variable loadings across

mental health and physical activity models even when the shared variance between these two

constructs was accounted for, suggesting a potential overlap in brain network architecture

between these two constructs. Further analyses examining genetic correlations for mental health

and physical activity showed that negative affect exhibited significant genetic correlations with

several physical activity types, of which sleep duration demonstrated the strongest genetic

correlation that remained significant after controlling for BMI and sex effects. Together, these

results support the presence of shared multivariate brain and genetic architectures between

mental health and physical activity.

The three intrinsic brain networks, namely the DMN, SN and CEN, have been consistently

implicated in a wide range of psychiatric disorders including major depression and anxiety[10].

Interestingly, connections between or the configurations of these networks have also been

associated with physical exercises[71,72]. The current study therefore focused specifically on the

subnetworks from these large-scale networks and used the amplitude and connectivity strength

(i.e., subnetwork edge) to examine the associations of these networks with mental health and

with physical activity, respectively. In line with the literature, we observed significant

multivariate associations for all three networks with either mental health or physical activity, and

significant loadings on most of the subnetworks. In particular, the dorsal DMN and left CEN

showed the greatest involvement in the observed brain associations with both mental health and

physical activity (*Figure 2*). In this study, the dorsal DMN subnetwork primarily consisted of the

posterior cingulate cortex (PCC) and the ventromedial prefrontal cortex (vmPFC), the two brain

areas that are commonly considered as the core subsystem of the DMN[73]. Similarly, the

subnetwork of the CEN here included two most typical hub regions: the dorsolateral prefrontal

cortex (dlPFC) and posterior parietal cortex (PPC) for each individual hemisphere (*Figure S1*).

These major hubs of the DMN and CEN have been implicated in various mental illnesses

including depression and anxiety. For example, the PCC and mPFC have been suggested to

collectively support multiple cognitive functions including decision making and memory[74], the

impairment of which has often been reported in patients with major depression and anxiety

disorders[75–77]. Additionally, the dlPFC is known to be involved in emotion regulation [78–80] and

dysfunction of this region is often seen in abnormal processing of emotional experiences in

patients with depressive and anxiety symptoms[81,82]. As for physical activity, increased dlPFC

activity has been observed after acute physical exercises in participants with higher scores in the

Stroop test[83], whereas the involvement of the DMN subsystems in the medial temporal lobe

(MTL) including the hippocampus and its connection with the medial PFC are often observed in

relation to enhanced memory after physical exercises[20,84,85]. In our findings, the amplitude of

dorsal DMN and left CEN that indicated the magnitude of fluctuations of intra-network signal

(i.e., variance in the connections between PCC and vmPFC or between dlPFC and PCC), as well

as the connectivity strength between these two subnetworks, showed significantly high loadings

for the most critical association between brain measures and mental health measures (i.e., the

first canonical variate). These observations are in line with separate literature on mental health

and physical activity, and provide empirical evidence that mental health and physical activity

may share brain architecture involved in major cognitive functions.

Interestingly, mental health and physical activity also appear to have partially overlapping

genetic architectures. In line with previous reports that internalizing problems are negatively

genetically correlated with physical movement[45], we showed that a latent negative affect factor

capturing genetic covariance between subjective well-being, neuroticism, major depressive

disorder, and generalized anxiety disorder was negatively genetically correlated with overall

physical activity as well as a more fine-grained phenotype of moderate activity, and positively

genetically correlated with sleep duration. Protective effects of physical activity on mental health

have long been documented, as have negative health consequences of psychiatric disorders[5–7,86].

Here, we demonstrate that these relationships can be partially explained by shared genetic

predisposition, although results from our latent causal variable analysis indicate that these

associations do not reflect causal influences. Interestingly, with the adjustment of sex and BMI,

only the correlation with sleep duration remained significant. Sleep duration also showed the

lowest loading onto a latent physical activity genetic factor (*Figure S4*). These results suggest

that the sleep phenotype is somewhat distinct from the remaining physical activity phenotypes

and that the shared genetic architecture between negative affect and sleep duration is more

pronounced than that between negative affect and the degree of daily physical or sedentary

activity. This is in line with the frequent documentation of symptomatic sleep disturbances

across forms of psychopathology, including depression and anxiety[87,88], even in children[89].

Specifically, this aligns well with the literature showing reduced sleep duration in older adults

with depression and anxiety disorders [90–92].

Despite being the first to jointly investigate the shared brain network architecture and genetic

basis of mental health and physical activity in a large population cohort, our study has some

limitations. First, the brain measures in our study are derived from resting-state fMRI measures.

Although our choice reflected a rich literature that has implicated these measures in both mental

health and physical activity, addressing smaller-scale brain structures (e.g., specific regions) with

relevant hypotheses and inclusion of multimodal brain measures such as structural gray matter

volume, cortical thickness, and white matter integrity can provide complementary insights into

brain architecture in relation to mental health and physical activity, and thus may be of interest

for future investigations. Second, although the accelerometer recording took place prior to the

acquisition of resting-state fMRI and mental health assessment for most of the participants in our

study (i.e., 96%), the degrees of time difference between these measurements varied greatly at

the individual level (i.e., ranging between -473 to 2281 days). This time discrepancy was

accounted for in all CCA models as a covariate and cautions should be taken when interpreting

the observed brain associations with reference to time effects. It should also be noted that the

mean age of the current sample is relatively high as the UKB cohort comprises predominantly

middle-to-late aged individuals. Our findings therefore should be interpreted carefully in the

relevant context. Lastly, socioeconomic variables including education attainment can be relevant

to mental health and physical activity phenotypes and inclusion of these variables as

confounding factors may be considered in future investigations.

In conclusion, our study jointly analyzed resting-state network measures and genetic correlations

in a large cohort to test the hypothesis of a shared neurobiological basis of mental health and

physical activity. Our findings revealed that multivariate patterns of brain correlates were highly

similar between mental health and physical activity and highlighted genetic correlations between

mental health (negative affect) and overall physical activity, moderate activity levels, and sleep

duration. Taken together, these findings point towards neural and genetic mechanisms that may

subserve the protective influence of physical exercise and sleep on mental health.

Acknowledgements
JDB is supported by the NIH (1 R34 NS118618-01) and the McDonnell Center for Systems
Neuroscience. SEP is supported by the NIH (1 F31 AA029934-01).
646

647 **Conflict of Interest**
648 The authors report no biomedical financial interests or potential conflicts of interest.
649
650
651
652

Reference

Zhang Z, Chen W. A Systematic Review of the Relationship Between Physical Activity and Happiness. J. Happiness Stud. 2019. doi:10.1007/s10902-018-9976-0.

Fernandes J, Arida RM, Gomez-Pinilla F. Physical exercise as an epigenetic modulator of brain plasticity and cognition. Neurosci. Biobehav. Rev. 2017. doi:10.1016/j.neubiorev.2017.06.012.

Kays JL, Hurley RA, Taber KH. The dynamic brain: Neuroplasticity and mental health. *J Neuropsychiatry Clin Neurosci* 2012. doi:10.1176/appi.neuropsych.12050109.

Millan MJ, Agid Y, Brüne M, Bullmore ET, Carter CS, Clayton NS *et al.* Cognitive dysfunction in psychiatric disorders: Characteristics, causes and the quest for improved therapy. Nat. Rev. Drug Discov. 2012. doi:10.1038/nrd3628.

Biddle S. Physical activity and mental health: Evidence is growing. World Psychiatry. 2016. doi:10.1002/wps.20331.

Deslandes A, Moraes H, Ferreira C, Veiga H, Silveira H, Mouta R *et al.* Exercise and mental health: Many reasons to move. Neuropsychobiology. 2009. doi:10.1159/000223730.

Heinze K, Cumming J, Dosanjh A, Palin S, Poulton S, Bagshaw AP *et al.* Neurobiological evidence of longer-term physical activity interventions on mental health outcomes and cognition in young people: A systematic review of randomised controlled trials. Neurosci. Biobehav. Rev. 2021. doi:10.1016/j.neubiorev.2020.10.014.

Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J *et al.* UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. *PLoS Med* 2015. doi:10.1371/journal.pmed.1001779.

Woodward ND, Cascio CJ. Resting-state functional connectivity in psychiatric disorders. *JAMA Psychiatry* 2015; **72**: 743–744.

Menon V. Large-scale brain networks and psychopathology: A unifying triple network model. *Trends Cogn Sci* 2011; **15**: 483–506.

Tozzi L, Zhang X, Chesnut M, Holt-Gosselin B, Ramirez CA, Williams LM. Reduced functional connectivity of default mode network subsystems in depression: Meta-analytic evidence and relationship with trait rumination. *NeuroImage Clin* 2021. doi:10.1016/j.nicl.2021.102570.

Kaiser RH, Andrews-Hanna JR, Wager TD, Pizzagalli DA. Large-Scale Network Dysfunction in Major Depressive Disorder. *JAMA Psychiatry* 2015. doi:10.1001/jamapsychiatry.2015.0071.

Xu J, Van Dam NT, Feng C, Luo Y, Ai H, Gu R *et al.* Anxious brain networks: A coordinate-based activation likelihood estimation meta-analysis of resting-state functional connectivity studies in anxiety. Neurosci. Biobehav. Rev. 2019. doi:10.1016/j.neubiorev.2018.11.005.

Servaas MN, Geerligs L, Renken RJ, Marsman JBC, Ormel J, Riese H *et al.* Connectomics and neuroticism: An altered functional network organization. *Neuropsychopharmacology* 2015. doi:10.1038/npp.2014.169.

Colcombe SJ, Kramer AF, Erickson KI, Scalf P, McAuley E, Cohen NJ *et al.* Cardiovascular fitness, cortical plasticity, and aging. *Proc Natl Acad Sci U S A* 2004. doi:10.1073/pnas.0400266101.

Herold F, Aye N, Lehmann N, Taubert M, Müller NG. The contribution of functional magnetic resonance imaging to the understanding of the effects of acute physical exercise on cognition. *Brain Sci* 2020. doi:10.3390/brainsci10030175.

Ikuta T, Loprinzi PD. Association of cardiorespiratory fitness on interhemispheric hippocampal and parahippocampal functional connectivity. *Eur J Neurosci* 2019. doi:10.1111/ejn.14366.

Talukdar T, Nikolaidis A, Zwilling CE, Paul EJ, Hillman CH, Cohen NJ *et al.* Aerobic fitness explains individual differences in the functional brain connectome of healthy young adults. *Cereb Cortex* 2018. doi:10.1093/cercor/bhx232.

Weng TB, Pierce GL, Darling WG, Falk D, Magnotta VA, Voss MW. The Acute Effects of Aerobic Exercise on the Functional Connectivity of Human Brain Networks. *Brain Plast* 2016. doi:10.3233/bpl-160039.

Voss MW, Prakash RS, Erickson KI, Basak C, Chaddock L, Kim JS *et al.* Plasticity of brain networks in a randomized intervention trial of exercise training in older adults. *Front Aging Neurosci* 2010. doi:10.3389/fnagi.2010.00032.

Greeley B, Chau B, Jones CB, Neva JL, Kraeutner SN, Campbell KL *et al.* Multiple bouts of high-intensity interval exercise reverse age-related functional connectivity disruptions without affecting motor learning in older adults. *Sci Rep* 2021. doi:10.1038/s41598-021-96333-4.

Schmitt A, Upadhyay N, Martin JA, Rojas S, Strüder HK, Boecker H. Modulation of Distinct Intrinsic Resting State Brain Networks by Acute Exercise Bouts of Differing Intensity. *Brain Plast* 2019. doi:10.3233/bpl-190081.

Voss MW, Weng TB, Narayana-Kumanan K, Cole RC, Wharff C, Reist L *et al.* Acute exercise effects predict training change in cognition and connectivity. *Med Sci Sports Exerc* 2020. doi:10.1249/MSS.0000000000002115.

Nutt DJ, Wilson S, Paterson L. Sleep disorders as core symptoms of depression. *Dialogues Clin Neurosci* 2008. doi:10.31887/dcns.2008.10.3/dnutt.

Riemann D, Krone LB, Wulff K, Nissen C. Sleep, insomnia, and depression. Neuropsychopharmacology. 2020. doi:10.1038/s41386-019-0411-y.

Cox RC, Olatunji BO. Sleep in the anxiety-related disorders: A meta-analysis of subjective and objective research. Sleep Med. Rev. 2020. doi:10.1016/j.smrv.2020.101282.

Mellman TA. Sleep and Anxiety Disorders. Psychiatr. Clin. North Am. 2006. doi:10.1016/j.psc.2006.08.005.

Rumble ME, White KH, Benca RM. Sleep Disturbances in Mood Disorders. Psychiatr. Clin. North Am. 2015. doi:10.1016/j.psc.2015.07.006.

Chennaoui M, Arnal PJ, Sauvet F, Léger D. Sleep and exercise: A reciprocal issue? Sleep Med. Rev. 2015. doi:10.1016/j.smrv.2014.06.008.

Wainberg M, Jones SE, Beaupre LM, Hill SL, Felsky D, Rivas MA *et al.* Association of accelerometer-derived sleep measures with lifetime psychiatric diagnoses: A cross-sectional study of 89,205 participants from the UK Biobank. *PLOS Med* 2021; **18**: e1003782.

Full KM, Malhotra A, Gallo LC, Kerr J, Arredondo EM, Natarajan L *et al.* Accelerometer-Measured Sleep Duration and Clinical Cardiovascular Risk Factor Scores in Older Women. *Journals Gerontol - Ser A Biol Sci Med Sci* 2020. doi:10.1093/gerona/glz201.

De Havas JA, Parimal S, Soon CS, Chee MWL. Sleep deprivation reduces default mode network connectivity and anti-correlation during rest and task performance. *Neuroimage* 2012. doi:10.1016/j.neuroimage.2011.08.026.

Sämann PG, Tully C, Spoormaker VI, Wetter TC, Holsboer F, Wehrle R *et al.* Increased sleep pressure reduces resting state functional connectivity. *Magn Reson Mater Physics, Biol Med* 2010. doi:10.1007/s10334-010-0213-z.

Yeo BTT, Tandi J, Chee MWL. Functional connectivity during rested wakefulness predicts vulnerability to sleep deprivation. *Neuroimage* 2015. doi:10.1016/j.neuroimage.2015.02.018.

Khalsa S, Mayhew SD, Przezdzik I, Wilson R, Hale J, Goldstone A *et al.* Variability in cumulative habitual sleep duration predicts waking functional connectivity. *Sleep* 2016. doi:10.5665/sleep.5324.

Bijsterbosch J, Harrison S, Duff E, Alfaro-Almagro F, Woolrich M, Smith S. Investigations into within- and between-subject resting-state amplitude variations. *Neuroimage* 2017. doi:10.1016/j.neuroimage.2017.07.014.

Anderson E, Shivakumar G. Effects of exercise and physical activity on anxiety. *Front Psychiatry* 2013. doi:10.3389/fpsyt.2013.00027.

Blumenthal JA, Babyak MA, Doraiswamy PM, Watkins L, Hoffman BM, Barbour KA *et al.* Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med* 2007. doi:10.1097/PSY.0b013e318148c19a.

Choi KW, Zheutlin AB, Karlson RA, Wang MJ, Dunn EC, Stein MB *et al.* Physical activity offsets genetic risk for incident depression assessed via electronic health records in a biobank

762   cohort study. *Depress Anxiety* 2020. doi:10.1002/da.22967.

763   40   Dennison CA, Legge SE, Bracher-Smith M, Menzies G, Escott-Price V, Smith DJ *et al.*
764   Association of genetic liability for psychiatric disorders with accelerometer-assessed physical
765   activity in the UK Biobank. *PLoS One* 2021. doi:10.1371/journal.pone.0249189.

766   41   Anttila V, Bulik-Sullivan B, Finucane HK, Walters RK, Bras J, Duncan L *et al.* Analysis of shared
767   heritability in common disorders of the brain. *Science (80-)* 2018. doi:10.1126/science.aap8757.

768   42   Lee SH, Ripke S, Neale BM, Faraone S V., Purcell SM, Perlis RH *et al.* Genetic relationship
769   between five psychiatric disorders estimated from genome-wide SNPs. *Nat Genet* 2013.
770   doi:10.1038/ng.2711.

771   43   Taylor MJ, Martin J, Lu Y, Brikell I, Lundström S, Larsson H *et al.* Association of Genetic Risk
772   Factors for Psychiatric Disorders and Traits of These Disorders in a Swedish Population Twin
773   Sample. *JAMA Psychiatry* 2019. doi:10.1001/jamapsychiatry.2018.3652.

774   44   Smoller JW, Andreassen OA, Edenberg HJ, Faraone S V., Glatt SJ, Kendler KS. Psychiatric
775   genetics and the structure of psychopathology. Mol. Psychiatry. 2019. doi:10.1038/s41380-017-
776   0010-4.

777   45   Grotzinger AD, Mallard TT, Akingbuwa WA, Ip HF, Adams MJ, Lewis CM *et al.* Genetic
778   architecture of 11 major psychiatric disorders at biobehavioral, functional genomic, and molecular
779   genetic levels of analysis. medRxiv. 2020. doi:10.1101/2020.09.22.20196089.

780   46   Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J *et al.* Multimodal
781   population brain imaging in the UK Biobank prospective epidemiological study. *Nat Neurosci*
782   2016. doi:10.1038/nn.4393.

783   47   van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate imputation by chained equations in R.
784   *J Stat Softw* 2011. doi:10.18637/jss.v045.i03.

785   48   Doherty A, Smith-Byrne K, Ferreira T, Holmes M V., Holmes C, Pulit SL *et al.* GWAS identifies
786   14 loci for device-measured physical activity and sleep duration. *Nat Commun* 2018; **9**.
787   doi:10.1038/s41467-018-07743-4.

788   49   Griffanti L, Douaud G, Bijsterbosch J, Evangelisti S, Alfaro-Almagro F, Glasser MF *et al.* Hand
789   classification of fMRI ICA noise components. *Neuroimage* 2017.
790   doi:10.1016/j.neuroimage.2016.12.036.

791   50   Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD. Decoding Subject-Driven Cognitive
792   States with Whole-Brain Connectivity Patterns. *Cereb Cortex* 2012; **22**: 158–165.

793   51   Voss MW, Nagamatsu LS, Liu-Ambrose T, Kramer AF. Exercise, brain, and cognition across the
794   life span. J. Appl. Physiol. 2011. doi:10.1152/japplphysiol.00210.2011.

795   52   Wallman-Jones A, Perakakis P, Tsakiris M, Schmidt M. Physical activity and interoceptive
796   processing: Theoretical considerations for future research. Int. J. Psychophysiol. 2021.
797   doi:10.1016/j.ijpsycho.2021.05.002.

798   53   Sha Z, Wager TD, Mechelli A, He Y. Common Dysfunction of Large-Scale Neurocognitive
799   Networks Across Psychiatric Disorders. *Biol Psychiatry* 2019.
800   doi:10.1016/j.biopsych.2018.11.011.

801   54   Sha Z, Xia M, Lin Q, Cao M, Tang Y, Xu K *et al.* Meta-connectomic analysis reveals commonly
802   disrupted functional architectures in network modules and connectors across brain disorders.
803   *Cereb Cortex* 2018. doi:10.1093/cercor/bhx273.

804   55   Kolesar TA, Bilevicius E, Wilson AD, Kornelsen J. Systematic review and meta-analyses of
805   neural structural and functional differences in generalized anxiety disorder and healthy controls
806   using magnetic resonance imaging. NeuroImage Clin. 2019. doi:10.1016/j.nicl.2019.102016.

807   56   Nickerson LD, Smith SM, Öngür D, Beckmann CF. Using dual regression to investigate network
808   shape and amplitude in functional connectivity analyses. *Front Neurosci* 2017; **11**.
809   doi:10.3389/fnins.2017.00115.

810   57   Winkler AM, Renaud O, Smith SM, Nichols TE. Permutation inference for canonical correlation
811   analysis. *Neuroimage* 2020. doi:10.1016/j.neuroimage.2020.117065.

58   HOTELLING H. RELATIONS BETWEEN TWO SETS OF VARIATES. *Biometrika* 1936.

doi:10.1093/biomet/28.3-4.321.

59 Helmer M, Warrington S, Mohammadi-Nejad A-R, Ji LJ, Howell A, Rosand B *et al.* On stability
of Canonical Correlation Analysis and Partial Least Squares with application to brain-behavior
associations. *bioRxiv* 2020.

60 Nagel M, Jansen PR, Stringer S, Watanabe K, De Leeuw CA, Bryois J *et al.* Meta-analysis of
genome-wide association studies for neuroticism in 449,484 individuals identifies novel genetic
loci and pathways. *Nat Genet* 2018. doi:10.1038/s41588-018-0151-7.

61 Levey DF, Gelernter J, Polimanti R, Zhou H, Cheng Z, Aslan M *et al.* Reproducible Genetic Risk
Loci for Anxiety: Results From ~200,000 Participants in the Million Veteran Program. *Am J*
Psychiatry 2020. doi:10.1176/appi.ajp.2019.19030256.

62 Okbay A, Baselmans BML, De Neve JE, Turley P, Nivard MG, Fontana MA *et al.* Genetic
variants associated with subjective well-being, depressive symptoms, and neuroticism identified
through genome-wide analyses. *Nat Genet* 2016. doi:10.1038/ng.3552.

63 Howard DM, Adams MJ, Clarke TK, Hafferty JD, Gibson J, Shirali M *et al.* Genome-wide meta-
analysis of depression identifies 102 independent variants and highlights the importance of the
prefrontal brain regions. *Nat Neurosci* 2019. doi:10.1038/s41593-018-0326-7.

64 Levey DF, Stein MB, Wendt FR, Pathak GA, Zhou H, Aslan M *et al.* Bi-ancestral depression
GWAS in the Million Veteran Program and meta-analysis in >1.2 million individuals highlight
new therapeutic directions. *Nat Neurosci* 2021. doi:10.1038/s41593-021-00860-2.

65 Karlsson Linnér R, Biroli P, Kong E, Meddens SFW, Wedow R, Fontana MA *et al.* Genome-wide
association analyses of risk tolerance and risky behaviors in over 1 million individuals identify
hundreds of loci and shared genetic influences. *Nat Genet* 2019. doi:10.1038/s41588-018-0309-3.

66 Grotzinger AD, Rhemtulla M, de Vlaming R, Ritchie SJ, Mallard TT, Hill WD *et al.* Genomic
structural equation modelling provides insights into the multivariate genetic architecture of
complex traits. *Nat Hum Behav* 2019. doi:10.1038/s41562-019-0566-x.

67 O'Connor LJ, Price AL. Distinguishing genetic correlation from causation across 52 diseases and
complex traits. *Nat Genet* 2018. doi:10.1038/s41588-018-0255-0.

68 Jia H, Zack MM, Gottesman II, Thompson WW. Associations of Smoking, Physical Inactivity,
Heavy Drinking, and Obesity with Quality-Adjusted Life Expectancy among US Adults with
Depression. *Value Heal* 2018. doi:10.1016/j.jval.2017.08.002.

69 Metse AP, Clinton-Mcharg T, Skinner E, Yogaraj Y, Colyvas K, Bowman J. Associations between
suboptimal sleep and smoking, poor nutrition, harmful alcohol consumption and inadequate
physical activity ('snap risks'): A comparison of people with and without a mental health
condition in an australian community survey. *Int J Environ Res Public Health* 2021.
doi:10.3390/ijerph18115946.

70 R Core Team (2021). R: A Language and Environment for Statistical Computing. R Found. Stat.
Comput. 2021.

71 Baniqued PL, Gallen CL, Voss MW, Burzynska AZ, Wong CN, Cooke GE *et al.* Brain network
modularity predicts exercise-related executive function gains in older adults. *Front Aging*
Neurosci 2018. doi:10.3389/fnagi.2017.00426.

72 Chaddock-Heyman L, Weng TB, Kienzler C, Weisshappel R, Drollette ES, Raine LB *et al.* Brain
Network Modularity Predicts Improvements in Cognitive and Scholastic Performance in Children
Involved in a Physical Activity Intervention. *Front Hum Neurosci* 2020.
doi:10.3389/fnhum.2020.00346.

73 Smallwood J, Bernhardt BC, Leech R, Bzdok D, Jefferies E, Margulies DS. The default mode
network in cognition: a topographical perspective. Nat. Rev. Neurosci. 2021. doi:10.1038/s41583-
021-00474-4.

74 Wang S, Tepfer LJ, Taren AA, Smith D V. Functional parcellation of the default mode network: a
large-scale meta-analysis. *Sci Rep* 2020. doi:10.1038/s41598-020-72317-8.

75 Mirza SS, Ikram MA, Bos D, Mihaescu R, Hofman A, Tiemeier H. Mild cognitive impairment and
risk of depression and anxiety: A population-based study. *Alzheimer's Dement* 2017.

doi:10.1016/j.jalz.2016.06.2361.

76   Perini G, Ramusino MC, Sinforiani E, Bernini S, Petrachi R, Costa A. Cognitive impairment in
depression: Recent advances and novel treatments. *Neuropsychiatr Dis Treat* 2019.
doi:10.2147/NDT.S199746.

77   Rock PL, Roiser JP, Riedel WJ, Blackwell AD. Cognitive impairment in depression: A systematic
review and meta-analysis. Psychol. Med. 2014. doi:10.1017/S0033291713002535.

78   Delgado MR, Nearing KI, LeDoux JE, Phelps EA. Neural Circuitry Underlying the Regulation of
Conditioned Fear and Its Relation to Extinction. *Neuron* 2008. doi:10.1016/j.neuron.2008.06.029.

79   Morawetz C, Bode S, Baudewig J, Kirilina E, Heekeren HR. Changes in Effective Connectivity
Between Dorsal and Ventral Prefrontal Regions Moderate Emotion Regulation. *Cereb Cortex*
2016. doi:10.1093/cercor/bhv005.

80   Ochsner KN, Gross JJ. The cognitive control of emotion. Trends Cogn. Sci. 2005.
doi:10.1016/j.tics.2005.03.010.

81   Koenigs M, Grafman J. The functional neuroanatomy of depression: Distinct roles for
ventromedial and dorsolateral prefrontal cortex. Behav. Brain Res. 2009.
doi:10.1016/j.bbr.2009.03.004.

82   Nejati V, Majdi R, Salehinejad MA, Nitsche MA. The role of dorsolateral and ventromedial
prefrontal cortex in the processing of emotional dimensions. *Sci Rep* 2021. doi:10.1038/s41598-
021-81454-7.

83   Yanagisawa H, Dan I, Tsuzuki D, Kato M, Okamoto M, Kyutoku Y *et al.* Acute moderate exercise
elicits increased dorsolateral prefrontal activation and improves cognitive performance with
Stroop test. *Neuroimage* 2010. doi:10.1016/j.neuroimage.2009.12.023.

84   Li MY, Huang MM, Li SZ, Tao J, Zheng GH, Chen LD. The effects of aerobic exercise on the
structure and function of DMN-related brain regions: a systematic review. Int. J. Neurosci. 2017.
doi:10.1080/00207454.2016.1212855.

85   Tao J, Liu J, Egorova N, Chen X, Sun S, Xue X *et al.* Increased hippocampus-medial prefrontal
cortex resting-state functional connectivity and memory function after Tai Chi Chuan practice in
elder adults. *Front Aging Neurosci* 2016. doi:10.3389/fnagi.2016.00025.

86   De Hert M, Correll CU, Bobes J, Cetkovich-Bakmas M, Cohen DAN, Asai I *et al.* Physical illness
in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in
health care. World Psychiatry. 2011. doi:10.1002/j.2051-5545.2011.tb00014.x.

87   Fang H, Tu S, Sheng J, Shao A. Depression in sleep disturbance: A review on a bidirectional
relationship, mechanisms and treatment. J. Cell. Mol. Med. 2019. doi:10.1111/jcmm.14170.

88   Cox RC, Olatunji BO. A systematic review of sleep disturbance in anxiety and related disorders. J.
Anxiety Disord. 2016. doi:10.1016/j.janxdis.2015.12.001.

89   Cheng W, Rolls E, Gong W, Du J, Zhang J, Zhang XY *et al.* Sleep duration, brain structure, and
psychiatric and cognitive problems in children. *Mol Psychiatry* 2021. doi:10.1038/s41380-020-
0663-2.

90   Brenes GA, Miller ME, Stanley MA, Williamson JD, Knudson M, McCall WV. Insomnia in older
adults with generalized anxiety disorder. *Am J Geriatr Psychiatry* 2009.
doi:10.1097/JGP.0b013e3181987747.

91   Lippman S, Gardener H, Rundek T, Seixas A, Elkind MSV, Sacco RL *et al.* Short sleep is
associated with more depressive symptoms in a multi-ethnic cohort of older adults. *Sleep Med*
2017. doi:10.1016/j.sleep.2017.09.019.

92   Li Y, Wu Y, Zhai L, Wang T, Sun Y, Zhang D. Longitudinal Association of Sleep Duration with
Depressive Symptoms among Middle-aged and Older Chinese. *Sci Rep* 2017. doi:10.1038/s41598-
017-12182-0.

Figure Legends
914

Figure 1.   Overview of variables and analyses. Separate analyses were conducted for brain (left panel) and
genetic associations (right panel). For brain associations, canonical correlation analysis (CCA) was employed for
mental health (MH) and physical activity (PA) separately. Simple and unique CCA models only differed in whether
the shared variance between MH and PA was accounted for. For genetic associations, GWAS summary statistics for
5 mental health phenotypes (MHP_1-MHP_5) and for 5 physical activity phenotypes (PAP_1-PAP_5) were leveraged into
pairwise linkage disequilibrium score regression (LDSR) analyses in MHP and PAP models separately. Genomic
structural equation modeling (gSEM) was also employed to identify genetic associations between a latent factor
from MH phenotypes and each of PA phenotype, followed by latent causal variable analysis (LCV) that allows for
inferring causal genetic relationships among MH and PA phenotypes.
924

Figure 2. Canonical loadings of brain measures on the first canonical variates. These loadings represent the
linear correlation between the original brain measures (Y) and the first canonical variate (U) per model. Color
coding was made for brain variable names along the Y axis (i.e., subnetwork edges in gray with "-" between
subnetwork names and amplitude in orange), and for the bars representing canonical loadings (i.e., significance in
cyan, insignificance in yellow). Simple and unique models differ in whether the model accounted for the shared
variance in brain measures between mental health and physical activity. vDMN = ventral default mode network; R/L
CEN = right/ left central executive network; PCu = precuneus; a/p SN = anterior/posterior salience network.
933

935 **Figure 3. Heatmap of bivariate genetic correlations.** Pairwise genetic correlations were calculated separately for
936 phenotypes of mental health (A) and those of physical activity (B). rg = genetic correlation coefficient, GAD =
937 Generalized Anxiety Disorder, MDD = Major Depressive Disorder, Risk = Risk Tolerance, SWB = Subjective Well-
938 Being.
939

941 **Figure 4. Genetic correlations between negative affect and physical activity phenotypes.** Significant genetic
942 correlations with negative affect were observed for moderate (A), overall activities (B), and sleep duration (C). The
943 latent negative affect factor loaded positively on neuroticism, major depression disorder (MDD) and general anxiety
944 disorder (GAD) phenotypes, but negatively on the subjective well-being (SWB) phenotype.
945
946
947
948

949 **Tables**
950
951 **Table 1.** Correlations between mental health and physical activity measures.

Physical Activity		Mental Health						
		PC1	PC2	PC3	PC4	PC5	PC6	PC7
Overall	Sleep Duration	-0.08*	0.04*	0.01	0.00	0.04*	-0.01	0.03
	Sedentary	-0.03*	0.07*	0.07*	-0.02	-0.02	-0.03	0.03*
	Light Tasks	0.02	-0.02	0.01	-0.01	-0.04*	-0.01	0.00
	Moderate	-0.02	-0.06*	0.02	0.01	0.03*	0.04*	0.01
	Walking	0.01	-0.01	0.02	-0.03*	-0.06*	0.00	0.04*
	Overall Activity	-0.06*	0.04*	0.03*	-0.02	0.04*	0.01	0.08*
	MET	-0.06*	0.10*	0.07*	-0.03	-0.01	-0.05*	0.02
Weekdays	Sleep Duration	-0.08*	0.04*	0.02	0.00	0.03*	-0.01	0.03*
	Sedentary	-0.05*	0.08*	0.08*	-0.01	-0.04*	-0.02	0.04*
	Light Tasks	0.01	-0.02	0.03	-0.03	-0.03*	-0.01	0.01
	Moderate	-0.02	-0.05*	0.03	0.01	0.02	0.03*	0.03
	Walking	0.02	0.00	0.04*	-0.04*	-0.09*	-0.01	0.04*
	Overall Activity	-0.05*	0.04*	0.06*	-0.04*	-0.01	0.00	0.07*
	MET	-0.06*	0.11*	0.08*	-0.03	-0.07*	-0.04*	0.02
Weekend	Sleep Duration	-0.06*	0.04*	0.00	0.00	0.02	-0.01	0.02
	Sedentary	-0.02	0.05*	0.06*	-0.02	-0.01	-0.02	0.02
	Light Tasks	0.02	-0.01	0.00	-0.01	-0.05*	-0.02	-0.02
	Moderate	-0.02	-0.04*	0.01	0.00	0.01	0.04*	0.00
	Walking	0.02	0.00	0.02	-0.03*	-0.06*	0.00	0.04*
	Overall Activity	-0.05*	0.04*	0.03	-0.02	0.03*	0.01	0.06*
	MET	-0.05*	0.08*	0.05*	-0.02	0.01	-0.04*	0.02

952 Phenotypes of mental health were represented by the first seven principal components (i.e., PC1-PC7)
953 that altogether explained 51.05% variance. Significant correlations are indicated by * after FDR
954 corrections (i.e., all corrected p's≤0.045) and negative correlation coefficients are highlighted with light
955 grey shading
956
957
958 **Table 2. Summary statistics from GWAS studies.**

Reference	Phenotype	Dataset	Sample Size	SNP h^2
Nagel et al 2018[60]	Neuroticism	UK Biobank & PGC	390,279	0.10
Levey et al 2020[61]	GAD	Million Veteran Program	199,611	0.056
Okbay et al 2016[62]	SWB	Meta-analysis across 59 cohorts	298,420	0.040
Howard et al 2019[63]	MDD	UK Biobank & PGC	500,199	0.11
Levey et al 2021[64]		Million Veteran Program	250,215	
Linnér et al 2019[65]	Risk-taking	UK Biobank	431,126	0.050

Doherty et al 2018[48]	Physical activities	UK Biobank	91,105	0.10 – 0.21

GAD = general anxiety disorder; SWB = subjective well-being; MDD = major depressive disorder; PGC = Psychiatric Genetics Consortium.

Table 3. Genetic correlations between mental health and physical activity phenotypes.

Mental Health Phenotype	**Physical Activity Phenotype**	*rg*	*p*	*pFDR*
Risk Tolerance	Sleep Duration	-0.029	0.433	0.447
	Sedentary	-0.028	0.447	0.447
	Moderate	0.081	0.092	0.131
	Walking	**0.115**	4.95e-3	0.012
	Overall	**0.074**	0.020	0.040
Negative Affect	Sleep Duration	**0.121**	1.25e-5	1.25e-4
	Sedentary	*-0.062*	0.036	0.060
	Moderate	**-0.117**	1.33e-3	4.43e-3
	Walking	-0.029	0.383	0.447
	Overall	**-0.085**	7.70e-4	3.85e-3

Note, Significant results after FDR corrections were highlighted in bold. *rg = genetic correlation, pFDR = FDR-corrected p-value.*