Running Title: Shared brain and genetic architectures between mental health and physical activity

Wei Zhang[1], Sarah E. Paul[2], Anderson Winkler[3], Ryan Bogdan[2], Janine D. Bijsterbosch[1]

[1]Radiology Department, Washington University School of Medicine, St. Louis
[2]Department of Psychological & Brain Sciences, Washington University in St. Louis
[3]National Institute of Mental Health / National Institutes of Health

Abstract

Physical activity is correlated with, and effectively treats various forms of psychopathology. However, whether biological correlates of physical activity and psychopathology are shared remains unclear. Here, we examined the extent to which the neural and genetic architecture of physical activity and mental health are shared. Using data from the UK Biobank (N=6,389), canonical correlation analysis was applied to estimate associations between the amplitude and connectivity strength of sub-networks of three major neurocognitive networks (default mode, DMN; salience, SN; central executive networks, CEN) with accelerometer-derived measures of physical activity and self-reported mental health. We estimated the genetic correlation between mental health and physical activity measures, as well as putative causal relationships by applying linkage disequilibrium score regression, genomic structural equational modeling, and latent causal variable analysis to genome-wide association summary statistics (GWAS N=91,105-500,199). Physical activity and mental health were associated with connectivity strength and amplitude of the DMN, SN, and CEN (r's≥0.12, p's<0.048). These neural correlates exhibited highly similar loading patterns across mental health and physical activity models even when accounting for their shared variance. This suggests a largely shared brain network architecture between mental health and physical activity. Mental health and physical activity were also genetically correlated (|rg|=0.085-0.121), but we found no evidence for causal relationships between them. Collectively, our findings provide empirical evidence that mental health and physical activity have shared brain and genetic architectures and suggest potential candidate sub-networks for future studies on brain mechanisms underlying beneficial effects of physical activity on mental health.

Keywords: mental health, physical exercise, sleep, canonical correlation, multivariate analysis, genetic correlation

1. Introduction

Mental health and physical activity have both been linked to emotion, cognition, and brain correlates[1–4]. Furthermore, physical activity is known to improve various psychiatric conditions[5–7]. Yet, it remains elusive whether mental health and physical activity overlap in neural and genetic architectures.

1.1 Neural correlates of mental health

In addition to structural alterations such as cortical thickness and volumetric sizes[8–10], functional alterations in large-scale brain networks have been consistently implicated in a wide range of psychiatric disorders[11]. Dysfunctional configuration of neurocognitive networks such as the default mode, salience, and central executive networks has been hypothesized to characterize major psychiatric disorders including depression and anxiety (DMN, SN, CEN)[12]. In line with this hypothesis, findings from meta-analyses have shown that core cognitive and affective abnormalities in major depression can be accounted for by hypo-connectivity within the CEN and hyper-connectivity within the DMN[13], together with hypo-connectivity between the control systems (i.e., CEN) and salience, emotion processing systems (i.e., SN)[14]. Similarly, a recent meta-analysis suggests that anxiety disorders are characterized by hypo-connectivity between subcortical limbic circuits that partially overlap with the SN, CEN, and DMN, as well as decoupling between the CEN and DMN[15]. Furthermore, the personality trait neuroticism, which is considered a risk marker for psychopathology, has also been linked to alterations in functional brain networks[16].

1.2 Genetic associations with mental health

Evidence from genome-wide association studies (GWAS) and family-based studies have suggested overlapping genetic architectures across various psychiatric disorders, including anxiety and depression[17–19], which is in line with the highly polygenic nature of psychiatric phenotypes [20]. As such, latent genetic factors appear to capture shared variance across clusters of psychiatric symptoms and improve identification of independent genomic loci[21]. For instance, genomic contributions to disorders such as depression and anxiety are captured by an internalizing genetic factor that is negatively genetically correlated with other relevant health phenotypes, including adverse health outcomes and physical movement patterns[22].

1.3 Neural correlates of physical activity

Physical activity is known to promote resilience to various psychiatric conditions, alleviating

symptoms of depression, anxiety, and negative mood[23,24]. At the neural level, physical activity

has been shown to promote brain plasticity, leading to structural changes in cerebellum, motor

cortex, and hippocampus among other structures [25–27]. Furthermore, fMRI studies of physical

activity have reported changes in activity of and functional connectivity between brain areas

including hippocampus, parahippocampus, dorsal anterior cingulate cortex, and ventromedial

prefrontal cortex that primarily subserve executive functions such as memory, attention, and

inhibition[28–32]. At the network level, a 12-month aerobic walking intervention was found to

increase resting-state functional connectivity between DMN nodes[33], whereas connectivity in the

CEN was found to increase after multiple sessions of high intensity interval training[34].

Interestingly, the intensity of physical exercise appeared to modulate functional connectivity

changes in the hub regions of the CEN[35], as well as in the DMN subsystems acutely and after 3

months of training[36]. Additionally, although sleep might not be considered a type of physical

activity for its inactive nature, it is closely connected to mental health such that sleep problems

have been considered as a risk factor for subsequent development of depressive symptoms[37,38],

and disrupted sleep is often seen in patients with various types of mood and anxiety disorders[39–

41]. Sleep duration can be accurately measured with ecological sampling approaches such as

wrist-worn accelerometers, which have been adopted in medical and mental health research[42,43].

At the neural network level, sleep deprivation has been linked to reduction in functional

connectivity within the DMN [44–46], whereas increased sleep duration has been linked to

strengthened connectivity within the DMN but reduced connectivity between DMN and SN[47].

Fluctuations in arousal (i.e., indication of drowsiness) during resting-state scan acquisitions have

also been linked to the fMRI signal amplitude in sensorimotor networks[48].

1.4 Genetic associations of physical activity

Physical activity has been found to be heritable in twin and family studies, with genetic factors

explaining roughly 20-30% variance in physical activity or inactivity[49–52]. This heritability

appears to be even higher (i.e., up to 60%) when physical activity is measured by an objective

device such as acceleromete[53]. Genetic factors may explain different degrees of variance in

physical activity depending on sex[54] and activity duration[55]. GWAS and candidate gene studies

have also identified specific genomic loci implicated in physical activity, with a recent GWAS

conducted in the UK Biobank identifying 14 independent loci and gene pathways enriched in

neurological disease, brain structure, and cognitive function[52,56–58].

1.5 Shared brain and genetic architectures?

The studies reviewed above showed that mental health and physical activity both involve large-

scale brain networks such as the DMN, SN and CEN. Additionally, mental health and physical

activity may have partially overlapping genetic architectures, with preliminary evidence showing

a significant genetic correlation between internalizing problems and physical movement[22]. As

mental health and physical activity are also tightly related at the behavioral level (e.g., emotion,

cognition), it is reasonable to speculate that these two constructs may partially overlap in the

underlying neurobiological mechanisms. In this study, we aim to determine whether mental

health and physical activity have shared variance in brain and genetic architectures.

2. Methods and Materials

2.1 Participants

The UK Biobank is an openly accessible population dataset with neuroimaging data collection, in addition to extensive demographic, behavioral, lifestyle, and cognitive measures[59,60]. An initial sample of N=8,378 participants from the UK Biobank (UKB) was considered for this study. These participants had participated in accelerometer-based physical activity evaluations and visited the assessment center, where the resting-state fMRI and mental health data were acquired. Data quality assurance resulted in exclusion of N=64 participants for insufficient accelerometer data (see details below in section 2.2.2), and N=1,925 participants with considerable missing data in the mental health questionnaire (see details below in section 2.2.1). The final sample had N=6,389 participants with 2,994 (46.9%) females (sample mean age=63.74 ± SD 7.53). All participants provided informed consent. UK Biobank has ethical approval from the North West Multi-Centre Research Ethics Committee (MREC). Data access was obtained under UK Biobank application ID 47267.

2.2 Data Acquisition and Preprocessing

2.2.1 General mental health measures

The UKB general mental health questionnaire consisting of 41 items was conducted on the same day as fMRI data acquisition (https://biobank.ndph.ox.ac.uk/showcase/label.cgi?id=100060). This self-reported questionnaire primarily measured depression, anxiety, and neuroticism, as well as subjective well-being. These variables had varying degrees of missing data points partly due to responses such as "do not know", "prefer not to answer", or "none of above," and partly due to question dependencies. To ensure robust model estimation while maximizing statistical power, individual variables or questions that had more than 30% missing values were excluded (N=10; see full descriptions for each included individual question in *Table S1*). A multivariate imputation procedure was then leveraged to handle the missingness in the remaining data. By default, this procedure implements multiple imputations with separate imputation models for each incomplete variable[61]. Predictive mean matching (PMM) approach was employed for imputing continuous variables, which first estimates a linear regression model for the target variable (e.g., Y) from all other variables in the data (e.g., non-Y variables) with complete observations. New coefficients are then drawn from the posterior predictive distribution of the

estimated regression coefficients and used to calculate the predicted values for the missing

entries in Y. The predictive values for the observed Y are also calculated using the estimated

regression coefficients. Finally, a small set of candidate donors is formed from the observed Y

(i.e., usually 3 or 5 donors) that have the closest predicted values to the missing Y, and the

observed value from one donor will be randomly selected to replace the missing value[61]. Using

PMM, 20 iterations was performed for each incomplete variable of mental health, and the final

imputed value for any given missing entry was averaged across all iterations.

As the current mental health questionnaire covers a broad range of multiple constructs, including

depression and anxiety symptoms, neuroticism, and subjective well-being, we performed data

decomposition using principal component analysis (PCA) on the imputed data to extract the most

relevant information about general psychopathology, using a R package (see details below in

section 2.4). We retained the top principal components that collectively explained more than

50% variance of the data in the subsequent statistical analyses. PCA loadings of each individual

question per component can be found in *Table S1*.

164
165
2.2.2   Physical activity measures

Accelerometer data were acquired for a subset of UKB participants during a seven-day

monitoring period (https://biobank.ndph.ox.ac.uk/ukb/label.cgi?id=1008). This enabled real-time

measuring of physical activity for the participants throughout the entire week. Following a

recommendation for quality control [58], data from participants who had less than 72h device

wearing time or had no data in each one-hour period of the 24h cycle were excluded (N=64).

Using a publicly available machine learning algorithm, we extracted measures of five types of

physical activity including sleep, sedentary, walking, light task, and moderate activities[58]. This

algorithm applied random forest and hidden Markov models to a 126-dimental vector that

represented a range of time and frequency domain features for every non-overlapping 30-sec

epoch. The resulting probability of each physical activity was then defined as the count of

predicted activity type per 30-sec epoch divvied by the number of epochs[58]. In addition to these

probability measures, the average acceleration magnitude, and metabolic equivalents of task

(MET) to were included to indicate overall activity intensity. The mean values of these features

were calculated across weekdays and weekends, respectively, as well as across the entire

monitoring period (i.e., average over weekdays and weekends). To account for variation in each

individual physical activity measure at different time points, all mean values were standardized

by standard deviations for each participant. In total, 21 standardized physical activity measures

were included in our analyses.

2.2.3 Resting-state fMRI data and brain network measure

Resting-state fMRI was acquired using a multiband sequence with an acceleration factor of 8

(TR=0.735; voxel size=2.4x2.4x2.4mm3). Preprocessing steps included motion correction,

grand-mean intensity normalization, high-pass temporal filtering, unwarping and ICA-FIX

denoising (Alfaro-Almagro et al., 2018). Full details can be found in UK Biobank Brain Imaging

Documentation (https://biobank.ctsu.ox.ac.uk/crystal/crystal/docs/brain_mri.pdf).

For this study, we used IDPs (imaging-derived phenotypes) that were generated and released by

the UKB (Miller et al., 2016). Specifically, partial connectivity matrices from ICA with

dimensionalities of 100 and 25 were considered (i.e., ICA100 and ICA25). Based on the

literature, seven nodes or functional parcels in total were selected from the Stanford FIND atlas

[63] to construct canonical spatial maps for the three brain networks of interest: the DMN, SN and

CEN. Fifty-five and twenty-one signal components derived from ICA100 and ICA25

respectively were examined to identify the best matching components corresponding to each of

seven selected network nodes in terms of spatial correlations. In contrast to ICA25, components

from ICA100 exhibited higher spatial correlations with the atlas on average (mean r = 0.37) with

less spatial overlap between the components. Thus, seven ICA components from ICA100 were

identified as the network nodes in our study sample. Pairwise partial correlations between (i.e.,

node edges), and amplitudes (i.e., temporal fluctuations) of these 7 nodes were used as the brain

network variables, resulting in 21 node edges and 7 amplitudes (*see the selected nodes in Figure*

S1).

2.3   Statistical Analysis

Separate statistical analyses were performed to examine the shared brain network architecture

and genetic architecture between mental health and physical activity (see *Figure 1* for an

overview for the relevant variables and analyses).

212
213 **Figure 1. Overview of variables and analyses.** Different sets of variables were involved in independent models for
214 testing brain and genetic associations, respectively. CCA: canonical correlation analysis; LDSR: linkage
215 disequilibrium score regression; gSEM: genomic structural equation modeling; LCV: latent causal variable analysis.
216

2.3.1 Shared variance between mental health and physical activity

Pearson's correlation was used to identify the shared variance between phenotypes of mental

health and physical activity. As we decomposed the data of mental health into principal

components, all correlations were performed using the individual-specific component scores. We

further calculated the false discovery rate (FDR) to account for multiple testing on all pairwise

correlations between mental health and physical activity phenotypes.

2.3.2 Brain Associations with mental health and physical activity

Canonical Correlation Analysis (CCA) has been recognized as a key tool for population

neuroimaging that allows for investigating associations between imaging and non-imaging

variables[64]. Here in this study, CCA was performed to investigate the associations of brain

measures with physical activity and with mental health separately (i.e., simple CCA models).

Specifically, CCA finds a linear combination of brain measures that is maximally correlated with

a linear combination of mental health or physical activity variables respectively, as defined in $Y *$

$A = U \sim V = X * B$[65] where Y is the set of brain measures, X the set of mental health or physical

activity measures, A and B are the linear weights, and U and V the canonical variables or
canonical variate pair. The canonical correlation for each pair of canonical variates is defined as
the correlation between U and V. Canonical loadings that indicate the shared variance between
the original observations and canonical variables are calculated as the correlations between Y and
U, or between X and V.

To further identify unique brain associations with physical activity and with mental health
respectively, variance in brain network data explained by one set of variables was partialled out
in the CCA model for the other set of variables (i.e., the unique CCA model included physical
activity measures as covariates in the model for assessing brain-mental health associations and
vice versa). Confounding variables were included in all four CCA models and statistical
inference for CCA results was made via 1,000 permutations (i.e., breaking correspondence of
participant identity with brain measures and mental health/physical activity measures), as
implemented in the permCCA package[64]. Notably, the CCA model with the largest number of
non-confounding variables included 28 brain measures and 21 physical activity measures,
resulting in a ratio of approximately 130 observations (i.e., individuals) per feature. This is
expected to ensure sufficient stability for our study[66].

To further investigate whether the patterns of brain measures in relation to mental health and
physical activity overlap, post-hoc analyses were carried out to test the significance of canonical
loadings for each individual brain variable. Specifically, we aimed to determine whether the
same brain measures contributed significantly to the canonical associations both with mental
health and physical activity, and thus could indicate a shared brain basis. These analyses were
conducted only for significant canonical variates within each individual CCA model using
permutation testing, where correspondence between brain measures and mental health/physical
activity measures for each individual participant is shuffled. Canonical loading for each brain
variable was recorded per permutation, which resulted in separate null distributions of loadings
for each brain variable. The loadings from the true (unpermuted) CCA were then compared
against the matching null distributions for each individual brain variable. Statistical significance
was determined as the proportion of permuted loadings equal or higher than the observed
loadings from the unpermuted analysis, divided by the total number of 1,000 permutations.
These permutation-derived p values were further corrected for the number of significant
canonical variates within each model (i.e., record the permuted loadings across canonical

variates). To compare brain variable patterns across different CCA models, we matched the first

significant canonical variates from each model based on the correlations between the canonical

variate for the brain measures (i.e., correlating the vector □ obtained from the mental health

CCA models with the vector □ obtained from the physical activity CCA models).

To characterize the individual mental health questions and physical activity types in relation to

the tested brain associations, we further examined the loading patterns of each individual

question and physical activity type for the first canonical variate from all models, without testing

for statistical significance.

2.3.3   Shared genetic architecture between mental health and physical activity

Genetic correlations between mental health and physical activity were examined by leveraging

GWAS summary statistics for the relevant phenotypes.

2.3.3.1 Summary statistics

The mental health questionnaire used in this study includes items that measure neuroticism,

anxiety, subjective well-being, depression, and risk taking. We therefore sought to obtain

summary statistics for these psychopathological phenotypes. First, we extracted summary

statistics for *Neuroticism* from a GWAS meta-analysis of self-reported neuroticism in the UKB

(using the same questions as in our study) and Psychiatric Genetics Consortium (using the NEO-

FFI personality inventory)[67]. For *Generalized Anxiety Disorder,* we leveraged summary statistics

from a GWAS of self-reported Generalized Anxiety Disorder 2-item scale scores in the Million

Veteran Program[68]. We further obtained summary statistics for *Subjective Well-Being* from a

GWAS meta-analysis of life satisfaction, positive affect, or both life satisfaction and positive

affect across 59 cohorts[69]. For *Major Depressive Disorder*, we meta-analyzed summary statistics

from case-control GWAS in the UK Biobank and Psychiatric Genomics Consortium[70] and the

Million Veteran Program[71] (see *Supplementary Results* for further details). Lastly, we obtained

summary statistics for *Risk Taking* from a GWAS study using the UKB data, which included the

same question of risk taking as in our study[72].

Summary statistics for accelerometer data-derived physical activity phenotypes, including

moderate activity, overall activity, sedentary activity, walking, and sleep duration were derived

from a GWAS of N=91,105 participants of European ancestry in the UK Biobank[58]. These
phenotypes were also included in our brain association analyses.

Please refer to *Table 1* for an overview for all the summary statistics used in this study.

Table 1. Summary statistics from GWAS studies.

Reference	Phenotype	Dataset	Sample Size
Nagel et al 2018[67]	Neuroticism	UK Biobank & Psychiatric Genetics Consortium	390,279
Levey et al 2020[68]	GAD	Million Veteran Program	199,611
Okbay et al 2016[69]	SWB	Meta-analysis across 59 cohorts	298,420
Howard et al 2019[70]	MDD	UK Biobank & Psychiatric Genomics Consortium	500,199
Levey et al 2021[71]		Million Veteran Program	250,215
Linnér et al 2019[72]	Risk-taking	UK Biobank	431,126
Doherty et al 2018[58]	Physical activities	UK Biobank	91,105

GAD = general anxiety disorder; SWB = subjective well-being; MDD = major depressive
disorder

2.3.3.2 Genetic correlations

We used linkage disequilibrium score regression (LDSR) and genomic structural equation
modeling (gSEM) to test whether the genomic architecture associated with general mental
health is shared with physical activity. LDSR leverages GWAS summary statistics to estimate
genetic correlations by regressing the SNP statistics on the SNP linkage disequilibrium (LD)
scores, or correlations between nearby genomic loci due to population stratification (i.e.,
systematic differences in allele frequencies due to differences in ancestry). gSEM characterizes
the latent genetic architecture across phenotypes based on the LDSR-derived genetic correlation
matrices [21]. To this end, we first applied LDSR to existing GWAS summary statistics of
psychopathological phenotypes (i.e., neuroticism, generalized anxiety disorder, subjective well-
being, major depressive disorder, and risk taking) and physical activity phenotypes (i.e., overall
activity, moderate activity, sedentary activity, sleep duration, and walking), respectively, to
estimate pairwise genetic correlations within each construct (i.e., within mental health and within
physical activity respectively). We also examined genetic correlations between mental health

phenotypes and physical activity phenotypes adjusted for sex and BMI. We then applied gSEM
to the covariance matrix of psychopathology and that of physical activity separately, allowing
one single latent factor to load freely within each model. Metrics indicating model fit (i.e., CFI,
comparative fit index; SRMR, standardized root mean squared residual) and factor loadings from
each of these models were used to determine whether one common genetic factor fit the physical
activity and mental health data well, respectively.

Because our results indicated poor model fit for some gSEM analyses (see details below in
Results section 3.3), we focused on the model of mental health, where "risk-taking" was
excluded to generate a latent factor of "negative affect" across other phenotypes. Specifically, we
explored genetic correlations between the latent factor of "negative affect" (i.e., without risk-
taking) and each of the five physical activity phenotypes. In addition, we examined genetic
correlations between "risk-taking" alone and each individual physical activity phenotype, using
LDSR. False discovery rate correction was used to correct for multiple testing (N=10 tests). We
also repeated analyses that returned significant results, with adjustment for sex and BMI.
Adjusted summary statistics from Doherty and colleagues (2018) were used in these analyses.

2.3.3.3 Causal relationships

To examine plausible causal associations between physical activity phenotypes and negative
affect, we conducted Latent Causal Variable Analysis (LCV). This approach finds a latent
variable that mediates the genetic correlation between two traits, such as negative affect and
sleep duration. Generally, if the latent variable has a stronger genetic correlation with trait 1
(e.g., sleep duration) than with trait 2 (e.g., negative affect), part of the genetic component of
trait 1 is thought to be causal for trait 2. This partial causality can be quantified using the genetic
causality proportion (GCP) of trait 1 (sleep duration) on trait 2 (negative affect), which can range
between 0 (no partial genetic causality) and 1 (full genetic causality; O'Connor and Price, 2018).

2.4   Confounding variables

Based on the literature, BMI, smoking and drinking status were included as confounding factors
in all statistical analyses for brain associations[74,75]. Age, sex, head motion during rs-fMRI
acquisition (i.e., mean frame-wise displacement), time difference in days between accelerometer
recording (i.e., start date) and assessment center visit date (i.e., acquisition date for both mental

health questionnaire and rs-fMRI), as well as the scanning site were further included. Due to

varying degrees of missingness in the confound variables (i.e., up to 22%), the same imputation

procedure, as described in *Section 2.2.2*, was performed except that additional usage of

multinomial logistic regression was employed to impute categorical data with more than two

levels (i.e., smoking and drinking status). Complete observations from all variables including the

confounding variables were used in the imputation procedure. When calculating the final

predicted values to replace missing data, the predicted values for continuous variables were

averaged across 20 iterations and the level with the highest count across iterations was selected.

All categorical variables after imputation were dummy coded for subsequent statistical analyses.

In the genetic correlation analyses, to ensure robust test effects, we repeated models with

significant results including sex and BMI as covariates.

2.5   Data and Code Availability

The UK Biobank data used in this study can be accessed by researchers upon application

(https://www.ukbiobank.ac.uk/register-apply). A machine learning algorithm shared on github

was   used   to   extract   physical   activity   measures

(https://github.com/activityMonitoring/biobankAccelerometerAnalysis). The Matlab code for

running permutation inference for CCA are also available on github, including the new extension

for testing canonical variable loadings (https://github.com/andersonwinkler/PermCCA). We

performed Pearson's correlations and imputations in R (version 4.1.0; R Core Team, 2021),

using basic *stat* function and *MICE* package (van Buuren and Groothuis-Oudshoorn, 2011),

respectively. Genetic analyses including LDSR and gSEM were performed using the

GenomicSEM   package   also   in   R   [21],   with   code   available   on   github

(https://github.com/GenomicSEM/GenomicSEM). LCV was performed using the open-source

LCV software (https://github.com/lukejoconnor/LCV).

3. Results

3.1   Correlations between mental health and physical activity measures

In this sample, phenotypic measures of mental health and physical activity demonstrated overall

small but significant correlations, with coefficients ranging between –0.08 and 0.11 (FDR-

corrected p-values $\leq$ 0.045). The overall coefficient patterns were largely consistent for

correlations between physical activity and mental health measures despite of minor differences

across different time windows (i.e., overall, weekdays, weekend; see full correlations in Table

S2).

3.2   Shared neural correlates of mental health and physical activity

3.2.1   Brain associations with mental health and physical activity

In simple CCA models (i.e., without accounting for the potentially shared variance in brain

measures between physical activity and mental health), 2 and 3 significant canonical variate pairs

were observed respectively for brain-mental health (r_1=0.16, p_1=0.001; r_2=0.12, p_2=0.001) and

brain-physical activity associations (r_1=0.23, p_1=0.001; r_2=0.15, p_2=0.001; r_3=0.13, p_3=0.047).

IWhen controlling for the shared variance in brain measures between mental health and physical

activity (i.e., unique models), we found 2 significant canonical variates for both brain-mental

health (r_1=0.14, p_1=0.001; r_2=0.12, p_2=0.002) and brain-physical activity associations (r_1=0.21,

p_1=0.001; r_2=0.15, p_2=0.001), with slightly decreased canonical correlation coefficients

comparing to those from the simple models. The canonical variates for brain measures after

accounting for the shared variance (i.e., from the unique models) mapped well with those from

the simple models, as indicated by correlation coefficients between brain canonical variables

(i.e., □; all r's > 0.96; see full results in *Table S3*).

3.2.2   Canonical loadings of brain measures

Findings from the post-hoc analyses on the significant canonical variates showed considerable

overlap between the brain measures with significant loadings associated with mental health and

with physical activity. Overall, the amplitude of nodes that indicates the averaged signal

fluctuations in each node over time loaded higher than the edges, with the highest loadings on

the amplitude of dorsal DMN or left CEN for the first canonical variates in both the mental

health and physical activity models (*permuted p-values* $\leq$ 0.001; *Figure 2*). The similar loading

pattern of all brain measures were largely retained even when the shared variance was partialled

out for mental health and for physical activity, respectively, in the unique models (*permuted* p-

values ≤ 0.035). In addition to amplitude, connectivity between the dorsal DMN and left CEN

(i.e., node edge) also exhibited statistically significant loadings in the first canonical variates of

both the mental health and the physical activity models (simple and unique models; *permuted* p-

values ≤ 0.001; *Figure 2*). The brain variable loadings for the second canonical variate were

highly similar between simple and unique models for either the mental health or physical activity

models, with the amplitude and node edges showing evenly important involvement. Yet, the

patterns of these loadings differed between the mental health and physical activity model (*Figure*

S2).

Figure 2. Canonical loadings of brain measures on the first canonical variates. These loadings represent the
linear correlation between the original brain measures (Y) and the first canonical variate (U) per model. Color
coding was made for brain variable names along the Y axis (i.e., node edges in gray with "-" between node names
and amplitude in orange), and for the bars representing canonical loadings (i.e., significance in cyan, insignificance
in yellow). Simple and unique models differ in whether the model accounted for the shared variance in brain

measures between mental health and physical activity. vDMN = ventral default mode network; R/L CEN = right/
left central executive network; PCu = precuneus; a/p SN = anterior/posterior salience network.
425
426

3.2.3   Canonical loadings of mental health questions and physical activities
Canonical loadings of both individual mental health questions and physical activity types for the first
canonical variate also exhibited similar patterns between the simple and unique models (*Figure S3*). In the
models for brain-mental health associations, "risk taking" and "ever irritable/ argumentative for 2 days"
had the highest loadings in both simple and unique models, whereas in the models for brain-physical
activity associations, "walking" in all time windows (i.e., overall, weekdays, weekend) showed the
greatest importance across models. Interestingly, among all physical activity types, only "sleep" showed
the opposite direction in canonical loadings (*Figure S3*).
435

3.3   Genetic correlations
3.3.1   Genetic correlations for individual phenotypes
Pairwise LDSR was performed separately for psychopathological phenotypes and physical
activity phenotypes. All results were significant after controlling for multiple comparisons,
except for the genetic correlations between neuroticism and risk-taking ($rg = 0.010$, $p = 0.633$),
and between generalized anxiety disorder and risk-taking ($rg = 0.0677$, $p = 0.0472$, *FDR-*
corrected p-value = 0.0524; *Figure 3*). Specifically, among psychopathological phenotypes, the
smallest significant genetic correlation was observed for subjective well-being and risk-taking
($rg = -0.117$, $p = 1.13e-03$, *FDR-corrected p-value* = 1.41e-03), and the largest effect was
observed for MDD and generalized anxiety disorder ($rg = 0.768$, $p = 2.62e-90$, *FDR-corrected p-*
value = 1.31e-89). For physical activity, genetic correlations ranged from -0.217 ($p = 1.03e-04$,
FDR-corrected p-value = 1.03e-04) for sleep duration and walking to 0.796 ($p = 1.77e-31$, *FDR-*
corrected p-value = 8.85e-31) for moderate and overall activity (*Figure 3*). Physical activity
phenotypes adjusted for sex and BMI largely recapitulated these results, with genetic correlations
ranging from -0.209 ($p = 2.49e-04$, *FDR-corrected p-value* = 2.49e-04) for sleep duration and
walking, to 0.780 ($p = 2.12e-23$, *FDR-corrected p-value* = 1.06e-22) for moderate and overall
activities.
453

Figure 3. Heatmap of bivariate genetic correlations. Pairwise genetic correlations were calculated separately for

phenotypes of mental health (A) and those of physical activity (B). rg = genetic correlation coefficient, GAD =

Generalized Anxiety Disorder, MDD = Major Depressive Disorder, Risk = Risk Tolerance, SWB = Subjective Well-

Being.

3.3.2   Genetic correlations for latent factors

Analyses using gSEM across all mental health phenotypes returned results indicating suboptimal

estimations with a low factor loading on risk taking from the model for a general

psychopathology latent factor (standardized loading = 0.12), which likely reflected its

conceptually distinct construct from all other phenotypes (i.e., depression, anxiety, neuroticism,

and subjective well-being). Additionally, initial gSEM results for physical activity phenotypes

indicated overall poor model fit (CFI = 0.569, SRMR = 0.160; *Figure S4*). Thus, we performed

gSEM for psychopathological phenotypes grouped as one "negative affect" latent factor after

excluding "risk-taking", with an effective sample size of 571,170 and good model fit (CFI =

0.987, SRMR = 0.0535; *Figure S5*). After correction for multiple testing, genetic variance in

negative affect was significantly and positively correlated with sleep duration (*rg* = 0.121, *p* =

1.25e-05, *FDR-corrected p* = 1.25e-04), and negatively correlated with moderate (*rg* = -0.117, *p*

= 1.33e-03, *FDR-corrected p* = 4.43e-03) and overall activity (*rg* = -0.085, *p* = 7.70e-04, *FDR-*

corrected p = 3.85e-03; *Figure 4;* also see the results for a latent negative affect factor including

"risk-taking" in *the Supplementary Results*). No significant genetic correlation was observed

between negative affect and sedentary activity or walking after multiple comparison corrections

($|rgs| \leq 0.062$, p-values ≥ 0.036, FDR-corrected p-values ≥ 0.076; *Table 2*). Post-hoc analyses

revealed that after adjustment for sex and BMI, only the genetic correlation between negative

affect and sleep duration remained significant ($rg = 0.122$, $p = 9.38e\text{-}06$). Using LDSR, we found

no significant correlations between risk-taking and any physical activity phenotype ($|rgs| \leq 0.074$,

$ps \geq 0.0511$).

Figure 4. Genetic correlations between negative affect and physical activity phenotypes. Significant genetic
correlations with negative affect were observed for moderate (A), overall activities (B), and sleep duration (C). The
latent negative affect factor loaded positively on neuroticism, major depression disorder (MDD) and general anxiety
disorder (GAD) phenotypes, but negatively on the subjective well-being (SWB) phenotype.

Table 2. Genetic correlations between mental health and physical activity phenotypes.

Mental Health Phenotype	Physical Activity Phenotype	*rg*	*p*	*pFDR*
Risk Tolerance	Moderate	0.072	0.088	0.126
	Overall	0.031	0.311	0.346
	Sedentary	*0.074*	0.038	0.076
	Sleep Duration	-0.067	0.051	0.085
	Walking	0.041	0.300	0.346
Negative Affect	Moderate	**-0.117**	1.33e-3	4.43e-3
	Overall	**-0.085**	7.70e-4	3.85e-3
	Sedentary	*-0.062*	0.036	0.076
	Sleep Duration	**0.121**	1.25e-5	1.25e-4
	Walking	-0.029	0.383	0.383

Note, Significant results after FDR corrections were highlighted in bold. *rg* = genetic correlation,
pFDR = FDR-corrected p-value.

3.3.3 Latent Causal Variable Analysis.

Results from the LCV analyses did not indicate causal relationships between any physical

activity phenotypes and negative affect, in either direction ($|GCPs| < 0.425$; ps > 0.163).

4. Discussion

In this study, we investigated whether mental health and physical activity have shared brain and

genetic architectures using the UK Biobank cohort. Our findings showed significant associations

of mental health and physical activity separately with a set of brain measures that represent the

connectivity strength and amplitude of nodes from the DMN, SN and CEN. Critically, these

significant associations exhibited highly similar patterns of brain variable loadings across mental

health and physical activity models even when the shared variance between these two constructs

was accounted for, suggesting a potential overlap in brain network architecture between these

two constructs. Further analyses examining genetic correlations for mental health and physical

activity showed that negative affect exhibited significant genetic correlations with several

physical activity types, of which sleep duration demonstrated the strongest genetic correlation

that remained significant after controlling for BMI and sex effects. Together, these results

support the presence of shared multivariate brain and genetic architectures between mental health

and physical activity.

The three intrinsic brain networks, namely the DMN, SN and CEN, have been consistently

implicated in a wide range of psychiatric disorders including major depression and anxiety[12].

Interestingly, connections between or the configurations of these networks have also been

associated with physical exercises[76,77]. The current study therefore focused specifically on the

subnetworks or nodes from these large-scale networks and used the amplitude and connectivity

strength (i.e., node edge) to examine the associations of these network nodes with mental health

and with physical activity, respectively. In line with the literature, we observed significant

multivariate associations for all three networks with either mental health or physical activity, and

significant loadings on most of the network nodes. In particular, the dorsal DMN and left CEN

showed the greatest involvement in the obsecrated brain associations with both mental health and

physical activity (*Figure 2*). In this study, the dorsal DMN subnetwork primarily consisted of the

posterior cingulate cortex (PCC) and the ventromedial prefrontal cortex (vmPFC), the two brain

areas that are commonly considered as the core subsystem of the DMN[78]. Similarly, the CEN

node here included two most typical hub regions: the dorsolateral prefrontal cortex (dlPFC) and

posterior parietal cortex (PPC) for each individual hemisphere (*Figure S1*). These major DMN

and CEN nodes have been implicated in various mental illnesses including depression and

anxiety. For example, the PCC and mPFC have been suggested to collectively support multiple

cognitive functions including decision making and memory[79], the impairment of which has often

been reported in patients with major depression and anxiety disorders[80–82]. Additionally, the

dlPFC is known to be involved in emotion regulation [83–85] and dysfunction of this region is often

seen in abnormal processing of emotional experiences in patients with depressive and anxiety

symptoms[86,87]. As for physical activity, increased dlPFC activity has been observed after acute

physical exercises in participants with higher scores in the Stroop test[88], whereas the involvement

of the DMN subsystems in the medial temporal lobe (MTL) including the hippocampus and its

connection with the medial PFC are often observed in relation to enhanced memory after

physical exercises[33,89,90]. In our findings, the amplitude of dorsal DMN and left CEN that

indicated the temporal fluctuations of intra-node interactions (i.e., variance in the connections

between PCC and vmPFC or between dlPFC and PCC), as well as the connectivity strength

between these two nodes, showed significantly high loadings for the most critical association

between brain measures and mental health measures (i.e., the first canonical variate). These

observations are in line with separate literature on mental health and physical activity, and

provide empirical evidence that mental health and physical activity may share brain architecture

involved in major cognitive functions.

Interestingly, mental health and physical activity also appear to have partially overlapping

genetic architectures. In line with previous reports that internalizing problems are negatively

genetically correlated with physical movement[22], we showed that a latent negative affect factor

capturing genetic covariance between subjective well-being, neuroticism, major depressive

disorder, and generalized anxiety disorder was negatively genetically correlated with overall

physical activity as well as a more fine-grained phenotype of moderate activity, and positively

genetically correlated with sleep duration. Protective effects of physical activity on mental health

have long been documented, as have negative health consequences of psychiatric disorders[5–7,91].

Here, we demonstrate that these relationships can be partially explained by shared genetic

predisposition, although results from our latent causal variable analysis indicate that these

associations do not reflect causal influences. Interestingly, with the adjustment of sex and BMI,

only the correlation with sleep duration remained significant. Sleep duration also showed the

lowest loading onto a latent physical activity genetic factor (*Figure S4*). These results suggest

that the sleep phenotype is somewhat distinct from the remaining physical activity phenotypes

and that the shared genetic architecture between negative affect and sleep duration is more

pronounced than that between negative affect and the degree of daily physical or sedentary

activity. This is in line with the frequent documentation of symptomatic sleep disturbances

across forms of psychopathology, including depression and anxiety[92,93], even in children[94].

Despite being the first to jointly investigate the shared brain network architecture and genetic

basis of mental health and physical activity in a large population cohort, our study has some

limitations. First, the brain measures in our study are derived from resting-state fMRI measures.

Although our choice reflected a rich literature that has implicated these measures in both mental

health and physical activity, inclusion of multimodal brain measures such as structural gray

matter volume, cortical thickness, and white matter integrity can provide multiple aspects of

brain architecture in relation to mental health and physical activity, and thus may be of interest

for future investigations. Second, although the accelerometer recording took place prior to the

acquisition of resting-state fMRI and mental health assessment for most of the participants in our

study (i.e., 96%), the degrees of time difference between these measurements varied greatly at

the individual level (i.e., ranging between -473 to 2281 days). This time discrepancy was

accounted for in all CCA models as a covariate and cautions should be taken when interpreting

the observed brain associations with reference to time effects.

In conclusion, our study jointly analyzed resting-state network measures and genetic correlations

in a large cohort to test the hypothesis of a shared neurobiological basis of mental health and

physical activity. Our findings revealed that multivariate patterns of brain correlates were highly

similar between mental health and physical activity and highlighted genetic correlations between

mental health (negative affect) and overall physical activity, moderate activity levels, and sleep

duration. Taken together, these findings point towards neural and genetic mechanisms that may

subserve the protective influence of physical exercise and sleep on mental health.

Acknowledgements
JDB is supported by the NIH (1 R34 NS118618-01) and the McDonnell Center for Systems
Neuroscience. SEP is supported by the NIH (1 F31 AA029934-01).
589
590
591

Reference

1. Zhang Z, Chen W. A Systematic Review of the Relationship Between Physical Activity and Happiness. J. Happiness Stud. 2019. doi:10.1007/s10902-018-9976-0.

2. Fernandes J, Arida RM, Gomez-Pinilla F. Physical exercise as an epigenetic modulator of brain plasticity and cognition. Neurosci. Biobehav. Rev. 2017. doi:10.1016/j.neubiorev.2017.06.012.

3. Kays JL, Hurley RA, Taber KH. The dynamic brain: Neuroplasticity and mental health. *J Neuropsychiatry Clin Neurosci* 2012. doi:10.1176/appi.neuropsych.12050109.

4. Millan MJ, Agid Y, Brüne M, Bullmore ET, Carter CS, Clayton NS *et al.* Cognitive dysfunction in psychiatric disorders: Characteristics, causes and the quest for improved therapy. Nat. Rev. Drug Discov. 2012. doi:10.1038/nrd3628.

5. Biddle S. Physical activity and mental health: Evidence is growing. World Psychiatry. 2016. doi:10.1002/wps.20331.

6. Deslandes A, Moraes H, Ferreira C, Veiga H, Silveira H, Mouta R *et al.* Exercise and mental health: Many reasons to move. Neuropsychobiology. 2009. doi:10.1159/000223730.

7. Heinze K, Cumming J, Dosanjh A, Palin S, Poulton S, Bagshaw AP *et al.* Neurobiological evidence of longer-term physical activity interventions on mental health outcomes and cognition in young people: A systematic review of randomised controlled trials. Neurosci. Biobehav. Rev. 2021. doi:10.1016/j.neubiorev.2020.10.014.

8. Besteher B, Gaser C, Nenadić I. Brain Structure and Subclinical Symptoms: A Dimensional Perspective of Psychopathology in the Depression and Anxiety Spectrum. Neuropsychobiology. 2020. doi:10.1159/000501024.

9. Patel Y, Parker N, Shin J, Howard D, French L, Thomopoulos SI *et al.* Virtual Histology of Cortical Thickness and Shared Neurobiology in 6 Psychiatric Disorders. *JAMA Psychiatry* 2021. doi:10.1001/jamapsychiatry.2020.2694.

10. Van Tol MJ, Van Der Wee NJA, Van Den Heuvel OA, Nielen MMA, Demenescu LR, Aleman A *et al.* Regional brain volume in depression and anxiety disorders. *Arch Gen Psychiatry* 2010. doi:10.1001/archgenpsychiatry.2010.121.

11. Woodward ND, Cascio CJ. Resting-state functional connectivity in psychiatric disorders. *JAMA Psychiatry* 2015; **72**: 743–744.

12. Menon V. Large-scale brain networks and psychopathology: A unifying triple network model. *Trends Cogn Sci* 2011; **15**: 483–506.

13. Tozzi L, Zhang X, Chesnut M, Holt-Gosselin B, Ramirez CA, Williams LM. Reduced functional connectivity of default mode network subsystems in depression: Meta-analytic evidence and relationship with trait rumination. *NeuroImage Clin* 2021. doi:10.1016/j.nicl.2021.102570.

14. Kaiser RH, Andrews-Hanna JR, Wager TD, Pizzagalli DA. Large-Scale Network Dysfunction in Major Depressive Disorder. *JAMA Psychiatry* 2015. doi:10.1001/jamapsychiatry.2015.0071.

15. Xu J, Van Dam NT, Feng C, Luo Y, Ai H, Gu R *et al.* Anxious brain networks: A coordinate-based activation likelihood estimation meta-analysis of resting-state functional connectivity studies in anxiety. Neurosci. Biobehav. Rev. 2019. doi:10.1016/j.neubiorev.2018.11.005.

16. Servaas MN, Geerligs L, Renken RJ, Marsman JBC, Ormel J, Riese H *et al.* Connectomics and neuroticism: An altered functional network organization. *Neuropsychopharmacology* 2015. doi:10.1038/npp.2014.169.

17. Anttila V, Bulik-Sullivan B, Finucane HK, Walters RK, Bras J, Duncan L *et al.* Analysis of shared heritability in common disorders of the brain. *Science (80-)* 2018. doi:10.1126/science.aap8757.

18. Lee SH, Ripke S, Neale BM, Faraone S V., Purcell SM, Perlis RH *et al.* Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nat Genet* 2013. doi:10.1038/ng.2711.

19. Taylor MJ, Martin J, Lu Y, Brikell I, Lundström S, Larsson H *et al.* Association of Genetic Risk Factors for Psychiatric Disorders and Traits of These Disorders in a Swedish Population Twin Sample. *JAMA Psychiatry* 2019. doi:10.1001/jamapsychiatry.2018.3652.

650   20   Smoller JW, Andreassen OA, Edenberg HJ, Faraone S V., Glatt SJ, Kendler KS. Psychiatric
651   genetics and the structure of psychopathology. Mol. Psychiatry. 2019. doi:10.1038/s41380-017-
652   0010-4.

653   21   Grotzinger AD, Rhemtulla M, de Vlaming R, Ritchie SJ, Mallard TT, Hill WD *et al.* Genomic
654   structural equation modelling provides insights into the multivariate genetic architecture of
655   complex traits. *Nat Hum Behav* 2019. doi:10.1038/s41562-019-0566-x.

656   22   Grotzinger AD, Mallard TT, Akingbuwa WA, Ip HF, Adams MJ, Lewis CM *et al.* Genetic
657   architecture of 11 major psychiatric disorders at biobehavioral, functional genomic, and molecular
658   genetic levels of analysis. medRxiv. 2020. doi:10.1101/2020.09.22.20196089.

659   23   Anderson E, Shivakumar G. Effects of exercise and physical activity on anxiety. *Front Psychiatry*
660   2013. doi:10.3389/fpsyt.2013.00027.

661   24   Blumenthal JA, Babyak MA, Doraiswamy PM, Watkins L, Hoffman BM, Barbour KA *et al.*
662   Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med*
663   2007. doi:10.1097/PSY.0b013e318148c19a.

664   25   Thomas AG, Dennis A, Rawlings NB, Stagg CJ, Matthews L, Morris M *et al.* Multi-modal
665   characterization of rapid anterior hippocampal volume increase associated with aerobic exercise.
666   *Neuroimage* 2016. doi:10.1016/j.neuroimage.2015.10.090.

667   26   Thomas AG, Dennis A, Bandettini PA, Johansen-Berg H. The effects of aerobic activity on brain
668   structure. Front. Psychol. 2012. doi:10.3389/fpsyg.2012.00086.

669   27   Sexton CE, Betts JF, Dennis A, Doherty A, Leeson P, Holloway C *et al.* The effects of an aerobic
670   training intervention on cognition, grey matter volumes and white matter microstructure. *Physiol*
671   *Behav* 2020. doi:10.1016/j.physbeh.2020.112923.

672   28   Colcombe SJ, Kramer AF, Erickson KI, Scalf P, McAuley E, Cohen NJ *et al.* Cardiovascular
673   fitness, cortical plasticity, and aging. *Proc Natl Acad Sci U S A* 2004.
674   doi:10.1073/pnas.0400266101.

675   29   Herold F, Aye N, Lehmann N, Taubert M, Müller NG. The contribution of functional magnetic
676   resonance imaging to the understanding of the effects of acute physical exercise on cognition.
677   *Brain Sci* 2020. doi:10.3390/brainsci10030175.

678   30   Ikuta T, Loprinzi PD. Association of cardiorespiratory fitness on interhemispheric hippocampal
679   and parahippocampal functional connectivity. *Eur J Neurosci* 2019. doi:10.1111/ejn.14366.

680   31   Talukdar T, Nikolaidis A, Zwilling CE, Paul EJ, Hillman CH, Cohen NJ *et al.* Aerobic fitness
681   explains individual differences in the functional brain connectome of healthy young adults. *Cereb*
682   *Cortex* 2018. doi:10.1093/cercor/bhx232.

683   32   Weng TB, Pierce GL, Darling WG, Falk D, Magnotta VA, Voss MW. The Acute Effects of
684   Aerobic Exercise on the Functional Connectivity of Human Brain Networks. *Brain Plast* 2016.
685   doi:10.3233/bpl-160039.

686   33   Voss MW, Prakash RS, Erickson KI, Basak C, Chaddock L, Kim JS *et al.* Plasticity of brain
687   networks in a randomized intervention trial of exercise training in older adults. *Front Aging*
688   *Neurosci* 2010. doi:10.3389/fnagi.2010.00032.

689   34   Greeley B, Chau B, Jones CB, Neva JL, Kraeutner SN, Campbell KL *et al.* Multiple bouts of high-
690   intensity interval exercise reverse age-related functional connectivity disruptions without affecting
691   motor learning in older adults. *Sci Rep* 2021. doi:10.1038/s41598-021-96333-4.

692   35   Schmitt A, Upadhyay N, Martin JA, Rojas S, Strüder HK, Boecker H. Modulation of Distinct
693   Intrinsic Resting State Brain Networks by Acute Exercise Bouts of Differing Intensity. *Brain Plast*
694   2019. doi:10.3233/bpl-190081.

695   36   Voss MW, Weng TB, Narayana-Kumanan K, Cole RC, Wharff C, Reist L *et al.* Acute exercise
696   effects predict training change in cognition and connectivity. *Med Sci Sports Exerc* 2020.
697   doi:10.1249/MSS.0000000000002115.

698   37   Nutt DJ, Wilson S, Paterson L. Sleep disorders as core symptoms of depression. *Dialogues Clin*
699   *Neurosci* 2008. doi:10.31887/dcns.2008.10.3/dnutt.

700   38   Riemann D, Krone LB, Wulff K, Nissen C. Sleep, insomnia, and depression.

701   Neuropsychopharmacology. 2020. doi:10.1038/s41386-019-0411-y.

702 39 Cox RC, Olatunji BO. Sleep in the anxiety-related disorders: A meta-analysis of subjective and
703   objective research. Sleep Med. Rev. 2020. doi:10.1016/j.smrv.2020.101282.

704 40 Mellman TA. Sleep and Anxiety Disorders. Psychiatr. Clin. North Am. 2006.
705   doi:10.1016/j.psc.2006.08.005.

706 41 Rumble ME, White KH, Benca RM. Sleep Disturbances in Mood Disorders. Psychiatr. Clin. North
707   Am. 2015. doi:10.1016/j.psc.2015.07.006.

708 42 Wainberg M, Jones SE, Beaupre LM, Hill SL, Felsky D, Rivas MA *et al.* Association of
709   accelerometer-derived sleep measures with lifetime psychiatric diagnoses: A cross-sectional study
710   of 89,205 participants from the UK Biobank. *PLOS Med* 2021; **18**: e1003782.

711 43 Full KM, Malhotra A, Gallo LC, Kerr J, Arredondo EM, Natarajan L *et al.* Accelerometer-
712   Measured Sleep Duration and Clinical Cardiovascular Risk Factor Scores in Older Women.
713   *Journals Gerontol - Ser A Biol Sci Med Sci* 2020. doi:10.1093/gerona/glz201.

714 44 De Havas JA, Parimal S, Soon CS, Chee MWL. Sleep deprivation reduces default mode network
715   connectivity and anti-correlation during rest and task performance. *Neuroimage* 2012.
716   doi:10.1016/j.neuroimage.2011.08.026.

717 45 Sämann PG, Tully C, Spoormaker VI, Wetter TC, Holsboer F, Wehrle R *et al.* Increased sleep
718   pressure reduces resting state functional connectivity. *Magn Reson Mater Physics, Biol Med* 2010.
719   doi:10.1007/s10334-010-0213-z.

720 46 Yeo BTT, Tandi J, Chee MWL. Functional connectivity during rested wakefulness predicts
721   vulnerability to sleep deprivation. *Neuroimage* 2015. doi:10.1016/j.neuroimage.2015.02.018.

722 47 Khalsa S, Mayhew SD, Przezdzik I, Wilson R, Hale J, Goldstone A *et al.* Variability in cumulative
723   habitual sleep duration predicts waking functional connectivity. *Sleep* 2016.
724   doi:10.5665/sleep.5324.

725 48 Bijsterbosch J, Harrison S, Duff E, Alfaro-Almagro F, Woolrich M, Smith S. Investigations into
726   within- and between-subject resting-state amplitude variations. *Neuroimage* 2017.
727   doi:10.1016/j.neuroimage.2017.07.014.

728 49 Choh AC, Demerath EW, Lee M, Williams KD, Towne B, Siervogel RM *et al.* Genetic analysis of
729   self-reported physical activity and adiposity: The Southwest Ohio Family Study. *Public Health*
730   *Nutr* 2009. doi:10.1017/S1368980008003583.

731 50 Moore-Harrison T, Lightfoot JT. Driven to Be Inactive?—The Genetics of Physical Activity. In:
732   *Progress in Molecular Biology and Translational Science*. 2010 doi:10.1016/B978-0-12-375003-
733   7.00010-8.

734 51 Sallis JF, Patterson TL, Buono MJ, Atkins CJ, Nader PR. Aggregation of physical activity habits
735   in Mexican-American and Anglo families. *J Behav Med* 1988. doi:10.1007/BF00846167.

736 52 Simonen RL, Pérusse L, Rankinen T, Rice T, Rao DC, Bouchard C. Familial aggregation of
737   physical activity levels in the Québec family study. *Med Sci Sports Exerc* 2002.
738   doi:10.1097/00005768-200207000-00014.

739 53 Butte NF, Cai G, Cole SA, Comuzzie AG. Viva la Familia Study: Genetic and environmental
740   contributions to childhood obesity and its comorbidities in the Hispanic population. *Am J Clin*
741   *Nutr* 2006. doi:10.1093/ajcn/84.3.646.

742 54 Stubbe JH, Boomsma DI, Vink JM, Cornes BK, Martin NG, Skythe A *et al.* Genetic influences
743   on exercise participation in 37.051 twin pairs from seven countries. *PLoS One* 2006.
744   doi:10.1371/journal.pone.0000022.

745 55 Duncan GE, Goldberg J, Noonan C, Moudon AV, Hurvitz P, Buchwald D. Unique environmental
746   effects on physical activity participation: A twin study. *PLoS One* 2008.
747   doi:10.1371/journal.pone.0002019.

748 56 Aasdahl L, Nilsen TIL, Meisingset I, Nordstoga AL, Evensen KAI, Paulsen J *et al.* Genetic
749   variants related to physical activity or sedentary behaviour: a systematic review. Int. J. Behav.
750   Nutr. Phys. Act. 2021. doi:10.1186/s12966-020-01077-5.

751 57 De Moor MHM, Liu YJ, Boomsma DI, Li J, Hamilton JJ, Hottenga JJ *et al.* Genome-wide

752   association study of exercise behavior in dutch and american adults. *Med Sci Sports Exerc* 2009.
753   doi:10.1249/MSS.0b013e3181a2f646.

754   58   Doherty A, Smith-Byrne K, Ferreira T, Holmes M V., Holmes C, Pulit SL *et al.* GWAS identifies
755   14 loci for device-measured physical activity and sleep duration. *Nat Commun* 2018; **9**.
756   doi:10.1038/s41467-018-07743-4.

757   59   Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J *et al.* UK Biobank: An Open Access
758   Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old
759   Age. *PLoS Med* 2015. doi:10.1371/journal.pmed.1001779.

760   60   Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J *et al.* Multimodal
761   population brain imaging in the UK Biobank prospective epidemiological study. *Nat Neurosci*
762   2016. doi:10.1038/nn.4393.

763   61   van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate imputation by chained equations in R.
764   *J Stat Softw* 2011. doi:10.18637/jss.v045.i03.

765   62   Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J *et al.* Multimodal
766   population brain imaging in the UK Biobank prospective epidemiological study. *Nat Neurosci*
767   2016. doi:10.1038/nn.4393.

768   63   Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD. Decoding Subject-Driven Cognitive
769   States with Whole-Brain Connectivity Patterns. *Cereb Cortex* 2012; **22**: 158–165.

770   64   Winkler AM, Renaud O, Smith SM, Nichols TE. Permutation inference for canonical correlation
771   analysis. *Neuroimage* 2020. doi:10.1016/j.neuroimage.2020.117065.

772   65   HOTELLING H. RELATIONS BETWEEN TWO SETS OF VARIATES. *Biometrika* 1936.
773   doi:10.1093/biomet/28.3-4.321.

774   66   Helmer M, Warrington S, Mohammadi-Nejad A-R, Ji LJ, Howell A, Rosand B *et al.* On stability
775   of Canonical Correlation Analysis and Partial Least Squares with application to brain-behavior
776   associations. *bioRxiv* 2020.

777   67   Nagel M, Jansen PR, Stringer S, Watanabe K, De Leeuw CA, Bryois J *et al.* Meta-analysis of
778   genome-wide association studies for neuroticism in 449,484 individuals identifies novel genetic
779   loci and pathways. *Nat Genet* 2018. doi:10.1038/s41588-018-0151-7.

780   68   Levey DF, Gelernter J, Polimanti R, Zhou H, Cheng Z, Aslan M *et al.* Reproducible Genetic Risk
781   Loci for Anxiety: Results From ~200,000 Participants in the Million Veteran Program. *Am J*
782   *Psychiatry* 2020. doi:10.1176/appi.ajp.2019.19030256.

783   69   Okbay A, Baselmans BML, De Neve JE, Turley P, Nivard MG, Fontana MA *et al.* Genetic
784   variants associated with subjective well-being, depressive symptoms, and neuroticism identified
785   through genome-wide analyses. *Nat Genet* 2016. doi:10.1038/ng.3552.

786   70   Howard DM, Adams MJ, Clarke TK, Hafferty JD, Gibson J, Shirali M *et al.* Genome-wide meta-
787   analysis of depression identifies 102 independent variants and highlights the importance of the
788   prefrontal brain regions. *Nat Neurosci* 2019. doi:10.1038/s41593-018-0326-7.

789   71   Levey DF, Stein MB, Wendt FR, Pathak GA, Zhou H, Aslan M *et al.* Bi-ancestral depression
790   GWAS in the Million Veteran Program and meta-analysis in >1.2 million individuals highlight
791   new therapeutic directions. *Nat Neurosci* 2021. doi:10.1038/s41593-021-00860-2.

792   72   Karlsson Linnér R, Biroli P, Kong E, Meddens SFW, Wedow R, Fontana MA *et al.* Genome-wide
793   association analyses of risk tolerance and risky behaviors in over 1 million individuals identify
794   hundreds of loci and shared genetic influences. *Nat Genet* 2019. doi:10.1038/s41588-018-0309-3.

795   73   O'Connor LJ, Price AL. Distinguishing genetic correlation from causation across 52 diseases and
796   complex traits. *Nat Genet* 2018. doi:10.1038/s41588-018-0255-0.

797   74   Jia H, Zack MM, Gottesman II, Thompson WW. Associations of Smoking, Physical Inactivity,
798   Heavy Drinking, and Obesity with Quality-Adjusted Life Expectancy among US Adults with
799   Depression. *Value Heal* 2018. doi:10.1016/j.jval.2017.08.002.

800   75   Metse AP, Clinton-Mcharg T, Skinner E, Yogaraj Y, Colyvas K, Bowman J. Associations between
801   suboptimal sleep and smoking, poor nutrition, harmful alcohol consumption and inadequate
802   physical activity ('snap risks'): A comparison of people with and without a mental health

condition in an australian community survey. *Int J Environ Res Public Health* 2021.
doi:10.3390/ijerph18115946.

76 Baniqued PL, Gallen CL, Voss MW, Burzynska AZ, Wong CN, Cooke GE *et al.* Brain network
modularity predicts exercise-related executive function gains in older adults. *Front Aging*
Neurosci 2018. doi:10.3389/fnagi.2017.00426.

77 Chaddock-Heyman L, Weng TB, Kienzler C, Weisshappel R, Drollette ES, Raine LB *et al.* Brain
Network Modularity Predicts Improvements in Cognitive and Scholastic Performance in Children
Involved in a Physical Activity Intervention. *Front Hum Neurosci* 2020.
doi:10.3389/fnhum.2020.00346.

78 Smallwood J, Bernhardt BC, Leech R, Bzdok D, Jefferies E, Margulies DS. The default mode
network in cognition: a topographical perspective. Nat. Rev. Neurosci. 2021. doi:10.1038/s41583-
021-00474-4.

79 Wang S, Tepfer LJ, Taren AA, Smith D V. Functional parcellation of the default mode network: a
large-scale meta-analysis. *Sci Rep* 2020. doi:10.1038/s41598-020-72317-8.

80 Mirza SS, Ikram MA, Bos D, Mihaescu R, Hofman A, Tiemeier H. Mild cognitive impairment and
risk of depression and anxiety: A population-based study. *Alzheimer's Dement* 2017.
doi:10.1016/j.jalz.2016.06.2361.

81 Perini G, Ramusino MC, Sinforiani E, Bernini S, Petrachi R, Costa A. Cognitive impairment in
depression: Recent advances and novel treatments. *Neuropsychiatr Dis Treat* 2019.
doi:10.2147/NDT.S199746.

82 Rock PL, Roiser JP, Riedel WJ, Blackwell AD. Cognitive impairment in depression: A systematic
review and meta-analysis. Psychol. Med. 2014. doi:10.1017/S0033291713002535.

83 Delgado MR, Nearing KI, LeDoux JE, Phelps EA. Neural Circuitry Underlying the Regulation of
Conditioned Fear and Its Relation to Extinction. *Neuron* 2008. doi:10.1016/j.neuron.2008.06.029.

84 Morawetz C, Bode S, Baudewig J, Kirilina E, Heekeren HR. Changes in Effective Connectivity
Between Dorsal and Ventral Prefrontal Regions Moderate Emotion Regulation. *Cereb Cortex*
2016. doi:10.1093/cercor/bhv005.

85 Ochsner KN, Gross JJ. The cognitive control of emotion. Trends Cogn. Sci. 2005.
doi:10.1016/j.tics.2005.03.010.

86 Koenigs M, Grafman J. The functional neuroanatomy of depression: Distinct roles for
ventromedial and dorsolateral prefrontal cortex. Behav. Brain Res. 2009.
doi:10.1016/j.bbr.2009.03.004.

87 Nejati V, Majdi R, Salehinejad MA, Nitsche MA. The role of dorsolateral and ventromedial
prefrontal cortex in the processing of emotional dimensions. *Sci Rep* 2021. doi:10.1038/s41598-
021-81454-7.

88 Yanagisawa H, Dan I, Tsuzuki D, Kato M, Okamoto M, Kyutoku Y *et al.* Acute moderate exercise
elicits increased dorsolateral prefrontal activation and improves cognitive performance with
Stroop test. *Neuroimage* 2010. doi:10.1016/j.neuroimage.2009.12.023.

89 Li MY, Huang MM, Li SZ, Tao J, Zheng GH, Chen LD. The effects of aerobic exercise on the
structure and function of DMN-related brain regions: a systematic review. Int. J. Neurosci. 2017.
doi:10.1080/00207454.2016.1212855.

90 Tao J, Liu J, Egorova N, Chen X, Sun S, Xue X *et al.* Increased hippocampus-medial prefrontal
cortex resting-state functional connectivity and memory function after Tai Chi Chuan practice in
elder adults. *Front Aging Neurosci* 2016. doi:10.3389/fnagi.2016.00025.

91 De Hert M, Correll CU, Bobes J, Cetkovich-Bakmas M, Cohen DAN, Asai I *et al.* Physical illness
in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in
health care. World Psychiatry. 2011. doi:10.1002/j.2051-5545.2011.tb00014.x.

92 Fang H, Tu S, Sheng J, Shao A. Depression in sleep disturbance: A review on a bidirectional
relationship, mechanisms and treatment. J. Cell. Mol. Med. 2019. doi:10.1111/jcmm.14170.

93 Cox RC, Olatunji BO. A systematic review of sleep disturbance in anxiety and related disorders. J.
Anxiety Disord. 2016. doi:10.1016/j.janxdis.2015.12.001.

94   Cheng W, Rolls E, Gong W, Du J, Zhang J, Zhang XY *et al.* Sleep duration, brain structure, and
psychiatric and cognitive problems in children. *Mol Psychiatry* 2021. doi:10.1038/s41380-020-
0663-2.
857
858