Early signals of significantly increased vaccine breakthrough, decreased hospitalization rates, and less severe disease in patients with COVID-19 caused by the Omicron variant of SARS-CoV-2 in Houston, Texas

Paul A. Christensen,[*,†,‡,§] Randall J. Olsen,[*,†,‡,§] S. Wesley Long,[*,†,‡,§] Richard Snehal,[†]

James J. Davis,[¶,**] Matthew Ojeda Saavedra,[†] Kristina Reppond,[†] Madison N. Shyer,[†]

Jessica Cambric,[†] Ryan Gadd,[†] Rashi M. Thakur,[†] Akanksha Batajoo,[†] Regan

Mangham,[†] Sindy Pena,[†] Trina Trinh,[†] Jacob C. Kinskey,[†] Guy Williams,[†] Robert

Olson,[¶,**] Jimmy Gollihar,[‡] James M. Musser[†,‡,§,††]

*Contributed equally;

[†]Laboratory of Human Molecular and Translational Human Infectious Diseases and

[‡]Laboratory of Antibody Discovery and Accelerated Protein Therapeutics, Center for

Infectious Diseases, Houston Methodist Research Institute and Department of

Pathology and Genomic Medicine, Houston Methodist Hospital, Houston, Texas;

[§]Department of Pathology and Laboratory Medicine, Weill Cornell Medical College, New

York, New York;

¶*Consortium for Advanced Science and Engineering, University of Chicago, Chicago,*

Illinois;

**Computing, Environment and Life Sciences, Argonne National Laboratory, Lemont,*

Illinois;

††*Address correspondence to James M. Musser, M.D., Ph.D., Department of Pathology*

and Genomic Medicine, Houston Methodist Research Institute, 6565 Fannin Street,

Suite B490, Houston, Texas 77030. Tel: 713.441.5890, E-mail:

jmmusser@houstonmethodist.org

Running head: Omicron variant in Houston, Texas

Disclosures: None.

Funding: This project was supported by the Houston Methodist Academic Institute

Infectious Diseases Fund; and in part with funds from the National Institute of Allergy

and Infectious Diseases, National Institutes of Health, Department of Health and Human

Services, under Contract No. 75N93019C00076 (J.J.D.).

Number of text pages: 13

Number of figures: 2

Number of tables: 2

Abstract

Genetic variants of SARS-CoV-2 continue to dramatically alter the landscape of the

COVID-19 pandemic. The recently described variant of concern designated Omicron

(B.1.1.529) has rapidly spread worldwide and is now responsible for the majority of

COVID-19 cases in many countries. Because Omicron was recognized very recently,

many knowledge gaps exist about its epidemiology, clinical severity, and disease

course. A comprehensive genome sequencing study of SARS-CoV-2 in the Houston

Methodist healthcare system identified 1,313 symptomatic patients with infections

caused by Omicron from late November 2021 through December 20, 2021. Omicron

very rapidly increased in only three weeks to cause 90% of all new COVID-19 cases.

Compared to patients infected with either Alpha or Delta variants in our healthcare

system, Omicron patients were significantly younger, had significantly increased

vaccine breakthrough rates, and were significantly less likely to be hospitalized.

Omicron patients required less intense respiratory support and had a shorter length of

hospital stay, consistent with decreased disease severity. Although the number of

Omicron patients we studied is relatively small, in the aggregate the data document the

unusually rapid spread and increased occurrence of COVID-19 caused by the Omicron

variant in metropolitan Houston, and provide information about disease character.

63   [Introduction]
64
65
Over the last 14 months, the Alpha and Delta variants of concern (VOCs) of SARS-

CoV-2 have caused two distinct COVID-19 disease surges in the United States,

Southeast Asia, Europe, and elsewhere (https://www.cdc.gov/coronavirus/2019-

ncov/cases-updates/variant-surveillance/variant-info.html, last accessed December 30,

2021; https://www.gov.uk/government/collections/new-sars-cov-2-variant, last accessed

December 30, 2021)[1], and remodeled the landscape of human behavior and many

societies. Delta replaced the Alpha variant as the cause of virtually all COVID-19 in

many countries (https://www.who.int/publications/m/item/weekly-epidemiological-

update-on-covid-19---13-july-2021, last accessed August 18, 2021;

https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditions

anddiseases/bulletins/coronaviruscovid19infectionsurveypilot/9july2021, last accessed

August 18, 2021).

At the start of the pandemic almost two years ago, the Houston Methodist

healthcare system instituted a comprehensive and integrated population genomics

project designed to sequence all SARS-CoV-2 samples causing COVID-19 in patients

cared for at our facilities, which include eight hospitals located throughout the

metroplex. The project was implemented when the initial Houston Methodist COVID-19

case was diagnosed at the end of February 2020, and has continued unabated[2-6]. This

project was facilitated by the existence of a single large diagnostic laboratory that

serves the entire system and is seamlessly integrated with a research institute with

extensive genomics expertise and capacity. A key goal was to comprehensively map

the population genomics, trajectory, and other features of the pandemic in metropolitan

Houston with a population size of approximately 7.2 million. Houston is the fourth

largest city in the United States, the most ethnically diverse metropolitan area in the

country, and is a major port of entry. To date, SARS-CoV-2 genomes have been

sequenced from greater than 70,000 patient samples. Many features of four distinct

SARS-CoV-2 waves in Houston have been described[2-6].

The successes of rapid SARS-CoV-2 vaccine development and documented

efficacy, coupled with the significant downturn of the disease wave caused by Delta in

Houston and elsewhere in fall, 2021[6], suggested that the pandemic was abating.

However, the identification of a new VOC designated B.1.1.529 and commonly known

as Omicron that has spread rapidly in South Africa and the UK has tempered this

optimism[7-9]. Inasmuch as Omicron was recognized very recently, and much is not

known about its epidemiology and clinical characteristics and course, we used our

integrated infrastructure in an effort to address some pertinent knowledge gaps.

Genome sequencing identified 1,313 COVID-19 patients with symptomatic disease

caused by Omicron in the Houston Methodist healthcare system beginning in late

November 2021 and ending December 20, 2021. In three weeks Omicron spread

throughout the Houston metropolitan region and became the cause of 90% of new

COVID-19 cases. Although the number of our Omicron patients is relatively small,

compared to patients infected with either Alpha or Delta variants and cared for in our

system, significantly fewer Omicron patients were hospitalized, and those who were

hospitalized required significantly less intense respiratory support and had a shorter

length of stay. We cautiously interpret our findings to be consistent with decreased

disease severity among Houston Methodist Omicron patients. Many factors undoubtedly

have contributed, including but not limited to increased vaccination uptake, population

immunity, and patient demographics such as younger age. The extent to which our

findings translate to other cities and other patient populations, including children, is

unknown. Our study was not designed to address possible intrinsic differences in

virulence of Omicron compared to Alpha, Delta, and other VOCs. We believe this topic

remains an open question that warrants further investigation.

Materials and Methods

Patient Specimens

Specimens were obtained from patients registered at Houston Methodist facilities (e.g.,

hospitals and urgent care centers), and institutions in the Houston metropolitan region

that use our laboratory services. The great majority of individuals had signs or

symptoms consistent with COVID-19 disease. For analyses focusing on patients with

COVID-19 caused by the Omicron variant, samples obtained from November 27, 2021

through December 20, 2021 were used. This time frame was chosen because it

represents the period during which an Omicron variant was first identified in our

healthcare system and the last date of specimen collection used to generate genome

sequence data for this manuscript. For analyses comparing features of patients infected

with the Omicron VOC and Alpha and Delta VOCs, all patients documented to be

infected with these variants in the Houston Methodist system were studied. The study

included 36,793 unique patients identified in this time frame for whom we had SARS-

CoV-2 genome sequences. The work was approved by the Houston Methodist

Research Institute Institutional Review Board (IRB1010-0199).

SARS-CoV-2 Molecular Diagnostic Testing

Specimens obtained from symptomatic patients with a suspicion for COVID-19 disease

were tested in the Molecular Diagnostics Laboratory at Houston Methodist Hospital

using assays granted Emergency Use Authorization (EUA) from the FDA

(https://www.fda.gov/medical-devices/emergency-situations-medical-devices/faqs-

diagnostic-testing-sars-cov-2#offeringtests, last accessed June 7, 2021). Multiple

molecular testing platforms were used, including the COVID-19 test or RP2.1 test with

BioFire Film Array instruments, the Xpert Xpress SARS-CoV-2 test using Cepheid

GeneXpert Infinity or Cepheid GeneXpert Xpress IV instruments, the Cobas SARS-

CoV-2 & Influenza A/B Assay using the Roche Liat system, the SARS-CoV-2 Assay

using the Hologic Panther instrument, the Aptima SARS-CoV-2 Assay using the Hologic

Panther Fusion system, the Cobas SARS-CoV-2 test using the Roche 6800 system,

and the SARS-CoV-2 assay using Abbott Alinity m instruments. Virtually all tests were

performed on material obtained from nasopharyngeal swabs immersed in universal

transport media (UTM); oropharyngeal or nasal swabs, bronchoalveolar lavage fluid, or

sputum treated with dithiothreitol (DTT) were sometimes used. Standardized specimen

collection methods were used (https://vimeo.com/396996468/2228335d56, last

accessed June 7, 2021).

SARS-CoV-2 Genome Sequencing, Genome Analysis, and Identification of

Variants

We sequenced the SARS-CoV-2 genome of >90% of all positive cases in the Houston

Methodist healthcare system during the period studied. Libraries for whole SARS-CoV-2

genome sequencing were prepared according to version 4

(https://community.artic.network/t/sars-cov-2-version-4-scheme-release/312, last

accessed August 19, 2021) of the ARTIC nCoV-2019 sequencing protocol. The semi-

automated workflow used has been described previously[2-6]. Sequence reads were

generated with an Illumina NovaSeq 6000 instrument.

Viral genomes were assembled with the BV-BRC SARS-Cov2 assembly service

(https://www.bv-brc.org/app/ComprehensiveSARS2Analysis, last accessed June 7,

2021, requires registration). The pipeline currently uses seqtk version 1.3-r117 for

sequence trimming (https://github.com/lh3/seqtk.git) and minimap version 2.17 for

aligning reads against the Wuhan-Hu-1 (NC_045512.2) reference genome. Samtools

version 1.11 was used for sequence and file manipulation, where maximum depth and

minimum depth parameters in mpileup were set to 8,000 and 3, respectively. iVar

version 1.3.1 was used for primer trimming and variant calling. Genetic lineages,

VOCs, and variants of interest (VOIs) were identified based on genome sequence data

and designated by Pangolin v. 3.1.17 with pangoLEARN module 2021-12-06

(https://cov-lineages.org/resources/pangolin.html, last accessed December 12, 2021).

Genome data used in this study have been deposited to GISAID www.gisaid.org.

S-Gene Target-Failure Assay

An S-gene target-failure (SGTF) assay (TaqPath COVID-19 Combo Kit Thermo Fisher,

Inc.), was used as a surrogate marker for the Omicron VOC for some specimens

collected between December 18, 2021 and December 22, 2021. From November 1,

2021 onward, only Delta and Omicron were circulating in metropolitan Houston, as

documented by whole-genome sequence data. Thus, samples with S-gene target failure

were classified as Omicron, whereas samples yielding amplification of the S-gene were

classified as a Delta variant.

Patient Metadata and Geospatial Analysis

Patient metadata were acquired from the electronic medical record by standard

informatics methods. Figures showing geospatial distribution of spread for Omicron

were generated with Tableau version 2021.2.7 (Tableau Software, LLC, Seattle, WA)

using patient home address zip codes. A vaccination breakthrough case was defined as

a PCR-positive sample from a patient obtained greater than 14 days after full

vaccination (e.g., both doses of the Pfizer or Moderna mRNA vaccines) was completed.

A booster vaccination breakthrough case was defined as a PCR-positive sample from a

patient obtained greater than 14 days after receiving a third vaccine dose. For some

cases, manual chart review was conducted to resolve discrepancies or clarify

ambiguities.

Results

Omicron Epidemiologic Wave

The first Houston Methodist patient infected with an Omicron variant was identified at

the end of November 2021, a time when the Delta VOC was responsible for all COVID-

19 cases in metropolitan Houston[6]. During this period, the metropolitan area was

experiencing a steady decrease in total number of new COVID-19 cases (**Figure 1,**

Figure 2).

Omicron increased in frequency unusually rapidly over a three-week period in

December (**Figure 1, Figure 2).** By December 20, the genome sequence data showed

that Omicron accounted for >90% of all new COVID-19 cases in our healthcare system

(**Figure 2**). This represents the fifth wave of COVID-19 cases in metropolitan Houston

(**Figure 1**). The estimated case doubling time during this three-week period was

approximately 1.8 days (**Figure 2**), which means that Omicron increased in relative

frequency approximately three times faster than Delta had increased in our area[6], an

unprecedented trajectory for SARS-COV-2 infections.

Consistent with extensive infections caused by Omicron in southern Africa and

elsewhere (https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-

classifications.html, last accessed December 28, 2021;

https://www.gov.uk/government/collections/new-sars-cov-2-variant, last accessed

December 28, 2021), several patients had very recent travel histories to countries with a

high prevalence of this VOC, suggesting acquisition of virus by some cases from abroad

and importation into Houston. However, the great majority of Omicron patients had no

documented travel outside the US and undoubtedly acquired the infection domestically,

either in Houston or elsewhere.

To understand the geospatial distribution of Omicron in metropolitan Houston,

patient metadata were acquired from the electronic medical record by standard

informatics methods, and home address zip codes were used to visualize virus spread

(**Figure 2**). The 1,313 Houston Methodist patients infected with Omicron during this

period were distributed widely throughout metropolitan Houston, with 200 different zip

codes represented (**Figure 2**). The widespread distribution of Omicron in the Houston

metroplex in an extremely short period of time reflects the ability of this variant to spread

unusually rapidly and effectively between individuals, and cause symptomatic disease.

Comparison of Omicron, Alpha, and Delta COVID-19 Cases

There is a considerable lack of detailed information about patients in the United States

and elsewhere with COVID-19 caused by the Omicron VOC. We compared available

metadata for all Houston Methodist patients infected with Omicron, Alpha, and Delta

VOCs (**Table 1, Table 2**). The populations differed significantly in many characteristics,

including median age, hospital admission rates, maximum respiratory support, rate of

vaccine breakthrough, and median length of stay (**Table 1, Table 2**).

Patients infected with Omicron were significantly younger than Alpha and Delta

patients (**Table 1, Table 2**). Importantly, Omicron patients were hospitalized significantly

less frequently than patients infected with either the Alpha or Delta variants, and had a

significantly shorter median hospital length of stay (**Table 1, Table 2**).

We next analyzed Omicron vaccine breakthrough cases (**Table 1, Table 2**). We

found 675 of the 1313 total Omicron patients (51.4%) for whom we have whole genome

sequence data met the CDC definition of vaccine breakthrough cases (**Table 1, Table

2**). There was no simple relationship between the time elapsed since administration of

the second vaccination dose and the date of vaccination breakthrough. These 675

patients received either two doses of the Pfizer-BioNTech BNT162b2 ($n = 485$, 72%) or

Moderna mRNA-1273 ($n = 157$, 23%), or one dose of J&J/Janssen JNJ-78436735 ($n =$

33, 5%) vaccine. This distribution reflects the majority use of BNT162b2 vaccination

doses in our health system. Compared to either Alpha or Delta patients, a significantly

greater percentage of patients with breakthrough cases was caused by the Omicron

VOC (51.4% compared to 3.2% and 24.3% for Alpha and Delta VOCs, respectively)

(**Table 1, Table 2**). We next analyzed individuals with breakthrough cases after

receiving a third (booster) dose of either the Pfizer-BioNTech BNT162b2 or Moderna

mRNA-1273 vaccine. We found that 140 (10.7%) of the 1,313 Omicron patients met this

criteria. Consistent with Omicron causing a significantly increased number of vaccine

breakthrough cases, it has been reported that this variant has reduced sensitivity to

antibody neutralization *in vitro*, likely in large part due to the extensive number of amino

acid and other structural changes occurring in Omicron spike protein[10-31].

An Omicron Sublineage With Spike Protein Polymorphism L452R

The L452R amino acid change in spike protein of SARS-CoV-2 has arisen by

convergent evolution in multiple genetic lineages throughout the course of the

pandemic, most notably in the Delta VOC. L452R is located in the receptor binding

domain of spike protein and has been associated with decreased antibody

neutralization, escape from cell mediated immunity, and increased affinity for the ACE2

receptor[32, 33]. L452R was present in samples of Omicron collected in South Africa as

early as November 11, 2021. We found that all 43 specimens collected over the first

nine days after we identified Omicron in Houston Methodist patients had the 452R

amino acid replacement. Our first L452 isolate was collected on day 10. By day 16, over

half the Omicron isolates collected were L452, and the proportion of 452R isolates

continued to diminish. Among the 65,109 Omicron samples deposited in GISAID as of

28 December 2021, only 727 (1.1%) have the 452R amino acid replacement and over

half of these were from Houston Methodist patients. The initial high prevalence of

Omicron containing 452R in spike protein in our population may represent a founder

effect of the original genotype introduced into Houston.

Spike-Gene Target-Failure Assay

To estimate Omicron variant frequency in patient samples not yet sequenced, we

performed the TaqPath COVID-19 Combo Kit assay (ThermoFisher) on 657 samples

collected from symptomatic patients between December 18, 2021 and December 22,

2021. In total, 604 (91.9%) of patient samples yielded a RT-PCR result with S-gene

target-failure indicative of the Omicron variant. These data are consistent with the

increasing frequency of new cases of COVID-19 caused by Omicron in our population

(**Figure 2**).

Lack of Omicron Variant BA.2 in Houston Methodist Samples

The BA.2 Omicron sublineage was first identified in November 2021 in Australia in a

patient who had traveled to South Africa (https://github.com/cov-lineages/pango-

designation/issues/359; last accessed December 30, 2021). This sublineage does not

have the full set of polymorphisms characteristic of BA.1 (B.1.1.529) and also has

additional mutations unique to it (https://github.com/cov-lineages/pango-

designation/issues/361; last accessed December 30, 2021). One important difference is

that sublineage BA.2 lacks the spike protein deletion at amino acid 69/70 which means

that it will not be detected by the SGTF assay. We inspected all full genome sequences

present in our large database and this sublineage was not present. This includes all

specimens taken from symptomatic patients or asymptomatic individuals. Thus, it is

likely that routine use of an SGTF assay in our population will identify all or virtually all

patients infected with Omicron.

Discussion

In this work, we describe information relevant to the surging Omicron wave in metropolitan Houston. In three weeks, Omicron was first identified in our population and rapidly increased to cause 90% of all new COVID-19 cases, with an estimated case doubling time of 1.8 days. The study was based on genome sequence analysis of 1,313 Omicron samples taken from socioeconomically, geographically, and ethnically diverse symptomatic patients. Several key findings were made, including (i) the Omicron VOC rapidly increased as a cause of COVID-19 and spread throughout the metroplex in an unusually short period of time, far faster than any other SARS-CoV-2 variant; (ii) Omicron caused significantly more vaccine breakthrough cases than the Alpha or Delta VOCs; (iii) Omicron patients were significantly younger than Alpha or Delta patients; (iv) significantly fewer Omicron patients required hospitalization compared to Alpha and Delta patients; (v) the median length of stay for hospitalized Omicron patients was significantly shorter than for Alpha and Delta patients, and consistent with this observation, the maximum respiratory support required for Omicron patients was significantly less than for Alpha or Delta patients. Our findings are largely consistent with many aspects of Omicron data reported from the UK, South Africa, and Canada[7-9, 34-36] and are consistent with experimental animal infection data suggesting that Omicron causes less severe disease in mice and hamsters[37-40]. This study was facilitated by a comprehensive and integrated population genomics and epidemiology project[2-6] implemented at the end of February, 2020, when the initial COVID-19 case was diagnosed in the Houston Methodist healthcare system.

Because we sequence the genome of approximately 90% of SARS-CoV-2

causing COVID-19 in our diverse Houston Methodist patient population, and have done

so for almost two years, this means we are continuously monitoring the composition of

this virus in a major US metroplex. This affords us the opportunity to rapidly assess

changes in SARS-CoV-2 population genomic structure in the fourth largest city in the

US. However, our study has several limitations. Although we sequenced the genomes

of SARS-CoV-2 causing 90% of all Houston Methodist COVID-19 cases in the study

period, this sample represents only approximately 5% of cases reported in the

metropolitan region. Our patient population will underrepresent some demographic

groups, for example homeless individuals and pediatric patients. The samples

sequenced in this study were obtained from symptomatic individuals, which means that

it is possible that we failed to identify Omicron subvariants or features preferentially

represented in asymptomatic individuals. It is likely that our study included some

patients where Omicron was detected on hospital admission but was incidental to the

primary cause of admission.

In the aggregate, our data add critical new information to features of Omicron

genomic epidemiology and patient characteristics in the US. Further, the present study

highlights the importance of analyzing SARS-CoV-2 genome data integrated with

patient metadata and stresses the need to continue to do this in near-real time as the

Omicron surge continues, the virus evolves, and new variants with potentially altered

fitness and biomedically relevant phenotypes are generated. Analyses of this type are

also important in the context of vaccine formulation and long COVID, an increasing

health and economic problem globally. Finally, the strategy we have used in this and

previous studies[2-6] are readily applicable to future infectious diseases problems that

warrant special attention.

Acknowledgments

We thank Drs. Marc Boom and Dirk Sostman for their ongoing support, and Dr. Sasha

M. Pejerrey for editorial contributions. The research was supported by the Houston

Methodist Academic Institute Infectious Diseases Fund and many generous Houston

philanthropists. The funders had no role in the design and conduct of the study;

collection, management, analysis, and interpretation of the data; preparation, review, or

approval of the manuscript; and decision to submit the manuscript for publication.

We declare that we have no conflict of interest.

Author Contributions

P.A.C., R.J.O., S.W.L., and J.M.M. had full access to all study data and take

responsibility for the integrity of the data and the accuracy of the data analysis; concept

and design by J.M.M., P.A.C., R.J.O., and S.W.L; data acquisition, analysis, or

interpretation by all authors; drafting of the manuscript by all authors; statistical analysis

by P.A.C.; funding obtained by J.M.M. and J.J.D.; and overall supervision by J.M.M.

P.A.C., R.J.O., and S.W.L. contributed equally and are co-first authors.

References

[1] Dhar MS, Marwal R, Radhakrishnan V, Ponnusamy K, Jolly B, Bhoyar RC, et al.:
Genomic characterization and Epidemiology of an emerging SARS-CoV-2 variant in
Delhi, India. medRxiv 2021. [Preprint]. doi:2021.2006.2002.21258076

[2] Long SW, Olsen RJ, Christensen PA, Bernard DW, Davis JJ, Shukla M, Nguyen M,
Saavedra MO, Yerramilli P, Pruitt L, Subedi S, Kuo HC, Hendrickson H, Eskandari G,
Nguyen HAT, Long JH, Kumaraswami M, Goike J, Boutz D, Gollihar J, McLellan JS,
Chou CW, Javanmardi K, Finkelstein IJ, Musser JM: Molecular Architecture of Early
Dissemination and Massive Second Wave of the SARS-CoV-2 Virus in a Major
Metropolitan Area. mBio 2020, 11

[3] Musser JM, Olsen RJ, Christensen PA, Long SW, Subedi S, Davis JJ, Gollihar J:
Rapid, widespread, and preferential increase of SARS-CoV-2 B.1.1.7 variant in
Houston, TX, revealed by 8,857 genome sequences. medRxiv 2021. [Preprint].
doi:2021.2003.2016.21253753

[4] Olsen RJ, Christensen PA, Long SW, Subedi S, Hodjat P, Olson R, Nguyen M,
Davis JJ, Yerramilli P, Saavedra MO, Pruitt L, Reppond K, Shyer MN, Cambric J, Gadd
R, Thakur RM, Batajoo A, Finkelstein IJ, Gollihar J, Musser JM: Trajectory of Growth of
Severe Acute Respiratory (SARS-CoV-2) Syndrome Coronavirus 2 Variants in Houston,
Texas, January through May 2021, Based on 12,476 Genome Sequences. Am J Pathol
2021, Oct;191(10):1754-1773.

[5] Long SW, Olsen RJ, Christensen PA, Subedi S, Olson R, Davis JJ, Saavedra MO,
Yerramilli P, Pruitt L, Reppond K, Shyer MN, Cambric J, Finkelstein IJ, Gollihar J,
Musser JM: Sequence Analysis of 20,453 SARS-CoV-2 Genomes from the Houston

Metropolitan Area Identifies the Emergence and Widespread Distribution of Multiple

Isolates of All Major Variants of Concern. Am J Pathol 2021, Jun;191(6):983-992

[6] Christensen PA, Olsen RJ, Long SW, Subedi S, Davis JJ, Hodjat P, Walley DR,

Kinskey JC, Saavedra MO, Pruitt L, Reppond K, Shyer MN, Cambric J, Gadd R, Thakur

RM, Batajoo A, Mangham R, Pena S, Trinh T, Yerramilli P, Nguyen M, Olson R, Snehal

R, Gollihar J, Musser JM: Delta Variants of SARS-CoV-2 Cause Significantly Increased

Vaccine Breakthrough COVID-19 Cases in Houston, Texas. Am J Pathol 2021, Nov

11;S0002-9440(21)00480-6.

[7] Viana R, Moyo S, Amoako DG, Tegally H, Scheepers C, Lessells RJ, et al.: Rapid

epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. medRxiv

2021. [Preprint]. doi:2021.2012.2019.21268028

[8] Elliott P, Bodinier B, Eales O, Wang H, Haw D, Elliott J, Whitaker M, Jonnerby J,

Tang D, Walters CE, Atchison C, Diggle PJ, Page AJ, Trotter AJ, Ashby D, Barclay W,

Taylor G, Ward H, Darzi A, Cooke GS, Chadeau-Hyam M, Donnelly CA: Rapid increase

in Omicron infections in England during December 2021: REACT-1 study. medRxiv

2021. [Preprint]. doi:2021.2012.2022.21268252

[9] Sheikh AK, Steven; Woolhouse, Mark; McMenamin, Jim; Robertson, Chris. : Severity

of Omicron variant of concern and vaccine effectiveness against symptomatic disease:

national cohort with nested test negative design study in Scotland. The University of

Edinburgh 2021

[10] Meng B, Ferreira I, Abdullahi A, Kemp SA, Goonawardane N, Papa G, Fatihi S,

Charles O, Collier D, Collaboration C-NBC-, Consortium TGtPJ, Choi J, Hyeon Lee J,

Mlcochova P, James L, Doffinger R, Thukral L, Sato K, Gupta RK: SARS-CoV-2

Omicron spike mediated immune escape, infectivity and cell-cell fusion. bioRxiv 2021.

[Preprint]. doi:2021.2012.2017.473248

[11] Zeng C, Evans JP, Qu P, Faraone J, Zheng Y-M, Carlin C, Bednash JS, Zhou T,

Lozanski G, Mallampalli R, Saif LJ, Oltz EM, Mohler P, Xu K, Gumina RJ, Liu S-L:

Neutralization and Stability of SARS-CoV-2 Omicron Variant. bioRxiv 2021. [Preprint].

doi:2021.2012.2016.472934

[12] Jacobsen H, Strengert M, Maass H, Ynga Durand MA, Kessel B, Harries M, Rand

U, Abassi L, Kim Y, Lueddecke T, Hernandez P, Ortmann J, Heise J-K, Castell S,

Gornyk D, Gloeckner S, Melhorn V, Lange B, Dulovic A, Haering J, Junker D,

Schneiderhan-Marra N, Poehlmann S, Hoffmann M, Krause G, Cicin-Sain L: Diminished

neutralization responses towards SARS-CoV-2 Omicron VoC after mRNA or vector-

based COVID-19 vaccinations. medRxiv 2021. [Preprint].

doi:2021.2012.2021.21267898

[13] Eggink D, Andeweg SP, Vennema H, van Maarseveen N, Vermaas K, Vlaemynck

B, Schepers R, van Gageldonk-Lafeber AB, van den Hof S, Reusken CBEM, Knol MJ:

Increased risk of infection with SARS-CoV-2 Omicron compared to Delta in vaccinated

and previously infected individuals, the Netherlands, 22 November to 19 December

2021. medRxiv 2021. [Preprint]. doi:2021.2012.2020.21268121

[14] Edara V-V, Manning KE, Ellis M, Lai L, Moore KM, Foster SL, Floyd K, Davis-

Gardner ME, Mantus G, Nyhoff LE, Bechnack S, Alaaeddine G, Naji A, Samaha H, Lee

M, Bristow L, Hussaini L, Ciric CR, Nguyen P-V, Gagne M, Roberts-Torres J, Henry AR,

Godbole S, Grakoui A, Sexton M, Piantadosi A, Waggoner JJ, Douek DC, Anderson EJ,

Rouphael N, Wrammert J, Suthar MS: mRNA-1273 and BNT162b2 mRNA vaccines

have reduced neutralizing activity against the SARS-CoV-2 Omicron variant. bioRxiv

2021. [Preprint]. doi:2021.2012.2020.473557

[15] Zou j, Xia H, Xie X, Kurhade C, Machado RR, Weaver SC, Ren P, Shi P-Y:

Neutralization against Omicron SARS-CoV-2 from previous non-Omicron infection.

bioRxiv 2021. [Preprint]. doi:2021.2012.2020.473584

[16] Ikemura N, Hoshino A, Higuchi Y, Taminishi S, Inaba T, Matoba S: SARS-CoV-2

Omicron variant escapes neutralization by vaccinated and convalescent sera and

therapeutic monoclonal antibodies. medRxiv 2021. [Preprint].

doi:2021.2012.2013.21267761

[17] Dejnirattisai W, Shaw RH, Supasa P, Liu C, Stuart AS, Pollard AJ, Liu X, Lambe T,

Crook D, Stuart DI, Mongkolsapaya J, Nguyen-Van-Tam JS, Snape MD, Screaton GR,

group tC-Cs: Reduced neutralisation of SARS-COV-2 Omicron-B.1.1.529 variant by

post-immunisation serum. medRxiv 2021. [Preprint]. doi:2021.2012.2010.21267534

[18] Cameroni E, Saliba C, Bowen JE, Rosen LE, Culap K, Pinto D, et al.: Broadly

neutralizing antibodies overcome SARS-CoV-2 Omicron antigenic shift. bioRxiv 2021.

[Preprint]. doi:2021.2012.2012.472269

[19] Liu L, Iketani S, Guo Y, Chan JF-W, Wang M, Liu L, Luo Y, Chu H, Huang Y, Nair

MS, Yu J, Chik KK-H, Yuen TT-T, Yoon C, To KK-W, Chen H, Yin MT, Sobieszczyk ME,

Huang Y, Wang HH, Sheng Z, Yuen K-Y, Ho DD: Striking Antibody Evasion Manifested

by the Omicron Variant of SARS-CoV-2. bioRxiv 2021. [Preprint].

doi:2021.2012.2014.472719

[20] Planas D, Saunders N, Maes P, Benhassine FG, Planchais C, Porrot F, Staropoli I,

Lemoine F, Pere H, Veyer D, Puech J, Rodary J, Bolland WH, Buchrieser J, Baele G,

Dellicour S, Raymenants J, Gorissen S, Geenen C, Vanmechelen B, Wawina T, Marti J,

Cuypers L, Seve A, Hocqueloux L, Prazuck T, Loriere ES, REY F, Bruel T, Mouquet H,

Andre E, Schwartz O: Considerable escape of SARS-CoV-2 variant Omicron to

antibody neutralization. bioRxiv 2021. [Preprint]. doi:2021.2012.2014.472630

[21] Andrews N, Stowe J, Kirsebom F, Toffa S, Rickeard T, Gallagher E, Gower C, Kall

M, Groves N, O'Connell A-M, Simons D, Blomquist PB, Zaidi A, Nash S, Aziz NIBA,

Thelwall S, Dabrera G, Myers R, Amirthalingam G, Gharbia S, Barrett JC, Elson R,

Ladhani SN, Ferguson N, Zambon M, Campbell CN, Brown K, Hopkins S, Chand M,

Ramsay M, Bernal JL: Effectiveness of COVID-19 vaccines against the Omicron

(B.1.1.529) variant of concern. medRxiv 2021. [Preprint]. doi:2021.2012.2014.21267615

[22] Yu X, Wei D, Xu W, Li Y, Li X, Zhang X, Qu J, Yang Z, Chen E: Reduced sensitivity

of SARS-CoV-2 Omicron variant to booster-enhanced neutralization. medRxiv 2021.

[Preprint]. doi:2021.2012.2017.21267961

[23] Cele S, Jackson L, Khan K, Khoury DS, Moyo-Gwete T, Tegally H, Scheepers C,

Amoako D, Karim F, Bernstein M, Lustig G, Archary D, Smith M, Ganga Y, Jule Z,

Reedoy K, Cromer D, San JE, Hwa S-H, Giandhari J, Blackburn JM, Gosnell BI, Karim

SSA, Hanekom W, NGS-SA, Team C-K, von Gottberg A, Bhiman J, Lessells RJ, Moosa

M-YS, Davenport MP, de Oliveira T, Moore PL, Sigal A: SARS-CoV-2 Omicron has

extensive but incomplete escape of Pfizer BNT162b2 elicited neutralization and requires

ACE2 for infection. medRxiv 2021. [Preprint]. doi:2021.2012.2008.21267417

[24] Cao Y, Wang J, Jian F, Xiao T, Song W, Yisimayi A, Huang W, Li Q, Wang P, An

R, Wang J, Wang Y, Niu X, Yang S, Liang H, Sun H, Li T, Yu Y, Cui Q, Liu S, Yang X,

Du S, Zhang Z, Hao X, Shao F, Jin R, Wang X, Xiao J, Wang Y, Xie XS: B.1.1.529

escapes the majority of SARS-CoV-2 neutralizing antibodies of diverse epitopes.

bioRxiv 2021. [Preprint]. doi:2021.2012.2007.470392

[25] Hansen CH, Schelde AB, Moustsen-Helms IR, Emborg H-D, Krause TG, Moelbak

K, Valentiner-Branth P, Institut TIDPGaSS: Vaccine effectiveness against SARS-CoV-2

infection with the Omicron or Delta variants following a two-dose or booster BNT162b2

or mRNA-1273 vaccination series: A Danish cohort study. medRxiv 2021. [Preprint].

doi:2021.2012.2020.21267966

[26] Syed AM, Ciling A, Khalid MM, Sreekumar B, Kumar GR, Silva I, Milbes B, Kojima

N, Hess V, Shacreaw M, Lopez L, Brobeck M, Turner F, Spraggon L, Taha TY, Tabata

T, Chen IP, Ott M, Doudna JA: Omicron mutations enhance infectivity and reduce

antibody neutralization of SARS-CoV-2 virus-like particles. medRxiv 2021. [Preprint].

doi:2021.2012.2020.21268048

[27] Sheward DJ, Kim C, Ehling RA, Pankow A, Castro Dopico X, Martin DP, Reddy ST,

Dillner J, Karlsson Hedestam GB, Albert J, Murrell B: Variable loss of antibody potency

against SARS-CoV-2 B.1.1.529 (Omicron). bioRxiv 2021. [Preprint].

doi:2021.2012.2019.473354

[28] Haveri A, Solastie A, Ekström N, Österlund P, Nohynek H, Nieminen T, Palmu AA,

Melin M: Neutralizing antibodies to SARS-CoV-2 Omicron variant after 3rd mRNA

vaccination in health care workers and elderly subjects and response to a single dose in

previously infected adults. medRxiv 2021. [Preprint]. doi:2021.2012.2022.21268273

[29] Arien KK, Heyndrickx L, Michiels J, Vereecken K, Van Lent K, Coppens S, Pannus

P, Martens GA, Van Esbroeck M, Goossens ME, Marchant A, Bartholomeeusen K,

Desombere I: Three doses of the BNT162b2 vaccine confer neutralising antibody

capacity against the SARS-CoV-2 B.1.1.529 (Omicron) variant of concern. medRxiv

2021. [Preprint]. doi:2021.2012.2023.21268316

[30] Willett BJ, Grove J, MacLean O, Wilkie C, Logan N, De Lorenzo G, et al.: The

hyper-transmissible SARS-CoV-2 Omicron variant exhibits significant antigenic change,

vaccine escape and a switch in cell entry mechanism. medRxiv 2022. [Preprint].

doi:2022.2001.2003.21268111

[31] Boschi C, Colson P, Bancod A, Moal V, La Scola B: Omicron variant escapes

therapeutic mAbs contrary to eight prior main VOC. bioRxiv 2022. [Preprint].

doi:2022.2001.2003.474769

[32] Motozono C, Toyoda M, Zahradnik J, Saito A, Nasser H, Tan TS, Ngare I, Kimura I,

Uriu K, Kosugi Y, Yue Y, Shimizu R, Ito J, Torii S, Yonekawa A, Shimono N, Nagasaki

Y, Minami R, Toya T, Sekiya N, Fukuhara T, Matsuura Y, Schreiber G, Ikeda T,

Nakagawa S, Ueno T, Sato K: SARS-CoV-2 spike L452R variant evades cellular

immunity and increases infectivity. Cell Host Microbe 2021, 29:1124-1136.e1111

[33] Li Q, Wu J, Nie J, Zhang L, Hao H, Liu S, Zhao C, Zhang Q, Liu H, Nie L, Qin H,

Wang M, Lu Q, Li X, Sun Q, Liu J, Zhang L, Li X, Huang W, Wang Y: The Impact of

Mutations in SARS-CoV-2 Spike on Viral Infectivity and Antigenicity. Cell 2020,

182:1284-1294 e1289

[34] Ulloa AC, Buchan SA, Daneman N, Brown KA: Early estimates of SARS-CoV-2

Omicron variant severity based on a matched cohort study, Ontario, Canada. medRxiv

2021. [Preprint]. doi:2021.2012.2024.21268382

[35] Wolter N, Jassat W, Walaza S, Welch R, Moultrie H, Groome M, Amoako DG,

Everatt J, Bhiman JN, Scheepers C, Tebeila N, Chiwandire N, du Plessis M, Govender

N, Ismail A, Glass A, Mlisana K, Stevens W, Treurnicht FK, Makatini Z, Hsiao N-y,

Parboosing R, Wadula J, Hussey H, Davies M-A, Boulle A, von Gottberg A, Cohen C:

Early assessment of the clinical severity of the SARS-CoV-2 Omicron variant in South

Africa. medRxiv 2021. [Preprint]. doi:2021.2012.2021.21268116

[36] SARS-CoV-2 variants of concern and variants under investigation in England

Technical briefing: Update on hospitalisation and vaccine effectiveness for Omicron

VOC-21NOV-01 (B.1.1.529) UK Health Security: UK Health Security, 2021.

[37] Bentley EG, Kirby A, Sharma P, Kipar A, Mega DF, Bramwell C, Penrice-Randal R,

Prince T, Brown JC, Zhou J, Screaton GR, Barclay WS, Owen A, Hiscox JA, Stewart

JP: SARS-CoV-2 Omicron-B.1.1.529 Variant leads to less severe disease than Pango B

and Delta variants strains in a mouse model of severe COVID-19. bioRxiv 2021.

[Preprint]. doi:2021.2012.2026.474085

[38] Abdelnabi R, Foo CS, Zhang X, Lemmens V, Maes P, Slechten B, Raymenants J,

André E, Weynand B, Dallemier K, Neyts J: The omicron (B.1.1.529) SARS-CoV-2

variant of concern does not readily infect Syrian hamsters. bioRxiv 2021. [Preprint].

doi:2021.2012.2024.474086

[39] Diamond M, Peter H, Tadashi M, Kiyoko I-H, Shun I, Maki K, et al.: The SARS-

CoV-2 B.1.1.529 Omicron virus causes attenuated infection and disease in mice and

hamsters Nature Portfolio 2022. 29 December 2021. [Preprint].

doi:https://doi.org/10.21203/rs.3.rs-1211792/v1

[40] McMahan K, Giffin V, Tostanoski L, Chung B, Siamatu M, Suthar M, Halfmann P,

Kawaoka Y, Piedra-Mora C, Martinot A, Kar S, Andersen H, Lewis MG, Barouch DH:

Reduced Pathogenicity of the SARS-CoV-2 Omicron Variant in Hamsters. bioRxiv 2022.

[Preprint]. doi:2022.2001.2002.474743

Figure 1 Epidemiologic curve showing five COVID-19 disease waves in Houston

Methodist patients. Number of new COVID-19 cases (y-axis) totals are shown as a +/-

three-day moving average. Each of the five waves is shown in a different color. The first

and second waves were composed of a heterogenous array of SARS-CoV-2 genotypes.

The Alpha VOC shown in the third wave, the Delta VOC shown in the fourth, and the

Omicron VOC shown in the fifth wave indicate their numeric prominence in those

waves. The figure should not be interpreted to mean that all cases in the third, fourth,

and fifth waves were caused by Alpha, Delta, and Omicron VOCs, respectively. Rather,

they are the dominant single VOCs causing disease in Houston Methodist system

patients in those waves. The fifth wave shown includes data through December 20,

2021. The figure was generated with Tableau version 2021.2.7 (Tableau Software, LLC,

Seattle, WA), and is a modified version of one presented recently[6]. The curve is

essentially superimposable on COVID-19 activity in all metropolitan Houston, Texas.

Figure 2 Increase in Omicron frequency over time and distribution in metropolitan

Houston. The study time frame was November 27, 2021 through December 20, 2021.

A: Omicron logistic growth model. The estimated case doubling time is 1.8 days. **B:**

Cumulative increase in Omicron during the study period; y-axis is the cumulative

number of new COVID-19 Omicron cases. At the end of the study period, Omicron

caused 90% of all COVID-19 cases. **C – E:** Geospatial distribution of Omicron based on

home address zip code for each patient. **C**: November 27 – December 3; **D**: November

27 – December 10; **E**: November 27 – December 20. Note differences in heat map

scale for each panel. Figures were generated using Tableau version 2021.2.7. (Tableau

Software, LLC, Seattle, WA).

Table 1. Summary of pertinent patient metadata for 4,470 unique patients infected with Omicron or Alpha variants.

	Omicron Variant	Alpha Variant	Total	Statistical Analysis
No. (%) with data	1313 (29.4%)	3157 (70.6%)	4470	
Patient Characteristics				
Median Age (Years)	39.2	50.0	46.8	*P*<0.0001 Mann-Whitney
Female	763 (58.1%)	1622 (51.4%)	2385 (53.4%)	*P*<0.0001
Male	550 (41.9%)	1535 (48.6%)	2085 (46.6%)	Fisher's exact test
Ethnicity				
Caucasian	406 (31.0%)	1241 (39.5%)	1647 (37.0%)	*P*<0.0001
Hispanic or Latino	230 (17.6%)	945 (30.1%)	1175 (26.4%)	Chi-square
Black	544 (41.6%)	732 (23.3%)	1276 (28.7%)	
Asian	49 (3.7%)	123 (3.9%)	172 (3.9%)	
Other	7 (0.5%)	32 (1.0%)	39 (0.9%)	
Unavailable	77 (5.9%)	84 (2.7%)	161 (3.6%)	
BMI				
Median BMI	28.6	30.5	30.0	*P*<0.0001 Mann-Whitney
Admission Data				
Admitted	193 (14.7%)	1721 (54.5%)	1914 (42.8%)	*P*<0.0001
Not Admitted	1120 (85.3%)	1436 (45.5%)	2556 (57.2%)	Fisher's exact test Odds Ratio: 0.144 (95% CI 0.122-0.170)
Median LOS (Days) (Discharged patients only)	3.0	5.1	5.0	*P*<0.0001 Mann-Whitney
Max Respiratory Support				
ECMO	1 (0.5%)	7 (0.4%)	8 (0.4%)	*P*<0.0001
Mechanical Ventilation	10 (5.2%)	144 (8.4%)	154 (8.0%)	Chi-square

Non-Invasive Ventilation	16 (8.3%)	164 (9.5%)	180 (9.4%)	
High Flow Oxygen	18 (9.3%)	364 (21.2%)	382 (20.0%)	
Low Flow Oxygen	60 (31.1%)	722 (42.0%)	782 (40.9%)	
Room Air	88 (45.6%)	320 (18.6%)	408 (21.3%)	
Mortality				
Alive	1302 (99.2%)	2987 (94.6%)	4289 (96.0%)	*P*<0.0001 Fisher's exact test Odds Ratio: 0.148 (95% CI 0.080-0.274)
Deceased	11 (0.8%)	170 (5.4%)	181 (4.0%)	
Median PCR Cycle Threshold				
Abbott Alinity	19.2 n=519	22.4 n=1050	n=1569	*P*<0.0001 Mann-Whitney
Hologic Panther	22.0 n=88	24.2 n=359	n=447	*P*=.0950 Mann-Whitney
Vaccine				
Not Fully Vaccinated	638 (48.6%)	3056 (96.8%)	3694 (82.6%)	*P*<0.0001 Fisher's exact test Odds Ratio: 32.012 (95% CI 25.541-40.123)
Fully Vaccinated	675 (51.4%)	101 (3.2%)	776 (17.4%)	

599   BMI: body mass index; CI: confidence interval; ECMO: extracorporeal membrane oxygenation; LOS: length of stay

Table 2. Summary of pertinent patient metadata for 16,995 unique patients

infected with Omicron or Delta variants.

	Omicron Variant	Delta Variant	Total	Statistical Analysis
No. (%) with data	1313 (7.7%)	15682 (92.3%)	16995	
Patient Characteristics				
Median Age (Years)	39.2	48.2	47.5	*P*<0.0001 Mann-Whitney
Female	763 (58.1%)	8109 (51.7%)	8872 (52.2%)	*P*=0.0001
Male	550 (41.9%)	7573 (48.3%)	8123 (47.8%)	Fisher's exact test
Ethnicity				
Caucasian	406 (30.9%)	6874 (43.8%)	7280 (42.8%)	*P*<0.0001
Hispanic or Latino	230 (17.5%)	4166 (26.6%)	4396 (25.9%)	Chi-square
Black	544 (41.4%)	3450 (22.0%)	3994 (23.5%)	
Asian	49 (3.7%)	527 (3.4%)	576 (3.4%)	
Other	7 (0.5%)	112 (0.7%)	119 (0.7%)	
Unavailable	77 (5.9%)	553 (3.5%)	630 (3.7%)	
BMI				
Median BMI	28.6	29.5	29.5	*P*=0.0001 Mann-Whitney
Admission Data				
Admitted	193 (14.7%)	6749 (43.0%)	6942 (40.8%)	*P*<0.0001
Not Admitted	1120 (85.3%)	8933 (57.0%)	10053 (59.2%)	Fisher's exact test Odds Ratio: 0.228 (95% CI 0.195-0.267)
Median LOS (Days) (Discharged patients only)	3.0	5.4	5.3	*P*<0.0001 Mann-Whitney
Max Respiratory Support				
ECMO	1 (0.5%)	19 (0.3%)	20 (0.3%)	*P*<0.0001
Mechanical Ventilation	10 (5.2%)	723 (10.7%)	733 (10.6%)	Chi-square

Non-Invasive Ventilation	16 (8.3%)	638 (9.5%)	654 (9.4%)	
High Flow Oxygen	18 (9.3%)	1785 (26.4%)	1803 (26.0%)	
Low Flow Oxygen	60 (31.1%)	2282 (33.8%)	2342 (33.7%)	
Room Air	88 (45.6%)	1302 (19.3%)	1390 (20.0%)	
Mortality				
Alive	1302 (99.2%)	14855 (94.7%)	16157 (95.1%)	*P*<0.0001 Fisher's exact test
Deceased	11 (0.8%)	827 (5.3%)	838 (4.9%)	Odds Ratio: 0.152 (95% CI 0.083-0.276)
Median PCR Cycle Threshold				
Abbott Alinity	19.2 n=519	21.43 n=5102	n=5621	*P*<0.0001 Mann-Whitney
Hologic Panther	22.0 n=88	22.6 n=1295	n=1383	*P*=.8987 Mann-Whitney
Vaccine				
No vaccine	584 (44.5%)	11387 (72.6%)	11971 (70.4%)	*P*<0.0001 Chi-square
>7 days past 1st Vaccine	54 (4.1%)	492 (3.1%)	546 (3.2%)	
>14 days past 2nd Vaccine	535 (40.7%)	3665 (23.4%)	4200 (24.7%)	
>14 days past 3rd Vaccine	140 (10.7%)	138 (0.9%)	278 (1.6%)	

602   BMI: body mass index; CI: confidence interval; ECMO: extracorporeal membrane oxygenation; LOS: length of stay