

The hyper-transmissible SARS-CoV-2 Omicron variant exhibits significant antigenic change, vaccine escape and a switch in cell entry mechanism

Brian J. Willett*, Joe Grove, Oscar A. MacLean, Craig Wilkie, Nicola Logan, Giuditta De Lorenzo, Wilhelm Furnon, Sam Scott, Maria Manali, Agnieszka Szemiel, Shirin Ashraf, Elen Vink, William T. Harvey, Chris Davis, Richard Orton, Joseph Hughes, Poppy Holland, Vanessa Silva, David Pascall, Kathryn Puxty, Ana da Silva Filipe, Gonzalo Yebra, Sharif Shaaban, Matthew T. G. Holden, Rute Maria Pinto, Rory Gunson, Kate Templeton, Pablo R. Murcia, Arvind H. Patel, The COVID-19 Genomics UK (COG-UK) Consortium§, John Haughney, David L. Robertson, Massimo Palmarini, Surajit Ray & Emma C. Thomson*

§Full list of consortium names and affiliations are listed in the Supplementary information.

***Corresponding authors:** Professor Emma Thomson and Professor Brian Willett, MRC-University of Glasgow Centre for Virus Research, UK, G61 1QH.

emma.thomson@glasgow.ac.uk and brian.willett@glasgow.ac.uk

Abstract

Vaccines based on the spike protein of SARS-CoV-2 are a cornerstone of the public health response to COVID-19. The emergence of hypermutated, increasingly transmissible variants of concern (VOCs) threaten this strategy. Omicron, the fifth VOC to be described, harbours 30 amino acid mutations in spike including 15 in the receptor-binding domain. Here, we demonstrate substantial evasion of neutralisation by Omicron *in vitro* using sera from vaccinated individuals. Importantly, these data are mirrored by a substantial reduction in real-world vaccine effectiveness that is partially restored by booster vaccination. We also demonstrate that Omicron does not induce cell syncytia and favours a TMPRSS2-independent endosomal entry pathway. Such marked changes in antigenicity and replicative biology may underlie the rapid global spread and altered pathogenicity of the Omicron variant.

Main

Vaccination against SARS-CoV-2 is based primarily on vaccines that induce immunity to the spike glycoprotein. These vaccines have become the cornerstone of the global public health response to SARS-CoV-2¹. However, their effectiveness is now being threatened by the emergence of Variants of Concern (VOC) displaying enhanced transmissibility and evasion of host immunity². Of the five VOCs that have emerged, the Beta variant (B.1.351) and Gamma (P.1) variants were associated primarily with immune evasion; they spread locally but never dominated globally. In contrast, the Alpha (B.1.1.7) and Delta (B.1.617.2) VOCs spread globally and were responsible for significant waves of infections and an increase in reproduction number (R_0). The Alpha and Delta variants harbour mutations within the polybasic cleavage site in spike (a H681 in Alpha and R681 in Delta) that enhance cleavage by furin; changes that are associated with enhanced cell entry and may contribute to increased transmissibility. While the Alpha variant spread rapidly, it was in turn replaced by the Delta variant that combined augmented transmissibility with significant immune evasion^{2, 3, 4, 5}.

Omicron (lineage B.1.1.529) is the fifth variant to be named as a VOC by the World Health Organisation (WHO) and was first detected in mid-November 2021 in Botswana, South Africa⁶ and in quarantined travellers in Hong Kong⁷. It has since split into three divergent sublineages (BA.1, BA.2 and BA.3) of which BA.1 now dominates worldwide.

Emerging data indicate that the Omicron variant evades neutralisation by sera obtained from people vaccinated with 1 or 2 doses of vaccine, especially when antibody titres are waning. Indicative studies have shown that 3 doses of spike-based vaccines may provide only partial protection from infection with this variant, including unpublished data made available as a press release from Pfizer. Immune evasion by Omicron may have contributed to the extremely high transmission rates in countries with high vaccination rates or natural immunity (R_0 of 3-5 in the U K ^{8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18}).

In this study, we investigate the antigenic and biological properties of the Omicron variant that might underlie immune evasion and increased transmission of the virus using both *in vitro* assays and real-life population data.

Results

Omicron displays substantial changes within spike predicted to affect antigenicity and furin cleavage

Omicron is characterised by significant changes within the RBD of the spike glycoprotein, regions targeted by class 1,2 and 3 RBD-directed antibodies, and within the N-terminal domain (NTD) supersite (Fig.1a). The G339D, N440K, S477N, T478K, Q498R and N501Y mutations enhance binding of spike to the human ACE2 receptor, while combinations such as Q498R and N501Y may enhance ACE2 binding additively¹⁹. Overall, the Omicron RBD binds to the human ACE2 with approximately double the affinity (x2.4) of the Wuhan RBD⁸. Deep mutational scanning (DMS) estimates at mutated sites are predictive of substantially reduced monoclonal and polyclonal antibody binding and altered binding to human ACE2 (Fig.1b)²⁰. Fourteen mutations (K417N, G446S, E484A, Q493R, G496S, Q498R and to a lesser extent, G339D, S371L, S373P, N440K, S477N, T478K, N501Y and Y505H) may affect antibody binding based on a calculated escape fraction (a quantitative measure of the extent to which a mutation reduces polyclonal antibody binding by DMS). Seven Omicron RBD mutations (K417N, G446S, E484A, Q493R, G496S, Q498R and N501Y) have been shown previously to be associated with decreased antibody binding, importantly falling in epitopes corresponding to three major classes of RBD-specific neutralising antibodies (nAbs). The mutations present in spike also involve key structural epitopes targeted by several monoclonal antibodies in current clinical use. Of these, bamlanivimab, cilgavimab, casirivimab, etesevimab, imdevimab, regdanvimab and tixagevimab bind to the RBM, and neutralisation of Omicron has been shown to be negligible or absent. In contrast, sotrovimab, targets a conserved epitope common to SARS-CoV-1 and SARS-CoV-2 that is outside the RBM and has only a small reduction (x3) in neutralisation potency^{21, 22, 23}. N679K and P681H mutations at the furin cleavage site (FCS) are predicted individually to increase furin cleavage, although the combination of these changes and an adjacent change (H655Y, also present in the Gamma VOC) in the vicinity of the FCS is unknown²⁴.

Omicron bears three deletions (amino acids 69-70, 143-145 and 211) and an insertion (site 214) in the NTD of spike. The 69-70 deletion is also found in the Alpha and Eta (B.1.525) variants and is associated with enhanced fusogenicity and incorporation of cleaved spike into

virions²⁵. This 69-70 deletion is a useful proxy for estimates of Omicron prevalence in the population by S-gene target failure (SGTF) using the TaqPath™ (Applied Biosystems, Pleasanton, CA) diagnostic assay. Deletions in the vicinity of amino acids 143-145 have been shown to affect a range of NTD-specific nAbs^{26, 27}.

Emergence of the Omicron variant in the UK

Despite high vaccination rates and levels of natural immunity following previous exposure in the UK, Omicron has rapidly become dominant. The evolutionary relationships of SARS-CoV-2 variants at a global level are shown in Fig.1c. The first 8 cases of Omicron were detected in the UK on the 27th and 28th November 2021 (2 in England and 6 in Scotland). Due to the rapid spread of Omicron, early genome sequences were highly related with an average genetic divergence between 1 and 7 single nucleotide polymorphisms (SNPs) (Fig.1d). The phylogenetic relationship to Omicron sequences from other countries was consistent with multiple introductions associated with travel to South Africa followed by community transmission.




Fig. 1: Spike amino acid changes typifying the Omicron variant. **a**, Spike homotrimer in open conformation with locations of Omicron amino acid substitutions, deletions (Δ), or insertions (ins) highlighted as spheres with opaque surface representation. Colouring highlights mutations at residues with substitutions impacting RBD-specific antibodies of classes 1 (green), 2 (yellow), and 3 (blue)²⁸, or that belong to the NTD antibody supersite (magenta)²⁶, or that belong to the FCS (orange), with the remainder in grey. These are annotated on the monomer with an 'up' receptor-binding domain. The substitution D614G which is shared by common descent by all lineage B.1 descendants is italicised. The visualisation is made using a complete spike model²⁹ which is in turn based upon a partial cryo-EM structure (RCSB Protein Data Bank (PDB) ID: 6VSB³⁰). **b**, Aligned heatmaps showing properties of amino acid residues or of the specific amino acid substitution present in the Omicron variant, as appropriate (insertion not shown). Structure-based epitope scores³¹ for residues in the structure of the original genotype spike in closed and open conformations are shown. For RBD residues, the results of deep mutational scanning (DMS) studies show the escape fraction (that is, a quantitative measure of the extent to which a mutation reduced polyclonal antibody binding) for each mutant averaged across plasma ('plasma average') and for the most sensitive plasma ('plasma max')²⁰. Each mutation is classified as having evidence for mutations affecting neutralisation by either mAbs^{27, 32, 33, 34, 35} or antibodies in convalescent plasma from previously infected or vaccinated individuals^{20, 34, 35, 36}. Membership of the furin cleavage site is shown. The distance to ACE2-contacting residues that form the receptor-binding site (RBS) is shown (RBS defined as residues with an atom $<4\text{\AA}$ of an ACE2 atom in the structure of RBD bound to ACE2 (RCSB PDB ID: 6MOJ³⁷). Finally, ACE2 binding scores representing the binding constant ($\Delta\log_{10}$ KD) relative to the wild-type reference amino acid from DMS experiments³⁸. **c**, Inferred evolutionary relationships of SARS-CoV-2 from NextStrain (<https://nextstrain.org/ncov/gisaid/global>) with the Variants of Concern labelled. The colours of the tree tips correspond to the number of mutations causing Spike amino acid substitutions relative to the SARS-CoV-2 original genotype (OG) reference strain Wuhan-Hu-1. **d**, Inferred evolutionary relationships of the first 111 Omicron sequences in Scotland with NHS Scottish Health boards denoted: AA, Ayrshire and Arran; FF, Fife; FV, Forth Valley; GC, Great Glasgow and Clyde; GR, Grampian; HG, Highlands; LN, Lanarkshire; LO, Lothian; TY, Tayside, see key.

Neutralising responses to Omicron (BA.1) are substantially reduced following double and partially restored following triple vaccination

Levels of nAbs in patient sera correlate strongly with protection from infection^{39, 40, 41, 42}, and reductions in neutralising activity against the Alpha and Delta variants are consistent with a reduction in vaccine effectiveness^{2, 3, 4, 5, 43}. To predict the effect of the mutations within the Omicron spike glycoprotein on vaccine effectiveness, sera collected from healthy volunteers at more than 14 days post-2nd dose vaccination with either BNT162b2, ChAdOx1 or mRNA-1273 were sorted into three age-matched groups (n=24 per group, mean age 45 years). Sera were first screened by electrochemiluminescence (MSD-ECL) assay for reactivity with SARS-CoV-2 antigens (Spike, RBD, NTD or nucleoprotein (N)). The antibody responses to RBD and NTD were significantly higher ($p < 0.0001$) in the sera from individuals vaccinated with BNT162b2 or mRNA-1273 in comparison with the ChAdOx1 vaccinees (Fig. 2a and

Supplementary Table 1). In contrast, antibody responses to endemic human coronaviruses (HCoVs) (Extended data Fig. 1 and Supplementary Table 2) or influenza (Extended data Fig. 2 and Supplementary Table S3) were similar, with the exception of coronavirus OC43, where responses in BNT162b2 and ChAdOx1 vaccinees differed significantly, perhaps suggesting modulation (back-boosting) of pre-existing OC43 responses by BNT162b2 vaccination.

Next, the nAb responses against SARS-CoV-2 pseudotypes expressing the spike glycoprotein from either Wuhan-Hu-1, or Omicron (BA.1) were compared (Fig. 2b). Vaccination with mRNA-1273 elicited the highest nAb titres (mean titre Wuhan=21,118, Omicron=285), in comparison with those elicited by vaccination with either BNT162b2 (Wuhan=4978, Omicron=148.3) or ChAdOx1 (Wuhan=882.3, Omicron=61.9). Neutralising antibody titres against Wuhan differed significantly between the three study groups. Activity against Omicron was markedly reduced in comparison with Wuhan, reduced by 33-fold for BNT162b2, 14-fold for ChAdOX1 and 74-fold for mRNA-1273 (Supplementary Table 4). While the fold change in neutralisation was lowest in recipients of the ChAdOx1 vaccine and highest in recipients of the mRNA-1273 vaccine, absolute neutralisation values were highest in mRNA-1273 followed by BNT162b2 and ChAdOx1. Neutralisation was lowest in the ChAdOx1 group, however it is important to note that this was given to older patients during early vaccine rollout in the UK, especially to vulnerable patients in nursing homes and was not recommended in young adults less than 40 years of age.




Fig. 2 - Antibody responses elicited by SARS-CoV-2 vaccination. Antibody responses were studied in three groups of individuals (n=24 per group) receiving primary vaccination with either BNT162b2, ChAdOx1 or mRNA-1273 by **a**, MSD-ECL assay or **b**, pseudotype-based neutralisation assay. **a**, Responses were measured against full-length spike glycoprotein (Spike), receptor binding domain (RBD), N-terminal domain (NTD) and nucleoprotein (N) and are expressed as arbitrary units (AU/ml). **b**, NAb responses were quantified against Wuhan or Omicron spike glycoprotein bearing HIV (SARS-CoV-2) pseudotypes. Each point represents the mean of three replicates, bar represents the group mean. To assess the effect of booster vaccines antibody responses were studied in two groups of individuals primed with two doses of either BNT162b2 or ChAdOx1 and boosted with either BNT162b2 or mRNA-1273. Reactivity against SARS-CoV-2 antigens was measured by **c**, MSD-ECL assay while neutralising activity **d**, & **e**, was measured using HIV (SARS-CoV-2) pseudotypes, as above. Green data points represent those boosted with mRNA-1273, all others received BNT162b2. In panel **d**, & **e**, % Omicron neutralising refers to the proportion of serum samples that displayed neutralising activity against Omicron pseudotypes.

Next, samples were analysed from vaccine recipients at least 14 days post booster vaccination (third dose). Participants had been primed with two doses of either ChAdOx1 or BNT162b2, followed by a third dose of either BNT162b2 (full dose) or mRNA-1273 (half dose; 50µg). All sera reacted strongly with SARS-CoV-2 antigens by MSD-ECL, with no significant differences between the four groups (Fig. 2c and Supplementary Table 5). Antibody responses to HCoVs (Extended data Fig. 3 and Supplementary Table 6) or influenza (Extended data Fig. 4 and Supplementary Table 7) were similar, with the exception of influenza Michigan H1, where responses in ChAdOx1-primed and BNT162b2 or mRNA-1273-boosted groups differed significantly, likely reflecting co-administration of influenza booster vaccines during the booster campaign. Two vaccine recipients boosted with BNT162b2 displayed weak reactivity with nucleocapsid (Fig. 2c), suggesting previously undetected exposure to SARS-CoV-2. Sera from vaccine recipients primed with BNT162b2 and boosted with either BNT162b2 or mRNA-1273 displayed similar titres of nAb against Wuhan to the samples collected post-dose 2 (Fig. 2d). In contrast, vaccination of individuals primed with ChAdOx1 with a booster dose of either BNT162b2 or mRNA-1273 resulted in a marked increase in antibody titre (9.3-fold increase) against Wuhan relative to the low titres after dose 2 (Fig. 2e and Supplementary Table 8). The marked increase in antibody titre in ChAdOx1-primed individuals (Extended data Fig. 5) emphasises the importance of the third dose booster in this population. Indeed, following boost with either BNT162b2 or mRNA-1273, anti-Wuhan nAb titres in the ChAdOx1-primed group were not significantly different from those primed with BNT162b2 (Supplementary Table 8). NAb titres against Omicron were lower in both booster study groups and did not differ significantly in titre (Supplementary Table 8). However, absolute numbers displaying measurable Omicron neutralising activity were higher in the ChAdOx1-primed group (13/21, 62%) compared with the BNT162p2 primed group (5/20, 25%) (Fig. 2d, Fig. 2e).

Vaccine effectiveness against the Omicron variant is reduced compared to the Delta variant

A logistic additive model with a test negative case control design was used to estimate relative vaccine effectiveness against becoming a confirmed case with Delta (4911 cases) and/or Omicron (6166 cases) in a population of 1.2 million people in the largest health board in Scotland, NHS GG&C, between 22nd and 28th December 2021. Demographic data are shown

in Supplementary Table 9. The timing of first doses of vaccination are shown in Fig.3a and the occurrence of sequenced/confirmed infections with different variants in vaccine recipients over time is shown in Fig.3b. Infection status for Omicron and Delta was modelled by number and product type of vaccine doses, previous infection status, sex, SIMD quartile, and age (to control for demographic bias). Immunosuppressed individuals were removed from the analysis to ensure case-positivity could be attributed to vaccine escape rather than an inability to mount a vaccine response, with immunosuppression status derived from a list of those in GG&C shielding due to immunosuppression or listed as being given immunosuppressant drugs. Age and time since vaccination were each modelled as single smooth effects using thin plate regression splines⁴⁴.

The protection from vaccine-acquired and infection-acquired immunity were estimated as being markedly reduced against Omicron compared with Delta. Estimates of vaccine effectiveness in recent recipients (at 14 days post-dose) were only 11.44% for full primary courses of ChAdOx1 against Omicron and 78.92% against Delta. For two doses of mRNA vaccines, vaccine effectiveness was significantly lower for Omicron than Delta; BNT162b2 (26.02% versus 83.55%) and mRNA-1273 (23.70% versus 87.82%) (Fig.3c). These responses were similar following a third booster dose of BNT162b2 or mRNA-1273 against Delta (85.94% and 86.53%, respectively), but increased significantly against Omicron (43.22% and 46.38%, respectively). These estimates are in keeping with those reported recently against symptomatic infection in England where vaccine effectiveness was estimated as 71.4% and 75.5% for ChAdOx1 and BNT162b2 primary course recipients boosted with BNT162b2, respectively⁹.

Next, we estimated the additive protective effect of previous natural infection. Infection-acquired immunity directed against other VOCs may be broader in nature and may wane more slowly than that induced by vaccines^{45, 46, 47}. The level of protection following previous infection was 33.9% for Omicron, and 77.4% for Delta. This level of protection was greater than that observed following two doses of vaccine for Omicron but did not reach the levels attained by those who had never had natural infection and had received third dose boosters for either Omicron or Delta. Collectively, these results emphasise the importance of booster

vaccines. Importantly, vaccine-mediated protection against severe disease is likely to be more durable than that against detected infection⁴⁸.




Fig. 3 - Vaccine deployment and vaccine effectiveness estimates **a**, Denominator (violin) plot showing populations of test positive and test negative cohorts in NHS GG&C, with the widths of the grey bands represent the populations in each group at each time point. VOC classification of sequenced cases are overlaid as a dotplot, with points coloured by their VOC and a random jitter applied along the x-axis for visual clarity. **b**, Boxplots of date of first administered vaccine dose by vaccine product for the population of NHS Greater Glasgow and Clyde (NHS GG&C) aged 18 and older. The box limits are the quartiles and the centre line is the median, with whisker length of 1.5 times the interquartile range. Outliers are shown as dots outwith the whisker range. Data points are overlaid as a dotplot with points shown as black dots, with a random jitter along the x-axis applied for visual clarity. **c**, Error bar plot of estimated vaccine effectiveness against testing positive for Delta and Omicron SARS-CoV-2 infection in the population of over 18s in NHS GG&C who were tested between 6th and 12th December 2021. The points and corresponding text represent the estimated vaccine effectiveness (%) for each group, for each variant, with the error bar endpoints representing the endpoints of the corresponding 95% CIs. The additive effect of infection-acquired immunity was calculated for the entire population and plotted for the unvaccinated cohort.

Absence of syncytia in Omicron-infected cells

Our data demonstrate that antigenic change in Omicron permits evasion of vaccine induced immunity, however, the constellation of spike mutations in Omicron suggest that functional change may also contribute to its rapid transmission (Fig.1a). Therefore, we investigated the virological properties of live Omicron isolated from a patient sample. SARS-CoV-2 particles can achieve membrane fusion at the cell surface following proteolytic activation of spike by the plasma membrane protease TMPRSS2. This property also permits spike-mediated fusion of SARS-CoV-2 infected cells with adjacent cells resulting in syncytia⁴⁹; a feature that has been associated with severe disease⁵⁰. Moreover, the Delta variant has been shown to exhibit enhanced fusion compared to the Alpha and Beta variants⁵¹.

A split GFP cell-cell fusion system⁵² was used to quantify syncytia formation by Omicron, Delta and first wave Wuhan D614G virus (Fig. 4a). Cells expressing split GFP were infected with Wuhan-D614G, Delta or Omicron and the levels of the reconstituted GFP signal following cell-cell fusion was determined in real time (Fig. 4c). In addition, infected cells were probed by indirect immunofluorescence assay to assess viral replication by the detection of the viral nucleocapsid protein (Fig. 4b). The Delta variant exhibited the highest levels of cell fusion followed by Wuhan D614G. In contrast, the Omicron variant failed to form syncytia. This failure was not due to lack of infection as immunofluorescent detection of nucleocapsid protein confirmed viral replication by Omicron, Wuhan D614G and Delta¹⁸.

Reduced replication kinetics of Omicron in lung epithelial cells

The replication of Omicron, Delta and Wuhan D614G was compared in Calu-3, a human lung epithelial cell line. Wuhan D614G and Delta displayed comparable replication kinetics over a period of 72 hours, with visible CPE between 48-72hpi (Fig. 4d). In contrast, the titres of Omicron were at least an order of magnitude lower at each time point compared to Wuhan D614G and Delta. These observations are consistent with attenuated replication of Omicron in lower respiratory tissues as recently reported^{18, 53}.








Fig. 4. Omicron exhibits reduced syncytia formation and has switched entry route preference a,

Schematic representation of the split GFP system, used in this study to quantify virus induced cell fusion. This system is based on co-culture of two different cell lines (GFP-10 and GFP-11) expressing split GFP molecules. Upon virus-induced cell fusion, the intact GFP molecule is reconstituted, and the resulting signal can be detected and quantified. **b,** GFP-10 and GFP-11 A549 ACE2 TMPRSS2 cells were co-cultured and infected with Wuhan D614G, Delta and Omicron and incubated in a CLARIOStar Plus (BMG LABTECH) at 37°C / 5% CO₂. Micrographs display cells 22 hours post infection: reconstituted GFP (green), N viral nucleocapsid (red), and DAPI nuclear counter stain (blue). Scale bars 100µm. **c,** To quantify, GFP signal was measured every 30 min for 20h. Omicron infected cells showed only background levels of GFP signal. **d,** Calu-3 cells were infected with Wuhan D614G, Delta and Omicron and supernatants were collected at the indicated times and assessed by RT-qPCR. Omicron display reduced replication kinetics compared to Wuhan D614G and Omicron. **e,** SARS CoV-2 entry can occur via two routes. Route 1 permits rapid fusion at the cell surface following proteolytic processing by TMPRSS2. In Route 2 fusion occurs following endocytosis after processing by cathepsin B or L. Route 1 and 2 can be specifically inhibited using the protease inhibitors Camostat and E64d, respectively. **f,** SARS-CoV-2 pseudotype infection of the stated cell lines, data represent mean luciferase values from one representative experiment. In Calu-3 cells Route 1 entry predominates whereas HEK exclusively support Route 2, A549 ACE2 TMPRSS2 cells permit both routes. Pangolin CoV spike is included as a control; it can only achieve entry via Route 2. Pseudotypes without viral glycoproteins (No) are included as a negative control. **g,** Relative SARS-CoV-2 pseudotype infection (compared to untreated control) of cells treated with 10µM protease inhibitors. Data represent mean of four replicates, error bars indicate standard error of the mean, asterisks indicate statistical significance (ANOVA). **h,** Titration of Camostat and E64d against Delta, Omicron and Pangolin CoV in A549 ACE2 TMPRSS2 cells, data points represent mean relative infection, compared to untreated control.

Omicron spike has switched entry route preference

Entry of SARS-CoV-2, and related coronaviruses, can proceed via two routes⁵⁴. Cell surface fusion following proteolysis by TMPRSS2, as described above (“Route 1”, Fig. 4e), or fusion from the endosome after endocytosis and activation by the endosomal proteases Cathepsin B or L (“Route 2”, Fig. 4e). The ability of SARS-CoV-2 to achieve cell surface fusion is dependent on its S1/S2 polybasic cleavage site; this is absent from most closely related sarbecoviruses, which are confined to endosomal fusion^{55, 56, 57}. Given the reduced fusogenicity and replication kinetics of Omicron, HIV pseudotypes were used to evaluate entry route preference. We tested Wuhan D614G, Alpha, Delta and Omicron spike, while Pangolin CoV (Guangdong isolate) spike was included as a control. Pangolin CoV spike exhibits high affinity interactions with human ACE2 but lacks a polybasic cleavage site and, therefore, enters via the endosome only^{58, 59, 60, 61}.

Calu-3 cells support cell surface (Route 1) fusion predominantly, owing to their high endogenous expression of TMPRSS2^{56, 62}; in these cells, Delta yielded the highest infection, being ~4 fold higher than Omicron (Fig. 4f). Pangolin CoV infection was low, indicating that Calu-3 cells do not support robust endosomal entry. In contrast, HEK only support endosomal entry and in these cells Pangolin CoV had high infection. Notably, Omicron also achieved high infection in HEK cells, producing ~10-fold greater signal than Delta. This suggests that Omicron, like Pangolin CoV, is optimised for endosomal entry. All pseudotypes exhibited robust infection in A549-ACE2-TMPRSS2, where both entry routes are available^{63, 64}.

Entry pathway preference was further investigated using protease inhibitors targeting either TMPRSS2 (Camostat) or cathepsins (E64d)⁵⁵. In Calu-3 cells, all SARS-CoV-2 pseudotypes were inhibited by Camostat, whereas only Omicron exhibited E64d sensitivity, indicating that a component of infection occurs via endosomal entry (Fig. 4g). In HEK cells, all pseudotypes were inhibited by E64d, whereas Camostat was non-inhibitory; this confirms that only endosomal entry is available in these cells. Inhibitor treatment in A549 ACE2 TMPRSS2 provided the clearest evidence of altered entry by Omicron. D614G, Alpha and Delta were potently inhibited by Camostat, but not E64d. For Omicron, and Pangolin CoV, this pattern was reversed, suggesting a binary switch from cell surface to endosomal fusion; this conclusion was supported by titration of either inhibitor in A549 ACE2 TMPRSS2 cells (Fig. 4h).

These data indicate that, whilst Delta is optimised for fusion at the cell surface, Omicron preferentially achieves entry through endosomal fusion; this biological about-face may impact transmission, cellular tropism and pathogenesis. Moreover, this switch away from TMPRSS2-mediated activation offers a mechanistic explanation for reduced syncytia formation by Omicron infected cells.

Discussion

The Omicron variant represents a major change in biological function and antigenicity of SARS-CoV-2. In this study, we demonstrate substantial immune escape of the Omicron variant. We present clear evidence of vaccine failure in dual-vaccinated individuals and partial restoration of immunity following a third booster dose of mRNA vaccine. In addition, we demonstrate a shift in the SARS-CoV-2 entry pathway from cell surface fusion, triggered by TMPRSS2, to cathepsin-dependent fusion within the endosome. This fundamental biological shift may affect the pathogenesis and severity of disease and requires further evaluation in population-based studies.

Using sera from double vaccine recipients, Omicron was found to be associated with a drop in neutralisation greater in magnitude than that reported in all other variants of concern (including Beta and Delta). Boosting enhanced neutralising responses to both the vaccine strain (Wuhan) and Omicron, particularly in recipients of ChAdOx1, but did not completely overcome the inherent immune escape properties of Omicron. Importantly, we did not assess the impact of vaccination on clinical severity of disease which is likely to be much higher than detection of infection. Protection against severe disease is longer lasting than prevention of infection. We also did not measure the impact of vaccination on T cell immunity which may be better preserved as only 14% of CD8+ and 28% of CD4+ epitopes are predicted to be affected by key Omicron mutations¹².

The probability of infection with Omicron versus infection with the preceding Delta variant was significantly higher in double vaccine recipients, in keeping with the neutralisation data. A third dose of mRNA vaccine substantially reduced the probability of infection but did not restore immunity fully.

The emergence of a highly transmissible variant that is associated with escape from vaccine-induced immune responses means that over time, Omicron-specific vaccines may be required if disease severity was high, either directed at the general population or vulnerable groups. Early indications in young people are that Omicron infection is 40-70% less severe than Delta infection^{65, 66} similar calculations in the most vulnerable part of the population over the age of 40 years are awaited.

Genotypic changes in new variants have previously been shown to alter viral phenotype by modulating innate immune responses as well as evasion of the adaptive immune response^{67, 68}. Additionally, mutations can alter spike functionality to impact transmission and pathogenesis²⁴. Such changes may have provided emergent viruses with a selective advantage in lung cells and primary human airway epithelial cells. Enhanced spike activation by the plasma membrane protease TMPRSS2, may have enabled more rapid cell surface fusion⁵⁴. In this study, we found that the Omicron variant has switched entry pathway to use TMPRSS2-independent endosomal fusion preferentially, a major change in the biological behaviour of the virus. This switching in the mechanism of fusion activation also manifests in reduced syncytia formation in infected cells, likely to reduce the cell-to-cell transmission characteristics of other variants. These properties have the potential to change cellular tropism and disease pathogenesis significantly. However, even a variant that is less virulent with a very high transmission rate may continue to present a substantial risk to older people and those with co-morbidities, especially those with immunosuppression. Moreover, our work demonstrates that SARS-CoV-2 exhibits high antigenic and functional plasticity; further fundamental shifts in transmission and disease should be anticipated.

Methods

Cells. Calu-3 are human lung adenocarcinoma epithelial cells. Caco-2 are an immortalized cell line derived from human colorectal adenocarcinoma, primarily used as a model of the intestinal epithelial barrier. A549 cells, a human alveolar adenocarcinoma line, were modified to stably express human ACE-2 and TMPRSS2. Human embryonic kidney (HEK293T) cells were used in pseudotype production. Baby Hamster Kidney clone 21 cells and Vero ACE-2 TMPRSS2 cells were used in the isolation of live Omicron SARS-CoV-2. All cell lines were maintained at 37°C and 5% CO₂ in DMEM supplemented with 10% foetal bovine serum (FBS), except for Calu-3 cells which were supplemented with 20% FBS.

Generation of cell line expressing human ACE2 receptor. Lentiviral vectors encoding human *ACE2* (GenBank NM_001371415.1) were produced as described previously⁶⁹ and BHK-21 transduced

cells were selected with 200µg/ml of hygromycin B.

Generation of cell lines used for fusion assays. Retrovirus vectors were produced by transfecting HEK-293T cells with plasmid pQCXIP-GFP1-10 (Addgene #68715) or pQCXIP-BSR-GFP11 (Addgene #68716)⁵² and packaging vectors expressing MLV gal-pol and VSV-G using Lipofectamine 3000 (Invitrogen) according to manufacturer's instructions. Cell supernatants were harvested 24-48h post-transfection, pooled, clarified by centrifugation and filtered. One mL of each supernatant was used to transduce A549-Ace2-TMPRSS2 (AAT) cells⁶⁹ in presence of Polybrene (Merck). Two days post-transduction, the supernatant was replaced with selection medium (DMEM 10% FBS 1µg/mL puromycin) and cells incubated until complete death of the untransduced control cells were observed. The resulting puromycin-resistant cells (termed AAT-GFP1-10 and AAT-BSR-GFP11) were used in fusion assays.

Virus isolation from clinical samples. Nasopharyngeal swabs of patients infected with Omicron were collected with biorepository ethical approval (reference 20/ES/0061) in virus transport medium and resuspended in serum-free DMEM supplemented with 10 µg/ml gentamicin, 100 units/ml penicillin-streptomycin and 2.5µg/ml amphotericin B to a final volume of 1.5ml. Virus isolation was attempted in BHK-21 cells stably expressing the human ACE2 protein (BHK-hACE2) and VERO cells stably expressing ACE2 and TMPRSS2 (VAT⁶⁹). The

infected cells were incubated at 37°C and monitored for signs of cytopathic effect (CPE) and the presence of viral progeny in the medium by RT-qPCR. While no CPE was observed in any of the infected cells, RT-qPCR at 5 days post-infection (dpi) confirmed the presence of the virus derived from two of the five samples (referred to hereafter as 204 and 205) in the medium of BHK-hACE2, but not VAT cells (Extended data Fig. 7a). An aliquot of the clarified medium containing approximately 4×10^4 viral genomes of the P0 stocks of samples 204 and 205 was used to infect VAT, BHK-ACE2 and Calu-3 cells. No CPE was observed in the infected cells but once again, virus replication was confirmed in BHK-hACE2 and Calu-3 by RT-qPCR. Supernatants (termed P1) from infected Calu-3 cells at 3 dpi were collected and virus titrated by both focus forming assay and RT-qPCR. The virus reached more than 100-fold higher titres in Calu-3 cells compared to BHK-hACE2 (Extended data Fig. 7b). Further passage of sample 205-derived P1 virus in both Calu-3 and Caco-2 yielded equivalent genome copy numbers in both cell lines (Extended data Fig. 7b). CPE was observed at 3 dpi in both Calu-3 and Caco-2 cells (not shown). The medium (termed P2) of infected Calu-3 and Caco-2 cells was collected at 4 dpi, titrated and used in subsequent experiments.

Measurement of SARS-CoV-2, HCoV-229E, HCoV-OC43, HCoV-NL63, HCoV-HKU1 and influenza antibody response by electrochemiluminescence. IgG antibody titres were measured quantitatively against SARS-CoV-2 trimeric spike (S) protein, N-terminal domain (NTD), receptor binding domain (RBD) or nucleocapsid (N), human seasonal coronaviruses (HCoVs) 229E, OC43, NL63 and HKU1; and influenza A (Michigan H1, Hong Kong H3 and Shanghai H7) and B (Phuket HA and Brisbane) using MSD V-PLEX COVID-19 Coronavirus Panel 2 (K15369) and Respiratory Panel 1 (K15365) kits. Multiplex Meso Scale Discovery electrochemiluminescence (MSD-ECL) assays were performed according to manufacturer instructions. Briefly, 96-well plates were blocked for one hour. Plates were then washed, samples were diluted 1:5000 in diluent and added to the plates along with serially diluted reference standard (calibrator) and serology controls 1.1, 1.2 and 1.3. After incubation, plates were washed and SULFO-TAG detection antibody added. Plates were washed and were immediately read using a MESO Sector S 600 plate reader. Data were generated by Methodological Mind software and analysed using MSD Discovery Workbench (v4.0). Results are expressed as MSD arbitrary units per ml (AU/ml). Reference

plasma samples yielded the following values: negative pool - spike 56.6 AU/ml, NTD 119.4 AU/ml, RBD 110.5 AU/ml and nucleocapsid 20.7 AU/ml; SARS-CoV-2 positive pool - spike 1331.1 AU/ml, NTD 1545.2 AU/ml, RBD 1156.4 AU/ml and nucleocapsid 1549.0 AU/ml; NIBSC 20/130 reference - spike 547.7 AU/ml, NTD 538.8 AU/ml, RBD 536.9 AU/ml and nucleocapsid 1840.2 AU/ml.

Measurement of virus neutralising antibodies using viral pseudotypes. Pseudotype-based neutralisation assays were carried out as described previously^{70 71 72}. Briefly, HEK293, HEK293T, and 293-ACE2⁷¹ cells were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% FBS, 200mM L-glutamine, 100µg/ml streptomycin and 100 IU/ml penicillin. HEK293T cells were transfected with the appropriate SARS-CoV-2 S gene expression vector (wild type or variant) in conjunction with p8.91⁷³ and pCSFLW⁷⁴ using polyethylenimine (PEI, Polysciences, Warrington, USA). HIV (SARS-CoV-2) pseudotypes containing supernatants were harvested 48 hours post-transfection, aliquoted and frozen at -80°C prior to use. S gene constructs bearing the WUHAN (D614G) and Omicron (B.1.1.529) S genes were based on the codon-optimised spike sequence of SARS-CoV-2 and generated by GeneArt (ThermoFisher). Constructs bore the following mutations relative to the Wuhan-Hu-1 sequence (GenBank: MN908947): WUHAN (D614G) – D614G; Omicron (BA.1, B.1.1.529) - A67V, Δ69-70, T95I, G142D/Δ143-145, Δ211/L212I, ins214EPE, G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H, T547K, D614G, H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H, N969K, L981F. 293-ACE2 target cells were maintained in complete DMEM supplemented with 2µg/ml puromycin.

Neutralising activity in each sample was measured by a serial dilution approach. Each sample was serially diluted in triplicate from 1:50 to 1:36450 in complete DMEM prior to incubation with HIV (SARS-CoV-2) pseudotypes, incubated for 1 hour, and plated onto 239-ACE2 target cells. After 48-72 hours, luciferase activity was quantified by the addition of SteadyLite Plus chemiluminescence substrate and analysis on a Perkin Elmer EnSight multimode plate reader (Perkin Elmer, Beaconsfield, UK). Antibody titre was then estimated by interpolating the point at which infectivity had been reduced to 50% of the value for the no serum control samples.

Protease inhibitor studies. To selectively inhibit either cell surface or endosomal fusion of SARS-CoV-2, cells were pre-treated for one hour with 10µM of either Camostat mesylate

(referred to hence forth as Camostat) or E64d prior to inoculation with pseudotype. In these studies, spike proteins from Alpha and Delta VOCs, and Guangdong isolate Pangolin coronavirus (GISAID ref EPI_ISL_410721) were used as controls.

Viral RNA extraction and RT-qPCR. Viral RNA was extracted from culture supernatants using the RNAdvance Blood kit (Beckman Coulter Life Sciences) following the manufacturer's recommendations. RNA was used as template to detect and quantify viral genomes by duplex RT-qPCR using a Luna® Universal Probe One-Step RT-qPCR Kit (New England Biolabs, E3006E). SARS-CoV-2 specific RNAs were detected by targeting the N1 gene from the CDC panel as part of the SARS-CoV-2 Research Use Only qPCR Probe Kit (Integrated DNA Technologies) and the ORF1ab gene using the following set of primers and probes: SARS-CoV-2_Orf1ab_Forward 5' GACATAGAAGTTACTGG&CGATAG 3', SARS-CoV-2_Orf1ab_Reverse 5' TTAATATGACGCGCACTACAG 3', SARS-CoV-2_Orf1ab_Probe ACCCCGTGACCTTGGTGCTTGT with HEX/ZEN/3IABkFQ modifications. SARS-CoV-2 RNA was used to generate a standard curve and viral genomes were quantified and expressed as number of Orf1ab RNA molecules /ml of supernatant. All runs were performed on the ABI7500 Fast instrument and results analysed with the 7500 Software v2.3 (Applied Biosystems, Life Technologies).

Genome Sequencing and analysis. Sequencing was carried out by the UK public health agencies (UKHSA/PHE, PHS, PHW and PHNI) and by members of the COG-UK consortium using the ARTIC protocol as previously described. Sequences were aligned by mapping to the SARS-CoV-2 reference Wuhan-Hu-1 using Minimap2 ⁷⁵. Prior to phylogenetic analysis 85 sites exhibiting high genetic variability due to data quality issues in overseas sequencing labs were excluded using a masking script in Phylopipe (<https://github.com/cov-ert/phylopipe>). The phylogenetic tree was constructed with the maximum likelihood method FastTree2 ⁷⁶ using a JC+CAT nucleotide substitution model.

Replication curve. Calu-3 cells were seeded in a 96-well plate at a cell density of 3.5×10^4 cells per well. Cells were infected with the indicated viruses using the equivalent of 2×10^4 Orf1ab genome copies/well in serum-free RPMI-1640 medium (Gibco). After one hour of incubation at 37°C, cells were washed three times and left in 20% FBS RPMI-1640 medium. Supernatants were collected at different times post-infection and viral RNA extracted and quantified as described above.

Fusion assay. AAT-GFP1-10 and AAT-BSR-GFP11 cells were trypsinized and mixed at a ratio of 1:1 to seed a total of 2×10^4 cells/well in black 96-well plate (Greiner) in FluoroBrite DMEM medium (Thermo Fischer Scientific) supplemented with 2% FBS. Next day, cells were infected with the indicated viruses using the equivalent of 10^6 Orf1a genome copies/well in FluoroBrite DMEM 2% FBS. GFP signal was acquired for the following 20 hours using a CLARIOStar Plus (BMG LABTECH) equipped with ACU to maintain 37°C and 5% CO₂. Data were analysed using MARS software and plotted with GraphPad prims 9 software. At 22 hs post-infection, cells were fixed in 8% formaldehyde, permeabilized with 0.1 % Triton X-100 and stained with sheep anti-SARS-CoV-2 N (1:500) antiserum ⁶⁹ followed by Alexa Fluor 594 Donkey anti-sheep IgG (H+L) (1:500, Invitrogen) and DAPI (1:4000, Sigma). Cell images were acquired using EVOS Cell Imaging Systems (Thermo Fischer Scientific).

Demographic data. Data for the EVADE study were available using the NHS Greater Glasgow and Clyde (NHS GG&C) SafeHaven platform and included vaccination status (dates and product names for each dose), demographic data (age, sex and Scottish Index of Multiple Deprivation (SIMD) quartile) comorbidity (shielding and immunosuppression status) and dates of positive and negative PCR tests, for 1.2 million inhabitants of the (NHS GG&C) area over 18 years of age, from 1st March 2020 up to 12th January 2022. Data were matched by CHI number and pseudonymised before analysis. Derogated ethical approval was granted by the NHS GG&C SafeHaven committee (GSH/21/IM/001).

Vaccine effectiveness. We used a logistic additive regression model to estimate relative vaccine effectiveness against the Omicron variant as it emerged in a population of 1.2 million people in NHS Greater Glasgow & Clyde, the largest health board in Scotland. Infection status for Omicron and Delta was modelled by number and product type of vaccine doses, previous infection status, sex, SIMD quartile, age on 31st October 2021 and time since most recent vaccination.

We identified Omicron infections using 3 data streams: confirmed S gene target failure (SGTF), allele specific PCR, and Pango lineage assignments from the sequencing data. SGTF samples with Delta lineage assignments were assigned as Delta infections. Samples for which the sequencing date was more than two weeks away from the first positive PCR were removed from the analysis.

We removed a small number of individuals who received ChAdOx1 as a third dose or had their third dose before the first of September 2021 on the assumption that the majority were part of the COV-BOOST clinical trial, the results of which are published elsewhere. We removed anyone with ambiguous vaccination status or whose brand was unknown due to data entry error. We removed those who were vaccinated during the study period. We removed individuals who tested positive in the 90 days before the study period. To control for the effect of missed vaccinations due to recent infection, we exclude those who were eligible for a second or third dose but had not taken this up (i.e had a first dose more than 8 weeks ago but no second dose, or a second dose more than 12 weeks ago but no third dose). Since those who tested positive during the study period could not subsequently be vaccinated within the study period, and those who changed vaccination status during the study period were excluded from the dataset, there would be an inflated proportion of people testing positive and having only a second dose of a vaccine. Specifically, those who were given the third dose of a vaccine during the study period would be excluded from the analysis, while those who would have been given a third dose of the vaccine but could not due to becoming infected would be included in the analysis. This would lead to reduced estimates of effectiveness of second doses if not accounted for appropriately. Due to the timing of the rollout of booster doses coinciding with high levels of infection, it is vital to account for this.

Serum samples. Serum samples were collected from healthy participants in the COVID-19 Deployed Vaccine Cohort Study (DOVE), a cross-sectional post-licensing cohort study to determine the immunogenicity of deployed COVID-19 vaccines against evolving SARS-CoV-2 variants. 308 adult volunteers aged at least 18 years and were recruited into the study 14 days or more after a second or third dose of vaccine. All participants gave written informed consent to take part in the study. The DOVE study was approved by the North-West Liverpool Central Research Ethics Committee (REC reference 21/NW/0073).

Structural modelling. The file 6vsb_1_1_1.pdb containing a complete model of the full-length glycosylated spike homotrimer in open conformation with one monomer having the receptor-binding domain in the 'up' position was obtained from the CHARMM-GUI Archive^{77, 78}. This model is itself generated based upon a partial spike cryo-EM structure (PDB ID: 6VSB). For visualisation, the model was trimmed to the ectodomain (residues 14-1164) and the signal

peptide (residues 1-13) and glycans were removed. Using this structural model and the closed conformation equivalent (6vxx_1_1_1.pdb). Residues belonging to the receptor-binding site were identified as those with an atom within 4Å of an ACE2 atom in the bound RBD-ACE2 structure (PDB ID: 6M0J⁷⁹) and Alpha carbon-to-Alpha carbon distances between these residues in the 'up' RBD and all other spike residues were calculated. Antibody accessibility scores for open and closed conformations were calculated using BEpro³¹. Figures were prepared using PyMol⁸⁰.

Epidemiological description of the emergence of the Omicron variant in the UK

On the 27th November 2021, the UK Health Security Agency detected 2 cases of Omicron in England, the following day 6 Scottish cases were detected by community (Pillar 2) sequencing. Over the next 10 days (to 8th December 2021) a further 95 genome sequences were obtained. Due to the rapid spread of Omicron and low genetic diversity, the genome sequences are highly related with mean genetic divergence of 1 single nucleotide polymorphisms (SNPs) and maximum 7 SNPs.

The phylogenetic relationship to Omicron sequences from other countries is consistent with multiple introductions associated with travel to South Africa followed by community transmissions within Scotland. Amongst the Scottish samples diverged from the tree backbone, there were a number identified that are genetically divergent, i.e., greater than 2 single nucleotide polymorphisms from the nearest Scottish sample (Fig. 1d). Moreover, comparison to the wider international collection of Omicron samples revealed that they were more closely related to genomes from other countries than other Scottish samples. These samples therefore likely represent independent introductions to Scotland, but without more detailed epidemiological data, the number of introductions is unknown. Where there are indistinguishable samples in the phylogeny from Scotland and elsewhere in world, importation cannot be ruled out as a source of these samples in Scotland, rather than transmission from an established population circulating in Scotland.

Within Scotland, cases are spread across 9 separate Health Boards and distributed throughout the phylogeny (Fig. 1d). Basal Scottish genomes were sampled in 7 different Health Boards, most of them from NHS Greater Glasgow & Clyde (47%) and NHS Lanarkshire (25%). Notably, amongst these earliest samples are cases that were epidemiologically linked to early

spreading events. All but one of these samples were found on this basal branch and are indistinguishable, and which is consistent with transmission at these events.

Data and materials availability: The experimental data that support the findings of this study are available on reasonable request but restrictions apply to the availability of clinical data, which were used under ethical approvals for the current study, and so are not publicly available.

Codes used in this analysis are available in the study's GitHub repository: <https://github.com/centre-for-virus-research/Omicron>.

Acknowledgments:

The authors would like to thank the participants of the DOVE study and Sister Therese McSorley and her nursing team at the NHS GG&C clinical research facility. The authors also thank Alison Hamilton, Laura Stirling and Charlie Mayor from the NHS GG&C SafeHaven team for their invaluable input in facilitating this study. We thank Paula Olmo for administrative support and Chris Robertson and Aziz Sheikh for statistical advice. The authors thank all of the researchers who have shared genome data openly via the Global Initiative on Sharing All Influenza Data (GISAID).

Funding:

Health Data Research UK (HDR UK) - the Evaluation of Variants Affecting Deployed COVID-19 Vaccine (EVADE) study (ECT, SR, OM, CW, BJW) (grant code: 2021.0155).

COG-UK is supported by funding from the Medical Research Council (MRC) part of UK Research & Innovation (UKRI), the National Institute of Health Research (NIHR) [grant code: MC_PC_19027], and Genome Research Limited, operating as the Wellcome Sanger Institute (RMB, DLR, ECT).

Medical Research Council (MRC) - the COVID-19 DeplOyed VaccinE (DOVE) study (grant code: MCUU1201412) and COG-UK (ECT).

Medical Research Council (MRC) (grant code: MC_UU_12014/12) (AF, JH, RO, JG, ECT and DLR)

Medical Research Council (MRC) (grant codes: MR/R024758/1 and MR/W005611/1) (WTH)
UK Research and Innovation (UKRI) - G2P-UK National Virology Consortium (MR/W005611/1) (MP, ECT, AP, DLR).

Wellcome Trust (grant code: 220977/Z/20/Z) (DLR)

Biotechnology and Biological Sciences Research Council (grant code: BBSRC, BB/R004250/1) (NL, BJW)

Wellcome Trust and Royal Society Sir Henry Dale Fellowship (grant code: 107653/Z/15/A) (JG)

Author contributions:

Conceptualization: ECT, JG, MP, DLR, BW

Methodology: BJW, JG, OAM, CW, NL, GL, WF, SS, MM, AS, WTH, CD, RO, JH, DP, KP, AF, GY, SS, RMP, PRM, AHP, JH, DLR, SR

Investigation: BJW, JG, OAM, CW, NL, GL, WF, SS, MM, AS, SA, EV, WTH, CD, RO, JH, PH, VS, DP, KP, AF, GY, SS, MTGH, RG, KT, AHP, JH, DLR, MP, SR

Visualization: BJW, JG, OAM, CW, GL, WTH, DLR

Funding acquisition: ECT, DLR, SR, BJW, JG

Project administration: ECT, DLR, SR, JH,

Supervision: BJW, DLR, MP, SR, JH, ECT

Writing – original draft: ECT, BJW, JG, GY

Writing – review & editing: BJW, JG, OAM, CW, SA, WTH, DLR, MP, ECT

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: The experimental data that support the findings of this study are available on reasonable request but restrictions apply to the availability of clinical data, which were used under ethical approvals for the current study, and so are not publicly available. Anonymised data are available with permission of the NHS Greater Glasgow & Clyde SafeHaven.

Codes used in this analysis are available upon request from the corresponding author.

Biological materials including cell lines are available on reasonable request from the authors.

Clinical samples are restricted for use under the ethical approvals obtained for their use.

References

1. WHO SPRP 2021 Mid-term Report - WHO Strategic Action Against COVID 19.
2. Davis, C. *et al.* Reduced neutralisation of the Delta (B.1.617.2) SARS-CoV-2 variant of concern following vaccination. *PLoS Pathog* **17**, e1010022 (2021).
3. Wall, E.C. *et al.* Neutralising antibody activity against SARS-CoV-2 VOCs B.1.617.2 and B.1.351 by BNT162b2 vaccination. *The Lancet* **397**, 2331-2333 (2021).
4. Wall, E.C. *et al.* AZD1222-induced neutralising antibody activity against SARS-CoV-2 Delta VOC. *Lancet* **398**, 207-209 (2021).
5. Lopez Bernal, J. *et al.* Effectiveness of Covid-19 Vaccines against the B.1.617.2 (Delta) Variant. *New England Journal of Medicine* **385**, 585-594 (2021).
6. Viana, R. *et al.* Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. *medRxiv*, 2021.2012.2019.21268028 (2021).
7. Gu, H. *et al.* Probable Transmission of SARS-CoV-2 Omicron Variant in Quarantine Hotel, Hong Kong, China, November 2021. *Emerg Infect Dis* **28** (2021).
8. Cameroni, E. *et al.* Broadly neutralizing antibodies overcome SARS-CoV-2 Omicron antigenic shift. *Nature* (2021).
9. Andrews, N. *et al.* Effectiveness of COVID-19 vaccines against the Omicron (B.1.1.529) variant of concern. *medRxiv*, 2021.2012.2014.21267615 (2021).
10. Aggarwal, A. *et al.* SARS-CoV-2 Omicron: evasion of potent humoral responses and resistance to clinical immunotherapeutics relative to viral variants of concern. *medRxiv*, 2021.2012.2014.21267772 (2021).
11. Basile, K. *et al.* Improved neutralization of the SARS-CoV-2 Omicron variant after Pfizer-BioNTech BNT162b2 COVID-19 vaccine boosting. *bioRxiv*, 2021.2012.2012.472252 (2021).

12. Ahmed, S.F., Quadeer, A.A. & McKay, M.R. SARS-CoV-2 T cell responses are expected to remain robust against Omicron. *bioRxiv*, 2021.2012.2012.472315 (2021).
13. Cao, Y. *et al.* Omicron escapes the majority of existing SARS-CoV-2 neutralizing antibodies. *Nature* (2021).
14. Cele, S. *et al.* SARS-CoV-2 Omicron has extensive but incomplete escape of Pfizer BNT162b2 elicited neutralization and requires ACE2 for infection. *medRxiv*, 2021.2012.2008.21267417 (2021).
15. Dejnirattisai, W. *et al.* Reduced neutralisation of SARS-CoV-2 omicron B.1.1.529 variant by post-immunisation serum. *The Lancet* (2021).
16. Doria-Rose, N.A. *et al.* Booster of mRNA-1273 Vaccine Reduces SARS-CoV-2 Omicron Escape from Neutralizing Antibodies. *medRxiv*, 2021.2012.2015.21267805 (2021).
17. Garcia-Beltran, W.F. *et al.* mRNA-based COVID-19 vaccine boosters induce neutralizing immunity against SARS-CoV-2 Omicron variant. *Cell* (2022).
18. Meng, B. *et al.* SARS-CoV-2 Omicron spike mediated immune escape, infectivity and cell-cell fusion. *bioRxiv*, 2021.2012.2017.473248 (2021).
19. Zahradník, J. *et al.* SARS-CoV-2 variant prediction and antiviral drug design are enabled by RBD in vitro evolution. *Nature Microbiology* **6**, 1188-1198 (2021).
20. Greaney, A.J. *et al.* Comprehensive mapping of mutations in the SARS-CoV-2 receptor-binding domain that affect recognition by polyclonal human plasma antibodies. *Cell host & microbe* **29**, 463-476 e466 (2021).
21. Burnett, D.L. *et al.* Immunizations with diverse sarbecovirus receptor-binding domains elicit SARS-CoV-2 neutralizing antibodies against a conserved site of vulnerability. *Immunity* **54**, 2908-2921.e2906 (2021).
22. Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody. *Nature* **583**, 290-295 (2020).
23. Rouet, R. *et al.* Potent SARS-CoV-2 binding and neutralization through maturation of iconic SARS-CoV-1 antibodies. *MAbs* **13**, 1922134 (2021).

24. Peacock, T.P. *et al.* The furin cleavage site in the SARS-CoV-2 spike protein is required for transmission in ferrets. *Nature Microbiology* **6**, 899-909 (2021).
25. Meng, B. *et al.* Recurrent emergence of SARS-CoV-2 spike deletion H69/V70 and its role in the Alpha variant B.1.1.7. *Cell Reports* **35**, 109292 (2021).
26. McCallum, M. *et al.* N-terminal domain antigenic mapping reveals a site of vulnerability for SARS-CoV-2. *Cell* **184**, 2332-2347.e2316 (2021).
27. McCarthy, K.R. *et al.* Recurrent deletions in the SARS-CoV-2 spike glycoprotein drive antibody escape. *Science*, eabf6950 (2021).
28. Greaney, A.J. *et al.* Mapping mutations to the SARS-CoV-2 RBD that escape binding by different classes of antibodies. *Nat. Commun.* **12**, 4196 (2021).
29. Woo, H. *et al.* Developing a Fully Glycosylated Full-Length SARS-CoV-2 Spike Protein Model in a Viral Membrane. *J Phys Chem B* **124**, 7128-7137 (2020).
30. Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* **367**, 1260-1263 (2020).
31. Sweredoski, M.J. & Baldi, P. PEPITO: improved discontinuous B-cell epitope prediction using multiple distance thresholds and half sphere exposure. *Bioinformatics* **24**, 1459-1460 (2008).
32. McCallum, M. *et al.* SARS-CoV-2 immune evasion by variant B.1.427/B.1.429. *bioRxiv*, 2021.2003.2031.437925 (2021).
33. Starr, T.N. *et al.* Prospective mapping of viral mutations that escape antibodies used to treat COVID-19. *Science* **371**, 850-854 (2021).
34. Weisblum, Y. *et al.* Escape from neutralizing antibodies by SARS-CoV-2 spike protein variants. *Elife* **9** (2020).
35. Liu, Z. *et al.* Identification of SARS-CoV-2 spike mutations that attenuate monoclonal and serum antibody neutralization. *Cell host & microbe* (2021).
36. Wang, Z. *et al.* mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. *bioRxiv*, 2021.2001.2015.426911 (2021).

37. Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature* **581**, 215-220 (2020).
38. Starr, T.N. *et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295-1310 e1220 (2020).
39. Cromer, D. *et al.* Neutralising antibody titres as predictors of protection against SARS-CoV-2 variants and the impact of boosting: a meta-analysis. *The Lancet Microbe* (2021).
40. Gilbert, P.B. *et al.* Immune Correlates Analysis of the mRNA-1273 COVID-19 Vaccine Efficacy Trial. *medRxiv* (2021).
41. Earle, K.A. *et al.* Evidence for antibody as a protective correlate for COVID-19 vaccines. *Vaccine* **39**, 4423-4428 (2021).
42. Khoury, D.S. *et al.* Neutralizing antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2 infection. *Nature Medicine* **27**, 1205-1211 (2021).
43. Voysey, M. *et al.* Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *Lancet* **397**, 99-111 (2021).
44. Wood, S.N. *Generalized additive models: an introduction with R*. Chapman and Hall/CRC, 2006.
45. Cho, A. *et al.* Anti-SARS-CoV-2 receptor-binding domain antibody evolution after mRNA vaccination. *Nature* **600**, 517-522 (2021).
46. Kim, P., Gordon, S.M., Sheehan, M.M. & Rothberg, M.B. Duration of SARS-CoV-2 Natural Immunity and Protection against the Delta Variant: A Retrospective Cohort Study. *Clinical Infectious Diseases* (2021).
47. Goldberg, Y. *et al.* Protection and waning of natural and hybrid COVID-19 immunity. *medRxiv*, 2021.2012.2004.21267114-21262021.21267112.21267104.21267114 (2021).

48. Feikin, D. *et al.* Duration of Effectiveness of Vaccines Against SARS-CoV-2 Infection and COVID-19 Disease: Results of a Systematic Review and Meta-Regression. *SSRN Electronic Journal* (2021).
49. Papa, G. *et al.* Furin cleavage of SARS-CoV-2 Spike promotes but is not essential for infection and cell-cell fusion. *PLoS pathogens* **17**, e1009246-e1009246 (2021).
50. Braga, L. *et al.* Drugs that inhibit TMEM16 proteins block SARS-CoV-2 Spike-induced syncytia. *Nature* **594**, 88-93 (2021).
51. Zhang, J. *et al.* Membrane fusion and immune evasion by the spike protein of SARS-CoV-2 Delta variant. *Science*, eabl9463-eabl9463 (2021).
52. Kodaka, M. *et al.* A new cell-based assay to evaluate myogenesis in mouse myoblast C2C12 cells. *Experimental cell research* **336**, 171-181 (2015).
53. Abdelnabi, R. *et al.* The omicron (B.1.1.529) SARS-CoV-2 variant of concern does not readily infect Syrian hamsters. *bioRxiv*, 2021.2012.2024.474086-472021.474012.474024.474086 (2021).
54. Jackson, C.B., Farzan, M., Chen, B. & Choe, H. Mechanisms of SARS-CoV-2 entry into cells. *Nature Reviews Molecular Cell Biology* **23**, 3-20 (2022).
55. Mykytyn, A.Z. *et al.* SARS-CoV-2 entry into human airway organoids is serine protease-mediated and facilitated by the multibasic cleavage site. *Elife* **10**, e64508-e64508 (2021).
56. Hoffmann, M. *et al.* SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *cell* **181**, 271-280 (2020).
57. Hoffmann, M., Kleine-Weber, H. & Pöhlmann, S. A multibasic cleavage site in the spike protein of SARS-CoV-2 is essential for infection of human lung cells. *Molecular cell* **78**, 779-784 (2020).
58. Nie, J. *et al.* Functional comparison of SARS-CoV-2 with closely related pangolin and bat coronaviruses. *Cell discovery* **7**, 1-12 (2021).
59. Wrobel, A.G. *et al.* Structure and binding properties of Pangolin-CoV spike glycoprotein inform the evolution of SARS-CoV-2. *Nat. Commun.* **12**, 1-6 (2021).

60. Belouzard, S., Chu, V.C. & Whittaker, G.R. Activation of the SARS coronavirus spike protein via sequential proteolytic cleavage at two distinct sites. *Proceedings of the National Academy of Sciences* **106**, 5871-5876 (2009).
61. Dicken, S.J. *et al.* Characterisation of B. 1.1. 7 and Pangolin coronavirus spike provides insights on the evolutionary trajectory of SARS-CoV-2. *bioRxiv* (2021).
62. Saccon, E. *et al.* Cell-type-resolved quantitative proteomics map of interferon response against SARS-CoV-2. *Science* **24**, 102420-102420 (2021).
63. Winstone, H. *et al.* The polybasic cleavage site in SARS-CoV-2 spike modulates viral sensitivity to type I interferon and IFITM2. *Journal of virology* **95**, e02422-02420 (2021).
64. Daniloski, Z. *et al.* Identification of required host factors for SARS-CoV-2 infection in human cells. *Cell* **184**, 92-105 (2021).
65. Sheikh, A., Kerr, S., Woolhouse, M., McMenamin, J. & Robertson, C. Severity of omicron variant of concern and vaccine effectiveness against symptomatic disease: national cohort with nested test negative design study in Scotland. (2021).
66. Report 50 - Hospitalisation risk for Omicron cases in England | Faculty of Medicine | Imperial College London.
67. Benvenuto, D. *et al.* Evolutionary analysis of SARS-CoV-2: how mutation of Non-Structural Protein 6 (NSP6) could affect viral autophagy. *Journal of Infection* **81**, e24-e27 (2020).
68. Thorne, L.G. *et al.* Evolution of enhanced innate immune evasion by SARS-CoV-2. *Nature*, 1-12 (2021).
69. Rihn, S.J. *et al.* A plasmid DNA-launched SARS-CoV-2 reverse genetics system and coronavirus toolkit for COVID-19 research. *PLoS biology* **19**, e3001091-e3001091 (2021).
70. Davis, C. *et al.* Reduced neutralisation of the Delta (B. 1.617. 2) SARS-CoV-2 variant of concern following vaccination. *PLoS pathogens* **17**, e1010022-e1010022 (2021).
71. Hughes, E.C. *et al.* SARS-CoV-2 serosurveillance in a patient population reveals differences in virus exposure and antibody-mediated immunity according to host demography and healthcare setting. *J Infect Dis* (2020).

72. Newman, J. *et al.* Neutralising antibody activity against SARS-CoV-2 variants, including Omicron, in an elderly cohort vaccinated with BNT162b2. *medRxiv*, 2021.2012.2023.21268293-21262021.21268212.21268223.21268293 (2021).
73. Zufferey, R., Nagy, D., Mandel, R.J., Naldini, L. & Trono, D. Multiply attenuated lentiviral vector achieves efficient gene delivery in vivo. *Nat.Biotechnol.* **15**, 871-875 (1997).
74. Zufferey, R. *et al.* Self-inactivating lentivirus vector for safe and efficient in vivo gene delivery. *J Virol* **72**, 9873-9880 (1998).
75. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094-3100 (2018).
76. Price, M.N., Dehal, P.S. & Arkin, A.P. FastTree 2—approximately maximum-likelihood trees for large alignments. *PloS one* **5**, e9490-e9490 (2010).
77. Woo, H. *et al.* Developing a fully glycosylated full-length SARS-CoV-2 spike protein model in a viral membrane. *The Journal of Physical Chemistry B* **124**, 7128-7137 (2020).
78. Liu, Z. *et al.* Identification of SARS-CoV-2 spike mutations that attenuate monoclonal and serum antibody neutralization. *Cell host & microbe* **29**, 477-488 (2021).
79. Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature* **581**, 215-220 (2020).
80. Pymol, T. The PyMOL molecular graphics system. *Version 1*, r3pre-r3pre (2010).

Acknowledgments

The authors would like to thank the participants of the DOVE study and Sister Therese McSorley and her nursing team at the NHS GG&C clinical research facility. The authors also thank Alison Hamilton, Laura Stirling and Charlie Mayor from the NHS GG&C SafeHaven team for their invaluable input in facilitating this study. We thank Paula Olmo for administrative support and Chris Robertson and Aziz Sheikh for statistical advice. The authors thank all of the researchers who have shared genome data openly via the Global Initiative on Sharing All Influenza Data (GISAID). Funding was provided by Health Data Research UK (HDR UK) for the Evaluation of Variants Affecting Deployed COVID-19 Vaccine (EVADE) study (ECT, SR, OM, CW, BJW, grant code: 2021.0155). This research is part of the Data and Connectivity National Core Study, led by Health Data Research UK in partnership with the Office for National Statistics and funded by UK Research and Innovation (grant ref MC_PC_20058). This work was also supported by The Alan Turing Institute via 'Towards Turing 2.0' EPSRC Grant Funding. COG-UK is supported by funding from the Medical Research Council (MRC, part of UK Research & Innovation (UKRI)), the National Institute of Health Research (NIHR) [grant code: MC_PC_19027], and Genome Research Limited, operating as the Wellcome Sanger Institute (RMB, DLR, ECT). Medical Research Council (MRC) provided funding for both the COVID-19 Deployed VaccinE (DOVE) study (grant code: MCUU1201412) and COG-UK (ECT). AF, JH, RO, JG, ECT, NL and DLR were funded by the Medical Research Council (MRC) (grant code: MC_UU_12014/12). W.T.H. was supported by Medical Research Council (MRC), grant codes MR/R024758/1 and MR/W005611/1. The G2P-UK National Virology Consortium was funded by UK Research and Innovation (UKRI) award MR/W005611/1 (MP, ECT, AP, DLR). D.L.R. was funded by Wellcome Trust (grant code: 220977/Z/20/Z). N.L. and B.J.W. were funded by the Biotechnology and Biological Sciences Research Council (grant codes: BBSRC, BB/R004250/1 & BB/R019843/1). J.G. was funded by a Wellcome Trust and Royal Society Sir Henry Dale Fellowship (grant code: 107653/Z/15/A).

Author information

These authors contributed equally: Brian J. Willett, Joe Grove, Oscar A. MacLean, Craig Wilkie, Nicola Logan, Giuditta De Lorenzo & Wilhelm Furnon

These authors contributed equally: John Haughney, David L. Robertson, Massimo Palmarini, Surajit Ray & Emma C. Thomson

Affiliations

MRC-University of Glasgow Centre for Virus Research, University of Glasgow, UK.

Brian J. Willett, Joe Grove, Oscar A. MacLean, Nicola Logan, Giuditta De Lorenzo , Wilhelm Furnon, Sam Scott, Maria Manali, Agnieszka Szemiel, Shirin Ashraf, Elen Vink, William T. Harvey, Chris Davis, Richard Orton, Joseph Hughes, Ana da Silva Filipe, Rute Maria Pinto, Pablo R. Murcia, Arvind H. Patel, David L. Robertson, Massimo Palmarini, Emma C. Thomson

School of Mathematics & Statistics, University of Glasgow, UK

Craig Wilkie, Surajit Ray

NHS Greater Glasgow & Clyde, Glasgow, UK

Poppy Holland, Vanessa Silva, Kathryn Puxty, Rory Gunson, John Haughney, Emma C. Thomson

MRC Biostatistics Unit, University of Cambridge, UK.

David Pascall

Public Health Scotland.

Gonzalo Yebra, Sharif Shaaban, Matthew T. G. Holden

School of Medicine, University of St Andrews, UK.

Matthew T. G. Holden

NHS Lothian, Edinburgh, UK.

Kate Templeton

London School of Hygiene and Tropical Medicine, London, UK

Emma C. Thomson

Contributions

Conceptualization: ECT, JG, MP, DLR, BW

Methodology: BJW, JG, OAM, CW, NL, GL, WF, SS, MM, AS, WTH, CD, RO, JH, DP, KP, AF, GY, SS, RMP, PRM, AHP, JH, DLR, SR

Investigation: BJW, JG, OAM, CW, NL, GL, WF, SS, MM, AS, SA, EV, WTH, CD, RO, JH, PH, VS, DP, KP, AF, GY, SS, MTGH, RG, KT, AHP, JH, DLR, MP, SR

Visualization: BJW, JG, OAM, CW, GL, WTH, DLR

Funding acquisition: ECT, DLR, SR, BJW, JG

Project administration: ECT, DLR, SR, JH

Supervision: BJW, DLR, MP, SR, JH, ECT

Writing – original draft: ECT, BJW, JG, GY

Writing – review & editing: BJW, JG, OAM, CW, SA, WTH, DLR, MP, ECT

Corresponding authors

Correspondence to Professor Emma Thomson (emma.thomson@glasgow.ac.uk) and Professor Brian Willett (brian.willett@glasgow.ac.uk), MRC-University of Glasgow Centre for Virus Research, UK, G61 1QH.

Ethics declarations

The authors declare that they have no competing interests.


Extended data

Extended data Figs. 1 to 7

Supplementary Tables 1 to 10


The COVID-19 Genomics UK (COG-UK) consortium - June 2021 V.1

Extended data Fig.1. HCoV reactivity following two doses of SARS-CoV-2 vaccine. Antibody responses were studied in three groups of individuals (n=24 per group) vaccinated with either BNT162b2, ChAdOx1 or mRNA-1273 by MSD-ECL assay. Responses were measured against full-length spike glycoprotein (Spike) from HCoVs 229E, OC43, NL63 and HKU1 and are expressed as MSD arbitrary units (AU/ml). The response to OC43 was significantly higher in BNT162b2 vaccinates than in ChAdOx1 vaccinates.




Extended data Fig. 2. Influenza reactivity following two doses of SARS-CoV-2 vaccine.


Antibody responses were studied in three groups of individuals (n=24 per group) vaccinated with either BNT162b2, ChAdOx1 or mRNA-1273 by MSD-ECL assay. Responses were measured against haemagglutinins from influenza viruses; influenza A Michigan H1, Hong Kong H3 and Shanghai H7, and influenza B Phuket HA and Brisbane and are expressed as MSD arbitrary units (AU/ml). No significant differences were detected between the vaccine groups for each of the antigens.




Extended data Fig. 3. HCoV reactivity following third dose of SARS-CoV-2 vaccine. Antibody responses were studied in four groups of individuals primed with two doses of either ChAdOx1 or BNT162b2, followed by a booster of BNT162b2 or mRNA-1273. Responses were measured by MSD-ECL assay against full-length spike glycoprotein (Spike) from HCoVs 229E, OC43, NL63 and HKU1 and are expressed as MSD arbitrary units (AU/ml).




Extended data Fig. 4. Influenza reactivity following third dose of SARS-CoV-2 vaccine. Antibody responses were studied in four groups of individuals primed with two doses of either ChAdOx1 or BNT162b2, followed by a booster of BNT162b2 or mRNA-1273. Responses were measured by MSD-ECL against haemagglutinins from influenza viruses; influenza A Michigan H1, Hong Kong H3 and Shanghai H7, and influenza B Phuket HA and Brisbane and are expressed as MSD arbitrary units (AU/ml). * Significantly different p=0.0413.



Extended Data Fig. 5. Effect of third dose of SARS-CoV-2 vaccine on neutralising antibody titres. Two groups of healthy volunteers vaccinated with two doses of either ChAdOx1 or BNT162b2, were sampled two weeks following a third dose of either BNT162b2 or mRNA-1273. Each point represents the mean of three replicates. Where dose 2 and dose 3 samples were available from the same individual, points are joined by a solid line.



Extended Data Fig.6. Isolation of Omicron in cell culture. **a**, Vero ACE2 TMPRSS2 (VAT) and BHK-hACE2 cells were inoculated with diluted clinical samples. Viral progeny was quantified in the medium 5 dpi by RT-qPCR. **b**, Aliquots of the medium from samples named 204 and 205 were used to generate a P1 in BHK-hACE2 and Calu-3 cells and, limited to sample 205, a P2 in Calu-3 and Caco2 cells. Viral stocks were quantified by RT-qPCR.



Supplementary Table 1. Comparison of SARS-CoV-2 antibody responses elicited by two doses of SARS-CoV-2 vaccine. Data were analyzed in GraphPad Prism v8.4.3, groups were compared by ordinary one-way ANOVA.

Tukey's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value
Spike BNT162b2 vs. ChAdOx1	2588	-278.1 to 5453	No	ns	0.1212
Spike BNT162b2 vs. mRNA-1273	-248.4	-3114 to 2617	No	ns	>0.9999
Spike ChAdOx1 vs. mRNA-1273	-2836	-5702 to 29.73	No	ns	0.0553
RBD BNT162b2 vs. ChAdOx1	4649	1783 to 7514	Yes	****	<0.0001
RBD BNT162b2 vs. mRNA-1273	-449.1	-3315 to 2417	No	ns	>0.9999
RBD ChAdOx1 vs. mRNA-1273	-5098	-7964 to -2232	Yes	****	<0.0001
NTD BNT162b2 vs. ChAdOx1	5755	2889 to 8620	Yes	****	<0.0001
NTD BNT162b2 vs. mRNA-1273	714.7	-2151 to 3581	No	ns	0.9996
NTD ChAdOx1 vs. mRNA-1273	-5040	-7906 to -2174	Yes	****	<0.0001
N BNT162b2vs. ChAdOx1	2.183	-2864 to 2868	No	ns	>0.9999
N BNT162b2 vs. mRNA-1273	0.4	-2865 to 2866	No	ns	>0.9999
N ChAdOx1 vs. mRNA-1273	-1.783	-2868 to 2864	No	ns	>0.9999

Supplementary Table 2. Comparison of HCoV antibody responses elicited by two doses of SARS-CoV-2 vaccine. Data were analyzed in GraphPad Prism v8.4.3, groups were compared by ordinary one-way ANOVA.

Tukey's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value
229E BNT162b2 vs. ChAdOx1	3296	-19673 to 26265	No	ns	>0.9999
229E BNT162b2 vs. mRNA-1273	5282	-17687 to 28251	No	ns	0.9998
229E ChAdOx1 vs. mRNA-1273	1986	-20982 to 24955	No	ns	>0.9999
OC43 BNT162b2 vs. ChAdOx1	39581	16612 to 62550	Yes	****	<0.0001
OC43 BNT162b2 vs. mRNA-1273	18396	-4573 to 41365	No	ns	0.2628
OC43 ChAdOx1 vs. mRNA-1273	-21185	-44154 to 1784	No	ns	0.1027
NL63 BNT162b2 vs. ChAdOx1	1317	-21652 to 24286	No	ns	>0.9999
NL63 BNT162b2 vs. mRNA-1273	696.9	-22272 to 23666	No	ns	>0.9999
NL63 ChAdOx1 vs. mRNA-1273	-620	-23589 to 22349	No	ns	>0.9999
HKU1 BNT162b2 vs. ChAdOx1	17914	-5055 to 40883	No	ns	0.3015
HKU1 BNT162b2 vs. mRNA-1273	5468	-17501 to 28437	No	ns	0.9998
HKU1 ChAdOx1 vs. mRNA-1273	-12446	-35415 to 10523	No	ns	0.8243

Supplementary Table 3. Comparison of influenza antibody responses elicited by two doses of SARS-CoV-2 vaccine. Data were analyzed in GraphPad Prism v8.4.3, groups were compared by ordinary one-way ANOVA.

Tukey's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value
Flu A Michigan H1 BNT162b2 vs. ChAdOx1	42355	-49090 to 133800	No	ns	0.9615
Flu A Michigan H1 BNT162b2 vs. mRNA-1273	34380	-57065 to 125825	No	ns	0.9943
Flu A Michigan H1 ChAdOx1 vs. mRNA-1273	-7975	-99420 to 83469	No	ns	>0.9999
Flu A Hong Kong H3 BNT162b2 vs. ChAdOx1	22853	-68592 to 114298	No	ns	>0.9999
Flu A Hong Kong H3 BNT162b2 vs. mRNA-1273	43959	-47486 to 135403	No	ns	0.948
Flu A Hong Kong H3 ChAdOx1 vs. mRNA-1273	21106	-70339 to 112550	No	ns	>0.9999
Flu A Shanghai H7 BNT162b2 vs. ChAdOx1	-3370	-94815 to 88074	No	ns	>0.9999
Flu A Shanghai H7 BNT162b2 vs. mRNA-1273	-5570	-97015 to 85874	No	ns	>0.9999
Flu A Shanghai H7 ChAdOx1 vs. mRNA-1273	-2200	-93645 to 89244	No	ns	>0.9999
Flu B Phuket HA BNT162b2 vs. ChAdOx1	54701	-36744 to 146145	No	ns	0.7707
Flu B Phuket HA BNT162b2 vs. mRNA-1273	83834	-7611 to 175279	No	ns	0.1132
Flu B Phuket HA ChAdOx1 vs. mRNA-1273	29133	-62311 to 120578	No	ns	0.999
Flu B Brisbane BNT162b2 vs. ChAdOx1	35122	-56323 to 126566	No	ns	0.993
Flu B Brisbane BNT162b2 vs. mRNA-1273	64342	-27102 to 155787	No	ns	0.5138
Flu B Brisbane ChAdOx1 vs. mRNA-1273	29220	-62224 to 120665	No	ns	0.999

Supplementary Table 4. Comparison of neutralising antibody titres elicited by two doses of SARS-CoV-2 vaccine. Neutralising antibody responses were quantified against Wuhan or Omicron spike glycoprotein-bearing HIV(SARS-CoV-2) pseudotypes. Data were analyzed in GraphPad Prism v8.4.3, groups were compared by ordinary one-way ANOVA.

Tukey's multiple comparisons test	Mean 1	Mean 2	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value
Wuhan BNT162b2 vs. ChAdOx1	4978	882.3	4096	733.8 to 7458	Yes	**	0.0075
Wuhan BNT162b2 vs. mRNA-1273	4978	21118	-16140	-19502 to -12778	Yes	****	<0.0001
Wuhan ChAdOx1 vs. mRNA-1273	882.3	21118	-20236	-23598 to -16874	Yes	****	<0.0001
Omicron BNT162b2 vs. ChAdOx1	148.3	61.9	86.39	-3276 to 3448	No	ns	>0.9999
Omicron BNT162b2 vs. mRNA-1273	148.3	285	-136.7	-3499 to 3225	No	ns	>0.9999
Omicron ChAdOx1 vs. mRNA-1273	61.9	285	-223.1	-3585 to 3139	No	ns	>0.9999
BNT162b2 Wuhan vs. Omicron	4978	148.3	4830	1468 to 8192	Yes	***	0.0008
ChAdOx12 Wuhan vs. Omicron	882.3	61.9	820.4	-2542 to 4182	No	ns	0.981
mRNA-1273 Wuhan vs. Omicron	21118	285	20833	17471 to 24195	Yes	****	<0.0001

Supplementary Table 5. Comparison of SARS-CoV-2 antibody responses elicited by a third dose of SARS-CoV-2 vaccine. Data were analyzed in GraphPad Prism v8.4.3, groups were compared by ordinary one-way ANOVA. P= BNT162b2, AZ = ChAdOx1, M = mRNA-1273.

Tukey's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value
Spike AZ+P vs. AZ+M	1163	-4354 to 6680	No	ns	>0.9999
Spike AZ+P vs. P+P	1007	-3360 to 5374	No	ns	>0.9999
Spike AZ+P vs. P+M	361.3	-8410 to 9133	No	ns	>0.9999
Spike AZ+M vs. P+P	-156	-5239 to 4927	No	ns	>0.9999
Spike AZ+M vs. P+M	-801.5	-9951 to 8348	No	ns	>0.9999
Spike P+P vs. P+M	-645.5	-9151 to 7860	No	ns	>0.9999
RBD AZ+P vs. AZ+M	2675	-2842 to 8192	No	ns	0.9453
RBD AZ+P vs. P+P	1439	-2928 to 5806	No	ns	0.9988
RBD AZ+P vs. P+M	37.5	-8734 to 8809	No	ns	>0.9999
RBD AZ+M vs. P+P	-1236	-6319 to 3846	No	ns	>0.9999
RBD AZ+M vs. P+M	-2638	-11787 to 6511	No	ns	0.9997
RBD P+P vs. P+M	-1401	-9906 to 7104	No	ns	>0.9999
NTD AZ+P vs. AZ+M	3368	-2149 to 8885	No	ns	0.7418
NTD AZ+P vs. P+P	4098	-269.4 to 8465	No	ns	0.0919
NTD AZ+P vs. P+M	3739	-5033 to 12510	No	ns	0.9822
NTD AZ+M vs. P+P	729.8	-4353 to 5813	No	ns	>0.9999
NTD AZ+M vs. P+M	370.8	-8778 to 9520	No	ns	>0.9999
NTD P+P vs. P+M	-359	-8864 to 8146	No	ns	>0.9999
N AZ+P vs. AZ+M	2.755	-5514 to 5520	No	ns	>0.9999
N AZ+P vs. P+P	-0.7066	-4368 to 4366	No	ns	>0.9999
N AZ+P vs. P+M	-3.095	-8775 to 8768	No	ns	>0.9999
N AZ+M vs. P+P	-3.461	-5086 to 5079	No	ns	>0.9999
N AZ+M vs. P+M	-5.85	-9155 to 9143	No	ns	>0.9999
N P+P vs. P+M	-2.389	-8508 to 8503	No	ns	>0.9999

P = BNT162b2; AZ = ChAdOx1; M = mRNA-1273

Supplementary Table 6. Comparison of HCoV antibody responses elicited by a third dose of SARS-CoV-2 vaccine. Data were analyzed in GraphPad Prism v8.4.3, groups were compared by ordinary one-way ANOVA. P = BNT162b2, AZ = ChAdOx1, M = mRNA-1273.

Tukey's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value
229E AZ+P vs. AZ+M	-11229	-47020 to 24562	No	ns	0.9993
229E AZ+P vs. P+P	-5670	-34000 to 22661	No	ns	>0.9999
229E AZ+P vs. P+M	4487	-52417 to 61391	No	ns	>0.9999
229E AZ+M vs. P+P	5560	-27414 to 38533	No	ns	>0.9999
229E AZ+M vs. P+M	15716	-43637 to 75069	No	ns	>0.9999
229E P+P vs. P+M	10156	-45019 to 65332	No	ns	>0.9999
OC43 AZ+P vs. AZ+M	-29568	-65359 to 6224	No	ns	0.2369
OC43 AZ+P vs. P+P	-121	-28451 to 28209	No	ns	>0.9999
OC43 AZ+P vs. P+M	-5301	-62205 to 51603	No	ns	>0.9999
OC43 AZ+M vs. P+P	29447	-3527 to 62420	No	ns	0.1382
OC43 AZ+M vs. P+M	24266	-35086 to 83619	No	ns	0.988
OC43 P+P vs. P+M	-5180	-60356 to 49996	No	ns	>0.9999
NL63 AZ+P vs. AZ+M	-11140	-46931 to 24651	No	ns	0.9994
NL63 AZ+P vs. P+P	-8074	-36404 to 20256	No	ns	0.9998
NL63 AZ+P vs. P+M	-16842	-73746 to 40062	No	ns	0.9996
NL63 AZ+M vs. P+P	3066	-29908 to 36040	No	ns	>0.9999
NL63 AZ+M vs. P+M	-5702	-65055 to 53651	No	ns	>0.9999
NL63 P+P vs. P+M	-8768	-63944 to 46407	No	ns	>0.9999
HKU1 AZ+P vs. AZ+M	9482	-26309 to 45273	No	ns	>0.9999
HKU1 AZ+P vs. P+P	8028	-20303 to 36358	No	ns	0.9998
HKU1 AZ+P vs. P+M	8678	-48226 to 65582	No	ns	>0.9999
HKU1 AZ+M vs. P+P	-1454	-34428 to 31519	No	ns	>0.9999
HKU1 AZ+M vs. P+M	-804.1	-60157 to 58549	No	ns	>0.9999
HKU1 P+P vs. P+M	650.4	-54525 to 55826	No	ns	>0.9999

P = BNT162b2; AZ = ChAdOx1; M = mRNA-1273

Supplementary Table 7. Comparison of influenza antibody responses elicited by a third dose of SARS-CoV-2 vaccine. Data were analyzed in GraphPad Prism v8.4.3, groups were compared by ordinary one-way ANOVA. P = BNT162b2, AZ = ChAdOx1, M = mRNA-1273.

Tukey's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value
Flu A Michigan H1 AZ+P vs. AZ+M	410671	6882 to 814460	Yes	*	0.0413
Flu A Michigan H1 AZ+P vs. P+P	279008	-40609 to 598625	No	ns	0.1757
Flu A Michigan H1 AZ+P vs. P+M	459436	-182547 to 1101420	No	ns	0.5271
Flu A Michigan H1 AZ+M vs. P+P	-131663	-503667 to 240341	No	ns	0.9993
Flu A Michigan H1 AZ+M vs. P+M	48766	-620842 to 718373	No	ns	>0.9999
Flu A Michigan H1 P+P vs. P+M	180429	-442054 to 802911	No	ns	>0.9999
Flu A Hong Kong H3 AZ+P vs. AZ+M	-64025	-467814 to 339763	No	ns	>0.9999
Flu A Hong Kong H3 AZ+P vs. P+P	-32804	-352421 to 286812	No	ns	>0.9999
Flu A Hong Kong H3 AZ+P vs. P+M	56571	-585413 to 698554	No	ns	>0.9999
Flu A Hong Kong H3 AZ+M vs. P+P	31221	-340783 to 403225	No	ns	>0.9999
Flu A Hong Kong H3 AZ+M vs. P+M	120596	-549012 to 790204	No	ns	>0.9999
Flu A Hong Kong H3 P+P vs. P+M	89375	-533107 to 711857	No	ns	>0.9999
Flu A Shanghai H7 AZ+P vs. AZ+M	6841	-396948 to 410630	No	ns	>0.9999
Flu A Shanghai H7 AZ+P vs. P+P	469.8	-319147 to 320087	No	ns	>0.9999
Flu A Shanghai H7 AZ+P vs. P+M	9363	-632620 to 651346	No	ns	>0.9999
Flu A Shanghai H7 AZ+M vs. P+P	-6371	-378375 to 365633	No	ns	>0.9999
Flu A Shanghai H7 AZ+M vs. P+M	2522	-667086 to 672130	No	ns	>0.9999
Flu A Shanghai H7 P+P vs. P+M	8893	-613589 to 631376	No	ns	>0.9999
Flu B Phuket HA AZ+P vs. AZ+M	-101482	-505271 to 302307	No	ns	>0.9999
Flu B Phuket HA AZ+P vs. P+P	-6309	-325926 to 313308	No	ns	>0.9999
Flu B Phuket HA AZ+P vs. P+M	65660	-576323 to 707643	No	ns	>0.9999
Flu B Phuket HA AZ+M vs. P+P	95173	-276831 to 467178	No	ns	>0.9999
Flu B Phuket HA AZ+M vs. P+M	167142	-502466 to 836750	No	ns	>0.9999
Flu B Phuket HA P+P vs. P+M	71969	-550513 to 694451	No	ns	>0.9999
Flu B Brisbane AZ+P vs. AZ+M	-2954	-406743 to 400834	No	ns	>0.9999
Flu B Brisbane AZ+P vs. P+P	24661	-294956 to 344278	No	ns	>0.9999
Flu B Brisbane AZ+P vs. P+M	122461	-519522 to 764444	No	ns	>0.9999
Flu B Brisbane AZ+M vs. P+P	27615	-344389 to 399620	No	ns	>0.9999
Flu B Brisbane AZ+M vs. P+M	125415	-544193 to 795023	No	ns	>0.9999
Flu B Brisbane P+P vs. P+M	97800	-524683 to 720282	No	ns	>0.9999

P = BNT162b2; AZ = ChAdOx1; M = mRNA-1273

Supplementary Table 8. Effect of third dose of SARS-CoV-2 vaccine on neutralising antibody titres.
Neutralising antibody responses were quantified against Wuhan or Omicron spike glycoprotein-bearing HIV (SARS-CoV-2) pseudotypes. Data were analyzed in GraphPad Prism v8.4.3, groups were compared by ordinary one-way ANOVA.

BNT162b2 prime (doses 1 & 2)						
Tukey's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value	
Wuhan (dose 2) vs. (dose 3)	-1186	-4381 to 2010	No	ns	0.7655	
Omicron (dose 2) vs. (dose 3)	56.56	-3139 to 3252	No	ns	>0.9999	
Test details	Mean 1	Mean 2	Mean Diff.	SE of diff.	n1	
Wuhan (dose 2) vs. (dose 3)	4978	6164	-1186	1219	24	
Omicron (dose 2) vs. (dose 3)	148.3	91.73	56.56	1219	24	
ChAdOx1 prime (doses 1 & 2)						
Tukey's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value	
Wuhan (dose 2) vs. (dose 3)	-7360	-11073 to -3647	Yes	****	<0.0001	
Omicron (dose 2) vs. (dose 3)	-203.7	-3916 to 3509	No	ns	0.9989	
Test details	Mean 1	Mean 2	Mean Diff.	SE of diff.	n1	
Wuhan (dose 2) vs. (dose 3)	882.3	8242	-7360	1416	24	
Omicron (dose 2) vs. (dose 3)	61.9	265.5	-203.7	1416	24	
Dose 3 titre comparison						
Tukey's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value	
Wuhan (BNT162b2 dose 2) vs. (ChAdOX1 dose 2)	-2079	-7008 to 2851	No	ns	0.6859	
Omicron (BNT162b2 dose 2) vs. (ChAdOX1 dose 2)	-173.8	-5103 to 4756	No	ns	0.9997	
Test details	Mean 1	Mean 2	Mean Diff.	SE of diff.	n1	
Wuhan (BNT162b2 dose 2) vs. (ChAdOX1 dose 2)	6164	8242	-2079	1877	20	
Omicron (BNT162b2 dose 2) vs. (ChAdOX1 dose 2)	91.73	265.5	-173.8	1877	20	

Supplementary Table 9: Demographics, SARS-CoV-2 positivity status and vaccination status for the study population. Population consisted of 101,310 people aged 18 and over, registered as living in NHS Greater Glasgow and Clyde and tested by PCR test for SARS-CoV-2 infection between 22nd and 28th December 2021, split by SARS-CoV-2 variant status.

	<i>Negative</i>	<i>Delta</i>	<i>Omicron</i>
Demographics			
Age on 31st October 2021			
<i>Minimum</i>	18.85	18.85	18.85
<i>1st Quartile</i>	40.57	30.17	29.75
<i>Median</i>	57.88	39.74	44.73
<i>Mean</i>	55.46	43.99	46.21
<i>3rd Quartile</i>	68.70	57.05	61.13
<i>Maximum</i>	108.56	107.32	98.66
Sex			
<i>Female</i>	53438 (59.22%)	2634 (53.63%)	3356 (54.43%)
<i>Male</i>	36795 (40.78%)	2277 (46.37%)	2810 (45.57%)
SIMD (2016) quartile			
<i>1</i>	33908 (37.58%)	2365 (48.16%)	2302 (37.33%)
<i>2</i>	16901 (18.73%)	895 (18.22%)	1184 (19.2%)
<i>3</i>	14855 (16.46%)	725 (14.76%)	1048 (17%)
<i>4</i>	23003 (25.49%)	831 (16.92%)	1526 (24.75%)
<i>Unknown</i>	1566 (1.74%)	95 (1.93%)	106 (1.72%)
SARS-CoV-2 positivity status			
Previous confirmed SARS-CoV-2 infection status			
<i>No previous infection</i>	83000 (91.98%)	4812 (97.98%)	5793 (93.95%)
<i>Had previous infection</i>	7233 (8.02%)	99 (2.02%)	373 (6.05%)
Vaccination status			
Vaccination schedule			
<i>Unvaccinated</i>	13017 (14.45%)	3017 (61.52%)	1901 (30.88%)
<i>1st dose AstraZeneca</i>	19 (0.02%)	<=5	<=5
<i>1st dose Pfizer</i>	437 (0.49%)	26 (0.53%)	45 (0.73%)
<i>1st dose Moderna</i>	295 (0.33%)	17 (0.35%)	30 (0.49%)
<i>2nd dose AstraZeneca</i>	468 (0.52%)	19 (0.39%)	45 (0.73%)
<i>2nd dose Pfizer</i>	2448 (2.72%)	71 (1.45%)	278 (4.52%)
<i>2nd dose Moderna</i>	761 (0.84%)	17 (0.35%)	90 (1.46%)
<i>2 x AstraZeneca, 1 x Pfizer</i>	23641 (26.25%)	581 (11.85%)	1130 (18.36%)
<i>2 x Moderna, 1 x Pfizer</i>	235 (0.26%)	10 (0.2%)	19 (0.31%)
<i>3 x Pfizer</i>	21704 (24.1%)	491 (10.01%)	1193 (19.38%)
<i>2 x AstraZeneca, 1 x Moderna</i>	15159 (16.83%)	394 (8.03%)	743 (12.07%)
<i>2 x Pfizer, 1 x Moderna</i>	11581 (12.86%)	254 (5.18%)	655 (10.64%)

3 x Moderna		296 (0.33%)			<=5			25 (0.41%)		
Time (days) since most recent dose by variant and dose number										
	<i>Negative</i>			<i>Delta</i>			<i>Omicron</i>			
	<i>1</i>	<i>2</i>	<i>3</i>	<i>1</i>	<i>2</i>	<i>3</i>	<i>1</i>	<i>2</i>	<i>3</i>	
<i>Minimum</i>	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.0	
<i>1st Quartile</i>	22.00	34.00	25.00	20.00	33.50	18.00	24.00	38.00	25.0	
<i>Median</i>	32.00	49.00	41.00	34.00	51.00	30.00	34.00	56.00	39.0	
<i>Mean</i>	31.15	48.32	42.04	30.29	47.53	35.44	32.71	53.15	41.5	
<i>3rd Quartile</i>	39.00	65.00	58.00	37.00	65.50	50.00	40.00	69.00	58.0	
<i>Maximum</i>	49.00	77.00	77.00	49.00	77.00	77.00	49.00	77.00	77.0	

The COVID-19 Genomics UK (COG-UK) consortium - June 2021 V.1

Funding acquisition, Leadership and supervision, Metadata curation, Project administration, Samples and logistics, Sequencing and analysis, Software and analysis tools, and Visualisation:
Dr Samuel C Robson ^{13, 84}

Funding acquisition, Leadership and supervision, Metadata curation, Project administration, Samples and logistics, Sequencing and analysis, and Software and analysis tools:
Dr Thomas R Connor ^{11, 74} and Prof Nicholas J Loman ⁴³

Leadership and supervision, Metadata curation, Project administration, Samples and logistics, Sequencing and analysis, Software and analysis tools, and Visualisation:
Dr Tanya Golubchik ⁵

Funding acquisition, Leadership and supervision, Metadata curation, Samples and logistics, Sequencing and analysis, and Visualisation:
Dr Rocio T Martinez Nunez ⁴⁶

Funding acquisition, Leadership and supervision, Project administration, Samples and logistics, Sequencing and analysis, and Software and analysis tools:
Dr David Bonsall ⁵

Funding acquisition, Leadership and supervision, Project administration, Sequencing and analysis, Software and analysis tools, and Visualisation:
Prof Andrew Rambaut ¹⁰⁴

Funding acquisition, Metadata curation, Project administration, Samples and logistics, Sequencing and analysis, and Software and analysis tools:
Dr Luke B Snell ¹²

Leadership and supervision, Metadata curation, Project administration, Samples and logistics, Software and analysis tools, and Visualisation:
Rich Livett ¹¹⁶

Funding acquisition, Leadership and supervision, Metadata curation, Project administration, and Samples and logistics:
Dr Catherine Ludden ^{20, 70}

Funding acquisition, Leadership and supervision, Metadata curation, Samples and logistics, and Sequencing and analysis:
Dr Sally Corden ⁷⁴ and Dr Eleni Nastouli ^{96, 95, 30}

Funding acquisition, Leadership and supervision, Metadata curation, Sequencing and analysis, and Software and analysis tools:
Dr Gaia Nebbia ¹²

Funding acquisition, Leadership and supervision, Project administration, Samples and logistics, and Sequencing and analysis:
Ian Johnston ¹¹⁶

Leadership and supervision, Metadata curation, Project administration, Samples and logistics, and Sequencing and analysis:

Prof Katrina Lythgoe⁵, Dr M. Estee Torok^{19,20} and Prof Ian G Goodfellow²⁴

Leadership and supervision, Metadata curation, Project administration, Samples and logistics, and Visualisation:

Dr Jacqui A Prieto^{97,82} and Dr Kordo Saeed^{97,83}

Leadership and supervision, Metadata curation, Project administration, Sequencing and analysis, and Software and analysis tools:

Dr David K Jackson¹¹⁶

Leadership and supervision, Metadata curation, Samples and logistics, Sequencing and analysis, and Visualisation:

Dr Catherine Houlihan^{96,94}

Leadership and supervision, Metadata curation, Sequencing and analysis, Software and analysis tools, and Visualisation:

Dr Dan Frampton^{94,95}

Metadata curation, Project administration, Samples and logistics, Sequencing and analysis, and Software and analysis tools:

Dr William L Hamilton¹⁹ and Dr Adam A Witney⁴¹

Funding acquisition, Samples and logistics, Sequencing and analysis, and Visualisation:

Dr Giselda Bucca¹⁰¹

Funding acquisition, Leadership and supervision, Metadata curation, and Project administration:

Dr Cassie F Pope^{40,41}

Funding acquisition, Leadership and supervision, Metadata curation, and Samples and logistics:

Dr Catherine Moore⁷⁴

Funding acquisition, Leadership and supervision, Metadata curation, and Sequencing and analysis:

Prof Emma C Thomson⁵³

Funding acquisition, Leadership and supervision, Project administration, and Samples and logistics:

Dr Ewan M Harrison^{116,102}

Funding acquisition, Leadership and supervision, Sequencing and analysis, and Visualisation:

Prof Colin P Smith¹⁰¹

Leadership and supervision, Metadata curation, Project administration, and Sequencing and analysis:

Fiona Rogan⁷⁷

Leadership and supervision, Metadata curation, Project administration, and Samples and logistics:

Shaun M Beckwith⁶, Abigail Murray⁶, Dawn Singleton⁶, Dr Kirstine Eastick³⁷, Dr Liz A Sheridan⁹⁸, Paul Randell⁹⁹, Dr Leigh M Jackson¹⁰⁵, Dr Cristina V Ariani¹¹⁶ and Dr Sónia Gonçalves¹¹⁶

Leadership and supervision, Metadata curation, Samples and logistics, and Sequencing and analysis:

Dr Derek J Fairley^{3,77}, Prof Matthew W Loose¹⁸ and Joanne Watkins⁷⁴

Leadership and supervision, Metadata curation, Samples and logistics, and Visualisation:

Dr Samuel Moses^{25,106}

Leadership and supervision, Metadata curation, Sequencing and analysis, and Software and analysis tools:

Dr Sam Nicholls⁴³, Dr Matthew Bull⁷⁴ and Dr Roberto Amato¹¹⁶

Leadership and supervision, Project administration, Samples and logistics, and Sequencing and analysis:

Prof Darren L Smith^{36,65,66}

Leadership and supervision, Sequencing and analysis, Software and analysis tools, and Visualisation:

Prof David M Aanensen^{14,116} and Dr Jeffrey C Barrett¹¹⁶

Metadata curation, Project administration, Samples and logistics, and Sequencing and analysis:

Dr Dinesh Aggarwal^{20,116,70}, Dr James G Shepherd⁵³, Dr Martin D Curran⁷¹ and Dr Surendra Parmar⁷¹

Metadata curation, Project administration, Sequencing and analysis, and Software and analysis tools:

Dr Matthew D Parker¹⁰⁹

Metadata curation, Samples and logistics, Sequencing and analysis, and Software and analysis tools:

Dr Catryn Williams⁷⁴

Metadata curation, Samples and logistics, Sequencing and analysis, and Visualisation:

Dr Sharon Glaysher⁶⁸

Metadata curation, Sequencing and analysis, Software and analysis tools, and Visualisation:

Dr Anthony P Underwood^{14,116}, Dr Matthew Bashton^{36,65}, Dr Nicole Pacchiarini⁷⁴, Dr Katie F Loveson⁸⁴ and Matthew Byott^{95,96}

Project administration, Sequencing and analysis, Software and analysis tools, and Visualisation:

Dr Alessandro M Carabelli²⁰

Funding acquisition, Leadership and supervision, and Metadata curation:

Dr Kate E Templeton^{56,104}

Funding acquisition, Leadership and supervision, and Project administration:

Dr Thushan I de Silva¹⁰⁹, Dr Dennis Wang¹⁰⁹, Dr Cordelia F Langford¹¹⁶ and John Sillitoe¹¹⁶

Funding acquisition, Leadership and supervision, and Samples and logistics:

Prof Rory N Gunson⁵⁵

Funding acquisition, Leadership and supervision, and Sequencing and analysis:

Dr Simon Cottrell ⁷⁴, Dr Justin O'Grady ^{75, 103} and Prof Dominic Kwiatkowski ^{116, 108}

Leadership and supervision, Metadata curation, and Project administration:

Dr Patrick J Lillie ³⁷

Leadership and supervision, Metadata curation, and Samples and logistics:

Dr Nicholas Cortes ³³, Dr Nathan Moore ³³, Dr Claire Thomas ³³, Phillipa J Burns ³⁷, Dr Tabitha W Mahungu ⁸⁰ and Steven Liggett ⁸⁶

Leadership and supervision, Metadata curation, and Sequencing and analysis:

Angela H Beckett ^{13, 81} and Prof Matthew TG Holden ⁷³

Leadership and supervision, Project administration, and Samples and logistics:

Dr Lisa J Levett ³⁴, Dr Husam Osman ^{70, 35} and Dr Mohammed O Hassan-Ibrahim ⁹⁹

Leadership and supervision, Project administration, and Sequencing and analysis:

Dr David A Simpson ⁷⁷

Leadership and supervision, Samples and logistics, and Sequencing and analysis:

Dr Meera Chand ⁷², Prof Ravi K Gupta ¹⁰², Prof Alistair C Darby ¹⁰⁷ and Prof Steve Paterson ¹⁰⁷

Leadership and supervision, Sequencing and analysis, and Software and analysis tools:

Prof Oliver G Pybus ²³, Dr Erik M Volz ³⁹, Prof Daniela de Angelis ⁵², Prof David L Robertson ⁵³, Dr Andrew J Page ⁷⁵ and Dr Inigo Martincorena ¹¹⁶

Leadership and supervision, Sequencing and analysis, and Visualisation:

Dr Louise Aigrain ¹¹⁶ and Dr Andrew R Bassett ¹¹⁶

Metadata curation, Project administration, and Samples and logistics:

Dr Nick Wong ⁵⁰, Dr Yusri Taha ⁸⁹, Michelle J Erkiert ⁹⁹ and Dr Michael H Spencer Chapman ^{116, 102}

Metadata curation, Project administration, and Sequencing and analysis:

Dr Rebecca Dewar ⁵⁶ and Martin P McHugh ^{56, 111}

Metadata curation, Project administration, and Software and analysis tools:

Siddharth Mookerjee ^{38, 57}

Metadata curation, Project administration, and Visualisation:

Stephen Aplin ⁹⁷, Matthew Harvey ⁹⁷, Thea Sass ⁹⁷, Dr Helen Umpleby ⁹⁷ and Helen Wheeler ⁹⁷

Metadata curation, Samples and logistics, and Sequencing and analysis:

Dr James P McKenna ³, Dr Ben Warne ⁹, Joshua F Taylor ²², Yasmin Chaudhry ²⁴, Rhys Izuagbe ²⁴, Dr Aminu S Jahun ²⁴, Dr Gregory R Young ^{36, 65}, Dr Claire McMurray ⁴³, Dr Clare M McCann ^{65, 66}, Dr Andrew Nelson ^{65, 66} and Scott Elliott ⁶⁸

Metadata curation, Samples and logistics, and Visualisation:

Hannah Lowe ²⁵

Metadata curation, Sequencing and analysis, and Software and analysis tools:

Dr Anna Price ¹¹, Matthew R Crown ⁶⁵, Dr Sara Rey ⁷⁴, Dr Sunando Roy ⁹⁶ and Dr Ben Temperton ¹⁰⁵

Metadata curation, Sequencing and analysis, and Visualisation:

Dr Sharif Shaaban ⁷³ and Dr Andrew R Hesketh ¹⁰¹

Project administration, Samples and logistics, and Sequencing and analysis:

Dr Kenneth G Laing ⁴¹, Dr Irene M Monahan ⁴¹ and Dr Judith Heaney ^{95, 96, 34}

Project administration, Samples and logistics, and Visualisation:

Dr Emanuela Pelosi ⁹⁷, Siona Silveira ⁹⁷ and Dr Eleri Wilson-Davies ⁹⁷

Samples and logistics, Software and analysis tools, and Visualisation:

Dr Helen Fryer ⁵

Sequencing and analysis, Software and analysis tools, and Visualization:

Dr Helen Adams ⁴, Dr Louis du Plessis ²³, Dr Rob Johnson ³⁹, Dr William T Harvey ^{53, 42}, Dr Joseph Hughes ⁵³, Dr Richard J Orton ⁵³, Dr Lewis G Spurgin ⁵⁹, Dr Yann Bourgeois ⁸¹, Dr Chris Ruis ¹⁰², Áine O'Toole ¹⁰⁴, Marina Gourtovaia ¹¹⁶ and Dr Theo Sanderson ¹¹⁶

Funding acquisition, and Leadership and supervision:

Dr Christophe Fraser ⁵, Dr Jonathan Edgeworth ¹², Prof Judith Breuer ^{96, 29}, Dr Stephen L Michell ¹⁰⁵ and Prof John A Todd ¹¹⁵

Funding acquisition, and Project administration:

Michaela John ¹⁰ and Dr David Buck ¹¹⁵

Leadership and supervision, and Metadata curation:

Dr Kavitha Gajee ³⁷ and Dr Gemma L Kay ⁷⁵

Leadership and supervision, and Project administration:

Prof Sharon J Peacock ^{20, 70} and David Heyburn ⁷⁴

Leadership and supervision, and Samples and logistics:

Katie Kitchman ³⁷, Prof Alan McNally ^{43, 93}, David T Pritchard ⁵⁰, Dr Samir Dervisevic ⁵⁸, Dr Peter Muir ⁷⁰, Dr Esther Robinson ^{70, 35}, Dr Barry B Vipond ⁷⁰, Newara A Ramadan ⁷⁸, Dr Christopher Jeanes ⁹⁰, Danni Weldon ¹¹⁶, Jana Catalan ¹¹⁸ and Neil Jones ¹¹⁸

Leadership and supervision, and Sequencing and analysis:

Dr Ana da Silva Filipe ⁵³, Dr Chris Williams ⁷⁴, Marc Fuchs ⁷⁷, Dr Julia Miskelly ⁷⁷, Dr Aaron R Jeffries ¹⁰⁵, Karen Oliver ¹¹⁶ and Dr Naomi R Park ¹¹⁶

Metadata curation, and Samples and logistics:

Amy Ash ¹, Cherian Koshy ¹, Magdalena Barrow ⁷, Dr Sarah L Buchan ⁷, Dr Anna Mantzouratou ⁷, Dr Gemma Clark ¹⁵, Dr Christopher W Holmes ¹⁶, Sharon Campbell ¹⁷, Thomas Davis ²¹, Ngee Keong Tan ²², Dr Julianne R Brown ²⁹, Dr Kathryn A Harris ^{29, 2}, Stephen P Kidd ³³, Dr Paul R Grant ³⁴, Dr Li Xu-McCrae ³⁵, Dr Alison Cox ^{38, 63}, Pinglawathee Madona ^{38, 63}, Dr Marcus Pond ^{38, 63}, Dr Paul A Randell ^{38, 63}, Karen T Withell ⁴⁸, Cheryl Williams ⁵¹, Dr Clive Graham ⁶⁰, Rebecca Denton-Smith ⁶², Emma Swindells ⁶², Robyn Turnbull ⁶², Dr Tim J Sloan ⁶⁷, Dr Andrew Bosworth ^{70, 35}, Stephanie Hutchings ⁷⁰, Hannah M Pymont ⁷⁰, Dr Anna Casey ⁷⁶, Dr Liz Ratcliffe ⁷⁶, Dr Christopher R Jones ^{79, 105}, Dr Bridget A Knight ^{79, 105}, Dr Tanzina Haque ⁸⁰, Dr Jennifer Hart ⁸⁰, Dr Dianne Irish-Tavares ⁸⁰, Eric Witele ⁸⁰, Craig Mower ⁸⁶, Louisa K Watson ⁸⁶, Jennifer Collins ⁸⁹, Gary Eltringham ⁸⁹, Dorian Crudgington ⁹⁸, Ben Macklin ⁹⁸, Prof Miren Iturriza-Gomara ¹⁰⁷, Dr Anita O Lucaci ¹⁰⁷ and Dr Patrick C McClure ¹¹³

Metadata curation, and Sequencing and analysis:

Matthew Carlile¹⁸, Dr Nadine Holmes¹⁸, Dr Christopher Moore¹⁸, Dr Nathaniel Storey²⁹, Dr Stefan Rooke⁷³, Dr Gonzalo Yebra⁷³, Dr Noel Craine⁷⁴, Malorie Perry⁷⁴, Dr Nabil-Fareed Alikhan⁷⁵, Dr Stephen Bridgett⁷⁷, Kate F Cook⁸⁴, Christopher Fearn⁸⁴, Dr Salman Goudarzi⁸⁴, Prof Ronan A Lyons⁸⁸, Dr Thomas Williams¹⁰⁴, Dr Sam T Haldenby¹⁰⁷, Jillian Durham¹¹⁶ and Dr Steven Leonard¹¹⁶

Metadata curation, and Software and analysis tools:

Robert M Davies¹¹⁶

Project administration, and Samples and logistics:

Dr Rahul Batra¹², Beth Blane²⁰, Dr Moira J Spyer^{30,95,96}, Perminder Smith^{32,112}, Mehmet Yavus^{85,109}, Dr Rachel J Williams⁹⁶, Dr Adhyana IK Mahanama⁹⁷, Dr Buddhini Samaraweera⁹⁷, Sophia T Girgis¹⁰², Samantha E Hansford¹⁰⁹, Dr Angie Green¹¹⁵, Dr Charlotte Beaver¹¹⁶, Katherine L Bellis^{116,102}, Matthew J Dorman¹¹⁶, Sally Kay¹¹⁶, Liam Prestwood¹¹⁶ and Dr Shavanthi Rajatileka¹¹⁶

Project administration, and Sequencing and analysis:

Dr Joshua Quick⁴³

Project administration, and Software and analysis tools:

Radoslaw Poplawski⁴³

Samples and logistics, and Sequencing and analysis:

Dr Nicola Reynolds⁸, Andrew Mack¹¹, Dr Arthur Morriss¹¹, Thomas Whalley¹¹, Bindi Patel¹², Dr Iliana Georgana²⁴, Dr Myra Hosmillo²⁴, Malte L Pinckert²⁴, Dr Joanne Stockton⁴³, Dr John H Henderson⁶⁵, Amy Hollis⁶⁵, Dr William Stanley⁶⁵, Dr Wen C Yew⁶⁵, Dr Richard Myers⁷², Dr Alicia Thornton⁷², Alexander Adams⁷⁴, Tara Annett⁷⁴, Dr Hibo Asad⁷⁴, Alec Birchley⁷⁴, Jason Coombes⁷⁴, Johnathan M Evans⁷⁴, Laia Fina⁷⁴, Bree Gatica-Wilcox⁷⁴, Lauren Gilbert⁷⁴, Lee Graham⁷⁴, Jessica Hey⁷⁴, Ember Hilvers⁷⁴, Sophie Jones⁷⁴, Hannah Jones⁷⁴, Sara Kumziene-Summerhayes⁷⁴, Dr Caoimhe McKerr⁷⁴, Jessica Powell⁷⁴, Georgia Pugh⁷⁴, Sarah Taylor⁷⁴, Alexander J Trotter⁷⁵, Charlotte A Williams⁹⁶, Leanne M Kermack¹⁰², Benjamin H Foulkes¹⁰⁹, Marta Gallis¹⁰⁹, Hailey R Hornsby¹⁰⁹, Stavroula F Louka¹⁰⁹, Dr Manoj Pohare¹⁰⁹, Paige Wolverson¹⁰⁹, Peijun Zhang¹⁰⁹, George MacIntyre-Cockett¹¹⁵, Amy Trebes¹¹⁵, Dr Robin J Moll¹¹⁶, Lynne Ferguson¹¹⁷, Dr Emily J Goldstein¹¹⁷, Dr Alasdair Maclean¹¹⁷ and Dr Rachael Tomb¹¹⁷

Samples and logistics, and Software and analysis tools:

Dr Igor Starinskij⁵³

Sequencing and analysis, and Software and analysis tools:

Laura Thomson⁵, Joel Southgate^{11,74}, Dr Moritz UG Kraemer²³, Dr Jayna Raghwanani²³, Dr Alex E Zarebski²³, Olivia Boyd³⁹, Lily Geidelberg³⁹, Dr Chris J Illingworth⁵², Dr Chris Jackson⁵², Dr David Pascall⁵², Dr Sreenu Vattipally⁵³, Timothy M Freeman¹⁰⁹, Dr Sharon N Hsu¹⁰⁹, Dr Benjamin B Lindsey¹⁰⁹, Dr Keith James¹¹⁶, Kevin Lewis¹¹⁶, Gerry Tonkin-Hill¹¹⁶ and Dr Jaime M Tovar-Corona¹¹⁶

Sequencing and analysis, and Visualisation:

MacGregor Cox²⁰

Software and analysis tools, and Visualisation:

Dr Khalil Abudahab^{14,116}, Mirko Menegazzo¹⁴, Ben EW Taylor MEng^{14,116}, Dr Corin A Yeats¹⁴, Afrida Mukaddas⁵³, Derek W Wright⁵³, Dr Leonardo de Oliveira Martins⁷⁵, Dr Rachel Colquhoun¹⁰⁴, Verity

Hill ¹⁰⁴, Dr Ben Jackson ¹⁰⁴, Dr JT McCrone ¹⁰⁴, Dr Nathan Medd ¹⁰⁴, Dr Emily Scher ¹⁰⁴ and Jon-Paul Keatley ¹¹⁶

Leadership and supervision:

Dr Tanya Curran ³, Dr Sian Morgan ¹⁰, Prof Patrick Maxwell ²⁰, Prof Ken Smith ²⁰, Dr Sahar Eldirdiri ²¹, Anita Kenyon ²¹, Prof Alison H Holmes ^{38,57}, Dr James R Price ^{38,57}, Dr Tim Wyatt ⁶⁹, Dr Alison E Mather ⁷⁵, Dr Timofey Skvortsov ⁷⁷ and Prof John A Hartley ⁹⁶

Metadata curation:

Prof Martyn Guest ¹¹, Dr Christine Kitchen ¹¹, Dr Ian Merrick ¹¹, Robert Munn ¹¹, Dr Beatrice Bertolusso ³³, Dr Jessica Lynch ³³, Dr Gabrielle Vernet ³³, Stuart Kirk ³⁴, Dr Elizabeth Wastnedge ⁵⁶, Dr Rachael Stanley ⁵⁸, Giles Idle ⁶⁴, Dr Declan T Bradley ^{69,77}, Dr Jennifer Poyner ⁷⁹ and Matilde Mori ¹¹⁰

Project administration:

Owen Jones ¹¹, Victoria Wright ¹⁸, Ellena Brooks ²⁰, Carol M Churcher ²⁰, Mireille Fragakis ²⁰, Dr Katerina Galai ^{20,70}, Dr Andrew Jermy ²⁰, Sarah Judges ²⁰, Georgina M McManus ²⁰, Kim S Smith ²⁰, Dr Elaine Westwick ²⁰, Dr Stephen W Attwood ²³, Dr Frances Bolt ^{38,57}, Dr Alisha Davies ⁷⁴, Elen De Lacy ⁷⁴, Fatima Downing ⁷⁴, Sue Edwards ⁷⁴, Lizzie Meadows ⁷⁵, Sarah Jeremiah ⁹⁷, Dr Nikki Smith ¹⁰⁹ and Luke Foulser ¹¹⁶

Samples and logistics:

Dr Themoula Charalampous ^{12,46}, Amita Patel ¹², Dr Louise Berry ¹⁵, Dr Tim Boswell ¹⁵, Dr Vicki M Fleming ¹⁵, Dr Hannah C Howson-Wells ¹⁵, Dr Amelia Joseph ¹⁵, Manjinder Khakh ¹⁵, Dr Michelle M Lister ¹⁵, Paul W Bird ¹⁶, Karlie Fallon ¹⁶, Thomas Helmer ¹⁶, Dr Claire L McMurray ¹⁶, Mina Odedra ¹⁶, Jessica Shaw ¹⁶, Dr Julian W Tang ¹⁶, Nicholas J Willford ¹⁶, Victoria Blakey ¹⁷, Dr Veena Raviprakash ¹⁷, Nicola Sheriff ¹⁷, Lesley-Anne Williams ¹⁷, Theresa Feltwell ²⁰, Dr Luke Bedford ²⁶, Dr James S Cargill ²⁷, Warwick Hughes ²⁷, Dr Jonathan Moore ²⁸, Susanne Stonehouse ²⁸, Laura Atkinson ²⁹, Jack CD Lee ²⁹, Dr Divya Shah ²⁹, Adela Alcolea-Medina ^{32,112}, Natasha Ohemeng-Kumi ^{32,112}, John Ramble ^{32,112}, Jasveen Sehmi ^{32,112}, Dr Rebecca Williams ³³, Wendy Chatterton ³⁴, Monika Pusok ³⁴, William Everson ³⁷, Anibolina Castigador ⁴⁴, Emily Macnaughton ⁴⁴, Dr Kate El Bouzidi ⁴⁵, Dr Temi Lampejo ⁴⁵, Dr Malur Sudhanva ⁴⁵, Cassie Breen ⁴⁷, Dr Graciela Sluga ⁴⁸, Dr Shazaad SY Ahmad ^{49,70}, Dr Ryan P George ⁴⁹, Dr Nicholas W Machin ^{49,70}, Debbie Binns ⁵⁰, Victoria James ⁵⁰, Dr Rachel Blacow ⁵⁵, Dr Lindsay Coupland ⁵⁸, Dr Louise Smith ⁵⁹, Dr Edward Barton ⁶⁰, Debra Padgett ⁶⁰, Garren Scott ⁶⁰, Dr Aidan Cross ⁶¹, Dr Mariyam Mirfenderesky ⁶¹, Jane Greenaway ⁶², Kevin Cole ⁶⁴, Phillip Clarke ⁶⁷, Nichola Duckworth ⁶⁷, Sarah Walsh ⁶⁷, Kelly Bicknell ⁶⁸, Robert Impey ⁶⁸, Dr Sarah Wyllie ⁶⁸, Richard Hopes ⁷⁰, Dr Chloe Bishop ⁷², Dr Vicki Chalker ⁷², Dr Ian Harrison ⁷², Laura Gifford ⁷⁴, Dr Zoltan Molnar ⁷⁷, Dr Cressida Auckland ⁷⁹, Dr Cariad Evans ^{85,109}, Dr Kate Johnson ^{85,109}, Dr David G Partridge ^{85,109}, Dr Mohammad Raza ^{85,109}, Paul Baker ⁸⁶, Prof Stephen Bonner ⁸⁶, Sarah Essex ⁸⁶, Leanne J Murray ⁸⁶, Andrew I Lawton ⁸⁷, Dr Shirelle Burton-Fanning ⁸⁹, Dr Brendan Al Payne ⁸⁹, Dr Sheila Waugh ⁸⁹, Andrea N Gomes ⁹¹, Maimuna Kimuli ⁹¹, Darren R Murray ⁹¹, Paula Ashfield ⁹², Dr Donald Dobie ⁹², Dr Fiona Ashford ⁹³, Dr Angus Best ⁹³, Dr Liam Crawford ⁹³, Dr Nicola Cumley ⁹³, Dr Megan Mayhew ⁹³, Dr Oliver Megram ⁹³, Dr Jeremy Mirza ⁹³, Dr Emma Moles-Garcia ⁹³, Dr Benita Percival ⁹³, Megan Driscoll ⁹⁶, Leah Ensell ⁹⁶, Dr Helen L Lowe ⁹⁶, Laurentiu Maftai ⁹⁶, Matteo Mondani ⁹⁶, Nicola J Chaloner ⁹⁹, Benjamin J Cogger ⁹⁹, Lisa J Easton ⁹⁹, Hannah Huckson ⁹⁹, Jonathan Lewis ⁹⁹, Sarah Lowdon ⁹⁹, Cassandra S Malone ⁹⁹, Florence Munemo ⁹⁹, Manasa Mutingwende ⁹⁹, Roberto Nicodemi ⁹⁹, Olga Podplomyk ⁹⁹, Thomas Somassa ⁹⁹, Dr Andrew Beggs ¹⁰⁰, Dr Alex Richter ¹⁰⁰, Claire Cormie ¹⁰², Joana Dias ¹⁰², Sally Forrest ¹⁰², Dr Ellen E Higginson ¹⁰², Mailis Maes ¹⁰², Jamie Young ¹⁰², Dr Rose K Davidson ¹⁰³, Kathryn A Jackson ¹⁰⁷, Dr Lance Turtle ¹⁰⁷, Dr Alexander J Keeley ¹⁰⁹, Prof Jonathan Ball ¹¹³, Timothy Byaruhanga ¹¹³, Dr

Joseph G Chappell ¹¹³, Jayasree Dey ¹¹³, Jack D Hill ¹¹³, Emily J Park ¹¹³, Arezou Fanaie ¹¹⁴, Rachel A Hilson ¹¹⁴, Geraldine Yaze ¹¹⁴ and Stephanie Lo ¹¹⁶

Sequencing and analysis:

Safiah Afifi ¹⁰, Robert Beer ¹⁰, Joshua Maksimovic ¹⁰, Kathryn McCluggage ¹⁰, Karla Spellman ¹⁰, Catherine Bresner ¹¹, William Fuller ¹¹, Dr Angela Marchbank ¹¹, Trudy Workman ¹¹, Dr Ekaterina Shelest ^{13,81}, Dr Johnny Debebe ¹⁸, Dr Fei Sang ¹⁸, Dr Marina Escalera Zamudio ²³, Dr Sarah Francois ²³, Bernardo Gutierrez ²³, Dr Tetyana I Vasylyeva ²³, Dr Flavia Flaviani ³¹, Dr Manon Ragonnet-Cronin ³⁹, Dr Katherine L Smollett ⁴², Alice Broos ⁵³, Daniel Mair ⁵³, Jenna Nichols ⁵³, Dr Kyriaki Nomikou ⁵³, Dr Lily Tong ⁵³, Ioulia Tsatsani ⁵³, Prof Sarah O'Brien ⁵⁴, Prof Steven Rushton ⁵⁴, Dr Roy Sanderson ⁵⁴, Dr Jon Perkins ⁵⁵, Seb Cotton ⁵⁶, Abbie Gallagher ⁵⁶, Dr Elias Allara ^{70,102}, Clare Pearson ^{70,102}, Dr David Bibby ⁷², Dr Gavin Dabrera ⁷², Dr Nicholas Ellaby ⁷², Dr Eileen Gallagher ⁷², Dr Jonathan Hubb ⁷², Dr Angie Lackenby ⁷², Dr David Lee ⁷², Nikos Manesis ⁷², Dr Tamyo Mbisa ⁷², Dr Steven Platt ⁷², Katherine A Twohig ⁷², Dr Mari Morgan ⁷⁴, Alp Aydin ⁷⁵, David J Baker ⁷⁵, Dr Ebenezer Foster-Nyarko ⁷⁵, Dr Sophie J Prosolek ⁷⁵, Steven Rudder ⁷⁵, Chris Baxter ⁷⁷, Sílvia F Carvalho ⁷⁷, Dr Deborah Lavin ⁷⁷, Dr Arun Mariappan ⁷⁷, Dr Clara Radulescu ⁷⁷, Dr Aditi Singh ⁷⁷, Miao Tang ⁷⁷, Helen Morcrette ⁷⁹, Nadua Bayzid ⁹⁶, Marius Cotic ⁹⁶, Dr Carlos E Balcazar ¹⁰⁴, Dr Michael D Gallagher ¹⁰⁴, Dr Daniel Maloney ¹⁰⁴, Thomas D Stanton ¹⁰⁴, Dr Kathleen A Williamson ¹⁰⁴, Dr Robin Manley ¹⁰⁵, Michelle L Michelsen ¹⁰⁵, Dr Christine M Sambles ¹⁰⁵, Dr David J Studholme ¹⁰⁵, Joanna Warwick-Dugdale ¹⁰⁵, Richard Eccles ¹⁰⁷, Matthew Gemmell ¹⁰⁷, Dr Richard Gregory ¹⁰⁷, Dr Margaret Hughes ¹⁰⁷, Charlotte Nelson ¹⁰⁷, Dr Lucille Rainbow ¹⁰⁷, Dr Edith E Vamos ¹⁰⁷, Hermione J Webster ¹⁰⁷, Dr Mark Whitehead ¹⁰⁷, Claudia Wierzbicki ¹⁰⁷, Dr Adrienn Angyal ¹⁰⁹, Dr Luke R Green ¹⁰⁹, Dr Max Whiteley ¹⁰⁹, Emma Betteridge ¹¹⁶, Dr Iraad F Bronner ¹¹⁶, Ben W Farr ¹¹⁶, Scott Goodwin ¹¹⁶, Dr Stefanie V Lensing ¹¹⁶, Shane A McCarthy ^{116,102}, Dr Michael A Quail ¹¹⁶, Diana Rajan ¹¹⁶, Dr Nicholas M Redshaw ¹¹⁶, Carol Scott ¹¹⁶, Lesley Shirley ¹¹⁶ and Scott AJ Thurston ¹¹⁶

Software and analysis tools:

Dr Will Rowe ⁴³, Amy Gaskin ⁷⁴, Dr Thanh Le-Viet ⁷⁵, James Bonfield ¹¹⁶, Jennifer Liddle ¹¹⁶ and Andrew Whitwham ¹¹⁶

1 Barking, Havering and Redbridge University Hospitals NHS Trust, **2** Barts Health NHS Trust, **3** Belfast Health & Social Care Trust, **4** Betsi Cadwaladr University Health Board, **5** Big Data Institute, Nuffield Department of Medicine, University of Oxford, **6** Blackpool Teaching Hospitals NHS Foundation Trust, **7** Bournemouth University, **8** Cambridge Stem Cell Institute, University of Cambridge, **9** Cambridge University Hospitals NHS Foundation Trust, **10** Cardiff and Vale University Health Board, **11** Cardiff University, **12** Centre for Clinical Infection and Diagnostics Research, Department of Infectious Diseases, Guy's and St Thomas' NHS Foundation Trust, **13** Centre for Enzyme Innovation, University of Portsmouth, **14** Centre for Genomic Pathogen Surveillance, University of Oxford, **15** Clinical Microbiology Department, Queens Medical Centre, Nottingham University Hospitals NHS Trust, **16** Clinical Microbiology, University Hospitals of Leicester NHS Trust, **17** County Durham and Darlington NHS Foundation Trust, **18** Deep Seq, School of Life Sciences, Queens Medical Centre, University of Nottingham, **19** Department of Infectious Diseases and Microbiology, Cambridge University Hospitals NHS Foundation Trust, **20** Department of Medicine, University of Cambridge, **21** Department of Microbiology, Kettering General Hospital, **22** Department of Microbiology, South West London Pathology, **23** Department of Zoology, University of Oxford, **24** Division of Virology, Department of Pathology, University of Cambridge, **25** East Kent Hospitals University NHS Foundation Trust, **26** East Suffolk and North Essex NHS Foundation Trust, **27** East Sussex Healthcare NHS Trust, **28** Gateshead Health NHS Foundation Trust, **29** Great Ormond Street Hospital for Children NHS Foundation Trust, **30** Great Ormond Street Institute of Child Health (GOS ICH), University College London (UCL), **31** Guy's and St. Thomas' Biomedical Research Centre, **32** Guy's and St. Thomas' NHS Foundation Trust, **33** Hampshire Hospitals NHS Foundation Trust, **34** Health Services Laboratories, **35** Heartlands Hospital, Birmingham, **36** Hub for Biotechnology in the Built Environment, Northumbria University, **37** Hull University Teaching Hospitals NHS Trust, **38** Imperial College Healthcare NHS Trust, **39** Imperial College London, **40** Infection Care Group, St George's University Hospitals NHS Foundation Trust, **41** Institute for Infection and Immunity, St George's University of London, **42** Institute of Biodiversity,

Animal Health & Comparative Medicine, **43** Institute of Microbiology and Infection, University of Birmingham, **44** Isle of Wight NHS Trust, **45** King's College Hospital NHS Foundation Trust, **46** King's College London, **47** Liverpool Clinical Laboratories, **48** Maidstone and Tunbridge Wells NHS Trust, **49** Manchester University NHS Foundation Trust, **50** Microbiology Department, Buckinghamshire Healthcare NHS Trust, **51** Microbiology, Royal Oldham Hospital, **52** MRC Biostatistics Unit, University of Cambridge, **53** MRC-University of Glasgow Centre for Virus Research, **54** Newcastle University, **55** NHS Greater Glasgow and Clyde, **56** NHS Lothian, **57** NIHR Health Protection Research Unit in HCAI and AMR, Imperial College London, **58** Norfolk and Norwich University Hospitals NHS Foundation Trust, **59** Norfolk County Council, **60** North Cumbria Integrated Care NHS Foundation Trust, **61** North Middlesex University Hospital NHS Trust, **62** North Tees and Hartlepool NHS Foundation Trust, **63** North West London Pathology, **64** Northumbria Healthcare NHS Foundation Trust, **65** Northumbria University, **66** NU-OMICS, Northumbria University, **67** Path Links, Northern Lincolnshire and Goole NHS Foundation Trust, **68** Portsmouth Hospitals University NHS Trust, **69** Public Health Agency, Northern Ireland, **70** Public Health England, **71** Public Health England, Cambridge, **72** Public Health England, Colindale, **73** Public Health Scotland, **74** Public Health Wales, **75** Quadram Institute Bioscience, **76** Queen Elizabeth Hospital, Birmingham, **77** Queen's University Belfast, **78** Royal Brompton and Harefield Hospitals, **79** Royal Devon and Exeter NHS Foundation Trust, **80** Royal Free London NHS Foundation Trust, **81** School of Biological Sciences, University of Portsmouth, **82** School of Health Sciences, University of Southampton, **83** School of Medicine, University of Southampton, **84** School of Pharmacy & Biomedical Sciences, University of Portsmouth, **85** Sheffield Teaching Hospitals NHS Foundation Trust, **86** South Tees Hospitals NHS Foundation Trust, **87** Southwest Pathology Services, **88** Swansea University, **89** The Newcastle upon Tyne Hospitals NHS Foundation Trust, **90** The Queen Elizabeth Hospital King's Lynn NHS Foundation Trust, **91** The Royal Marsden NHS Foundation Trust, **92** The Royal Wolverhampton NHS Trust, **93** Turnkey Laboratory, University of Birmingham, **94** University College London Division of Infection and Immunity, **95** University College London Hospital Advanced Pathogen Diagnostics Unit, **96** University College London Hospitals NHS Foundation Trust, **97** University Hospital Southampton NHS Foundation Trust, **98** University Hospitals Dorset NHS Foundation Trust, **99** University Hospitals Sussex NHS Foundation Trust, **100** University of Birmingham, **101** University of Brighton, **102** University of Cambridge, **103** University of East Anglia, **104** University of Edinburgh, **105** University of Exeter, **106** University of Kent, **107** University of Liverpool, **108** University of Oxford, **109** University of Sheffield, **110** University of Southampton, **111** University of St Andrews, **112** Viapath, Guy's and St Thomas' NHS Foundation Trust, and King's College Hospital NHS Foundation Trust, **113** Virology, School of Life Sciences, Queens Medical Centre, University of Nottingham, **114** Watford General Hospital, **115** Wellcome Centre for Human Genetics, Nuffield Department of Medicine, University of Oxford, **116** Wellcome Sanger Institute, **117** West of Scotland Specialist Virology Centre, NHS Greater Glasgow and Clyde, **118** Whittington Health NHS Trust