Mapping the antigenic diversification of SARS-CoV-2

Karlijn van der Straten[1-3], Denise Guerra[1-2], Marit J. van Gils[1-2], Ilja Bontjer[1-2],

Tom G. Caniels[1-2], Hugo D.G. van Willigen[1-2], Elke Wynberg[4], Meliawati Poniman[1-2], Judith A.

Burger[1-2], Joey H. Bouhuijs[1-2], Jacqueline van Rijswijk[1-2], Wouter Olijhoek[1-2], Marinus H.

Liesdek[1-3], A. H. Ayesha Lavell[2,5], Brent Appelman[6], Jonne J. Sikkens[2,5], Marije K. Bomers[2,5],

Alvin X. Han[1-2], Brooke E. Nichols[1-2,7], Maria Prins[3,4], Harry Vennema[8], Chantal Reusken[8],

Menno D. de Jong[1-2], Godelieve J. de Bree[3], Colin A. Russell[1-2*], Dirk Eggink[1-2,8*], Rogier W.

Sanders[1-2,9*]

[1] Amsterdam UMC location University of Amsterdam, Department of Medical Microbiology

and Infection prevention, Laboratory of Experimental Virology, Meibergdreef 9, 1105 AZ

Amsterdam, The Netherlands

[2] Amsterdam institute for Infection and Immunity, Infectious diseases, Amsterdam, The

Netherlands

[3] Amsterdam UMC location University of Amsterdam, Department of Internal Medicine,

Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

[4] Department of Infectious Diseases, Public Health Service of Amsterdam, GGD, 1018 WT

Amsterdam, The Netherlands

[5] Amsterdam UMC location VU University Amsterdam, Department of Internal Medicine,

Boelelaan 1117, 1081 HV Amsterdam, The Netherlands

[6] Amsterdam UMC location University of Amsterdam, Center for Experimental and

Molecular Medicine, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

[7] Department of Global Health, Boston University School of Public Health, Boston, USA

[8] Centre for Infectious Disease Control, National Institute for Public Health and the

Environment, 3721 MA Bilthoven, The Netherlands

[9] Department of Microbiology and Immunology, Weill Medical College of Cornell University,

New York, USA

* correspondence to:

c.a.russell@amsterdamumc.nl, dirk.eggink@rivm.nl, r.w.sanders@amsterdamumc.nl

Summary

Large-scale vaccination campaigns have prevented countless hospitalizations and deaths
due to COVID-19. However, the emergence of SARS-CoV-2 variants that escape from
immunity challenges the effectiveness of current vaccines. Given this continuing evolution,
an important question is when and how to update SARS-CoV-2 vaccines to antigenically
match circulating variants, similar to seasonal influenza viruses where antigenic drift
necessitates periodic vaccine updates. Here, we studied SARS-CoV-2 antigenic drift by
assessing neutralizing activity against variants-of-concern (VOCs) of a unique set of sera
from patients infected with a range of VOCs. Infections with D614G or Alpha strains induced
the broadest immunity, while individuals infected with other VOCs had more strain-specific
responses. Omicron BA.1 and BA.2 were substantially resistant to neutralization by sera
elicited by all other variants. Antigenic cartography revealed that Omicron BA.1 and BA.2 are
antigenically most distinct from D614G, associated with immune escape and likely requiring
vaccine updates to ensure vaccine effectiveness.

Keywords: SARS-CoV-2, variants of concern (VOCs), convalescent, vaccination,
neutralization, antibodies, antigenic cartography

Main text

The COVID-19 pandemic, caused by the SARS-CoV-2 virus, represents an enormous threat to

human health and a burden to healthcare systems and economies worldwide. The

unprecedented rapid development of efficacious vaccines fuelled hope of curtailing this

pandemic and permitting a return to a society without societal restrictions. However,

genetic drift of SARS-CoV-2 resulted in the emergence of multiple variants of concern (VOCs)

with a higher transmissibility compared to the ancestral strain, and that challenge the

effectiveness of public health measures, vaccines and/or therapeutics (World Health

Organization, 2021). Based on this definition, the WHO designated the Alpha (Pango lineage

B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2) and Omicron (B.1.1.529, sublineages

BA.1 and BA.2) variants as VOCs. The Alpha, Beta, Gamma and Delta VOCs have

approximately 7 to 12 mutations in the Spike protein (S), while Omicron BA.1 with 34

mutations, of which 3 deletions, and BA.2 with 28 mutations, differ substantially more from

the ancestral strain (Figure 1A)(World Health Organization, 2021). Approximately half of

Omicron's S mutations are located in the Receptor Binding Domain (RBD) and eight

mutations in the N-terminal domain (NTD), the two most important antigenic site of S.

Indeed, sera from COVID-19 patients infected with the ancestral strain and sera from

vaccinees show up to 7-fold and 4-fold reductions in neutralization activity against Beta and

Gamma, while 20 to 40-fold reductions are observed against Omicron BA.1 (Caniels et al.,

2021; Garcia-Beltran et al., 2021; van Gils et al., 2022; Wilhelm et al., 2021).

69   However, the precise antigenic relationships between these VOCs are only starting

to become clear. Understanding the differences between the serological antibody responses

elicited by these variants is important to assess the risk of re-infections after natural

infection and breakthrough infections after vaccination. For seasonal influenza viruses, this

type of antigenic data is combined with virus genetic and epidemiological data to quantify

the evolution of the virus and guide annual updates of the seasonal influenza virus vaccines.

Antigenic cartography can be used to visualize antigenic relationships between viral variants

(Fonville et al., 2014; Smith et al., 2004) and is routinely used in influenza virus vaccines

strain selection. Until recently, antigenic cartography for SARS-CoV-2 has only been applied

to cohorts of COVID-19 patients with uncertainty about their history of previous SARS-CoV-2

exposure and COVID-19 vaccinations, and without the usage of Omicron infected human

data (Liu et al., 2021; Mykytyn et al., 2022; Wilks et al., 2022). Here, we studied the (cross-
)neutralizing antibody responses in sera from a well-defined population of convalescent
individuals with a sequence confirmed, or high likelihood of, primary infection by the
D614G, Alpha, Beta, Gamma, Delta or Omicron BA.1 or BA.2 variants and used this data as
input for antigenic cartography to map the antigenic evolution of SARS-CoV-2.

We collected and analysed a unique set of serum samples from 66 COVID-19 patients with a
PCR-confirmed primary SARS-CoV-2 infection who did not receive any COVID-19
vaccinations. Blood was drawn 3 to 11 weeks after symptom onset (median 40 days, range
24 to 75 days), which corresponds with the peak of the antibody response (Table 1 and
Table S1)(Long et al., 2020). In total, n=20 D614G, n=11 Alpha, n=8 Beta, n=4 Gamma, n=11
Delta, n=8 Omicron BA.1 and n=4 Omicron BA.2 infected participants were included. Of
these participants, 39 had a sequence-confirmed VOC infection. The other 27 participants
met our inclusion criteria of a high likelihood of VOC infection (see STAR Methods section
and Table S1), of which 20 participants were assumed to be infected with the D614G strain
as they were sampled before the emergence of any VOC in the Netherlands, but after
D614G became dominant (Korber et al., 2020).

We assessed the neutralizing capacity of the convalescent sera in a lentiviral-based
pseudovirus neutralization assay against the D614G strain, the Alpha, two Beta, the Gamma,
Delta and Omicron BA.1 and BA.2 variants (Figure 1A). The two Beta subvariants differ from
each other in the NTD, where one Beta subvariant (L242H, R246I) is based on a very early
available sequence while the other (Δ242-244) is retrospectively more representative for
the predominant circulating strains.

The highest neutralization titres were generally measured against the homologous
virus, as might be expected (Figure 1A and Figure S1A). Only the Beta infected participants
showed higher cross-neutralization titres against the Gamma variant compared to
homologous neutralization, which is in line with other research(Wilks *et al.*, 2022). This
might be explained by the shared RBD mutations, as the RBDs of these variants only differ
by one amino acid (K417N in Beta *versus* K417T in Gamma). Our analyses suffer somewhat
from a disbalance in hospitalized versus non-hospitalized patients between the different
VOC groups(Table 1). However, when comparing only non-hospitalized patients which
generally have lower antibody levels compared to hospitalized patients, patients infected

with the Alpha variant showed the strongest homologous neutralization (1881 IU/mL, range
1658 to 2103 IU/mL), followed by individuals infected with the Gamma variant (median of
156 IU/mL, range 22 to 761 IU/mL), the D614G strain (median of 90 IU/mL, range 28 to 237
IU/mL) the Delta variant (median of 85 IU/mL, range 10 to 1635 IU/mL), the Omicron BA.2
(median of 64 IU/mL, range 10 to 95 IU/mL) and the Omicron BA.1 variant (median of 23
IU/mL, range 10 to 90 IU/mL) (Figure S1A). By contrast, none of the Beta infected
participants showed substantial homologous neutralization against either Beta subvariants
(Figure 1A, Figure S2).

Overall, the VOCs differed in their capacity to induce cross-neutralizing antibodies.
Individuals infected with the Alpha variant induced the broadest response, followed by
D614G strain-infected, Gamma-infected and Delta-infected patients (Figure 1A, Figure 1B
and Figure S1B), though there was substantial heterogeneity within all groups. Notably,
none of the patients infected with the Beta, Omicron BA.1 or Omicron BA.2 variants showed
substantial cross-neutralization activity.

Reductions in neutralizing activity against the two Omicron variants were substantial
in all groups (Figure 1A and 1B). Omicron neutralization dropped below the limit of
detection (10 IU/mL or an ID_{50} of 100) in 44/66 of the studied individuals for BA.1 and 37/66
for BA.2. The median fold-reduction of Omicron BA.1 neutralization versus homologous
neutralization was 9-fold (range 1 to 93-fold) when considering all patients, 10-fold (range 3
to 93-fold) for patients infected with a D614G strain, 52-fold (range 11 to 89-fold) for Alpha,
6-fold (range 1 to 22-fold) for Gamma, and 6-fold (range 1 to 51-fold) for Delta infected
patients. The median fold-reduction of Omicron BA.2 neutralization versus homologous
neutralization was 5.4-fold (range -3.7 to 134-fold) when considering all patients, 8-fold
(range 3 to 47-fold) for patients infected with D614G strain, 68-fold (range 18 to 134-fold)
for Alpha, 6-fold (range 1 to 22-fold) for Gamma, and 6-fold (range 1 to 60-fold) for Delta
infected patients.

To explore the antigenic relationships between the VOCs, we used the neutralization
data to construct a SARS-CoV-2 antigenic map (Figure 2A). In this map, homologous sera
tend to cluster around the infecting strain, reflecting that homologous neutralization is
dominant. The D614G and Alpha viruses cluster tightly together in the centre of the map,
while the Beta (L242H, R246I), Gamma, and Delta variants all lie within 2 antigenic units (1
unit = 2-fold change in neutralization titre) of the D614G strain suggesting a high degree of

antigenic similarity. For influenza viruses, variants are considered to be antigenically similar

in case of antigenic distances below 3 antigenic units, i.e. an 8-fold change in neutralization

titre, and different when above this threshold(Barr et al., 2014; Prevention, 2021). By

analogy, the D614G, Alpha, Beta (L242H, R246I), Gamma and Delta variants belong to one

antigenic cluster. Interestingly, the Beta (Δ242-244) subvariant is antigenically more distinct

from the D614G strain compared to Beta (L242H, R246I) (e.g. 3 to 4 units), implying that the

deletion at region 242-244 has a substantial effect on antigenicity and illustrates the

importance of the NTD as target of neutralizing antibodies and/or in modulating antigenicity

of other domains by allosteric means. The distance between the main antigenic cluster and

Omicron BA.1 and BA.2 variants is more than 4 antigenic units (>16-fold change in

neutralization) implying that Omicron BA.1 and BA.2 are the antigenically most distinct

SARS-CoV-2 variants (Figure 2A). One caveat is that it is unclear whether 2-fold changes in

pseudovirus neutralization titres are directly comparable to 2-fold changes in

hemagglutination inhibition assay titres used to define different antigenic clusters of

influenza viruses. However, the change in neutralization between Omicron BA.1 and BA.2

and other variants of SARS-CoV-2, including the D614G strain, is striking.

We next used neutralization data from sera of 109 COVID-19 naïve vaccinees

receiving either two Moderna (mRNA-1273, n=30), Pfizer/BioNTech (BNT162b2, n=49), or

AstraZeneca (AZD1222, n=30) vaccines, which are all based on the ancestral S sequence to

generate a second antigenic map (van Gils *et al.*, 2022). This map (Figure 2B) agreed well

with the infectee maps (Figure 2A), and corroborated that Omicron BA.1 represents a

distinct antigenic variant from viruses currently included in vaccines. Interestingly, while the

distributions of sera from recipients of different vaccines overlap, there is a skew of sera of

mRNA-1273 vaccinees towards Omicron BA.1, suggesting small differences in antigen

stimulation among vaccine formulations considered here.

We have started to define the antigenic SARS-CoV-2 landscape after two years of

antigenic drift, which should inform risk assessment of re-infections as well as strain

selection for COVID-19 vaccine updates. We can draw several conclusions. First,

homologous neutralization was usually stronger than heterologous neutralization. Second,

heterologous responses were broadest and most potent in individuals infected with Alpha

and D614G strains, while infection with Delta resulted in narrow-specificity responses. In

addition, the individuals infected with the Beta and Omicron BA.1 variant, and to lesser
extent Omicron BA.2 infected individuals, developed weak neutralizing responses against
any VOC, including the homologous strains, suggesting that the S proteins of Beta and both
Omicron variants are less immunogenic compared to the S of other VOCs. Interestingly, the
weak homologous and cross-neutralization levels of Beta variant infected individuals are in
contrast with the higher titres found by others(Cele et al., 2022; Liu *et al.*, 2021; Rossler et
al., 2022; Wilks *et al.*, 2022). It is unlikely that this weak homologous neutralization is caused
by a sequence mismatch between the strains causing the infection and the sequence used in
our pseudoviruses, as both Beta subvariant pseudoviruses escape homologous
neutralization of Beta sera, and the reduction of cross-neutralization of sera elicited by
other VOCs against the Beta variants are in line with previous studies (Cele *et al.*, 2022; Liu
et al., 2021; Rossler *et al.*, 2022; Wilks *et al.*, 2022). One contributing factor for these low
(cross-)neutralization titres by the Beta variant includes cohort specific differences between
studies, which is, however, hard to verify due to limited patient characteristic and
demographic information available in publications. The weak homologous and cross-
neutralization of Omicron BA.1 variant infected individuals is in line with other pre-print
data(Mykytyn *et al.*, 2022). Third, the D614G and Alpha strains are at the centre of our
antigenic map, which supports the use of the current COVID-19 vaccines based on the
ancestral strain, in case of circulation of the Alpha, Beta (L242H, R246I), Gamma and Delta
variants. Our data suggest that updated vaccines based on the Beta (L242H, R246I) or Delta
variants would not have been appreciably more effective than the ancestral virus-based
vaccines. However, the substantial reduction of neutralization in all groups against the Beta
Δ242-244, but especially against the Omicron variants indicates a high risk of re-infections
and vaccine breakthrough cases when exposed to these VOCs. The long antigenic distance
between Omicron variants and the preceding variants in the antigenic map indicates that
the current high rates of Omicron infections are at least partially associated with immune
escape and that a vaccine update is required. While finishing this study, several other efforts
to antigenically characterize VOCs became available (Mykytyn *et al.*, 2022; Wilks *et al.*,
2022). Our antigenic cartography is largely in accordance with these other studies.

As in the case of seasonal influenza viruses, the prospect of SARS-CoV-2 becoming an
endemic virus with recurring outbreaks implies the need for surveillance of antigenic drift
and possibly yearly administration of updated vaccines, especially for individuals at risk for

severe COVID-19. Antigenic cartography efforts such as those presented here, can inform
future vaccine updates.

Acknowledgements
We thank all public health services (GGD) in the Netherlands for their help in contacting
participants. We are also thankful to the study personnel and the participants of the COSCA,
RECoVERED and the S3-study for their contribution to this research.

Author Contributions
Conceptualization, K.vdS., D.G., M.J.vG., C.R., C.A.R., D.E., R.W.S.,
Methodology, K.vdS., D.G., M.J.vG., C.A.R., D.E., R.W.S.,
Validation, K.vdS., D.G., M.J.vG., I.B.,
Formal Analysis, K.vdS., A.X.H., B.E.N., C.A.R.,
Investigation, I.B., M.Po., J.A.B., J.H.B., J.vR., W.O.,
Resources, K.vdS., I.B., H.D.G.vW., E.W., M.H.L., A.H.A.L., B.A., J.L.S., M.K.B., M.Pr., H.V.,
C.R., M.D.dJ, G.J.dB,
Writing – original draft, K.vdS., D.G., A.X.H., C.A.R., D.E., R.W.S.,
Writing – review & editing, M.J.vG, I.B., T.G.C., H.D.G.vW., E.W., M.Po., J.A.B., J.H.B.,
J.vR., W.O., M.H.L., A.H.A.L., B.A., J.J.S., M.K.B., B.E.N., M.Pr., H.V., C.R., M.D.dJ.,
G.J.dB.
Visualization, K.vdS.,T.G.C., A.X.H., C.A.R., R.W.S.,
Supervision, C.A.R., D.E., R.W.S.,
Project administration, K.vdS., D.G.,
Funding acquisition, J.J.S., M.K.B., A.X.H., M.P., M.D.dJ., G.J.dB., C.A.R., R.W.S.,

Conflicts of interest
None of the authors have conflicts of interest related to this research.

Funding
R.W.S. and C.A.R. are recipients of Vici grants from the Netherlands Organization for
Scientific Research (NWO no. 91818627 for R.W.S.). C.A.R. and A.X.H. are also supported by
an ERC Consolidator Award. This work was supported by the NWO ZonMw grant agreement
no. 101500062010002 to M.D.dJ., and 10430072110003 to G.J. de Bree and the Public Health
Service of Amsterdam Research & Development grant number 21-14 to M. Prins

(RECoVERED). J.J.S. and M.K.B. are recipients of the NWO grant agreement no.
10430022010023 and 10430022010030.

Figure legends

Figure 1. SARS-CoV-2 genetic diversity and neutralization. A. Molecular models of SARS-
CoV-2 S, highlighting the locations of mutations in the D614G strain (blue), Alpha (green),
Beta (yellow), Gamma (orange), Delta (red), Omicron BA.1 (magenta) and Omicron BA.2
(pink) variants. Midpoint neutralization titres against the VOCs in International Units per mL
(IU/mL). The individuals are grouped per VOC and plotted accordingly. Median
neutralization titres are highlighted while the individual points are depicted with higher
transparency. The light grey bar (10 IU/ml) indicates the neutralization cut-off for all strains
except Omicron (cut-off 2 IU/mL, dark grey bar). Non-hospitalized patients are indicated
with dots and hospitalized patients with triangles. The individuals that were infected with an
Alpha strain that also included the E484K mutation are indicated in green squares. The two
individuals in the Omicron BA.1 group that may have been infected with BA.2 instead of
BA.1 are indicated in magenta diamonds (see also Table S1). The homologous neutralization
is highlighted using a light blue bar. The Wilcoxon signed rank test with Benjamini Hochberg
correction was used to compare cross-neutralization titres with the homologous
neutralization (see Table S3A for exact p-values). Only statistically significant differences are
indicated. * = $p < 0.005$, ** = $p < 0.01$, **** = $p < 0.0001$. **B.** Spider plot of the median
neutralization titre (IU/mL) of each group against all VOCs. A cut-off of 10 IU/mL is used for
all strains.

Figure 2. SARS-CoV-2 antigenic cartography. **A.** Antigenic map of SARS-CoV-2 VOCs
based on convalescent SARS-CoV-2 infection sera. SARS-CoV-2 variants are shown as circles
and sera are indicated as squares. Each square corresponds to sera of one individual and is
coloured by the infecting SARS-CoV-2 variant. Both axes of the map are antigenic distance
and each grid square (1 antigenic unit) represents a two-fold change in neutralization titre.
The distance between points in the map can be interpreted as a measure of antigenic
similarity, where the points more closely together show higher cross-neutralization and are
therefore antigenically more similar. The left panel included both Beta subvariants used in
this study. The right panel is without the Beta (Δ242-244) subvariant **B.** Antigenic map of

SARS-CoV-2 VOCs based on post-vaccination sera from individuals without prior SARS-CoV-2

infections. Each serum is coloured by the vaccine that individual received.

Table 1. Sociodemographics and clinical characteristics. A summary of the convalescent

SARS-CoV-2 patients included in this study. Table S1 contains a more comprehensive

overview   per   individual.

 STAR METHODS

 KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
DMEM media	Gibco	Cat#: 11966025
Opti-MEM I reduced serum media	Gibco	Cat#: 15392402
Phosphate-buffered saline (PBS)	Gibco	Cat#: 15326239
Bacterial and virus strains		
Chemically competent DH5α *Escherichia coli*	Thermo Fisher Scientific	Cat#: 12879416
Biological samples		
Human sera, convalescent SARS-Cov-2	This study	N/A
Human sera, post-COVID-19 vaccination	(van Gils *et al.*, 2022)	N/A
Chemicals, peptides, and recombinant proteins		
FastDigest SacI	Thermo Fisher Scientific	Cat#: 10324720
FastDigest ApaI	Thermo Fisher Scientific	Cat#: 10450280
Polyethylenimine hydrochloride (PEI) MAX	PolySciences	Cat#: 24765-1
Trypsin-EDTA	Gibco	Cat#: 25-200-056
GlutaMAX	Gibco	Cat#: 35050061
Glycylglycine, 99+%	Acros Organics/Thermo Scientific	Cat#: 120141000
Magnesium sulfate heptahydrate, 99.5%, for analysis, Thermo Scientific™	Acros Organics/ Thermo Scientific	Cat#: AC21311500
EGTA (ethylene glycol bis(2-aminoethyl ether)-N,N,N',N'-tetraacetic acid) ≥97%, Ultra Pure Grade	VWR	Cat#: 0732-100G
Triton™ X-100 (Electrophoresis), Fisher BioReagents™	Fisher BioReagents	Cat#: BP151-500
Poly-L-Lysine Hydrobromide	Sigma-Aldrich	Cat#: P1399
Critical commercial assays		
Gibson Assembly	New England BioLabs	E5510S
QuikChange Site-Directed Mutagenesis Kit	Agilent Technologies	Cat#: 200523
Experimental models: Cell lines		
HEK293T cells	American Type Culture Collection	CRL-11268
HEK-293T-hACE2	(Schmidt et al., 2020)	RRID:CVCL_A7UK
Oligonucleotides		
SARS-CoV-2 D614G spike gene fragment	Integrated DNA Technologies	GenBank:MT449663.1 Mutations in this paper
SARS-CoV-2 Alpha spike gene fragment	Integrated DNA Technologies	GenBank:MT449663.1 Mutations in this paper
SARS-CoV-2 Beta spike gene fragment	Integrated DNA Technologies	GenBank:MT449663.1 Mutations in this paper

SARS-CoV-2 Gamma spike gene fragment	Integrated DNA Technologies	GenBank:MT449663.1 Mutations in this paper
SARS-CoV-2 Delta spike gene fragment	Integrated DNA Technologies	GenBank:MT449663.1 Mutations in this paper
SARS-CoV-2 Omicron BA.1 spike gene fragment	Integrated DNA Technologies	GenBank:MT449663.1 Mutations in this paper
SARS-CoV-2 Omicron BA.2 spike gene fragments	Integrated DNA Technologies	GenBank:MT449663.1 Mutations in this paper
Recombinant DNA		
pCR3 SARS-CoV-2–S$_{\Delta19}$ expression plasmid	GenBank	ID: MT449663.1
pHIV-1$_{NL43}$ ΔEnv-NanoLuc reporter virus plasmid	(Schmidt *et al.*, 2020)	
Software and algorithms		
ACMACS antigenic cartography software	https://acmacs-web.antigenic-cartography.org	N/A
GraphPad Prism 8.3.0	GraphPad	RRID:SCR_002798; http://www.graphpad.com/
Microsoft Excel	Microsoft	RRID:SCR_016137; https://www.microsoft.com/en-gb/
Other		
Nano-Glo Luciferase Assay System	Promega	Cat#: N1130
GloMax system	Turner BioSystems	Cat#: 9101-002

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will

be fulfilled by the lead contact, Rogier W. Sanders (r.w.sanders@amsterdamumc.nl)

Materials availability

This study did not generate new unique reagents.

Data and code availability

Neutralisation and patient characteristic data have been deposited as supplementary tables

within this manuscript (Table S1 and Table S2) and are publicly available as of the date of

publication. Other accession numbers are listed in the key resources table. This paper does

not report original code. Any additional information required to reanalyse the data reported

in this paper is available from the lead contact upon request.

EXPERIMETNAL MODEL AND SUBJECT DETAILS

Study population

66 adults (aged 18 to 76) with a PCR proven primary SARS-CoV-2 infection were included in

the COSCA-study (NL 73281.018.20) or the RECOVERED study (NL73759.018.20) between

June 2020 and April 2022 at Amsterdam UMC and via the Dutch national SARS-CoV-2

sequence surveillance program as described previously(Grobben et al., 2021; Wynberg et

al., 2021). In short, 3-11 weeks after symptom onset, blood, patient demographics, time

between symptom onset and sampling, and admission status were collected (Table 1 and

Table S1). The diagnostic oropharyngeal swab was available for 39 participants and were

used to determine the SARS-CoV-2 strain causing the infection. The remaining 27 SARS-CoV-

2 infected participants fell within the following inclusion criteria: (1) ≥95% of circulating

strains at time of symptom onset belonged the suspected VOC of infection or (2) ≥75% of

circulating strains at the time of symptom onset belonged the suspected VOC of infection

AND a household member had a concurrent sequence confirmed infection with that

particular VOC. Prevalence data of CoVariants.org and the National Institute for Public

Health and the Environment were used to determine the current prevalence of a VOC

(CoVariants, 2022; Rijksinstituut voor Volksgezondheid en Milieu (RIVM)). Most individuals

of which no sequence confirmation of the infected strain was available, were presumed to

be infected with de D614G variant (n=20) as they were sampled before the emergence of

any VOC in the Netherlands and after D614G became predominant in the

Netherlands(Korber *et al.*, 2020). More details about the remaining n=7 individuals can be

found in the Table S1. The two Omicron individuals that may have been infected by either

BA.1 or BA.2 are indicated as diamonds in all graphs. Two of the individuals infected with an

Alpha strain harbouring the E484K mutation are indicated as squares in all graphs.

Neutralization data on COVID-19 naive vaccinee sera were kindly provided by the S3-

study of the Amsterdam UMC, The Netherlands(NL73478.029.20) (van Gils *et al.*, 2022). In

short, post-vaccination sera was obtained approximately four weeks after the second doses

of either Moderna (mRNA-1273), Pfizer/BioNTech (BNT162b2), or AstraZeneca (AZD1222).

Post-vaccination serum after Janssen (Ad26.COV2.S) were excluded from analysis because

they did not have enough non-threshold titres to be included in the map.

All above mentioned studies were conducted at the Amsterdam University Medical

Centres, the Netherlands, and approved by the local ethical committees. All individuals

provided written informed consent before participating.

Pseudovirus design

The D614G strain and the Alpha pseudovirus constructs contained the following mutations: D614G in D614G strain; deletion (Δ) of H69, V70 and Y144, N501Y, A570D, D614G, P681H, T716I, S982A, and D1118H in Alpha. The two Beta subvariants differ from each other in the NTD region 242-246, where one Beta subvariant (L242H, R246I) is based on a very early available sequence while the other (Δ242-244) is retrospectively more representative for the predominant circulating strains. These two Beta pseudovirus constructs contain therefore the following mutations: L18F, D80A, D215G, L242H, R246I, K417N, E484K, N501Y, D614G, and A701V in Beta (L242H, R246I); L18F, D80A, D215G, Δ242-244, K417N, E484K, N501Y, D614G, and A701V in Beta (Δ242-244). Only the D614G infected individuals showed statistically significant reduced neutralization against the Beta (Δ242-244) subvariant compared to the Beta (L242H, R246I) subvariant (Figure S2). The Gamma pseudovirus constructs contained the following mutations: L18F, T20N, P26S, D138Y, R190S, K417T, E484K, N501Y, D614G, H655Y, and T1027I in Gamma; This Gamma pseudovirus construct differs from the predominant strain in that it lacks a V1176F back bone mutation. However, it is not likely that this mutation, positioned at the S2 domain of the S, will affect escape of neutralization substantially. The Delta and Omicron BA.1 and BA.2 pseudovirus constructs contained the following mutations: T19R, G142D, E156G, Δ157-158, L452R, T478K, D614G, P681R and D950N in Delta; A67V, Δ69-70, T95I, G142D, Δ143-145, Δ211, L212I, ins214EPE, G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493K, G496S, Q498R, N501Y, Y505H, T547K, D614G, H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H, N969K, L981F in Omicron BA.1; and T19I, L24S, Δ125/127, G142D, V213G, G339D, S371F, S373P, S375F, T376A, D405N, R408S, K417N, N440K, S477N, T478K, E484A, Q493R, Q498R, N501Y, Y505H, D614G, H655Y, N679K, P681H, N764K, D796Y, Q954H, N969K in Omicron BA.2. The Omicron BA.1 strain used here harbors a Q493K mutation, while the predominant Omicron BA.1 harbors a Q493R mutation. This mutation did not impacted neutralization of several monoclonal SARS-CoV-2 antibodies tested (data not shown). The spike constructs were ordered as gBlock gene fragments (Integrated DNA Technologies) and cloned SacI and ApaI in the pCR3 SARS-CoV-2–S$_{\Delta19}$ expression plasmid (GenBank: MT449663.1) using Gibson Assembly (Thermo Fisher Scientific). The pseudovirus constructs were made using the QuikChange Site-Directed Mutagenesis Kit (Agilent Technologies) and verified by using Sanger sequencing. Pseudoviruses were procedures by cotransfecting HEK293T cells

(American Type Culture Collection, CRL-11268) with the pCR3 SARS-CoV-2-$S_{\Delta 19}$ expression

plasmid and the pHIV-1$_{NL43}$ΔEnv-NanoLuc reporter virus plasmid. Transfection takes place in

cell culture medium (DMEM), supplemented with 10% fetal bovine serum, penicillin (100

U/ml), streptomycin (100 g/ml). Medium is refreshed once 6-8 hours after transfection. 48

hours after the transfection, cell supernatants containing the pseudovirus were harvested

and stored at -80 °C until further use.

METHOD DETAILS

SARS-CoV-2 pseudovirus neutralization assay

The pseudovirus neutralization assay was performed as described previously(Caniels *et al.*,

2021). Shortly, HEK293T/ACE2 cells were kindly provided by P. Bieniasz(Schmidt *et al.*, 2020)

were seeded at a density of 20,000 cells per well in a 96-well plate coated with poly-lysine

(50 ug/ml) 1 day before the start of the neutralization assay. The next day, heat-inactivated

sera samples were in triplicate serial diluted in threefold steps, starting at 1:20 dilution to

test for Omicron BA.1 and BA.2 pseudovirus neutralization and 1:100 for all the other

variants. Sera was diluted in cell culture medium (DMEM), supplemented with 10% fetal

bovine serum, penicillin (100 U/ml), streptomycin (100 g/ml), and GlutaMAX (Gibco), mixed

in a 1:1 ratio with pseudovirus, and incubated for 1 hour at 37°C. These mixtures were then

added to the cells in a 1:1 ratio and incubated for 48 hours at 37°C, and lysis buffer was

added. The luciferase activity in cell lysates was measured using the Nano-Glo Luciferase

Assay System (Promega) and GloMax system (Turner BioSystems). Relative luminescence

units were normalized to those from cells infected with SARS-CoV-2 pseudovirus in the

absence of sera. The inhibitory neutralization titres (ID_{50}) were determined as the serum

dilution at which infectivity was inhibited by 50%, using a nonlinear regression curve fit

(GraphPad Prism software version 8.3). The International Standard for anti-SARS-CoV-2

immunoglobulins provided by the WHO(Kristiansen et al., 2021) were used to convert the

ID_{50} values into International Units per milliliters (IU/mL). Samples with IU/mL titres <10

were defined as having undetectable neutralization against the D614G, Alpha, Beta, Gamma

and Delta variant. For Omicron BA.1 and BA.2 neutralization, the start-dilution of 1:20

enables a cut-off of <2 IU/mL for all samples except for some Alpha infected individuals. A

limited amount of sera was available from the Alpha infected individuals, resulting in a start

dilution of 1:100 of n=7 samples against all variants including Omicron BA.1 and BA.2.

Neutralization data points of two Alpha infected individuals against BA.1 and BA.2 were

excluded from Figure 1A because a neutralization titres <10IU/mL (Table S2). This exclusion

did not impact the statistics as written below because a general cut-off of <10IU/mL were

used for neutralization against any variant, including Omicron BA.1 and BA.2.

Antigenic cartography

Antigenic maps were constructed as previously described (Fonville *et al.*, 2014; Smith *et al.*,

2004) using the antigenic cartography software from https://acmacs-web.antigenic-

cartography.org. In brief, this approach to antigenic mapping uses multidimensional scaling

to position antigens (viruses) and sera in a map to represent their antigenic relationships.

The maps here relied on the SARS-CoV-2 post-infection serology data and post-vaccination

serology data shown in Figure 1A and Table S2. The positions of antigens and sera were

optimized in the map to minimise the error between the target distances set by the

observed pairwise virus-serum combinations in the pseudovirus assay described above and

the resulting computationally derived map. Maps were constructed in 2, 3, 4, and 5

dimensions to investigate the dimensionality of the antigenic relationships. Both the

convalescent (Figure 2A) and post-vaccination datasets (Figure 2B) were strongly two-

dimensional with only small improvements in residual mean squared error of the maps as

map dimensionality increased.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data visualization and statistical analyses were performed in GraphPad Prism software

(version 8.3). Spider plots (Figure 1B) were made in Excel 2016. The antigenic maps were

produced using the antigenic cartography software mentioned above. Wilcoxon signed rank

test with Benjamini Hochberg correction was used to compare cross-neutralization titres

with the homologous neutralization (Figure 1A). Mann-Whitney test was used for non-

paired group comparisons (Figure S1). All statistics mentioned here were performed by

using a general neutralization cut-off of 10IU/mL against any variant of SARS-CoV-2.

References:

Barr, I.G., Russell, C., Besselaar, T.G., Cox, N.J., Daniels, R.S., Donis, R., Engelhardt, O.G.,
Grohmann, G., Itamura, S., Kelso, A., et al. (2014). WHO recommendations for the viruses
used in the 2013-2014 Northern Hemisphere influenza vaccine: Epidemiology, antigenic and

genetic characteristics of influenza A(H1N1)pdm09, A(H3N2) and B influenza viruses
collected from October 2012 to January 2013. Vaccine *32*, 4713-4725.
10.1016/j.vaccine.2014.02.014.
Caniels, T.G., Bontjer, I., van der Straten, K., Poniman, M., Burger, J.A., Appelman, B.,
Lavell, A.H.A., Oomen, M., Godeke, G.J., Valle, C., et al. (2021). Emerging SARS-CoV-2
variants of concern evade humoral immune responses from infection and vaccination. Sci
Adv *7*, eabj5365. 10.1126/sciadv.abj5365.
Cele, S., Karim, F., Lustig, G., San, J.E., Hermanus, T., Tegally, H., Snyman, J., Moyo-
Gwete, T., Wilkinson, E., Bernstein, M., et al. (2022). SARS-CoV-2 prolonged infection
during advanced HIV disease evolves extensive immune escape. Cell Host Microbe *30*, 154-
162 e155. 10.1016/j.chom.2022.01.005.
CoVariants (2022). Overview of Variants in Countries. https://covariants.org/per-country.
Fonville, J.M., Wilks, S.H., James, S.L., Fox, A., Ventresca, M., Aban, M., Xue, L., Jones,
T.C., Le, N.M.H., Pham, Q.T., et al. (2014). Antibody landscapes after influenza virus
infection or vaccination. Science *346*, 996-1000. 10.1126/science.1256427.
Garcia-Beltran, W.F., St Denis, K.J., Hoelzemer, A., Lam, E.C., Nitido, A.D., Sheehan, M.L.,
Berrios, C., Ofoman, O., Chang, C.C., Hauser, B.M., et al. (2021). mRNA-based COVID-19
vaccine boosters induce neutralizing immunity against SARS-CoV-2 Omicron variant.
medRxiv, 2021.2012.2014.21267755. 10.1101/2021.12.14.21267755.
Grobben, M., van der Straten, K., Brouwer, P.J., Brinkkemper, M., Maisonnasse, P.,
Dereuddre-Bosquet, N., Appelman, B., Lavell, A.A., van Vught, L.A., Burger, J.A., et al.
(2021). Cross-reactive antibodies after SARS-CoV-2 infection and vaccination. Elife *10*,
2021.2005.2026.21256092. 10.7554/eLife.70330.
Korber, B., Fischer, W.M., Gnanakaran, S., Yoon, H., Theiler, J., Abfalterer, W., Hengartner,
N., Giorgi, E.E., Bhattacharya, T., Foley, B., et al. (2020). Tracking Changes in SARS-CoV-2
Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. Cell *182*, 812-827
e819. 10.1016/j.cell.2020.06.043.
Kristiansen, P.A., Page, M., Bernasconi, V., Mattiuzzo, G., Dull, P., Makar, K., Plotkin, S.,
and Knezevic, I. (2021). WHO International Standard for anti-SARS-CoV-2 immunoglobulin.
Lancet *397*, 1347-1348. 10.1016/S0140-6736(21)00527-4.
Liu, C., Ginn, H.M., Dejnirattisai, W., Supasa, P., Wang, B., Tuekprakhon, A., Nutalai, R.,
Zhou, D., Mentzer, A.J., Zhao, Y., et al. (2021). Reduced neutralization of SARS-CoV-2
B.1.617 by vaccine and convalescent serum. Cell *184*, 4220-4236 e4213.
10.1016/j.cell.2021.06.020.
Long, Q.X., Liu, B.Z., Deng, H.J., Wu, G.C., Deng, K., Chen, Y.K., Liao, P., Qiu, J.F., Lin, Y.,
Cai, X.F., et al. (2020). Antibody responses to SARS-CoV-2 in patients with COVID-19. Nat
Med *26*, 845-848. 10.1038/s41591-020-0897-1.
Mykytyn, A.Z., Rissmann, M., Kok, A., Rosu, M.E., Schipper, D., Breugem, T.I., van den
Doel, P.B., Chandler, F., Bestebroer, T., de Wit, M., et al. (2022). Omicron BA.1 and BA.2
are antigenically distinct SARS-CoV-2 variants. bioRxiv, 2022.2002.2023.481644.
10.1101/2022.02.23.481644.
Prevention, C.o.D.C.a. (2021). Antigenic Characterization.
https://www.cdc.gov/flu/about/professionals/antigenic.htm.
Rijksinstituut voor Volksgezondheid en Milieu (RIVM). Resultaten kiemsurveillance.
https://www.rivm.nl/sites/default/files/2022-
04/Kiemsurveillance%2020220401%20tabel%20NL.pdf.
Rossler, A., Riepler, L., Bante, D., von Laer, D., and Kimpel, J. (2022). SARS-CoV-2
Omicron Variant Neutralization in Serum from Vaccinated and Convalescent Persons. N
Engl J Med *386*, 698-700. 10.1056/NEJMc2119236.
Schmidt, F., Weisblum, Y., Muecksch, F., Hoffmann, H.H., Michailidis, E., Lorenzi, J.C.C.,
Mendoza, P., Rutkowska, M., Bednarski, E., Gaebler, C., et al. (2020). Measuring SARS-
CoV-2 neutralizing antibody activity using pseudotyped and chimeric viruses. J Exp Med
217. 10.1084/jem.20201181.

Smith, D.J., Lapedes, A.S., de Jong, J.C., Bestebroer, T.M., Rimmelzwaan, G.F., Osterhaus,
A.D., and Fouchier, R.A. (2004). Mapping the antigenic and genetic evolution of influenza
virus. Science *305*, 371-376. 10.1126/science.1097211.
van Gils, M.J., Lavell, A., van der Straten, K., Appelman, B., Bontjer, I., Poniman, M., Burger,
J.A., Oomen, M., Bouhuijs, J.H., van Vught, L.A., et al. (2022). Antibody responses against
SARS-CoV-2 variants induced by four different SARS-CoV-2 vaccines in health care
workers in the Netherlands: A prospective cohort study. PLOS Medicine *19*, e1003991.
10.1371/journal.pmed.1003991.
Wilhelm, A., Widera, M., Grikscheit, K., Toptan, T., Schenk, B., Pallas, C., Metzler, M.,
Kohmer, N., Hoehl, S., Helfritz, F.A., et al. (2021). Reduced Neutralization of SARS-CoV-2
Omicron Variant by Vaccine Sera and monoclonal antibodies. medRxiv,
2021.2012.2007.21267432. 10.1101/2021.12.07.21267432.
Wilks, S.H., Mühlemann, B., Shen, X., Türeli, S., LeGresley, E.B., Netzl, A., Caniza, M.A.,
Chacaltana-Huarcaya, J.N., Daniell, X., Datto, M.B., et al. (2022). Mapping SARS-CoV-2
antigenic relationships and serological responses. bioRxiv, 2022.2001.2028.477987.
10.1101/2022.01.28.477987.
World   Health   Organization   (2021).   Tracking   SARS-CoV-2   variants
https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/.
Wynberg, E., van Willigen, H.D.G., Dijkstra, M., Boyd, A., Kootstra, N.A., van den Aardweg,
J.G., van Gils, M.J., Matser, A., de Wit, M.R., Leenstra, T., et al. (2021). Evolution of COVID-
19 symptoms during the first 12 months after illness onset. Clin Infect Dis.
10.1093/cid/ciab759.
502

Fig.1 A

Fig. 2 A

B

Variant of concern	N= (%)	Age (years) median (range)	Male	Hospital admission	Sequence confirmed	Time since symptom onset (days) median (range)
D614G	20 (30%)	53 (22-74)	9 (45%)	9 (45%)	0 (0%)	38 (30-45)
Alpha	11 (17%)	48 (25-76)	6 (55%)	9 (82%)	11 (100%)	39 (24-58)
Alpha + E484K	2/11	62 (48-76)	0	2/2	2/2	41 (24-58)
Beta	8 (12%)	41 (18-53)	3 (38%)	0	7 (88%)	39 (30-63)
Gamma	4 (6%)	35 (27-10)	2 (50%)	0	4 (100%)	40 (38-55)
Delta	11 (17%)	31 (19-63)	6 (55%)	1 (9%)	10 (91%)	46 (36-52)
Omicron BA.1	8 (12%)	31 (21-45)	4 (50%)	0	3 (38%)	44 (25-75)
Omicron BA.2	4 (6%)	51 (31-55)	4 (100%)	0	4 (100%)	40 (39-41)
Total	66	41 (18-76)	34 (52%)	21 (32%)	39 (59%)	40 (24-75)

Table 1. Sociodemographics and clinical characteristics