Surgical interventions targeting the nucleus caudalis for craniofacial pain: a systematic and historical review.

Brendan Santyr M.D.[1], Mohamad Abbass M.D.[1], Alan Chalil, M.D.[1], Amirti Vivekanandan, M.D.[1], Margaret Tindale, M.D.[2], Nicholas M. Boulis, M.D.[3], Jonathan C. Lau, M.D., Ph.D., FRCSC.[1]

[1]Division of Neurosurgery, Department of Clinical Neurological Sciences, Western University, London, Ontario, Canada.
[2]Division of Emergency Medicine, Western University, London, Ontario, Canada.
[3]Department of Neurosurgery, Emory University, Atlanta, Georgia, USA.

Conflict of interest:
The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Disclosure of funding:
No funding was received for the completion of this work.

Corresponding Author:
Jonathan C. Lau, M.D., Ph.D., FRCSC
Department of Clinical Neurological Sciences
University Hospital
339 Windermere Rd, London, Ontario, Canada
N6A 5A5
Tel. 519-685-8500 x33707
jonathan.lau@lhsc.on.ca

Keywords: Neurosurgery, Neuromodulation, Pain, Facial Pain, Nucleus Caudalis, Trigeminal Nucleus.

Abstract

Introduction: Craniofacial pain is a prevalent group of conditions and when refractory to conventional treatments poses a significant burden. The last decade has seen a renewed interest in the multimodal management of pain. Interventions targeting the nucleus caudalis (NC) of the trigeminocervical complex have been available as a treatment option since the 1930s, yet evidence for efficacy remains limited.

Methods: We present a systematic review of the literature providing a historical perspective on interventions targeting the NC leading up to the present. We examine the various intervention techniques, clinical indications, and procedural efficacy. A novel outcome reporting scheme was devised to enable comparison among studies due to historically variable reporting methods.

Results: A review of the literature revealed 33 retrospective studies published over the last 80 years reporting on 827 patients. The most common technique was the open NC dorsal root entry zone nucleotomy/tractotomy; however, there has been an emergence of novel approaches such as endoscopic and spinal cord stimulation in the last 10 years. Regardless of intervention technique or preoperative diagnosis, 87% of patients demonstrated improvement with treatment.

Conclusion: The literature surrounding NC intervention techniques is reviewed. Recent advancements and the wide range of craniofacial pain syndromes for which these interventions show potential efficacy is discussed. New and less invasive techniques continue to emerge as putative therapeutic options. However, prospective studies are lacking. Furthermore, the evidence supporting even well-established techniques remain of poor quality. Future work should be prospective, utilize standard outcome reporting, and address efficacy comparisons between intervention type and preoperative diagnosis.

Introduction

Refractory craniofacial pain is a heterogeneous group of disorders with multiple classification systems depending on clinical features that include anatomical distribution, anatomical origin, pathology, and symptomatology.[1] The prevalence of craniofacial pain is estimated to be 26%.[2] Epidemiological studies have yet to define the social and economic burden of this complex and debilitating group of diseases, however estimates may be made from the literature on trigeminal neuralgia (TN), the best studied of the chronic craniofacial pain syndromes outside the realm of headache. TN has been reported to impact multiple health status domains including activity, mood, ambulation, work, relationships, sleep, and life enjoyment.[3] More than one-third of patients with TN had employment impacted through work hour reduction, disability, or unemployment.[3] Despite advancements over the past two decades, many TN patients remain refractory to medical and conventional surgical options.[4] Conventional surgical approaches to treat TN have targeted the nerve root and ganglion with demonstrated efficacy in treating typical paroxysmal TN type 1 pain but are relatively ineffective at treating the background pain characteristic of TN type 2.[5] Worsening of constant background pain and the development of neuropathic pain can be seen with repeated treatments targeting the anterior system (distal to the nerve root).[6] Therefore, an investigation into alternative surgical approaches and treatment options for these syndromes remains crucial to improving pain management.

Nociceptive craniofacial sensation, temperature, and crude touch are carried by general somatic afferent first-order neurons of the Vth, VIIth, IXth, and Xth cranial nerves to the trigeminocervical nuclear complex, where they synapse to second-order neurons within the nucleus caudalis (NC).[7] The trigeminal nucleus forms ascending connections to structures involved in processing nociceptive stimuli, such as the thalamus, hypothalamus, locus coeruleus, and periaqueductal gray. Here nociceptive stimuli are modulated and pain is perceived.[8] There is partial functional overlap between the C2 and C3 substantia gelatinosa and the NC, and it is this close anatomical relationship that inspired the use of dorsal root entry zone (DREZ) lesioning techniques to treat craniofacial pain.[8]

Interruption of the trigeminocervical complex at the level of the brainstem was first reported in humans in 1938.[9] Introduced in 1968, DREZ procedures for chronic pain have progressed to include various methods of lesioning or modulating the DREZ at the spinal level of the pain, attempting to eliminate hyperactivity and spontaneous discharges that result in central perception of pain.[10–13] Favorable outcomes in DREZ operations for other pain syndromes and advancements in techniques for open and percutaneous trigeminal tractotomy helped provide motivation for NC DREZ procedures for craniofacial pain.[14–17] Similar to standard spinal DREZ procedures that target the dorsal horn of the spine, NC DREZ operations target the spinal trigeminal nucleus pars caudalis at the cervicomedullary junction, relieving pain through destruction of the second-order neurons.[10,17–19]

Although the first application of lesioning of the trigeminocervical complex dates back to the 1930s, prospective studies assessing its efficacy and indications remain lacking. Evidence supporting the use of surgical techniques targeting the NC is limited to small retrospective case

104 reports and case series. A comprehensive summary of the efficacy, indications, and variations
105 in intervention technique has yet to be reported. In the present study, we performed a
106 systematic review of the literature published regarding the surgical intervention of the NC,
107 providing a historical perspective for the development of NC-related interventions to the present
108 day.
109
110 **Methods**
111
Literature Search and Inclusion Criteria
We conducted a systematic review according to Preferred Reporting Items for Systematic
Reviews and Meta-Analyses (PRISMA) guidelines. Embase, MEDLINE, and Cochrane central
databases were queried using the search strategy indicated in Supplementary Materials 1 on
May 31, 2021. The search comprised of the following search terms (combination of subject
headings and keywords): "caudalis", "trigeminal nucleus", "trigeminal tractotomy", "trigeminal
nucleotractotomy", "trigeminal nucleotomy", "dorsal root entry zone", "ablation", "lesion",
"stimulation", and various combinations of the above.
120
The search strategies were modified for each database to include database-specific thesaurus
terms, syntax, and field names. Only original peer-reviewed clinical studies in humans published
in the English language were included. Studies were not restricted based on diagnosis or the
lesioning/stimulation method. Additional relevant studies that met our inclusion criteria were
identified through examination of bibliographies of relevant articles and selected reviews.
Conference abstracts and unpublished studies were excluded. Anatomical studies, descriptions
of technique, studies reporting on the same patient population, and reviews were excluded. Full-
text analysis excluded studies with fewer than 5 patients as a method to reduce bias from case
reports or small case series. The PICOS (Population, Intervention, Comparison, Outcomes, and
Study Design) format was used to depict study eligibility with prespecified inclusion and
exclusion criteria (Table 1).
132
133 *Quality Assessment*
134 Two reviewers (B.S., M.A.) independently screened each article title and abstract for relevance.
135 Disagreements were resolved by discussion and consensus in the presence of a third reviewer
136 (A.C.). All authors reviewed the preliminary results and the final analysis.
137
Data Collection and Outcome Measures
Of the included studies, information regarding study type, study size, mean follow-up duration,
interventional technique, preoperative diagnosis, and pre- and post-operative pain rating was
collected (Table 2). Postoperative pain rating was considered a primary outcome. Mean follow-
up duration was collected in years following intervention. All reported preoperative diagnoses
were collected and classified based on available information. Diagnostic classification included
trigeminal neuralgia or neuropathic pain, oncologic craniofacial pain, post-herpetic pain,
anesthesia dolorosa, traumatic craniofacial pain, glossopharyngeal neuralgia, post-stroke
craniofacial pain, headache or migraine, multiple sclerosis-related craniofacial pain, and
geniculate neuralgia. Most of the literature failed to distinguish between different classifications

or causes of craniofacial pain, thereby necessitating the combination of these
pathophysiological entities. Intervention techniques were classified into all procedural variations
of open procedures targeting the NC or trigeminocervical complex, including nucleotomy,
nucleotomy/tractotomy, and DREZ lesioning procedures; computed tomography (CT) guided
and free-hand percutaneous nucleotomy/tractotomy; endoscopic DREZ nucleotomy/tractotomy;
cervical spinal cord stimulation (SCS); and ultrasonic nucleotomy (Figure 1). Information
regarding total perioperative complications was collected. These were further subdivided into
postoperative facial dysfunction (pain, numbness, weakness), corneal anesthesia,
oropharyngeal dysfunction (dysarthria, dysphagia, vocal cord paralysis), ataxia, limb dysfunction
(pain, numbness, weakness, dysmetria), transient dysfunction (pain, numbness, weakness,
ataxia), reoperation rate, general medical complication, and mortality.
159
Due to the historical nature of this review spanning back to the first descriptions of interventional
procedures involving the NC, we observed a wide variability of methods for reporting pain in the
literature. Most commonly, qualitative and non-standardized methods were used to report pain
outcomes with only 5 studies reporting using the standardized Visual Analog Scale. To
accommodate this and facilitate the comparison of modern and classic studies, a composite
rating scale was employed in this review (Table 3). A composite pain score of 1 includes
patients that are pain-free, Visual Analog Scale (VAS) 0, or have had a 100% pain reduction. A
pain score of 2 includes those with a "good" response, VAS 1-3, or a >50% pain reduction. A
pain score of 3 includes those with a "satisfactory" or "fair" response, VAS 4-6, or a <50% pain
reduction. Finally, a pain score of 4 includes those who experienced no improvement, pain
worsening, or VAS 7-10.
171
The fraction of patients demonstrating any symptomatic improvement post-procedure (pain
score of 1-3) was calculated as a percentage of the total number of subjects included in the
study (Table 2). Patients with mortality prior to follow-up or if the outcome was not reported were
deemed outcome unknown. To determine the relative surgical success between intervention
techniques, the techniques were divided as follows: open focal lesioning techniques
(nucleotomy/tractotomy), DREZ (multipoint lesioning technique), percutaneous
nucleotomy/tractotomy, and neuromodulation (cervical SCS). Pain outcome distribution was
then compared between each using a Chi-Squared test. The fraction of patients with post-
procedural improvement, as a measure of relative procedural success, was assessed with
respect to publication date to determine the influence of procedural modification over time
(Supplementary Materials 2). This analysis was achieved using a linear model for all
intervention types as well as individually for open focal lesioning techniques, DREZ (multipoint
lesioning technique), and percutaneous procedures. Other procedures were excluded due to an
inadequate number of studies. Linear models, covariance, and correlation were implemented
using the R statistical language and environment and the functions lm(), var(), and cor()
respectively.[20] Event rates for composite scores and complications were pooled across studies,
and logit transformed with 95% confidence intervals calculated assuming a binomial distribution.
This was calculated using MATLAB R2019b.[21]
190
Results

A total of 782 articles were identified through the literature search and underwent title and
abstract review (Figure 2). Two separate reviewers (B.S. and M.A.) screened each article to
meet inclusion/exclusion criteria. Following removal of duplicates, review of bibliographies from
relevant articles, and exclusion of studies with n < 5 patients, 33 studies encompassing 827
patients were included for analysis as summarized in Table 2. Disagreement was resolved by a
third reviewer (A.C.). All identified studies for inclusion were case series published between
1940 and 2020. No randomized or prospective trials were identified. The results of the data
extraction for included studies are summarized in Table 2. The publications included in this work
represent an 80-year history of techniques targeting the NC for treatment of craniofacial pain
(Figure 3). The rate of publications continues to increase over time and there are emerging
techniques including endoscopic approaches and spinal cord stimulation (SCS) seen in the last
10 years (Figure 3)

Preoperative Diagnosis
Treatments targeting the nucleus caudalis have been used to ameliorate craniofacial pain in a
wide range of patient populations. Of the 827 patients included in this review, 41.5% (n=343)
had a diagnosis of trigeminal neuralgia or neuropathic pain (Figure 4). Oncologic craniofacial
pain comprised 19.7% (n=163), 8.5% (n=70) had post herpetic pain, 5.7% (n=47) had
anesthesia dolorosa, 5.6% (n=46) had traumatic craniofacial pain, 3.6% (n=30) had
glossopharyngeal neuralgia, 1.9% (n=16) had post-stroke craniofacial pain, 0.8% (n=7) had
headache or migraine, 0.7% (n=6) had multiple sclerosis related craniofacial pain, 0.6% (n=5)
had geniculate neuralgia, and 11.4% (n=94) had multiple diagnoses or unspecified craniofacial
pain (Figure 4).

Intervention Technique
Review of the included articles revealed 5 intervention techniques targeting the nucleus caudalis
for craniofacial pain (Figures 1 and 5). The most common approach was open NC DREZ
nucleotomy/tractotomy, reported in 21 studies and comprising 60.1% of patients (n=497). CT
guided and free-hand percutaneous nucleotomy/tractotomy is reported in 8 studies and 35.6%
of patients (n=294). Cervical SCS is reported in 2 studies and 2.1% of patients (n=17).
Ultrasonic nucleotomy is reported in one study and 1.7% of patients (n=14). Endoscopic DREZ
nucleotomy/tractotomy is reported in 1 study and 0.6% of patients (n=5). Although not included
in the analysis, Gamma Knife radiosurgery was reported in a single case series of 2 patients.[22]

Treatment Outcomes
As mentioned previously, various pain reporting methods are utilized in the studies reviewed. To
facilitate comparison of treatment outcomes between studies, a composite pain rating scale was
developed and outlined in Table 3. Preoperative pain is even less consistently reported. Of the
827 included patients, 142 (17.2%) have preoperative VAS reported with a mean of 8.6. Using
the implemented composite score developed to facilitate comparison between modern and
historical studies, interventions targeting the NC resulted in a pain score of 1 (i.e., pain freedom,
VAS 0, or 100% pain reduction) in 237 patients (28.7%), pain score of 2 (i.e., "good" response,
VAS 1-3, or a >50% pain reduction) in 215 patients (26.0%), pain score of 3 (i.e., "satisfactory"

or "fair" response, VAS 4-6, or a <50% pain reduction) in 35 patients (4.2%), pain score of 4
(i.e., no improvement, pain worsening, or VAS 7-10) in 49 patients (5.7%) (Figure 6). The
treatment outcome was unknown in 291 patients (35.2%). This includes patients lost to follow-
up, passed away, or inadequately reported. The mean follow-up duration was 2.52 years.
240
Focusing on subjects with clearly reported outcome values, the intervention types were
reclassified as open focal lesioning techniques (nucleotomy/tractotomy) (n patients = 154), open
DREZ (multipoint lesioning technique) (n patients = 141), percutaneous nucleotomy/tractotomy
(n patients = 224), and cervical SCS (n patients = 17) and outcome distribution were compared
between the types (Figure 7). Chi-squared demonstrates a significant difference in the outcome
distribution between the intervention types (χ^2 = 57.5322, p <0.001) (Supplemental Material 2).
247
Linear regression failed to demonstrate a trend toward improving postoperative outcomes in
interventions over the 80-year history (Supplemental Materials 3). A non-significant trend
towards improved outcomes is seen in open focal lesioning and percutaneous
nucleotomy/tractotomy when separated by intervention type. The correlation coefficients are
0.45 and 0.44, the covariances are 1.05 and 1.12, and p-values are 0.19 and 0.33 respectively
(Supplemental Materials 3).
254
Complications
A total of 251 complications are reported. The proportion of individual complications per
intervention type is presented in Figure 8. All complications except corneal anesthesia (n = 5),
oropharyngeal dysfunction (n = 11), and limb dysfunction (n = 14) demonstrate a statistically
significant difference in distribution between intervention types (focal lesioning techniques,
DREZ, percutaneous, and neuromodulation) (p < 0.001) (Supplemental Material 4).
261
262 **Discussion**
263
In this study, we performed a systematic review of the published literature regarding the surgical
interventions of the NC and investigated the efficacy and indications for these procedures. 87%
of subjects demonstrated some improvement from NC-targeted treatment. Furthermore, 44.2%
reported complete pain freedom postoperatively. We show percutaneous lesioning has a
relatively lower complication rate compared to other intervention types. Despite the increase in
described surgical options for craniofacial pain, up to 27% of patients undergoing surgical
treatment have insufficiently managed pain at 5 years.[2,23] The present study demonstrates that
there has been a recent increase in publication rate over 80 years since the original descriptions
of nucleus caudalis interventions. Contributing to this are not only the advancements in open
and percutaneous techniques but also the emergence of new approaches such as ultrasonic
lesioning, radiosurgery, and SCS.
275
The studies included in this review report on diverse diagnoses resulting in chronic craniofacial
pain including post-herpetic pain, trigeminal neuralgia or neuropathic pain, oncological pain,
migraine, multiple sclerosis-related craniofacial pain, and geniculate neuralgia. A detailed
description regarding the pathophysiology of pain associated with each of these conditions is

outside the scope of this review; however, common to each of these diseases is injury
(compression, denervation/deafferentation, demyelination) of the general somatic afferent fibers
of the Vth, VIIth, IXth, or Xth cranial nerves. These fibers synapse onto second order neurons
within the descending trigeminocervical complex, which extends from the pons to the upper
cervical spinal cord on the dorsal surface of the brainstem.[24] Based on anatomic and clinical
data, the nucleus caudalis was identified as the most important locus for modulation, integration,
and conduction of nociception from craniofacial structures to higher-order intracranial structures
for pain perception.[9,25–28] This provides a logical target for treatment of pain associated with
injury to cranial nerves V, VII, IX, and X. Intervention techniques such as the NC DREZ
nucleotomy/tractotomy utilize a small suboccipital craniectomy and C1-C2 laminectomy to
elevate the cerebellar tonsils and visualize the obex. The NC occupies the triangular area
between the dorsolateral sulcus and the emerging points of the accessory nerve. It tapers
caudally before joining the spinal DREZ of C2. During NC DREZ nucleotomy/tractotomy multiple
rostrocaudal and mediolateral lesions are made at a depth of 3-4mm covering 5mm below the
obex to the upper dorsal rootlets of C2.[24]
295	
The identified articles reveal progressive adaptation of intervention techniques targeting the NC.
Most studies (21 studies comprising 60.1% of patients) report on the open NC DREZ
nucleotomytractotomy. Open procedures in this region date back to the 1930s where Sjoqvist
demonstrated that sectioning of the trigeminocervical complex at the level of the medulla above
the obex resulted in ipsilateral thermoanalgesia of the face.[9] Although initially plagued by
complications, a near century of surgical advancement has considerably improved its safety and
efficacy, including the development of the more selective vertical trigeminal partial nucleotomy
in 1965.[29] The 1970s saw the emergence of stereotactic and percutaneous targeting
techniques, improving pain outcomes through patient cooperation and intraoperative monitoring
of lesion-induced analgesia under local anesthetic.[15] Surgical refinement and improved
anatomical understanding of the region in the 1970s and 80s resulted in specific open NC
DREZ lesioning procedures gaining popularity with higher immediate success rates.[30] In recent
decades a trend towards less invasive techniques is apparent, as morbidity related to injury of
nearby structures of the spinal cord and medulla from open procedures remain high; historically
5% encounter ataxia, 25% with paresthesias, and 8-24% with lesioning of the 10th nerve.[24]
Endoscopic NC DREZ nucleotomy/tractotomy procedures have been developed to minimize the
morbidity associated with large incisions and craniotomies.[31,32] Noninvasive techniques using
ultrasound[33] and Gamma Knife radiosurgery[22] have similarly been developed. Finally, the most
recent adaptation of techniques targeting the NC is in the form of spinal cord stimulation as a
titratable and reversible form of neuromodulation of the region.[34–38]
316	
It is clear that the techniques targeting the NC for ablation or neuromodulation represent an
additional option for the treatment of intractable craniofacial pain of diverse pathophysiological
origins. Despite the variations in intervention technique and preoperative diagnosis, 87% of
subjects demonstrated some improvement from treatment. Furthermore, 44.2% reported
complete pain freedom postoperatively, suggesting remarkable efficacy for a group of
treatments typically reserved for complex and otherwise surgically and medically refractory
craniofacial pain. Although this might be an overestimation of the overall effectiveness of these

procedures due to publication bias, it is nevertheless encouraging for continued prospective
study of these procedures. Also demonstrated here are significant differences in outcome
distributions between different procedural techniques. However, due to the inconsistencies in
outcome reporting from the inclusion of studies prior to standardization of reporting pain metrics,
efficacy comparisons between intervention type or preoperative diagnosis could not be reliably
made. There have been efforts in the literature to standardize pain outcome reporting, with
multiple well-validated pain rating scales available.[39] Here we present added evidence for the
need to continually report outcomes in a way that allows for robust comparisons. It is imperative
that future prospective studies address these lingering questions.
333
Analysis of reported complication rates demonstrates significant differences in complications
between the different intervention techniques. The data support percutaneous interventions
having a relatively lower rate of complications compared to others (Figure 8). These results
should be interpreted cautiously as they are certainly confounded by publication bias,
inadequate reporting of complications, and limited duration of follow-up (2.52 years). Further
investigation should be the subject of future prospective trials.
340
Open trigeminal tractotomy (focal lesioning technique) as a procedure has become largely
historical due to frequent partial/patchy craniofacial analgesia[40], high recurrence rates[41], and
frequency of morbidity and mortality.[24] Advancements in technology and understanding in the
1970s paved the way for the more common adaptation of these procedures seen today:
caudalis DREZ lesioning (multipoint lesioning technique) and percutaneous lesioning.
Determining when to pursue each technique and for whom remains a difficult question to
answer when prospective trials and comparison studies are lacking. The literature reviewed
here suggests that with caudalis DREZ lesioning pain improvement may be achieved in 70%
patients. Bernard et al. surmise that patient description of pain may help predict postoperative
outcome[30], patients with sharp/burning pain finding better relief than those with dull pain, 55%
and 24% respectively.[30] Better results were also achieved when pain is restricted to the
trigeminal territory and with fewer trigeminal distributions involved.[30] Good outcome is observed
in 75% of patients with pain involving one trigeminal distribution, 50% of patients with two, and
38% of patients where all three territories are involved.[30] Similarly percutaneous procedures
demonstrate pain improvement in 71% of patients. However, these procedures as described in
the literature tend to target afferent fiber tracts, second-order neurons, and internuncial neurons
and tracts in the spinal trigeminal complex eliminating abnormal discharging of deafferented
neurons and thereby providing better pain relief in patients with central or deafferentation pain
compared to caudalis DREZ lesioning.[18,30,33,42,43] Furthermore, we demonstrate a lower total
complication rate in percutaneous techniques with only a slightly higher reoperation/revision rate
compared to caudalis DREZ lesions. Historical literature and the results from this work support
the use of percutaneous procedures in providing an equivalent degree of craniofacial pain
reduction as caudalis DREZ lesioning, with better efficacy in deafferentation pain, and fewer
complications. Caudalis DREZ lesioning may be reserved for uncooperative patients or those
with pain refractory to other treatments, and those with sharp/burning pain in a limited
distribution.
367

Newer techniques such as endoscopic, ultrasound, gamma knife, and SCS report on too few
patients to make meaningful conclusions regarding their indications and relative efficacy,
however neuromodulation with cervical SCS is showing a recent rise in publication rate.[34–38] The
efficacy and safety of SCS in the treatment of chronic pain is well accepted and supported in the
literature.[44] Likewise, this efficacy is replicated in this review, where 88% of patients receiving
cervical SCS experience some degree of pain reduction. Velasquez et al. report a mean pain
reduction of 57.1% making it comparable to ablative procedures on a patient level.[37,45,46]
However large prospective studies are lacking and indication recommendations are limited. We
show a lower overall complication rate relative to caudalis DREZ lesioning with no reported
mortality or permanent neurological deficit such as ataxia, weakness, or paresthesia. There was
a slightly higher total complication rate compared to percutaneous techniques, likely secondary
to a higher reoperation/revision rate from infection, loss of efficacy, and lead malpositioning
rather than neurological deficit.[37] With a body of literature that is continually growing, cervical
SCS is presenting itself as a potential early, titratable, and reversible therapy with a lower
complication risk and comparable outcomes.

The need for ongoing innovation in chronic pain management is apparent and less invasive
methods are emerging. Of particular interest, recent reports have described both intraventricular
and intrathecal analgesia administration in cases of refractory craniofacial pain.[47,48] As the
evidence continues to evolve these procedures may become viable options for patients that
have failed other strategies or whose pain is bilateral.[47,48]

The first application of lesioning of the trigeminocervical complex dates back to the 1930s[9];
however, evidence supporting its use is limited to small retrospective case series and case
reports. As mentioned above the limitations encountered are a lack of prospective studies and
inadequate outcome reporting to strengthen comparisons between intervention types and
diagnosis. Historical studies, as encountered in this work, particularly suffer from a lack of
access to more recently devised outcome reporting schemes and classification systems. For
example, the contemporary classification of trigeminal neuralgia and as such differentiating
trigeminal neuropathic pain from trigeminal neuralgia could not be performed. Furthermore, the
heterogeneity of included studies with respect to intervention, diagnosis, and outcome reporting
measures necessitated grouping all these factors together and creating a custom post-
procedural pain reporting scale. This may limit the strength of efficacy conclusions; however, it
does contribute greatly to the generalizability of the included results and allowed comparison to
historical studies. Additionally, information regarding follow-up and treatment outcome durability
is almost universally lacking in the literature and only 64.8% of patients have reported outcome
data. Combining preoperative diagnoses further limits the conclusions able to be drawn as life
expectancy for some conditions, i.e. those with oncological facial pain, may be shorter than the
time it may take to achieve full benefit from the procedure.

Conclusion

This review summarizes 80 years of literature regarding interventions targeting the NC for
chronic refractory craniofacial pain. Recent advancements in intervention techniques are

highlighted, but also shortcomings in the literature impede widespread adoption. Only 64.8% of
included subjects have outcomes reported and due to the inclusion of studies prior to the
standardization of pain reporting metrics, there are inconsistencies in the way these outcomes
are reported. New and less invasive techniques continue to emerge, however prospective
studies remain absent in the literature. Additional studies addressing efficacy comparisons
between intervention type or preoperative diagnosis would help solidify these techniques in the
armamentarium of craniofacial pain treatments.
419

References

1. Zakrzewska JM. Facial pain: neurological and non-neurological. *Journal of neurology, neurosurgery, and psychiatry*. 2002;72 Suppl 2:ii27-ii32. doi:10.1136/jnnp.72.suppl_2.ii27

2. Macfarlane T v., Blinkhorn AS, Davies RM, Kincey J, Worthington H v. Oro-facial pain in the community: prevalence and associated impact. *Community Dentistry and Oral Epidemiology*. 2002;30(1):52-60. doi:10.1034/J.1600-0528.2002.300108.X

3. Tolle T, Dukes E, Sadosky A. Patient Burden of Trigeminal Neuralgia: Results from a Cross-Sectional Survey of Health State Impairment and Treatment Patterns in Six European Countries. *Pain Practice*. 2006;6(3):153-160. doi:10.1111/j.1533-2500.2006.00079.x

4. Cruccu G, Aziz TZ, Garcia-Larrea L, et al. EFNS guidelines on neurostimulation therapy for neuropathic pain. *European Journal of Neurology*. 2007;14(9):952-970. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed10&AN=47300786

5. Tyler-Kabara EC, Kassam AB, Horowitz MH, et al. Predictors of outcome in surgically managed patients with typical and atypical trigeminal neuralgia: comparison of results following microvascular decompression. *Journal of Neurosurgery*. 2002;96(3):527-531. doi:10.3171/jns.2002.96.3.0527

6. Broggi G, Ferroli P, Franzini A, Galosi L. The role of surgery in the treatment of typical and atypical facial pain. *Neurological Sciences*. 2005;26(SUPPL. 2). doi:10.1007/s10072-005-0418-5

7. Bullard DE, Nashold Jr. BS. The caudalis DREZ for facial pain. *Stereotactic & Functional Neurosurgery*. 1997;68(1-4 Pt 1):168-174. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med4&AN=9711711

8. Goadsby PJ, Holland PR, Martins-Oliveira M, Hoffmann J, Schankin C, Akerman S. Pathophysiology of Migraine: A Disorder of Sensory Processing. *Physiological Reviews*. 2017;97(2):553. doi:10.1152/PHYSREV.00034.2015

9. SJOQVIST, O. Studies of pain conduction in the trigeminal nerve. *Act Psychiat Neurol Scand Suppl*. 1938;17:1-139. Accessed October 26, 2021. https://ci.nii.ac.jp/naid/10008988203

10. Loeser JD, Ward AA, White LE. Chronic Deafferentation of Human Spinal Cord Neurons. *Journal of Neurosurgery*. 1968;29(1):48-50. doi:10.3171/JNS.1968.29.1.0048

11. Guenot M, Bullier J, Rospars JP, Lansky P, Mertens P, Sindou M. Single-Unit Analysis of the Spinal Dorsal Horn in Patients With Neuropathic Pain. *Journal of Clinical Neurophysiology*. 2003;20(2):143-150. doi:10.1097/00004691-200304000-00009

12. Sindou M. Drez lesions for brachial plexus injury. *Neurosurgery*. 1988;23(4):528. doi:10.1097/00006123-198810000-00025

13. Drake CG, Stavraky GW. AN EXTENSION OF THE "LAW OF DENERVATION" TO AFFERENT NEURONES. *Journal of Neurophysiology*. 1948;11(3):229-238. doi:10.1152/jn.1948.11.3.229

14. MCKENZIE KG. Trigeminal tractotomy. *Clinical neurosurgery*. 1954;2:50-69; discussion, 69-70. Accessed October 26, 2021. http://www.ncbi.nlm.nih.gov/pubmed/13270405

15. Hitchcock ER, Schvarcz JR. Stereotaxic trigeminal tractotomy for post-herpetic facial pain. *Journal of Neurosurgery*. 1972;37(4):412-417. doi:10.3171/JNS.1972.37.4.0412

16. Schvarcz JR. Spinal Cord Stereotactic Techniques re Trigeminal Nucleotomy and Extralemniscal Myelotomy. *Stereotactic and Functional Neurosurgery*. 1978;41(1-4):99-112. doi:10.1159/000102406

17. Gorecki JP, Nashold Jr. BS, Rubin L, Ovelmen-Levitt J. The Duke experience with nucleus caudalis DREZ coagulation. *Stereotactic & Functional Neurosurgery*. 1995;65(1-4):111-116. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med3&AN=8916338

18. Anderson LS, Black RG, Abraham J, Ward AA. Neuronal hyperactivity in experimental trigeminal deafferentation. *Journal of Neurosurgery*. 1971;35(4):444-452. doi:10.3171/JNS.1971.35.4.0444

19. Bernard Jr EJ, Nashold Jr BS, Caputi F, Moossy JJ. Nucleus caudalis DREZ lesions for facial pain. *British Journal of Neurosurgery*. 1987;1(1):81-92. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed3&AN=18068984

20. Bunn A, Korpela M. R: A language and environment for statistical computing. Published online 2013. doi:10.1016/j.dendro.2008.01.002

21. Lau JC, Kosteniuk SE, Walker T, Iansavichene A, Macdonald DR, Megyesi JF. Operative Complications with and without Image Guidance: A Systematic Review and Meta-Analysis of the Ommaya Reservoir Literature. *World Neurosurgery*. 2019;122:404-414. doi:10.1016/j.wneu.2018.11.036

22. Chang JW, Choi JY, Yoon Y, Park YG, Chung SS. Unusual causes of trigeminal neuralgia treated by gamma knife radiosurgery. Report of two cases. *Journal of Neurosurgery*. 2002;97(5 Suppl):533-535. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med4&AN=12507091

23. Al-Quliti KW. Update on neuropathic pain treatment for trigeminal neuralgia. *Neurosciences*. 2015;20(2):107-114. doi:10.17712/nsj.2015.2.20140501

24. Teixeira MJ, Fonoff ET. Technique of Trigeminal Nucleotractotomy. *Textbook of Stereotactic and Functional Neurosurgery*. Published online 2009:2097-2123. doi:10.1007/978-3-540-69960-6_124

25. Hu JW, Sessle BJ. Trigeminal nociceptive and non-nociceptive neurones: brain stem intranuclear projections and modulation by orofacial, periaqueductal gray and nucleus raphe magnus stimuli. *Brain Research*. 1979;170(3):547-552. doi:10.1016/0006-8993(79)90973-9

26. Hosobuchi Y, Rutkin B. Descending trigeminal tractotomy. Neurophysiological approach. *Archives of neurology*. 1971;25(2):115-125. doi:10.1001/ARCHNEUR.1971.00490020033004

27. Wall PD, Taub A. FOUR ASPECTS OF TRIGEMINAL NUCLEUS AND A PARADOX. *Journal of Neurophysiology*. 1962;25(1):110-126. doi:10.1152/jn.1962.25.1.110

28. Broton JG, Rosenfeld JP. Rostral trigeminal projections signal perioral facial pain. *Brain Research*. 1982;243(2):395-400. doi:10.1016/0006-8993(82)90269-4

29.   Kunc Z. Treatment of Essential Neuralgia of the 9th Nerve by Selective Tractotomy.
Journal of Neurosurgery. 1965;23(5):494-500. doi:10.3171/jns.1965.23.5.0494
30.   Bernard Jr EJ, Nashold Jr BS, Caputi F. Clinical review of nucleus caudalis dorsal root
entry zone lesions for facial pain. *Applied Neurophysiology*. 1988;51(2-5):218-224.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed4
&AN=18121840
31.   Teixeira MJ, de Almeida FF, de Oliveira YS, Fonoff ET. Microendoscopic stereotactic-
guided percutaneous radiofrequency trigeminal nucleotractotomy. *Journal of
Neurosurgery*. 2012;116(2):331-335.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med9&
AN=21999320
32.   Rahimpour S, Patel V. Endoscopic trigeminal nucleus caudalis doral root entry zone
lesioning for atypical facial pain. *Clinical Neurosurgery*. 2015;62 (Supplement 1):205.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed16
&AN=624269046
33.   Grigoryan Yu A, Slavin K v, Ogleznev Ky. Ultrasonic lesion of the trigeminal nucleus
caudalis for deafferentation facial pain. *Acta Neurochirurgica*. 1994;131(3-4):229-235.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med3&
AN=7754827
34.   Yearwood TL, Venkatesan L, Harrell CC. Resolution of primary cervicogenic headache
with sub-perception active recharge spinal cord stimulation (10721). *Neuromodulation*.
2016;19 (3):e150.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed17
&AN=72286353
35.   Richter B, Tomycz ND, Patterson M. C1 retrograde spinal cord stimulation for trigeminal
deafferentation pain. case study with long-term follow-up. *Neuromodulation*. 2017;20
(7):e165.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed18
&AN=619001349
36.   Patel K v, Lohani M. "Paresthesia-free relief: Higher frequency stimulation of high cervical
leads for patients with chronic facial pain." *Neuromodulation*. 2018;21 (3):e28.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed19
&AN=621997721
37.   Velásquez C, Tambirajoo K, Franceschini P, Eldridge PR, Farah JO. Upper Cervical
Spinal Cord Stimulation as an Alternative Treatment in Trigeminal Neuropathy. *World
Neurosurgery*. 2018;114:e641-e646. doi:10.1016/j.wneu.2018.03.044
38.   Jones MR, Rosenow JM. Importance of Laterality in Cervical Spinal Cord Stimulation for
Facial Pain: Case Report and Anatomic Review. *Operative Neurosurgery*.
2020;19(1):E83-E86. doi:10.1093/ons/opz381
39.   Younger J, McCue R, Mackey S. Pain outcomes: A brief review of instruments and
techniques. *Current Pain and Headache Reports*. 2009;13(1):39-43. doi:10.1007/s11916-
009-0009-x

40. Falconer MA. INTRAMEDULLARY TRIGEMINAL TRACTOTOMY AND ITS PLACE IN
THE TREATMENT OF FACIAL PAIN. *Journal of Neurology, Neurosurgery, and*
Psychiatry. 1949;12(4):297. doi:10.1136/JNNP.12.4.297

41. Moffie D. Late results of bulbar trigeminal tractotomy. Some remarks on recovery of
sensibility. *Journal of neurology, neurosurgery, and psychiatry*. 1971;34(3):270-274.
doi:10.1136/JNNP.34.3.270

42. Hitchcock ER, Tsukamoto Y. Physiological correlates in stereotactic spinal surgery. *Acta*
neurochirurgica. 1974;Suppl 21:119-123. doi:10.1007/978-3-7091-8355-7_16

43. Westrum LE, Black RG. Changes in the synapses of the spinal trigeminal nucleus after
ipsilateral rhizotomy. *Brain research*. 1968;11(3):706-709. doi:10.1016/0006-
8993(68)90163-7

44. Deer TR, Mekhail N, Provenzano D, et al. The Appropriate Use of Neurostimulation of the
Spinal Cord and Peripheral Nervous System for the Treatment of Chronic Pain and
Ischemic Diseases: The Neuromodulation Appropriateness Consensus Committee.
Neuromodulation: Technology at the Neural Interface. 2014;17(6):515-550.
doi:10.1111/NER.12208

45. Chivukula S, Tempel ZJ, Chen CJ, Shin SS, Gande A v, Moossy JJ. Spinal and Nucleus
Caudalis Dorsal Root Entry Zone Lesioning for Chronic Pain: Efficacy and Outcomes.
World Neurosurgery. 2015;84(2):494-504.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med12
&AN=25900792

46. Kanpolat Y, Tuna H, Bozkurt M, Elhan AH. Spinal and nucleus caudalis dorsal root entry
zone operations for chronic pain. *Neurosurgery*. 2008;62(3 Suppl 1):235-242; discussion
242-4.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med7&
AN=18424991

47. Dupoiron D, Dupoiron D. Targeted Drug Delivery (Intrathecal and Intracranial) for
Treatment of Facial Pain. *Progress in neurological surgery*. 2020;35:181-193.
doi:10.1159/000509623

48. Lee DJ, Gurkoff GG, Goodarzi A, Muizelaar JP, Boggan JE, Shahlaie K.
Intracerebroventricular opiate infusion for refractory head and facial pain. *World Journal*
of Clinical Cases : WJCC. 2014;2(8):351. doi:10.12998/WJCC.V2.I8.351

49. Grant F, Groff R, FH Lewy. Section of the descending spinal root of the fifth cranial nerve.
Archives of Neurology & Psychiatry. Published online 1940. Accessed November 1,
2021. https://jamanetwork.com/journals/archneurpsyc/article-
abstract/648387?casa_token=F6XWxXw0YTsAAAAA:vIllZs5646NFijw8tywI9eJce9dlhZn
wCtw74NXCQeT-zWpvuK6P9Fq0C9jj1YvNMdU2gCKoFw

50. GRANT FC. EXPERIENCES WITH INTRAMEDULLARY TRACTOTOMY. *Archives of*
Surgery. 1941;42(4):681. doi:10.1001/archsurg.1941.01210100041004

51. OLIVECRONA H. TRACTOTOMY FOR RELIEF OF TRIGEMINAL NEURALGIA.
Archives of Neurology And Psychiatry. 1942;47(4):544.
doi:10.1001/archneurpsyc.1942.02290040034002

52. HAMBY W, SHINNERS B, MARSH I. Trigeminal tractotomy: Observations on forty-eight
cases. *Archives of Surgery*. Published online 1948. Accessed November 2, 2021.

https://jamanetwork.com/journals/jamasurgery/article-
abstract/548105?casa_token=el4PG2XRRbIAAAAA:hB4eu_aqNL5sx8FJPrzDmRJXeH-
dvqeNBbFA8m8j1U-36GlVO2cLrXqReFcBMgG6j0ua-0-g

53.   Guidetti B. Tractotomy for the Relief of Trigeminal Neuralgia: Observations in 124 Cases.
Journal of Neurosurgery. 1950;7(6):499-508. doi:10.3171/JNS.1950.7.6.0499

54.   Raney R, Raney AA, Hunter CR. Treatment of major trigeminal neuralgia through section
of the trigeminospinal tract in the medulla. *The American Journal of Surgery*.
1950;80(1):11-17. doi:10.1016/0002-9610(50)90550-2

55.   Fox JL. Intractable Facial Pain Relieved by Percutaneous Trigeminal Tractotomy. *JAMA:*
The Journal of the American Medical Association. 1971;218(13):1940.
doi:10.1001/jama.1971.03190260054019

56.   Crue BL, Carregal EJA, Felsoory A. Percutaneous Stereotactic Radiofrequency
Trigeminal Tractotomy with Neurophysiological Recordings1. *Confin neurol*. 1972;34:389-
397.

57.   Fox JL. Percutaneous trigeminal tractotomy for facial pain. *Acta Neurochirurgica*.
1973;29(1-2):83-88. doi:10.1007/BF01414619

58.   Plangger CA, Fischer J, Grunert V, Mohsenipour I. Tractotomy and partial vertical
nucleotomy--for treatment of special forms of trigeminal neuralgia and cancer pain of face
and neck. *Acta Neurochirurgica - Supplementum*. 1987;39:147-150.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med2&
AN=3478977

59.   Rossitch Jr. E, Zeidman SM, Nashold Jr. BS. Nucleus caudalis DREZ for facial pain due
to cancer. *British Journal of Neurosurgery*. 1989;3(1):45-49.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med3&
AN=2789713

60.   Young JN, Nashold Jr. BS, Cosman ER. A new insulated caudalis nucleus DREZ
electrode. Technical note. *Journal of Neurosurgery*. 1989;70(2):283-284.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med3&
AN=2913226

61.   Morita M, Hosobuchi Y. Descending Trigeminal Tractotomy for Trigeminal Neuralgia after
Surgical Failure. *Stereotactic and Functional Neurosurgery*. 1992;59(1-4):52-55.
doi:10.1159/000098917

62.   Nashold BS, El-Naggar AO, Ovelmen-Levitt J, Abdul-Hak M. A new design of
radiofrequency lesion electrodes for use in the caudalis nucleus DREZ operation. *Journal*
of Neurosurgery. 1994;80(6):1116-1120. doi:10.3171/jns.1994.80.6.1116

63.   Kanpolat Y, Cosman ER. Special Radiofrequency Electrode System for Computed
Tomography-guided Pain-relieving Procedures. *Neurosurgery*. 1996;38(3):600-603.
doi:10.1097/00006123-199603000-00040

64.   Husain AM, Elliott SL, Gorecki JP, et al. Neurophysiological monitoring for the nucleus
caudalis dorsal root entry zone operation. *Neurosurgery*. 2002;50(4):822-828.
http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed7
&AN=36322808

65.   Delgado-Lopez P, Garcia-Salazar F, Mateo-Sierra O, Carrillo-Yague R, Llaurado G,
Lopez E. Trigeminal nucleus caudalis dorsal root entry zone radiofrequency

637   thermocoagulation for invalidating facial pain. *Neurocirugia (Asturias, Spain).*
638   2003;14(1):25-32; discussion 32.
639   http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med5&
640   AN=12655381
641 66. Teixeira MJ, Lepski G, Aguiar PHP, Cescato VAS, Rogano L, Alaminos AB. Bulbar
642   Trigeminal Stereotactic Nucleotractotomy for Treatment of Facial Pain. *Stereotactic and*
643   *Functional Neurosurgery.* 2003;81(1-4):37-42. doi:10.1159/000075102
644 67. Kanpolat Y, Kahilogullari G, Ugur HC, Elhan AH. Computed tomography-guided
645   percutaneous trigeminal tractotomy-nucleotomy. *Neurosurgery.* 2008;63(1 SUPPL.).
646   doi:10.1227/01.NEU.0000320139.27501.69
647 68. Raslan AM. Percutaneous computed tomography-guided radiofrequency ablation of
648   upper spinal cord pain pathways for cancer-related pain. *Neurosurgery.* 2008;62(3
649   SUPPL. 1):ONS226-ONS233.
650   http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed10
651   &AN=351572640
652
653

654 **Table 1:** Inclusion and exclusion criteria used for article screening presented in PICOS format.

	INCLUSION CRITERIA	EXCLUSION CRITERIA	DATA EXTRACTED
POPULATION	Patients undergoing NC intervention for craniofacial pain. Adults (18 years of age or older).	Patients undergoing intervention at anatomical site other than NC. Children, adolescents, or nonhuman subjects.	Total number of patients (n) Preoperative diagnosis
INTERVENTION	NC lesioning or modulation for craniofacial pain.	Anatomic studies or studies focused on description of technique or technological advancement.	Intervention technique utilized.
COMPARATOR	Single-arm studies reporting outcome data.	Studies published on the same patient population.	Intervention technique utilized.
OUTCOME	Pain-related outcomes. Procedure-related complications.	None.	Pre- and post-operative pain rating. Mean follow-up duration. Total number of perioperative complications, as well as postoperative facial dysfunction, corneal anesthesia, oropharyngeal dysfunction, ataxia, limb dysfunction, transient neurological dysfunction, reoperation rate, general medical complication, and mortality.
STUDY DESIGN	Any clinical study design: randomized control trial, observational (case-control; retrospective or prospective cohort), case series.	Nonclinical study. Conference abstract, review, editorial, letter, or commentary. Study cohort of < 5 patients.	Study design. Year of publication.

655 Abbreviations: PCIOS, Population, Intervention, Comparison, Outcome, and Study Design; NC,
656 Nucleus Caudalis.

657 **Table 2:** Summary of articles reporting interventions targeting the nucleus caudalis for
658 craniofacial pain.
659

Author (year)	Patients, #	Diagnoses (n)	Intervention Technique	Percent of patients improved
Grant (1940)[49]	12	TN (4) Oncologic (8)	Open DREZ nucleotomy/tractotomy	100%
Grant (1941)[50]	17	TN (6) Oncologic (11)	Open DREZ nucleotomy/tractotomy	88%
Olivecrona (1942)[51]	34	TN (34)	Open DREZ nucleotomy/tractotomy	91%
Hamby (1948)[52]	48	TN (35) Oncologic (13)	Open DREZ nucleotomy/tractotomy	67%
Falconer (1949)[40]	20	TN (13) Postherpetic (4) Anesthesia Dolorosa (2) Not specified (1)	Open DREZ nucleotomy/tractotomy	84%
Guidetti (1950)[53]	97	TN (97)	Open DREZ nucleotomy/tractotomy	Not reported
Raney (1950)[54]	59	TN (59)	Open DREZ nucleotomy/tractotomy	62%
Kunc (1965)[29]	6	Glossopharyngeal Neuralgia (6)	Open DREZ nucleotomy/tractotomy	100%
Fox (1971)[55]	12	TN (1) Oncologic (7) Postherpetic (2) Anesthesia Dolorosa (2)	Percutaneous nucleotomy/tractotomy	67%
Hosobuchi (1971)[26]	6	TN (1) Oncologic (3) Anesthesia Dolorosa (1) Not specified (1)	Open DREZ nucleotomy/tractotomy	100%
Crue (1972)[56]	12	TN (4) Oncologic (8)	Percutaneous nucleotomy/tractotomy	100%
Fox (1973)[57]	18	Oncologic (14) Postherpetic (2) Anesthesia Dolorosa (1) Not specified (1)	Percutaneous nucleotomy/tractotomy	61%
Schvarcz (1978)[16]	100	TN (19) Oncologic (31) Trauma (25) Anesthesia Dolorosa (14) Postherpetic (8) Glossopharyngeal Neuralgia (2) MS (1)	Percutaneous nucleotomy/tractotomy	77%

Plangger (1987)[58]	20	TN (12) Oncologic (6) Postherpetic (1) Anesthesia Dolorosa (1)	Open DREZ nucleotomy/tractotomy	100%
Bernard Jr. (1988)[30]	27	Not specified (27)	Open DREZ nucleotomy/tractotomy	95%
Rossitch Jr. (1989)[59]	5	Oncologic (5)	Open DREZ nucleotomy/tractotomy	100%
Young (1989)[60]	9	Not specified (9)	Open DREZ nucleotomy/tractotomy	Not reported
Morita (1992)[61]	7	TN (7)	Open DREZ nucleotomy/tractotomy	100%
Grigoryan (1994)[33]	14	Oncologic (10) Postherpetic (4)	Ultrasonic nucleotomy	93%
Nashold Jr. (1994)[62]	21	TN (8) Postherpetic (5) Anesthesia Dolorosa (2) Traumatic (1) Post Stroke (3) MS (1) Headache (1)	Open DREZ nucleotomy/tractotomy	57%
Gorecki (1995)[17]	46	Oncologic (1) Postherpetic (8) Anesthesia Dolorosa (15) Post Stroke (5) MS (1) Headache (4) Not specified (12)	Open DREZ nucleotomy/tractotomy	83%
Kanpolat (1996)[63]	19	TN (14) Oncologic (5)	Percutaneous nucleotomy/tractotomy	Not reported
Bullard (1997)[7]	25	TN (8) Oncologic (1) Postherpetic (1) Anesthesia Dolorosa (3) Traumatic (4) Post Stroke (1) MS (3) Not specified (4)	Open DREZ nucleotomy/tractotomy	100%
Husain (2002)[64]	5	Postherpetic (1) Traumatic (2) Headache (1) Not specified (1)	Open DREZ nucleotomy/tractotomy	80%
Delgado-Lopez (2003)[65]	6	Anesthesia Dolorosa (6)	Open DREZ nucleotomy/tractotomy	83%
Teixeira (2003)[66]	58	TN (2) Postherpetic (28) Oncologic (14)	Percutaneous nucleotomy/tractotomy	90%

		Post Stroke (7) Not specified (7)		
Kanpolat (2008a)[67]	65	TN (5) Oncologic (13) Postherpetic (3) Anesthesia Dolorosa (1) Glossopharyngeal Neuralgia (17) Geniculate Neuralgia (4) Headache (1) Not specified (21)	Percutaneous nucleotomy/tractotomy	95%
Kanpolat (2008b)[46]	11	TN (5) Oncologic (3) Glossopharyngeal Neuralgia (2) Geniculate Neuralgia (1)	Open DREZ nucleotomy/tractotomy	73%
Raslan (2008)[68]	10	Oncologic (10)	Percutaneous nucleotomy/tractotomy	80%
Chivukula (2015)[45]	16	TN (9) Postherpetic (3) Glossopharyngeal Neuralgia (3) Traumatic (1)	Open DREZ nucleotomy/tractotomy	100%
Rahimpour (2015)[32]	5	Not specified (5)	Endoscopic DREZ nucleotomy/tractotomy	80%
Patel (2018)[36]	5	Not specified (5)	Cervical SCS	100%
Velasquez (2018)[37]	12	Traumatic (12)	Cervical SCS	75%

660
661 Abbreviations: DREZ, dorsal root entry zone; MS; multiple sclerosis; SCS, spinal cord
662 stimulation; TN, Trigeminal Neuralgia/neuropathic pain.
663
664

665 **Table 3:** Composite pain reporting scale.

Composite pain score	Description
1	Pain free, VAS 0, 100% pain reduction
2	"Good" response, VAS 1-3, >50% pain reduction
3	"Satisfactory" or "fair" response, VAS 4-6, <50% pain reduction
4	No improvement, pain worsening, VAS 7-10

666   Abbreviations: VAS, visual analog scale.

Figure 1: Illustration of the dorsal aspect of the cervicomedullary junction, depicting the lesion site for nucleus caudalis interventions including DREZ nucleotomy/tractotomy, percutaneous nucleotomy/tractotomy, and cervical SCS. Abbreviations: DREZ, dorsal root entry zone; SCS, spinal cord stimulation.

673
Figure 2: Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)
flowchart outlining article selection. *Articles with fewer than 5 patients.

676
Figure 3: Cumulative number of studies examining nucleus caudalis targeting
interventions in the treatment of craniofacial pain. Over the last 80 years, 33 articles have
been published investigating techniques targeting the NC for treatment of craniofacial pain. The
1940s and 1970s saw the development of open (blue) and percutaneous approaches (orange)
respectively. The rate of publication continues to increase over time and novel techniques
continue to emerge including endoscopic approaches (light blue) and SCS (green) seen in the
last 10 years and ultrasonic (grey) previously. Abbreviations: DREZ, dorsal root entry zone; NC,
nucleus caudalis; SCS, spinal cord stimulation.
685

686
Figure 4: Distribution of preoperative diagnoses. Thirty-three articles reporting on 827
patients demonstrate a wide range of sources of craniofacial pain treated with NC interventions.
The most common being trigeminal neuralgia/neuropathic pain (n=343, 41.5%), oncological
(n=163, 19.7%), and postherpetic pain (n=70, 8.5%). Rarer causes of craniofacial pain are also
represented, such as headache (n=7, 0.8%), MS (n=6, 0.7%), and geniculate neuralgia (n=5,
0.6%). Abbreviations: MS, multiple sclerosis; NC, nucleus caudalis.

694
Figure 5: Distribution of intervention technique. The included studies encompass 5 main
intervention techniques targeting the NC. In order of decreasing prevalence: Open NC DREZ
nucleotomy/tractotomy (n=497, 60.1%), Percutaneous nucleotomy/tractotomy (n=294, 35.6%),
Cervical SCS (n=17, 2.1%), Ultrasonic nucleotomy (n=14, 1.7%), and Endoscopic DREZ
nucleotomy/tractotomy (n=5, 0.6). Abbreviations: DREZ, dorsal root entry zone; NC, nucleus
caudalis; SCS, spinal cord stimulation.
701

Figure 6: Post-intervention craniofacial pain outcome. Intervention efficacy was reported for
536 patients. A composite pain score was created to facilitate comparison between modern and
historical studies. Of those with reported outcome 44.2% (n=237) were pain free, 40.1% (n=215)
had a "good" response, 6.5% (n=35) had a "satisfactory" response, and 9.1% (n=49) had no
improvement or worsening pain. Abbreviation: VAS, visual analog scale.

Figure 7: Distribution of pain outcomes by intervention technique. The intervention types were reclassified as open focal lesioning techniques (nucleotomy/tractotomy) (n patients = 154), open DREZ (multipoint lesioning technique) (n patients = 141), percutaneous nucleotomy/tractotomy (n patients = 224), and cervical SCS (n patients = 17). Proportion of patients with a given outcome was reported with error bars representing 95% confidence intervals. Chi-squared demonstrates a significant difference in the outcome distribution between the intervention types (χ^2 = 57.5322, p <0.001). Abbreviations: DREZ, dorsal root entry zone; Perc, percutaneous; SCS, spinal cord stimulation.

719
Figure 8: Proportion of patients with a given complication arranged by intervention technique. All complications except corneal anesthesia (n = 5), oropharyngeal dysfunction (n = 11), and limb dysfunction (n = 14) demonstrate a statistically significant difference in distribution between intervention types. Proportion of patients with a given complication is reported with error bars representing 95% confidence intervals. See Supplemental Material 4 for data and statistical results. Abbreviations: DREZ, dorsal root entry zone; Perc, percutaneous; SCS, spinal cord stimulation.