

15 **Abstract**

16 **Background**

17 Leptospirosis is an emerging zoonosis of global importance. In South Africa, the
18 infection is an underreported public health concern, with limited information on its
19 incidence and distribution. This study aimed to determine the incidence of human
20 leptospirosis from 2010 to 2019 in Western Cape Province (WCP), and to compare
21 the incidence based on seasonal and demographic factors.

22 **Methods**

23 A retrospective study was conducted with data on leptospirosis diagnoses by sex, age,
24 season, and year in WCP obtained from the National Health Laboratory Services. With
25 the provincial population sizes as the denominator, the incidence of leptospirosis was
26 estimated and expressed as cases per 100,000 population. Negative binomial
27 regression was used to estimate the effect of sex, season, and year on the incidence
28 of leptospirosis.

29 **Results**

30 A total of 254 cases of leptospirosis were reported between 2010 and 2019, with the
31 highest number of cases being in 2015. The annual incidence ranged between 0.15
32 and 0.66/100,000 population with a 10-year average incidence of 0.40/100,000
33 population. The incidence was higher among males than in females (0.55 vs. 0.25/

34 100,000 population; incidence rate ratio (IRR) 2.2, 95% CI: 1.66,3.03). The 18-44 age
35 cohort and had the highest average incidence (0.56/100,000 population), while the
36 ≤ 17 age cohort had the lowest incidence (0.07/100,000 population). The 18-44 (IRR
37 8.0, 95% CI: 4.65,15.15) and ≥ 45 (IRR 7.4, 95% CI: 4.17,14.17) age cohorts were
38 more at risk of infection compared to ≤ 17 age cohort. The average incidence of the
39 infection was similar among seasons and there was no significant association between
40 season and incidence of leptospirosis.

41 **Conclusions**

42 The results highlight that leptospirosis is an important zoonosis within the province
43 disproportionately affecting males and the productive age demographic groups. These
44 findings should enhance targeted prevention and provoke further investigation on the
45 importance of environmental and socioeconomic factors on leptospirosis burden.

46 **Author Summary**

47 Leptospirosis is an emerging zoonotic bacterial disease of global importance. Despite
48 its wide distribution, the disease is largely underestimated because its clinical
49 manifestations mimic certain commonly known febrile illnesses such as malaria,
50 influenza, hepatitis, and yellow fever. Leptospirosis burden in South Africa has been
51 suggested to be moderately high however studies on the burden of the infection are
52 lacking. This study sought to determine the incidence and trends of leptospirosis in the
53 Western Cape Province, South Africa between 2010 and 2019. Overall, leptospirosis
54 average incidence was 0.4 cases per 100,000 population (ranging from 0.15 to 0.66
55 cases per 100,000 population). Leptospirosis incidence was highest among male
56 individuals and among those who were aged 18-years and above, however, the
57 incidence did not differ by seasons. These estimates highlight that leptospirosis is an
58 important zoonotic disease within the province and potentially disproportionately
59 affecting males and productive age demographic groups. Therefore, this indicates the
60 need for an all-encompassing One Health Approach to obtain all relevant information
61 concerning leptospirosis distribution and risk factors in South Africa and in Africa at
62 large to effectively enhance leptospirosis prevention strategies.

63 1. Introduction

64 Leptospirosis is an emerging, global zoonosis with an estimated 1.03 million cases
65 and 58,900 deaths occurring annually [1–5]. The disease is caused by pathogenic
66 spirochetes *Leptospira* species currently reported to comprise 66 different species with
67 more than 300 serovars [6]. The infection in humans is associated with varying clinical
68 manifestations ranging from a mild self-limiting febrile illness to severe illness
69 characterized by dysfunction of multiple organs such as liver, kidneys, lungs, and the
70 brain [7] potentially leading to pulmonary hemorrhagic syndrome and acute kidney
71 injury due to Weil's disease [5,8–10] . In addition, leptospirosis presents with
72 symptoms that can mimic commonly known infections that cause febrile illnesses such
73 as malaria, influenza, hepatitis, yellow fever, and viral hemorrhagic diseases among
74 others [11,12]. Consequently, the infection has often been underdiagnosed and
75 underreported [11]. Despite the life-threatening nature of leptospirosis, there is little
76 published data on morbidity associated with the infection [4,5,13] contributing to its
77 neglected status.

78 The risk of leptospirosis infection among humans occurs either through direct
79 exposure to urine or aborted tissues of wild or domestic reservoir animals such as
80 rodents and livestock/pets [4,14], or through indirect exposure such as contact with
81 contaminated water, soil, and food [3,14]. In South Africa, a recent prison outbreak of
82 leptospirosis was associated with exposure of inmates to rat urine in an overcrowded
83 prison. Due to the predominant modes of transmission, leptospirosis is regarded as
84 an occupational hazard affecting mainly farmers, sewer workers, veterinarians, and
85 military personnel [11,15–18]. However, the transmission patterns are changing

86 because there are increasing reports of leptospirosis in the general population. These
87 cases are due to recreational exposures such as camping, kayaking, adventure
88 travelling, hiking, cave exploration and other activities done in infected water
89 [11,16,18,19]. Due to the diverse avenues through which humans can be exposed to
90 the leptospires, occurrence of leptospirosis within the general population requires
91 consideration [7,20].

92 Globally, tropical regions as well as resource limited settings are reported to have a
93 higher burden of leptospirosis compared to temperate regions [5]. However, in Africa,
94 few countries have data and reports on human leptospirosis [15,20] and research
95 studies are scarce [2,5]. A previously published systematic review, of peer reviewed
96 studies conducted in Africa, reported a prevalence of acute human leptospirosis
97 ranging from 2.3% to 19.8% among hospital patients presenting with febrile illness and
98 an estimated total of 750,000 cases per annum [15]. It was further concluded in this
99 study that the morbidity of human leptospirosis in Africa is likely to be high relative to
100 other global regions [15]. The need of more studies on the occurrence of leptospirosis
101 in Africa was emphasized in another systematic review by De Vries et.al [20] to reliably
102 understand the extent of the problem.

103 The incidence of human leptospirosis in South Africa has been suggested to be
104 moderately high within the population [21–23]. Findings about the continued
105 circulation of highly pathogenic *Leptospira* spp. among rodents suggests that the
106 infection maybe an important underreported public health concern in the country
107 [21,22]. A study conducted between 2009 and 2011 reported a seroprevalence that
108 ranged from 9% to 12.5% among all clinical samples sent to the NHLS for IgM ELISA

109 testing from all over the country [23]. Environmental conditions in Western Cape South
110 Africa, (the coastal, and temperate conditions with wet winters and warmer summers)
111 have been suggested to favor transmission of pathogenic *leptospira* species *L.*
112 *borgpetersenii* and *L. interrogans* traditionally associated with rats [4,21,22]. The most
113 recent retrospective study conducted between 2005-2015, focused on patients
114 presenting at a tertiary referral hospital in Western Cape Province (WCP), and a
115 seroprevalence of 20% was reported [24].

116 Studies outside Africa have highlighted the influence of season, gender and age on
117 the distribution and occurrence of human leptospirosis [25–27]. Such associations
118 have scarcely been studied in Africa despite changes in climatic factors, demographic
119 shifts, urbanization, and globalization [2,11,20,28–30]. The risk conferred by rodent
120 infestation and overcrowding have been demonstrated in a recent leptospirosis
121 outbreak that occurred in a prison in WCP, South Africa. [22]. Despite this information,
122 studies describing leptospirosis occurrence in relation to demographic and seasonal
123 factors are lacking in WCP.

124 To address these knowledge gaps, a retrospective study on human leptospirosis
125 cases in the WCP, South Africa was conducted using data generated from diagnostic
126 tests done in a public health care setting in Cape Town, South Africa. The study aimed
127 at determining the incidence and trends of human leptospirosis over a 10-year period
128 (2010-2019), with respect to year and seasonality and to describe the demographic
129 characteristics of leptospirosis cases.

130 **2. Methods**

131 **2.1 Study setting**

132 WCP is located on the south-western coast of South Africa and is bordered by the
133 Northern Cape and Eastern Cape Provinces. WCP has a population of approximately
134 seven million inhabitants [31], three quarters of whom use public-sector healthcare
135 services [32]. The province is approximately 129,462 km² and has a population density
136 of 45 inhabitants per Km² [31]. Administratively, the province is divided into one
137 metropolitan municipality (City of Cape Town) and five district municipalities (Central
138 Karoo, Garden Route, Overberg, West Coast and Cape Winelands) [31]. The province
139 has a diverse climate but is dominated by a Mediterranean climate with a cool, wet
140 winter and a warm, dry summer. The inland daily maximum temperatures range from
141 20 °C in winter to 32 °C in summer and the mean annual rainfall is less than 380 mm.

142 **2.2 Study design and study population**

143 A retrospective study was conducted by using data provided by the data warehouse
144 of the National Health Laboratory Services (NHLS), on all human leptospirosis tests
145 conducted at public healthcare facilities in the WCP, South Africa. The NHLS does
146 serological testing on all serum samples of patients who are clinically suspected to be
147 having a leptospirosis infection. Specimens from suspected patients are collected from
148 different public healthcare facilities within the province and sent to a central NHLS
149 laboratory in Cape Town. In this analysis, data included are from patients who had
150 laboratory confirmed leptospirosis between 1st January 2010 and 31st December 2019.

151 **2.3 Data collection**

152 Human leptospirosis cases were considered for all ELISA IgM serological tests that
153 were positive for Leptospirosis during the study period (2010 to 2019). ELISA IgM
154 testing is the serological test conducted on all samples submitted to NHLS for
155 leptospirosis screening and is sensitive in detecting new onset of illness [11]. Patient
156 data extracted included the patient's age, sex, year, date of test (year, month, and
157 day) and the name of the health facility that submitted the specimen for testing.

158 The population sizes based on year, sex, and age in WCP for the 10-year period were
159 extracted from the Western Cape Department of Health population circular H102 of
160 2020 [33]. This data was used as the denominator when calculating the incidence of
161 leptospirosis infection.

162 **2.4 Statistical analysis**

163 All human leptospirosis cases were tabulated according to sex, age, season, and year
164 of occurrence (2010-2019) and their frequency and proportions were calculated. The
165 cases were not grouped based on their geographical location (district municipality)
166 because the patient address information was not part of received data. Furthermore,
167 the location of the health facility that submitted the specimen could not be assumed to
168 be the same as the patient's location because of the possibility of patients seeking
169 healthcare or being hospitalized in a facility outside of their actual district of residence.
170 The date when the test was done was used to assign the case to a particular year and
171 season. Season was categorized as Summer (1st December-28/29th February),

172 Autumn/Fall (1st March-31st May), Winter (1st June-31st August), and Spring (1st
173 September-30th November). The variable year was conceptualized as a consecutive
174 12-month period from 1st January to 31st December, thus there were ten categories for
175 year (2010 to 2019).

176 The incidence proportion of human leptospirosis for each year (2010-2019) was
177 estimated and expressed as leptospirosis cases per 100,000 population. The
178 incidence proportion by sex and age group as well as for each season of the year was
179 also estimated. The Kruskal-Wallis Rank Sum Test was used to compare the average
180 annual incidence between seasons of the year, age groups and between years. The
181 Wilcoxon Rank Sum Exact Test was used to compare the average annual incidence
182 between sex demographics. The negative binomial regression was used to estimate
183 the effect of sex, gender, year of occurrence and season of the year on the incidence
184 of human leptospirosis over the study period. This is a suitable regression model to
185 use instead of the traditional Poisson regression in situations where modelling involves
186 a count variable that is over-dispersed (the mean is less than the variance). The results
187 were presented as incidence rate ratios (IRR) with 95% confidence intervals (CI). The
188 cut-off value for statistical significance was 0.05. All statistical analysis was conducted
189 using R software version 1.2.5033.

190 **2.5 Ethical considerations**

191 Ethical approval was obtained from the University of Cape Town Human Research
192 Ethics Committee (UCT-HREC) (reference number: HREC REF: 303/2021). The
193 Western Cape Provincial approval was granted by Groote Schuur Hospital Research
194 Ethics Committee. The approval to use data from NHLS database was given by the
195 NHLS Academic Affairs and Research office (reference number: PR2118953).

196 **3. Results**

197 **3.1 Distribution of leptospirosis cases in WCP, 2010-2019**

198 During the 10-year study period (2010-2019), a total of 254 cases of human
199 leptospirosis were recorded by the National Health Laboratory Services (NHLS)
200 (**Table 1**Error! Reference source not found.) for persons in the Western Cape
201 Province, South Africa. Most of the cases reported had their specimen submitted from
202 healthcare facilities located within the City of Cape Town Metropolitan (S2 Table 1).
203 The highest number of cases was recorded in 2015 (42 cases, 16.5%) and lowest in
204 2012 (9 cases, 3.5%). When grouped by month of occurrence, the month of March (30
205 cases) had the highest number of cases over the 10-year period while September (16
206 cases) had the lowest number of the cases (**Fig 1**). When grouped into seasons, more
207 cases were observed in fall season (75 cases), followed by summer, spring, and winter
208 (64, 62, 53 cases respectively) (**Table 1**). More males were infected than females, with
209 the males accounting for 173 (68.1%) cases compared to 81 (31.9%) cases who were
210 females, and the overall male to female ratio was 2.136:1 (**Table 1**). The median age


211 of the observed cases was 37.0 (Inter Quantile Range (IQR) 28.0-48.0) years, with a
212 mean age of 38.12 years. Overall, 163 cases (64.2 %) were in the 18–44-year-old age
213 group, 78 cases (30.7%) were in ≥ 45 -year-old age group while 13 cases (5.1%) were
214 in the ≤ 17 -year-old age group over the 10-year study period (**Table 1**).

215 **Table 1: Distribution of leptospirosis cases by age, sex, season, and year of**
 216 **occurrence in Western Cape Province South Africa (2010-2019)**

217

Variables	All Cases (n=254) n (%)
Age (years) median (IQR)	37.0 (28.0,48.0)
Age group	
≤17	13 (5.1)
18-44	163 (64.2)
≥ 45	78 (30.7)
Sex	
Male	173 (68.1)
Female	81 (31.9)
Season of the Year	
Summer	64 (25.2)
Fall	75 (29.5)
Winter	53 (20.9)
Spring	62 (24.4)
Year of occurrence	
2010	34 (13.4)
2011	19 (7.5)
2012	9 (3.5)
2013	12 (4.7)
2014	28 (11.0)
2015	42 (16.5)
2016	37 (14.6)
2017	22 (8.7)
2018	31 (12.2)
2019	20 (7.9)


218 IQR: Inter Quantile Range; n: frequency.



219 **Fig 1: Monthly distribution of leptospirosis cases in Western Cape Province over**
220 **a 10-year period (2010-2019)**

221 **3.2 Incidence of leptospirosis in the WCP, 2010-2019**


222 The incidence of leptospirosis fluctuated widely across the 10 years. The annual
223 incidence ranged from 0.15 to 0.66 cases per 100,000 population, with an average
224 annual incidence of 0.40 cases per 100,000 population. The annual number of cases
225 and the incidence are plotted against the year of occurrence to show the trends across
226 the years (**Fig 2**). The incidence followed a downward trend between 2010 (0.59 cases
227 per 100,000 population) and 2012 (0.15 cases per 100,000 population). However,
228 there was an increase in the incidence from 2013 (0.20 cases per 100,000 population)
229 which peaked in 2015 (0.66 cases per 100,000 population). This was followed by a
230 gradual decrease in the incidence between 2015 and 2019 (0.66-0.27 cases per
231 100,000 population) except for year 2018 where the annual incidence was higher than
232 the one observed in 2017.




233 Fig 2: The number of cases and incidence of leptospirosis per year (2010 to
234 2019) in Western Cape Province, South Africa.

235 Throughout the study period, the incidence was highest among males compared to
236 females (**Fig 3**), with the average incidence among males being 0.55 cases per
237 100,000 population and 0.25 cases per 100,000 population among females (p-
238 values=0.004). During the entire study period, the incidence was highest in the 18-44-
239 year-old and ≥ 45 years-old age groups compared to the ≤ 17 -year-old age group (**Fig**
240 **4** and **Error! Reference source not found.**). The average incidence differed
241 substantially between the age groups with the highest average incidence being among
242 the 18-44-year-old age group (0.56 cases per 100,000 population) over the 10-year
243 period, followed by ≥ 45 years-old age group (0.49 cases per 100,000 population) and
244 the ≤ 17 -year-old age group had the lowest incidence (0.07 cases per 100,000
245 population) (**Error! Reference source not found.**).

246 The results indicate that on average, the annual incidence was higher during the fall
247 season (0.12 cases per 100,000 population) and lowest in the winter season (0.08
248 cases per 100,000 population) over the 10-year period (**Table 2**). However, the
249 observed difference in the average incidence of leptospirosis between the seasons of
250 the year was not statistically significant.



251 **Fig 3: Incidence of leptospirosis in males and females from 2010 to 2019 in**
252 **Western Cape Province.**



253 **Fig 4: Incidence of leptospirosis by age group from 2010 to 2019 in Western**
254 **Cape Province, South Africa.**

255 **Table 2: Seasonal incidence of leptospirosis for the years 2010-2019, in Western**
256 **Cape Province, South Africa.**

Year	Summer	Fall	Winter	Spring
	IP/100,000 (95% CI)	IP/100,000 (95% CI)	IP/100,000 (95% CI)	IP/100,000 (95% CI)
2010	0.10 (0.02,0.19)	0.28 (0.14,0.41)	0.12 (0.03,0.21)	0.09(0.01,0.16)
2011	0.12 (0.03,0.21)	0.03 (-0.01,0.08)	0.07 (0.001,0.13)	0.10 (0.02,0.18)
2012	0.07 (0.001,0.13)	0.03 (-0.01,0.08)	0.02 (-0.02,0.05)	0.03 (-0.01,0.08)
2013	0.05 (-0.01,0.10)	0.05 (-0.01,0.10)	0.07 (0.001,0.13)	0.03 (-0.01,0.08)
2014	0.02 (-0.01,0.05)	0.10 (0.02,0.17)	0.14 (0.05,0.24)	0.19 (0.08,0.30)
2015	0.20 (0.09,0.31)	0.17 (0.07,0.27)	0.16 (0.06,0.25)	0.13 (0.04,0.21)
2016	0.18 (0.08,0.29)	0.21 (0.10,0.33)	0.03 (-0.01,0.07)	0.14 (0.05,0.23)
2017	0.12 (0.03,0.18)	0.09 (0.02,0.16)	0.08 (0.01,0.14)	0.06 (0.001,0.12)
2018	0.12 (0.04,0.20)	0.15 (0.06,0.24)	0.09 (0.02,0.16)	0.10 (0.03,0.18)
2019	0.04(-0.01,0.09)	0.07 (0.01,0.14)	0.07 (0.01,0.14)	0.10 (0.03,0.18)
Average incidence	0.10 (0.03,0.18)	0.12 (0.04,0.20)	0.08 (0.02,0.15)	0.10 (0.02,0.17)

257 IP: incidence proportion; CI: confidence interval

258 **3.3 Effect of sex, age, year, and season on leptospirosis**

259 **incidence in WCP, 2010-2019**

260 **Table 3** shows the results from the Multivariable Negative Binomial Regression
261 (MNBR) model that was run to identify factors associated with the incidence of
262 leptospirosis during the 10-year period. The incidence rate of leptospirosis among the
263 male demographic was 2.2 (95% CI: 1.66,3.03) times higher than among females
264 during the entire study period. The model results indicated that on average the
265 incidence rate of leptospirosis among 18-44 and ≥ 45 age groups was 8.0 (95% CI:
266 4.65,15.15) and 7.4 (95% CI: 4.17,14.17) times higher than the incidence rate among
267 ≤ 17 years age group. The incidence was substantially lower in 2012 (IRR 0.25; 95%
268 CI 0.11-0.54), 2013 (IRR 0.34; 95% CI 0.16-0.68), 2017 (IRR 0.53; 0.28-0.98) and
269 2019 (IRR 0.47; 95% CI 0.24-0.87) compared to the incidence in 2010. From the model
270 results, the incidence rate in fall, summer and spring seasons was 1.4 (95% CI:
271 0.95,2.15), 1.2 (95% CI: 0.78,1.81) and 1.2 (95% CI: 0.77,1.78) times higher than what
272 was observed in winter season, however the observed difference was not statistically
273 significant.

274 **Table 3: MNBR analysis results between year, season, sex, and age group with**
275 **incidence of leptospirosis**

Variables	IRR	95% CI	P.value
Year of occurrence			
2010	Ref	–	–
2011	0.543	(0.281,1.026)	0.061
2012	0.25	(0.106,0.536)	0.001
2013	0.337	(0.157,0.684)	0.003
2014	0.73	(0.402,1.314)	0.294
2015	1.033	(0.597,1.796)	0.907
2016	0.956	(0.546,1.677)	0.874
2017	0.53	(0.282,0.98)	0.046
2018	0.748	(0.418,1.334)	0.323
2019	0.466	0.244,0.872	0.019
Season of the year			
Winter	Ref	–	–
Spring	1.166	(0.765,1.782)	0.475
Summer	1.184	(0.778,1.808)	0.431
Fall/autumn	1.422	0.947,2.147	0.091
Sex			
Female	Ref	–	–
Male	2.239	(1.664,3.034)	<0.001
Age group			
≤17	Ref	–	–
18-44	8.045	(4.654,15.147)	<0.001
≥45	7.391	(4.167,14.169)	<0.001

276 IRR: incidence rate ratio; CI: confidence interval; ref: reference category; MNBR:
277 Multivariable Negative Binomial Regression.

278 4. Discussion

279 This study provides a first description of the incidence of human leptospirosis in the
280 Western Cape Province (WCP), South Africa across a 10-year period. Here we report
281 an average incidence of 0.40 leptospirosis cases per 100,000 population, ranging from
282 0.15 in 2012 to 0.66 in 2015 cases per 100,000. There was no overall increase in the
283 annual incidence of leptospirosis, however the incidence fluctuated widely across the
284 10-year period. The incidence was significantly higher in males compared to female.
285 The 18-44-year-old and ≥ 45 years-old age groups had a higher incidence of the
286 infection compared to the ≤ 17 -year-old age group. The incidence did not vary based
287 on season over the 10-year period.

288 The annual incidence observed during this 10-year period lies within the range of the
289 incidence that has been reported globally to be occurring in temperate regions (0.1-1
290 cases per 100,000 population) [34]. The cold wet winters and warm dry summers
291 might allow transmission and survival of pathogenic *Leptospira* species [21]. WCP is
292 known for its harsh drought, and different types of floods such as river floods, flash
293 floods and these occur in each year at varying intensities [35]. Annual fluctuation in
294 such environmental conditions could partly explain the observed wide fluctuation in
295 the annual incidence of leptospirosis because increase in incidence is normally
296 reported after occurrence of extreme weather events such as heavy rains and high
297 temperatures [29,34,36]. In addition, rodent infestation, increased prevalence of
298 infected rodents and overcrowded settings were reported as risk factors for human
299 leptospirosis outbreak in a one of the prisons in WCP in 2015 [22]. This outbreak could
300 have contributed to the observed increase in the incidence in 2015. In a study

301 conducted after this outbreak, Naidoo et al. further highlighted the importance of
302 continued circulation of pathogenic *leptospira spp* among rodents in maintaining the
303 status of leptospirosis transmission particularly in informal settlements [22].

304 Previous studies have indicated a significant relationship between leptospirosis
305 incidence and seasonality [34,36,37]. In Western Cape, seasonality was not
306 associated with the incidence of leptospirosis between 2010 and 2019, however the
307 incidence was slightly higher in fall and summer as compared to winter and spring.
308 This is consistent with some reports from other countries in temperate regions [34,38].
309 Studies on correlation between leptospirosis incidence and seasonality require
310 consideration of other variables such as rodent seasonal population, temperature,
311 rainfall, and other climate related parameters to be included in the analysis [37]. This
312 could help in understanding the importance of the interaction of different environmental
313 and climatic factors in determining the incidence of leptospirosis in resource limited
314 settings.

315 The results showed that the overall incidence was significantly higher in males as
316 compared to females (0.55 cases vs 0.25 cases per 100,000 population) during the
317 study period. Similar results have been reported in both seroprevalence studies in
318 Africa and in incidence studies conducted using surveillance data globally
319 [26,27,34,38,39]. The higher incidence among males has been largely attributed to
320 occupational or environmental exposure, whereby males engage in activities that may
321 put them at higher risk of contracting the infection [36,40]. Epidemiological studies
322 collecting data on occupation, underlying medical conditions, and place of settlement

323 could aid in categorically establishing risk groups hence guiding policies on targeted
324 prevention of leptospirosis.

325 The incidence of leptospirosis was highest among those within the 18-44 and ≥ 45 -
326 year-old age groups compared to those below ≤ 17 years of age. These findings are in
327 line with a systematic review by Costa et al. that reported the highest incidence to be
328 occurring among adult males aged 20-49 years [5]. The observed results also
329 correspond with other studies, where incidence has been shown to be higher among
330 the active adult population [34,38]. People belonging to the 18-44 and ≥ 45 age groups
331 would theoretically have an increased environmental exposure compared to those
332 belonging to ≤ 17 age group, thus increasing their risk of being infected.

333 The data presented here are the first results to inform on the incidence trends and
334 distribution of leptospirosis in WCP. The data covers a 10-year period, hence could
335 help to guide designing of preventive strategies. However, there are some study
336 limitations that need to be considered. Firstly, the study utilized data collected from
337 passive surveillance and from a public health laboratory setting excluding data from
338 private laboratories, hence there is a great potential for underestimation of the real
339 occurrence of leptospirosis in WCP. Some bias may still be inherent in the analysis
340 due to ambiguity in clinical presentation; the latter contributes to underestimation of
341 actual incidence. Secondly, the analysis does not provide information on the
342 geographical distribution of the cases within the province as such information was not
343 included in provided data. Consideration of the geographical distribution of the
344 incidence would have given insights into which specific districts are at risk, hence
345 informing targeted interventions.

346 Findings from this analysis highlight the continued circulation of leptospirosis infection
347 within WCP with age and sex being significant risk factors for infection. However, there
348 is need of a good knowledge about the epidemiology of leptospirosis (such as the
349 geographical and seasonal patterns, the specific risk populations, circulating
350 *Leptospira* strains and the importance of reservoir animals) [13] within the province to
351 improve prevention strategies, prediction, and detection of leptospirosis burden and
352 outbreaks. For instance, future studies focusing on the impact of heavy rainfall, floods,
353 seasonal fluctuation in weather-related factors, occupation, geographical location, and
354 rodent dynamic parameters on leptospirosis incidence could help address the
355 knowledge gaps on the actual burden of this emerging zoonosis.

356 In conclusion, the incidence of leptospirosis in WCP fluctuated within the range that
357 has been reported to be occurring in temperate regions. The incidence was strongly
358 related to sex and age; however, it did not differ across the seasons of the year. These
359 results show that leptospirosis is an important zoonosis within the province and
360 potentially disproportionately affecting males and the productive age demographic
361 groups. The findings, therefore, can guide targeted intervention strategies within the
362 province to decrease the burden of human leptospirosis.

363 **5. Acknowledgement**

364 Special thanks to the National Health Laboratory Services (NHLS), South Africa for
365 providing all the necessary data on leptospirosis tests conducted in Western Cape
366 Province.

367 6. References

- 368 [1] Bharti AR, Nally JE, Ricaldi JN, Matthias MA, Diaz MM, Lovett MA, et al.
369 Leptospirosis: A zoonotic disease of global importance. *Lancet Infectious*
370 *Diseases* 2003;3:757–71. [https://doi.org/10.1016/S1473-3099\(03\)00830-2](https://doi.org/10.1016/S1473-3099(03)00830-2).
- 371 [2] Pappas G, Papadimitriou P, Siozopoulou V, Christou L, Akritidis N. The
372 globalization of leptospirosis: worldwide incidence trends. *International Journal*
373 *of Infectious Diseases* 2008;12:351–7.
374 <https://doi.org/10.1016/j.ijid.2007.09.011>.
- 375 [3] World Health Organisation. Human leptospirosis guidelines for Diagnosis,
376 Surveillance, and Control. International Leptospirosis Society Geneva 2003.
377 Available from: <https://apps.who.int/iris/handle/10665/42667> (accessed
378 November 11, 2021).
- 379 [4] Cosson J-F, Picardeau M, Mielcarek M, Tatarski C, Chaval Y, Suputtamongkol
380 Y, et al. Epidemiology of *Leptospira* Transmitted by Rodents in Southeast
381 Asia. *PLoS Neglected Tropical Diseases* 2014;8:e2902.
382 <https://doi.org/10.1371/journal.pntd.0002902>.
- 383 [5] Costa F, Hagan JE, Calcagno J, Kane M, Torgerson P, Martinez-Silveira MS,
384 et al. Global Morbidity and Mortality of Leptospirosis: A Systematic Review.
385 *PLoS Neglected Tropical Diseases* 2015;9.
386 <https://doi.org/10.1371/journal.pntd.0003898>.
- 387 [6] Caimi K, Ruybal P. *Leptospira* spp., a genus in the stage of diversity and
388 genomic data expansion. *Infection, Genetics and Evolution : Journal of*

- 389 Molecular Epidemiology and Evolutionary Genetics in Infectious Diseases
390 2020;81. <https://doi.org/10.1016/J.MEEGID.2020.104241>.
- 391 [7] Haake DA, Levett PN. Leptospirosis in Humans, 2015.
392 https://doi.org/10.1007/978-3-662-45059-8_5.
- 393 [8] Gouveia EL, Metcalfe J, de Carvalho ALF, Aires TSF, Villasboas-Bisneto JC,
394 Queirroz A, et al. Leptospirosis-associated Severe Pulmonary Hemorrhagic
395 Syndrome, Salvador, Brazil. Emerging Infectious Diseases 2008;14.
396 <https://doi.org/10.3201/eid1403.071064>.
- 397 [9] Papa A, Theoharidou D, Antoniadis A. Pulmonary Involvement and
398 Leptospirosis, Greece. Emerging Infectious Diseases 2009;15.
399 <https://doi.org/10.3201/eid1505.080270>.
- 400 [10] Lameire NH, Bagga A, Cruz D, de Maeseneer J, Endre Z, Kellum JA, et al.
401 Acute kidney injury: an increasing global concern. The Lancet 2013;382.
402 [https://doi.org/10.1016/S0140-6736\(13\)60647-9](https://doi.org/10.1016/S0140-6736(13)60647-9).
- 403 [11] Clem A, Izurieta R, Galwankar S. Leptospirosis: The “mysterious” mimic.
404 Journal of Emergencies, Trauma and Shock 2008;1:21.
405 <https://doi.org/10.4103/0974-2700.40573>.
- 406 [12] Crump JA, Morrissey AB, Nicholson WL, Massung RF, Stoddard RA, Galloway
407 RL, et al. Etiology of Severe Non-malaria Febrile Illness in Northern Tanzania:
408 A Prospective Cohort Study. PLoS Neglected Tropical Diseases 2013;7.
409 <https://doi.org/10.1371/journal.pntd.0002324>.

- 410 [13] Goarant C. Leptospirosis: risk factors and management challenges in
411 developing countries. *Research and Reports in Tropical Medicine* 2016;7:49.
412 <https://doi.org/10.2147/RRTM.S102543>.
- 413 [14] Levett PN. Leptospirosis. *Clinical Microbiology Reviews* 2001;14:296–326.
414 <https://doi.org/10.1128/CMR.14.2.296-326.2001>.
- 415 [15] Allan KJ, Biggs HM, Halliday JEB, Kazwala RR, Maro VP, Cleaveland S, et al.
416 Epidemiology of Leptospirosis in Africa: A Systematic Review of a Neglected
417 Zoonosis and a Paradigm for ‘One Health’ in Africa. *PLoS Neglected Tropical*
418 *Diseases* 2015;9. <https://doi.org/10.1371/journal.pntd.0003899>.
- 419 [16] Davis CP. Leptospirosis Vaccine, Symptoms, Treatment & Diagnosis 2021.
420 <https://www.medicinenet.com/leptospirosis/article.htm> (accessed January 25,
421 2021).
- 422 [17] Levett PN. Leptospirosis: A forgotten zoonosis? *Clinical and Applied*
423 *Immunology Reviews* 2004;4. <https://doi.org/10.1016/j.cair.2004.08.001>.
- 424 [18] Musso D, la Scola B. Laboratory diagnosis of leptospirosis: A challenge.
425 *Journal of Microbiology, Immunology and Infection* 2013;46:245–52.
426 <https://doi.org/10.1016/j.jmii.2013.03.001>.
- 427 [19] Monahan AM, Miller IS, Nally JE. Leptospirosis: risks during recreational
428 activities. *Journal of Applied Microbiology* 2009;107:707–16.
429 <https://doi.org/10.1111/J.1365-2672.2009.04220.X>.
- 430 [20] de Vries SG, Visser BJ, Nagel IM, Goris MGA, Hartskeerl RA, Grobusch MP.
431 Leptospirosis in Sub-Saharan Africa: A systematic review. *International*

- 432 Journal of Infectious Diseases 2014;28:e47–64.
433 <https://doi.org/10.1016/j.ijid.2014.06.013>.
- 434 [21] Moseley M, Naidoo K, Bastos A, Retief L, Freaan J, Telfer S, et al. Multi-locus
435 sequence analyses reveal a clonal *L. borgpetersenii* genotype in a
436 heterogeneous invasive *Rattus* spp. community across the City of
437 Johannesburg, South Africa. *Parasites & Vectors* 2020 13:1 2020;13:1–9.
438 <https://doi.org/10.1186/S13071-020-04444-0>.
- 439 [22] Naidoo K, Moseley M, McCarthy K, Chingonzoh R, Lawrence C, Setshedi GM,
440 et al. Fatal rodentborne leptospirosis in prison inmates, South Africa, 2015.
441 *Emerging Infectious Diseases* 2020;26:1033–55.
442 <https://doi.org/10.3201/eid2605.191132>.
- 443 [23] Saif A, Freaan J, Rossouw J, Trataris AN. Leptospirosis in South Africa.
444 *Onderstepoort Journal of Veterinary Research* 2012;79.
445 <https://doi.org/10.4102/ojvr.v79i2.478>.
- 446 [24] Mteshana Z. Epidemiology of leptospirosis in Groote Schuur Hospital
447 [dissertation]. 2020.
- 448 [25] Dhewantara PW, Mamun A al, Zhang WY, Yin WW, Ding F, Guo D, et al.
449 Geographical and temporal distribution of the residual clusters of human
450 leptospirosis in China, 2005–2016. *Scientific Reports* 2018;8.
451 <https://doi.org/10.1038/s41598-018-35074-3>.
- 452 [26] Benacer D, Thong KL, Min NC, Verasahib K bin, Galloway RL, Hartskeerl RA,
453 et al. Epidemiology of human leptospirosis in Malaysia, 2004-2012. *Acta
454 Tropica* 2016;157:162–8. <https://doi.org/10.1016/j.actatropica.2016.01.031>.

- 455 [27] Vieira ML, Gama-Simões MJ, Collares-Pereira M. Human leptospirosis in
456 Portugal: a retrospective study of eighteen years. *International Journal of*
457 *Infectious Diseases* 2006;10:378–86. <https://doi.org/10.1016/j.ijid.2005.07.006>.
- 458 [28] Durski KN, Jancloes M, Chowdhary T, Bertherat E. A global, multi-disciplinary,
459 multi-sectorial initiative to combat leptospirosis: Global Leptospirosis
460 Environmental Action Network (GLEAN). *International Journal of*
461 *Environmental Research and Public Health* 2014;11:6000–8.
462 <https://doi.org/10.3390/ijerph110606000>.
- 463 [29] Lau CL, Smythe LD, Craig SB, Weinstein P. Climate change, flooding,
464 urbanisation and leptospirosis: Fuelling the fire? *Transactions of the Royal*
465 *Society of Tropical Medicine and Hygiene* 2010;104:631–8.
466 <https://doi.org/10.1016/j.trstmh.2010.07.002>.
- 467 [30] Munoz-Zanzi C, Groene E, Morawski BM, Bonner K, Costa F, Bertherat E, et
468 al. A systematic literature review of leptospirosis outbreaks worldwide, 1970-
469 2012 2020. <https://doi.org/10.26633/RPSP.2020.78>.
- 470 [31] Municipalities of South Africa [Internet]. Western Cape Municipalities. [Cited
471 2021 Oct 26] 2021. Available from:
472 <https://municipalities.co.za/provinces/view/9/western-cape>.
- 473 [32] Boulle A, Heekes A, Tiffin N, Smith M, Mutemaringa T, Zinyakatira N, et al.
474 Data Centre Profile: The Provincial Health Data Centre of the Western Cape
475 Province, South Africa. *International Journal of Population Data Science*
476 2019;4. <https://doi.org/10.23889/IJPDS.V4I2.1143>.

- 477 [33] Western Cape DoH. Department of Health, Population Circular H102 of 2020
478 2020. Available from:
479 [https://www.westerncape.gov.za/assets/departments/health/h_102_2020_covi](https://www.westerncape.gov.za/assets/departments/health/h_102_2020_covid-19_population_data.pdf)
480 [d-19_population_data.pdf](https://www.westerncape.gov.za/assets/departments/health/h_102_2020_covid-19_population_data.pdf).
- 481 [34] Zhang H, Zhang C, Zhu Y, Mehmood K, Liu J, McDonough SP, et al.
482 Leptospirosis trends in China, 2007–2018: A retrospective observational study.
483 *Transboundary and Emerging Diseases* 2020;67:1119–28.
484 <https://doi.org/10.1111/tbed.13437>.
- 485 [35] Holloway Ailsa, Fortune Gillian, Chasi Vimbai. RADAR Western Cape 2010
486 Risk and Development Annual Review Disaster Mitigation for Sustainable
487 Livelihoods Programme 2010.
- 488 [36] Mohan ARM, Cumberbatch A, Adesiyun AA, Chadee DD. Epidemiology of
489 human leptospirosis in Trinidad and Tobago, 1996-2007: A retrospective
490 study. *Acta Tropica* 2009;112:260–5.
491 <https://doi.org/10.1016/j.actatropica.2009.08.007>.
- 492 [37] Desvars A, Jégo S, Chiroleu F, Bourhy P, Cardinale E, Michault A. Seasonality
493 of human leptospirosis in Reunion Island (Indian Ocean) and its association
494 with meteorological data. *PLoS ONE* 2011;6.
495 <https://doi.org/10.1371/journal.pone.0020377>.
- 496 [38] Habus J, Persic Z, Spicic S, Vince S, Stritof Z, Milas Z, et al. New trends in
497 human and animal leptospirosis in Croatia, 2009–2014. *Acta Tropica*
498 2017;168:1–8. <https://doi.org/10.1016/j.actatropica.2017.01.002>.

- 499 [39] Lau CL, Watson CH, Lowry JH, David MC, Craig SB, Wynwood SJ, et al.
500 Human Leptospirosis Infection in Fiji: An Eco-epidemiological Approach to
501 Identifying Risk Factors and Environmental Drivers for Transmission. PLOS
502 Neglected Tropical Diseases 2016;10.
503 <https://doi.org/10.1371/journal.pntd.0004405>.
- 504 [40] Puca E, Pipero P, Harxhi A, Abazaj E, Gega A, Puca E, et al. The role of
505 gender in the prevalence of human leptospirosis in Albania. The Journal of
506 Infection in Developing Countries 2018;12. <https://doi.org/10.3855/jidc.9805>.

507 **7. Supporting information:**

508 S1 Table 1: Summary of leptospirosis incidence distribution by age group from 2010

509 to 2019 in Western Cape Province, South Africa

510 S2 Table 1: Leptospirosis cases per healthcare facility between 2010-2019 in

511 Western Cape Province, South Africa.

512 S3 Fig 1: University of Cape Town – Human Research Ethics Committee approval

513 document.