Changes in foot progression angle during gait reduce the knee adduction moment and do not increase hip moments in individuals with knee osteoarthritis

Kirsten Seagers[1], Scott D. Uhlrich[1,2,3], Julie A. Kolesar[2,3], Madeleine Berkson[2], Janelle M. Kaneda[3], Gary S. Beaupre[1,2], Scott L. Delp[1,3,4]

[1]Department of Mechanical Engineering, Stanford University, Stanford, California, United States

[2]Department of Veterans Affairs Healthcare System, Palo Alto, California, United States

[3] Department of Bioengineering, Stanford University, Stanford, California, United States

[4]Department of Orthopaedic Surgery, Stanford University, Stanford, California, United States

Abstract

People with knee osteoarthritis who adopt a modified foot progression angle (FPA) during gait often benefit from a reduction in the knee adduction moment. It is unknown, however, whether changes in the FPA increase hip moments, a surrogate measure of hip loading, which will increase the mechanical demand on the joint. This study examined how altering the FPA affects hip moments. Individuals with knee osteoarthritis walked on an instrumented treadmill with their baseline gait, 10° toe-in gait, and 10° toe-out gait. A musculoskeletal modeling package was used to compute joint moments from the experimental data. Fifty participants were selected from a larger study who reduced their peak knee adduction moment with a modified FPA. In this group, participants reduced the first peak of the knee adduction moment by 7.6% with 10° toe-in gait and reduced the second peak by 11.0% with 10° toe-out gait. Modifying the FPA reduced the early-stance hip abduction moment, at the time of peak hip contact force, by 4.3% ± 1.3% for 10° toe-in gait (p=0.005, d=0.49) and by 4.6% ± 1.1% for 10° toe-out gait (p<0.001, d=0.59) without increasing the flexion and internal rotation moments (p>0.15). Additionally, 74% of individuals reduced their total hip moment at time of peak hip contact force with a modified FPA. In summary, when adopting a FPA modification that reduced the knee adduction moment, participants, on average, did not increase surrogate measures of hip loading.

Keywords: gait modifications, non-surgical intervention, joint loading, musculoskeletal modeling, osteoarthritis

1 Introduction

Of the 30 million Americans who suffer from osteoarthritis (OA) (Cisternas et al., 2016),

an estimated 14 million have symptomatic knee OA (Deshpande et al., 2016). Mechanical

loading, specifically joint forces, influence OA initiation and progression (Andriacchi et al.,

2004; Felson, 2013). Since direct measurement of compressive joint forces in a native joint is not

possible, joint moments, like the knee adduction moment (KAM), are often used as surrogate

measures of joint loading. The KAM is correlated with knee OA presence (Baliunas et al., 2002),

knee pain (Robbins et al., 2011; Thorp et al., 2007), knee cartilage degradation (Chehab et al.,

2014), and predicts knee OA progression (Miyazaki, 2002). People can change their KAM by

modifying features of their gait, such as trunk sway or walking speed (Hunt et al., 2008;

Mündermann et al., 2004).

Gait modifications are an attractive intervention because they can reduce surrogate

measures of knee loading, such as the KAM, without requiring surgery or external devices. One

such gait modification is changing the foot progression angle (FPA), i.e., toeing-in or toeing-out.

Altering the FPA requires changes to lower-body kinematics (Shull et al., 2013) and muscle

coordination (Charlton et al., 2018; Rutherford et al., 2010). Toeing-in generally reduces the first

peak of the KAM (Shull et al., 2013), whereas toeing-out reduces the second peak of the KAM

(Guo et al., 2007). Although the clinical efficacy of prescribing the same FPA modification to

all individuals remains unclear (Hunt et al., 2018), prescribing this intervention in a personalized

manner can increase the achievable reductions in KAM (Uhlrich et al., 2018) and avoid cases

where individuals increase their KAM. Felson et al., (2019) showed a personalized approach is

effective for other KAM-reducing interventions. If personalized FPA retraining proves to be

effective clinically, it could be deployed in the clinic with advancements in mobile sensors for

estimating the KAM (Wang et al., 2020; Boswell et al., 2021) and providing biofeedback to

change it (Xia et al., 2020).

As research on FPA modifications for people with knee OA progresses towards a

deployable intervention, it is important to understand their effects more comprehensively,

particularly at the hip. Studies that evaluate the efficacy of KAM-targeted gait modifications

focus primarily on the effects at the knee, although there is a need to evaluate the effects of these

modifications on other joints (Mündermann et al., 2005). People with knee OA have a 25%

lifetime risk of developing hip OA (Murphy et al., 2010) due to shared risk factors, which place

multiple joints at risk for OA (Felson et al., 2004; Issa et al., 2002). Although it is unknown how

gait modifications will affect the hip joint, increased hip moments in all planes will increase

mechanical demand on the soft tissues around the hip. Increased hip moments alter the

compressive loading environment in the joint (Simic et al., 2011) and are associated with hip OA

symptoms (Hall et al., 2019), presence (Maly et al., 2013), severity (Diamond et al., 2018), and

progression (Tateuchi et al., 2017). The total knee joint moment, which is a summary measure of

knee moments in all planes, has been used to evaluate gait modifications that have a multiplanar

effect (Asay et al., 2018). Computing a similar measure at the hip (i.e., the total hip moment)

may be a valuable way to estimate the net change in soft tissue demand at the hip from a gait

modification for each individual. Some studies have examined the effect of FPA gait

modifications on hip contact forces, kinetics, and kinematics to minimize torsional loading and

loosening of total hip replacements (Bowsher and Vaughan, 1995) and to reduce hip contact

forces for patients with hip pathology (Wesseling et al., 2015). Wesseling et al., (2015) found

that hip kinematic perturbations that mimic clinical gait patterns, which could result in a

modified FPA, reduced hip moments in a simulation study, but these results have not been

verified experimentally. Toe-out gait, which increased the KAM in asymptomatic individuals,

increased hip sagittal and frontal plane moments (Legrand et al., 2021); however, it is unclear

whether these effects will be the same in individuals with knee OA who reduce their KAM with

a modified FPA. Therefore, an important next step is to understand how interventions that target

knee loading affect hip kinetics, kinematics, and spatiotemporal parameters of gait in an

osteoarthritic population.

This study analyzes the biomechanical effects of a FPA modification on the hip in

individuals with medial compartment knee OA who reduce their KAM with the modification.

We hypothesize that changing the FPA will not increase hip moments or the total hip moment at

time of peak hip contact force compared to baseline gait. Additionally, we performed an

exploratory analysis to evaluate the effect of FPA on kinematics and spatiotemporal parameters.

Our results provide insight into whether gait modifications for knee osteoarthritis increase

mechanical demand at the hip.

Methods

Data Collection

Participant data were analyzed from a clinical study (clinicaltrials.gov NCT02767570)

that trained people with medial compartment knee OA to walk with a modified FPA to reduce

their peak KAM. One hundred seven people provided informed consent in agreement with a

Stanford University Institutional Review Board protocol. Inclusion criteria included medial

compartment knee OA grade between one and three on the Kellgren-Lawrence scale from

anterior-posterior weightbearing radiographs; medial knee pain of three or greater on an 11-point

numeric rating scale; the ability to walk safely on a treadmill without assistance for 25 minutes;

and BMI less than 35. Participants were excluded if they had symptomatic arthritis in lower limb

joints other than the knee that was more severe than their arthritic knee. The limb examined for

each participant was chosen based on the presence of radiographic and symptomatic medial knee

OA. If medial knee OA was present in both knees, the more symptomatic side was analyzed.

Experimental gait data were collected during a visit where different FPA modifications

were evaluated for their effect on KAM peaks. Nineteen reflective markers were placed

bilaterally on the 2^{nd} and 5^{th} metatarsal heads, calcanei and medial and lateral malleoli, medial

and lateral femoral epicondyles, anterior and posterior superior iliac spines, and the C7 vertebrae,

with eighteen additional markers for limb tracking. First, a standing calibration trial was

performed, followed by a functional hip joint center calibration trial during which participants

were asked to circumduct their hips (Piazza et al., 2004). Participants then walked naturally for

two minutes (i.e., baseline trial) at a self-selected speed on an instrumented split belt treadmill

(Bertec Corporation, Columbus, OH, USA) while ground reaction force data and optical marker

positions were recorded (Motion Analysis Corporation, Santa Rosa, CA, USA). Then, two-

minute FPA modification trials were performed where vibrotactile biofeedback (Engineering

Acoustics, Casselberry, FL, USA) was given to participants to toe-in and toe-out by 10° relative

to their natural FPA. The absolute FPA was calculated as the angle between the line connecting

the calcaneus and second metatarsal head marker positions and the forward direction of the

treadmill during 15-40% of stance (Rutherford et al., 2008).

Of the one hundred seven participants who completed this task in the clinical study, fifty

were included in this analysis (Table 1). Selection for this analysis required participants to

reduce the first or second peak KAM by at least 5%, a magnitude of KAM reduction that would

likely elicit an improvement in knee pain (Erhart-Hledick et al., 2012), with either a 10° toe-in or

a 10° toe-out gait modification as compared to baseline. Participants who did not reduce their

peak KAM would not be prescribed a FPA modification and were therefore excluded from this

analysis. The first and second peaks of the KAM during the baseline walking trial were defined

as the maximum values during 0-50% and 51-100% of the stance phase, respectively. The FPA

modification KAM peaks were identified within ±10% of the timing of the participant's baseline

peaks. Participants were excluded from this analysis if they used the treadmill handrails during

walking trials or had greater than an estimated 1 cm of soft tissue between the anterior superior

iliac spine marker and the anatomical bony landmark to ensure the fidelity of hip joint marker

data (Hicks et al., 2015; Leardini et al., 2005).

Musculoskeletal Modeling

We used a constrained musculoskeletal model (Rajagopal et al., 2016) with 25 degrees of

freedom for kinematic and kinetic analysis. This model includes a three degree of freedom hip

joint with Euler flexion, abduction, and rotation angles. We adapted the model published by

Rajagopal and colleagues, by locking the metatarsophalangeal joints, removing the arms, and

including a two degree of freedom knee joint (Lerner et al., 2015).

Scaling the generic musculoskeletal model to match the dimensions of each subject was a

two-step process. First, long-leg radiograph measurements were used to adjust frontal plane

lower limb alignment and pelvic width. The frontal plane lower limb alignment was found by

measuring the angle (ImageJ, National Institute of Mental Health, Bethesda, Maryland, USA)

between a line connecting the hip and knee joint centers and a line connecting the knee and ankle

joint centers (Lerner et al., 2015). The pelvic width was defined as the distance between the hip

joint centers identified on the radiograph. Second, the remainder of the model bodies were scaled

using locations of joint centers and anatomical markers in OpenSim 4.0 (Seth et al., 2018).

Inverse kinematics was performed to compute hip adduction, flexion, and internal

rotation angles. Results were filtered with a 4[th] order Butterworth filter and cutoff frequency of

15 Hz. These kinematics and ground reaction forces, also filtered at 15 Hz, were used in inverse

dynamics to calculate external hip abduction, extension, and internal rotation moments as well as

the KAM. Moments were reported in the joint coordinate reference frame.

Data and Statistical Analysis

For each gait trial (baseline, 10° toe-in, and 10° toe-out), the last 20 steps that achieved

the target FPA, within 2.5°, were analyzed. Each biomechanical metric of interest was calculated

for every step and averaged over each walking condition.

Hip moments were normalized by participant mass (Wesseling et al., 2015) and

compared between walking conditions for the primary kinetic metrics: the early-stance hip

moment, total hip moment, and angular impulse. To estimate the hip moment at the time of

greatest hip loading, the early-stance hip moment was computed as the average moment during

15-20% of stance phase, the time of first peak hip contact force (Bergmann et al., 2001;

Wesseling et al., 2015). To estimate the net effect of moment changes in all planes, the total hip

moment at time of peak hip contact force was calculated. This metric was defined as the square

root of the sum of squares of the early stance hip abduction, flexion, and internal rotation

moments (Asay et al., 2018). For analysis during the entire stance phase, the angular impulse

value was computed as the area under the absolute value of the hip moment curve during stance

phase. For kinematic analysis, joint angles were averaged during stance and compared between

walking conditions to capture global changes. Peak angles were also compared between

conditions. Spatiotemporal parameters were also determined for comparison, including FPA,

stride length, and step width. Stride length was defined as the difference in right and left foot

calcaneus marker positions at time of heel strike. Step width was defined as the distance between

the right and left foot center of pressure at 50% of stance (Donelan et al., 2001).

Statistical evaluations of 10° toe-in and 10° toe-out gait metrics compared to baseline

were performed in MATLAB R2020a (MathWorks Corporation, Natick, MA USA) using a

Wilcoxon signed-rank test ($\alpha = 0.05$), since a Kolmogorov-Smirnov test indicated the data were

not normally distributed. Bonferroni corrected p-values for multiple comparisons across FPA

conditions were reported to mitigate type I error. Additionally, the Cohen's d effect size was

calculated for further comparison of the hip moments between gait patterns, where d<0.20 is a

small effect and d>0.80 is a large effect.

Results

Participants adopted a FPA modification that reduced the peak KAM and did not increase

surrogate measures of hip loading. On average, 10° toe-in modifications reduced the first peak of

the KAM by 7.6% compared to baseline, and 10° toe-out reduced the second peak by 11.0%. The

early-stance hip abduction moment was significantly lower during 10° toe-in gait (-4.3% ± 1.3%,

[range: -31.7%, 11.0%], p=0.005, d=0.49) and during 10° toe-out gait (-4.6% ± 1.1%, [range: -

21.9%, 9.4%], p<0.001, d=0.59) compared to baseline. There were no significant differences

(p>0.05, d<0.25) in the early-stance hip extension moment or early-stance internal rotation

moment for 10° toe-in or toe-out conditions compared to baseline (Figure 1). The baseline

internal rotation moment and changes with FPA modifications were small in magnitude (Figure

2, Table 2), yet this results in large percent changes from baseline (Figure 1).

On average, there were no significant increases in the total hip moment at time of peak

hip contact force; however, on an individual basis when walking with a modified FPA that

maximally reduced their KAM, 37 individuals (74% of participants) reduced their total hip

moment, while 13 individuals (26% of participants) increased their total hip moment. When

walking with a toe-out gait, 46% of individuals increased this metric, and 40% increased it with

toe-in gait. The largest observed total hip moment increase from baseline was 9.2% and 12.4%

for 10° toe-in and 10° toe-out gait, respectively. There was no difference in the change in total

hip moment from baseline for 10° toe-in gait (-1.9% ± 0.8%, [range: -17.3%, 9.2%], p>0.05,

d=0.30) or 10° toe-out gait (-1.4% ± 0.8%, [range: -16.5%, 12.4%], p>0.05, d=0.22) compared to

baseline.

The frontal plane angular impulse was significantly reduced compared to baseline for 10°

toe-in gait (-2.3% ± 0.7%, [range: -11.7%, 7.4%], p<0.01, d=0.45) and for 10° toe-out gait (-

6.9% ± 0.8%, [range: -23.1%, 4.5%], p<0.001, d=1.23), but no significant differences (p>0.05,

d<0.18) were observed in the sagittal or transverse plane (Figure 3).

Changes in hip kinematics were minimal. Participants walked with a decreased *average* hip

adduction angle for 10° toe-in gait (-0.4° ± 1.1°, p=0.045) and 10° toe-out gait (-1.6° ± 1.2°,

p<0.001) compared to baseline (Figure 4). *Peak* hip adduction angles also decreased for both toe-

in (-0.7° ± 1.5°, p=0.009) and toe-out (-1.8° ± 1.5°, p=0.005) gait compared to baseline.

Participants walked with increased *average* hip flexion when toeing-in (2.7° ± 2.2°, p<0.001)

and toeing-out (1.7° ± 2.1°, p<0.001). Both modifications had a lower *peak* extension angle

compared to baseline (toe-in: -3.2° ± 2.7°, p<0.001; toe-out: -2.5° ± 2.7°, p<0.001). The *average*

hip internal rotation angle increased by an average of 4.4° ± 2.6° (p<0.001) when walking with a

10° toe-in gait and decreased by 5.7° ± 1.3° (p<0.001) with a 10° toe-out gait. Similar directional

changes were seen in the *peak* hip internal rotation angle compared to baseline (toe-in: 4.6° ±

2.7°, p<0.001; toe-out: -5.8° ± 1.6°, p<0.001). Changes in spatiotemporal parameters were also

observed. Stride length decreased with both FPA modifications (p<0.001), while step width

increased only for 10° toe-out gait (p<0.001) (Table 2).

Discussion

We analyzed how changing the FPA during walking affects hip moments in fifty

individuals with medial knee OA who successfully reduce their KAM with the modification.

Our hypothesis was confirmed, as hip moments did not increase, on average, when a modified

FPA was adopted. Both 10° toe-in and 10° toe-out modifications decreased the early-stance hip

abduction moment and angular impulse. On average, adopting a FPA modification that reduced

the KAM did not significantly increase hip moments at time of peak hip contact force, the total

hip moment at time of peak hip contact force, or the angular impulse. However, some

participants did increase their total hip moment at time of peak hip contact force with the FPA

modification. This suggests that for individuals with hip joint pathology or at risk of developing

hip OA, the effect of a FPA gait modification at the hip should be evaluated individually to

ensure that adopting a modification that improves knee loading does not cause a potentially

harmful increase in hip loading.

Our observed decrease in the hip abduction moment is consistent with other studies. The

4% decrease in the early-stance hip abduction moment when toeing-in and toeing-out was

smaller than the 16% decrease in hip abduction observed at the time of peak KAM in a healthy

population (Kettlety et al., 2018). This could be due to the varying participant population and

differences in gait cycle timing when the hip moments were analyzed (i.e., at peak hip contact

force versus peak KAM). Similarly, the reductions in early-stance hip abduction moment (10°

toe-in: 0.03 Nm/kg; 10° toe-out: 0.04 Nm/kg) agreed directionally with the 0.1 Nm/kg decrease

Wesseling et al., (2015) reported from simulated perturbations of frontal plane hip kinematics.

The magnitude differences are unsurprising when comparing simulation predictions to

experimental results, since participants may adopt other compensatory strategies to modify their

FPA that were not captured in these simulations. Additionally, our observed hip rotation moment

changes were small in magnitude, compared to the abduction and extension moments, resulting

in large percent changes.

Kinematic and spatiotemporal changes were also observed when participants modified

their FPA. Changes in the hip rotation angle were less than the change in FPA, which aligns with

results from Cibulka et al., (2016), who found that FPA modifications elicit changes at multiple

lower-extremity joints. In addition to the rotational kinematics changes, the average hip

adduction angle decrease relates to the increase in step width for toe-out gait. Other studies

(Kettlety et al., 2020; Uhlrich et al., 2018) have also observed that step width increases with a

modified FPA resulting from less hip adduction. While the decrease in hip adduction angle and

increase in step width are small, these changes increase medio-lateral stability (Meyer et al.,

2018). Additionally, participants decreased stride length, which can increase double-limb support

time, resulting in a more stable gait (Shkuratova et al., 2004). The changes to kinematics and

spatiotemporal parameters may attenuate over time as an individual becomes more comfortable

with the gait modification.

It is important to identify the limitations of our study. We estimated the effect of FPA

modifications on surrogate measures of hip loading. Electromyography was not used in this

study, but, in the future, validated electromyography-informed simulations could estimate

changes in muscle and hip contact forces. The surrogate measure of hip loading that we used,

joint moments, is related to muscle and joint contact forces, which provides an estimate of the

directional change in hip contact force. Meyer et al. (2018) reported that a decrease in the hip

abduction moment and hip adduction angle reduced the work generated by the hip abductor

muscles and decreased hip loading in people with hip OA. The relationship between the total hip

moment and hip OA pathogenesis has not been studied, but the total knee joint moment and the

change in relative contribution of its components (knee adduction, flexion, and rotation

moments) reflect gait changes associated with progressing knee OA and pain (Asay et al., 2018).

Similarly, the total hip moment is likely an effective summary metric of changes in muscular

demand in all planes at the hip and future work should investigate how it relates to hip OA

initiation and progression. Generally, little is known about the role of hip joint loading on the

initiation of hip OA; elucidating these relationships should be the topic of future work.

Several studies have shown that FPA modifications can provide meaningful reductions in

the KAM. Our results add to this growing body of evidence and show that, on average, FPA

modifications reduce frontal plane hip moments without increasing the moments in other planes.

However, some individuals did increase their total hip moment at time of peak hip contact force.

If an individual is at risk for hip OA, then hip moments will be an important biomechanical

parameter to monitor when prescribing gait modifications. Yet, 74% of people, who could

meaningfully reduce their KAM, reduced their total hip moment at time of peak hip contact force

with a modified FPA. In summary, on average and for most individuals, a FPA modification that

reduces knee loading does not increase surrogate measures of hip loading.

Acknowledgements

We would like to thank the participants involved in the study. This work was supported by the

Sang Samuel Wang Stanford Graduate Fellowship, NSF Graduate Research Fellowships (DGE-

1147470 and DGE-1656518), NIH grant P41EB027060, and by Merit Review Award I01

RX001811 from the United States Department of Veterans Affairs Rehabilitation Research and

Development Service.

References

Andriacchi, T.P., Mündermann, A., Smith, R.L., Alexander, E.J., Dyrby, C.O., Koo, S., 2004. A Framework for the in Vivo Pathomechanics of Osteoarthritis at the Knee. Annals of Biomedical Engineering 32, 447–457.

Asay, J.L., Erhart-Hledik, J.C., Andriacchi, T.P., 2018. Changes in the Total Knee Joint Moment in Patients with Medial Compartment Knee Osteoarthritis Over 5 years. Journal of Orthopaedic Research 36, 2373-2379.

Baliunas, A.J., Hurwitz, D.E., Ryals, A.B., Karrar, A., Case, J.P., Block, J.A., Andriacchi, T.P., 2002. Increased knee joint loads during walking are present in subjects with knee osteoarthritis. Osteoarthritis and Cartilage 10, 573–579.

Bergmann, G., Deuretzbacher, G., Heller, M., Graichen, F., Rohlmann, A., Strauss, J., Duda, G.N., 2001. Hip contact forces and gait patterns from routine activities. Journal of Biomechanics 34, 859–871.

Bowsher, K.A., Vaughan, C.L., 1995. Effect of foot-progression angle on hip joint moments during gait. Journal of Biomechanics 28, 759–762.

Charlton, J.M., Hatfield, G.L., Guenette, J.A., Hunt, M.A., 2018 Toe-in and toe-out walking require different lower limb neuromuscular patterns in people with knee osteoarthritis. Journal of Biomechanics 76, 112–118.

Chehab, E.F., Favre, J., Erhart-Hledik, J.C., Andriacchi, T.P., 2014. Baseline knee adduction and flexion moments during walking are both associated with 5-year cartilage changes in patients with medial knee osteoarthritis. Osteoarthritis and Cartilage 22, 1833–1839.

Cibulka, M.T., Winters, K., Kampwerth, T., McAfee, B., Payne, L., Roeckenhaus, T., Ross, S.A., 2016. Predicting Foot Progression Angle during Gait using Two Clinical Measures in Healthy Adults, a Preliminary Study. International Journal of Sports Physical Therapy 11, 400-408.

Cisternas, M.G., Murphy, L., Sacks, J.J., Solomon, D.H., Pasta, D.J., Helmick, C.G., 2016. Alternative Methods for Defining Osteoarthritis and the Impact on Estimating Prevalence in a US Population-Based Survey: OA Prevalence in a Population-Based Survey. Arthritis Care & Research 68, 574–580.

Deshpande, B.R., Katz, J.N., Solomon, D.H., Yelin, E.H., Hunter, D.J., Messier, S.P., Suter, L.G., Losina, E., 2016. Number of Persons With Symptomatic Knee Osteoarthritis in the US: Impact of Race and Ethnicity, Age, Sex, and Obesity: Symptomatic Knee OA in the US. Arthritis Care & Research 68, 1743–1750.

Diamond, L.E., Allison, K., Dobson, F., Hall, M., 2018. Hip joint loading during walking in people with hip osteoarthritis: a systematic review and meta-analysis. Osteoarthritis and Cartilage, Abstracts from the 2018 OARSI World Congress on Osteoarthritis 26, S370.

Donelan, J.M., Kram, R., Kuo, A.D., 2001. Mechanical and metabolic determinants of the preferred step width in human walking. Proceedings of the Royal Society: Biological Sciences 268, 1985-1992.

Erhart-Hledik, J.C., Elspas, B., Giori, N.J., Andriacchi, T.P., 2012. Effect of variable-stiffness walking shoes on knee adduction moment, pain, and function in subjects with medial compartment knee osteoarthritis after 1 year. Journal of Orthopedic Research 30, 514-21.

Felson, D.T., Parkes, M., Carter, S., Liu, A., Callaghan, M.J., Hodgson, R., Bowes, M., Jones, R.K., 2019. The Efficacy of a Lateral Wedge Insole for Painful Medical Knee Osteoarthritis After Prescreening: A Randomized Clinical Trial. Arthritis & Rheumatology 71, 908-915.

Felson, D.T., 2013. Osteoarthritis as a disease of mechanics. Osteoarthritis and Cartilage 21, 10–15.

Felson, D.T., 2004. Risk factors for osteoarthritis: understanding joint vulnerability. Clinical Orthopaedics and Related Research 427, S16-21.

Guo, M., Axe, M.J., Manal, K., 2007. The influence of foot progression angle on the knee adduction moment during walking and stair climbing in pain free individuals with knee osteoarthritis. Gait & Posture 26, 436–441.

Hall, M., Chabra, S., Shakoor, N., Leurgans, S.E., Demirtas, H., Foucher, K.C., 2019. Hip joint moments in symptomatic vs. asymptomatic people with mild radiographic hip osteoarthritis. Journal of Biomechanics 96, 109347.

Hicks, J.L., Uchida, T.K., Seth, A., Rajagopal, A., Delp, S.L., 2015. Is my model good enough? Best practices for verification and validation of musculoskeletal models and simulations of movement. Journal of Biomechanical Engineering 137, 0209051-02090524.

Hunt, M.A., Birmingham, T.B., Jenkyn, T.R., Giffin, J.R., Jones, I.C., 2008. Measures of frontal plane lower limb alignment obtained from static radiographs and dynamic gait analysis. Gait & Posture 27, 635–640.

Hunt, M.A., Charlton, J.M., Krowchuk, N.M., Tse, C.T.F, Hatfield, G.L., 2018. Clinical and biomechanical changes following a 4-month toe-out gait modification program for people with medial knee osteoarthritis: a randomized controlled trial. Osteoarthritis and Cartilage 26, 903-911.

Issa, R., Griffin, T., 2012. Pathobiology of obesity and osteoarthritis: integrating biomechanics and inflammation. Pathobiology of Aging & Age-related Diseases 2, 17470.

Kettlety, S., Lindsey, B., Eddo, O., Prebble, M., Caswell, S., Cortes, N., 2020. Changes in hip mechanics during gait modification to reduce knee adduction moment. Journal of Biomechanics 99, 1-7.

Leardini, A., Chiari L., Croce, U.D., Cappozzo, A., 2005. Human movement analysis using stereophotogrammetry. Part 3. Soft tissue artifact assessment and compensation. Gait & Posture 21, 212-225.

Legrand, T., Younesian, H., Equey, N., Campeau-Lecours, A., Turcot, K., 2021. Trunk lean and toe out gait strategies impact on lower limb joints. Journal of Biomechanics 129, 110740.

Lerner, Z.F., DeMers, M.S., Delp, S.L., Browning, R.C., 2015. How tibiofemoral alignment and contact locations affect predictions of medial and lateral tibiofemoral contact forces. Journal of Biomechanics 48, 644–650.

Maly, M.R., Robbins, S.M., Stratford, P.W., Birmingham, T.B., Callaghan, J.P., 2013. Cumulative knee adductor load distinguishes between healthy and osteoarthritic knees–A proof of principle study. Gait & Posture 37, 397–401.

Manal, K., Gardinier, E., Buchanan, T.S., Snyder-Mackler, L., 2015. A More Informed Evaluation of Medial Compartment Loading: the Combined Use of the Knee Adduction and Flexor Moments. Osteoarthritis and Cartilage 23(7), 1107-1111.

Meyer, C., Wesseling, M., Corten, K., Nieuwenhuys, A., Monari, D., Simon, J., Jonkers, I., Desloovere, K., 2018. Hip movement pathomechanics of patients with hip osteoarthritis aim at reducing hip joint loading on the osteoarthritic side. Gait & Posture, 59, 11-17.

Miyazaki, T., 2002. Dynamic load at baseline can predict radiographic disease progression in medial compartment knee osteoarthritis. Annals of the Rheumatic Diseases 61, 617–622.

Mündermann, A., Dyrby, C.O., Hurwitz, D.E., Sharma, L., Andriacchi, T.P., 2004. Potential strategies to reduce medial compartment loading in patients with knee osteoarthritis of

varying severity: Reduced walking speed: Relationship of Walking Speed to Maximum Knee Adduction Moment. Arthritis & Rheumatism 50, 1172–1178.

Mündermann, A., Dyrby, C.O., Andriacchi, T.P., 2005. Secondary gait changes in patients with medial compartment knee osteoarthritis: Increased load at the ankle, knee, and hip during walking. Arthritis & Rheumatism 52, 2835–2844.

Murphy, L.B., Helmick, C.G., Schwartz, T.A., Renner, J.B., Tudor, G., Koch, G.G., Dragomir, A.D., Kalsbeek, W.D., Luta, G., Jordan, J.M., 2010. One in four people may develop symptomatic hip osteoarthritis in his or her lifetime. Osteoarthritis and Cartilage 18, 1372–1379.

Piazza, S.J., Erdemir, A, Okita, N, Cavanagh, P.R., 2004. Assessment of the functional method of hip joint center location subject to reduced range of hip motion. Journal of Biomechanics 37, 349-356

Rajagopal, A., Dembia, C.L., DeMers, M.S., Delp, D.D., Hicks, J.L., Delp, S.L., 2016. Full-Body Musculoskeletal Model for Muscle-Driven Simulation of Human Gait. IEEE Transactions on Biomedical Engineering 63, 2068–2079.

Robbins, S.M., Birmingham, T.B., Callaghan, J.P., Jones, G.R., Chesworth, B.M., Maly, M.R., 2011. Association of pain with frequency and magnitude of knee loading in knee osteoarthritis. Arthritis Care & Research 63, 991–997.

Rutherford, D.J., Hubley-Kozey, C.L., Deluzio, K.J., Stanish, W.D., Dunbar, M., 2008. Foot progression angle and the knee adduction moment: a cross-sectional investigation in knee osteoarthritis. Osteoarthritis and Cartilage 16, 883-889.

Rutherford, D.J., Hubley-Kozey, C.L., Stanish, W.D., 2010. The neuromuscular demands of altering foot progression angle during gait in asymptomatic individuals and those with knee osteoarthritis. Osteoarthritis and Cartilage 18, 654–661.

Seth, A., Hicks, J.L, Uchida, T.K., Habib, A., Dembia, C.L., Dunne, J.J., Ong, C.F., DeMers, M.S., Rajagopal, A., Millard, M., Hamner, S.R., Arnold, E.M., Yong, J.R., Lakshimikanth, S.K., Sherman, M.A, Ku, J.P., Delp, S.L., 2018. OpenSim: Simulating musculoskeletal dynamics and neuromuscular control to study human and animal movement. PLoS Computational Biology 14, e1006223.

Shkuratova, N., Morris, M.E., Huxham, F., 2004. Effects of age on balance control during walking. Archives of Physical Medicine and Rehabilitation 85(4), 582-588.

Shull, P.B., Shultz, R., Silder, A., Dragoo, J.L., Besier, T.F., Cutkosky, M.R., Delp, S.L., 2013. Toe-in gait reduces the first peak knee adduction moment in patients with medial compartment knee osteoarthritis. Journal of Biomechanics 46, 122–128.

Simic, M., Hinman, R., Wrigley, T., Bennell, K., Hunt, M., 2011. Gait modification strategies for altering medial knee joint load: a systematic review. Arthritis Care & Research 63, 405–426.

Tateuchi, H., Koyama, Y., Akiyama, H., Goto, K., So, K., Kuroda, Y., Ichihashi, N., 2017. Daily cumulative hip moment is associated with radiographic progression of secondary hip osteoarthritis. Osteoarthritis and Cartilage 25, 1291–1298.

Thorp, L.E., Sumner, D.R., Wimmer, M.A., Block, J.A., 2007. Relationship between pain and medial knee joint loading in mild radiographic knee osteoarthritis. Arthritis Care & Research 57, 1254–1260.

Uhlrich, S.D., Silder, A., Beaupre, G.A., Shull, P.B., Delp, S.L., 2018. Subject-specific toe-in or toe-out gait modifications reduce the larger knee adduction moment peak more than a non-personalized approach. Journal of Biomechanics 66, 103–110.

Wang, C., Chan, P.P.K, Lam, B.M.F, Wang, S., Zhang, J.H., Chan, Z.Y.S., Chan, R.H.M, *Senior*
Member, IEEE, Ho, K.K.W., Cheung, R.T.H., 2020. Real-Time Estimation of Knee
Adduction Moment for Gait Retraining in Patients with Knee Osteoarthritis. IEEE
Transactions on Neural Systems and Rehabilitation Engineering 28, 888-894.
Wesseling, M., Groote, F. de, Meyer, C., Corten, K., Simon, J.P., Desloovere, K., Jonkers, I.,
2015. Gait alterations to effectively reduce hip contact forces. Journal of Orthopaedic
Research 33, 1094–1102.
Xia, H., Charlton, J.M., Shull, P.B., Hunt, M.A., 2020. Portable, automated foot progression
angle gait modification via a proof-of-concept haptic feedback-sensorized shoe. Journal of
Biomechanics 107, 109789

TABLES & FIGURES

Characteristic (N = 50)	Mean (SD)
Age (yr)	61 (7)
Height (m)	1.7 (0.1)
Mass (kg)	76.2 (15.4)
BMI (kg/m^2)	25.8 (3.8)
Gender	M: 31 F: 19
Static Knee Angle (°): Varus (-) Valgus (+)	Ipsilateral: -2.4 (3.6) Contralateral: -1.0 (3.6)
Participants who reduced peak KAM with a modified FPA	Peak 1 only: n =11 Peak 2 only: n =14 Peak 1 & Peak 2: n =25
Gait Speed (m/s)	1.2 (0.1)
Kellgren and Lawrence Grade	□: 12, □: 27, □: 11
Medial Knee Pain: 0 (no pain) - 10 (worst pain)	4 (2)

407 **Table 1:** Demographics of the fifty participants with symptomatic medial knee osteoarthritis

408 included in the analysis. Standard deviation (SD) is reported in parentheses.

Parameter	Baseline		10° Toe-in			10° Toe-out		
	mean	SD	mean	SD	p-value	mean	SD	p-value
FPA (°)	6.9	3.4	- 2.8	3.5	**< 0.001**	16.5	3.4	**< 0.001**
Stride Length (m)	1.33	0.14	1.29	0.13	**< 0.001**	1.29	0.13	**< 0.001**
Step Width (m)	0.18	0.04	0.19	0.04	0.863	0.22	0.05	**< 0.001**
Peak Hip Abduction Moment (Nm/kg)	0.97	0.12	0.93	0.13	**<0.001**	0.91	0.14	**<0.001**
Peak Hip Extension Moment (Nm/kg)	0.72	0.18	0.75	0.18	**0.003**	0.73	0.19	0.227
Peak Hip Flexion Moment (Nm/kg)	-0.64	0.17	-0.60	0.16	**<0.001**	-0.60	0.15	**<0.001**
Peak Hip Internal Rotation Moment (Nm/kg)	0.20	0.06	0.21	0.08	0.37	0.22	0.08	**0.025**
Peak Hip External Rotation Moment (Nm/kg)	-0.11	0.03	-0.09	0.04	**<0.001**	-0.08	0.04	**<0.001**
1st Peak KAM:								
(%BW*H)	3.69	1.17	3.41	1.11		3.80	1.23	
(Nm/kg)	0.62	0.20	0.57	0.19		0.64	0.21	
2nd Peak KAM:								
(%BW*H)	2.72	0.98	2.72	1.01		2.42	1.06	
(Nm/kg)	0.46	0.17	0.46	0.17		0.41	0.18	

Table 2: Spatiotemporal parameters, peak hip moments, and KAM for participants during

baseline gait, 10° toe-in and 10° toe-out. Note, p-values are not reported for the peak KAM

values since participants were selected based on their ability to reduce their peak KAM.

Figure 1: Participants reduced their early-stance hip abduction moment for 10° toe-in (blue) and

10° toe-out (red) gait as compared to baseline. No significant differences were observed

between baseline gait and toe-in or toe-out gait for the early-stance hip extension and internal

rotation moment. (*p<0.005)

Figure 2: Ensemble-averaged hip moment curves, normalized by bodyweight, for all participants

(n = 50) during stance phase. Baseline gait (black, dashed) hip moments are compared to 10°

toe-in gait (blue, solid) in the first row and 10° toe-out gait (red, solid) in the second row. The

shaded region indicates one standard deviation from the average moment.

427 **Figure 3:** Participants reduced their frontal plane angular impulse during 10° toe-in (blue) and

428 10° toe-out (red) gait compared to baseline. No significant differences were observed between

429 baseline and toe-in or toe-out gait for the angular impulse in the sagittal and transverse plane.

430 (*p<0.01)

Figure 4: Ensemble-averaged hip kinematics for all participants (n = 50) during stance phase. Baseline gait (black, dashed) hip angles are compared to 10° toe-in gait (blue, solid) in the first row and 10° toe-out gait (red, solid) in the second row. The shaded region indicates one standard deviation from the average hip angle. The peak hip adduction angle and the peak hip extension angle are significantly reduced with 10° toe-in gait and 10° toe-out gait compared to baseline (p<0.01). The peak hip internal rotation angle is significantly increased for 10° toe-in gait and reduced for 10° toe-out gait compared to baseline (p<0.001).