Infectious viral load in unvaccinated and vaccinated patients infected with SARS-CoV-2 WT, Delta and Omicron

Olha Puhach[1], Kenneth Adea[1], Nicolas Hulo[2], Pascale Sattonnet[1], Camille Genecand[3], Anne Iten[4], Frédérique Jacquérioz Bausch[5,6,7], Laurent Kaiser[5,8,9], Pauline Vetter[5,8,9,*,#], Isabella Eckerle[1,5,9,*,#], Benjamin Meyer[10,*,#]

Affiliations:

[1] Department of Microbiology and Molecular Medicine, Faculty of Medicine, University of Geneva, Geneva, Switzerland

[2] Service for Biomathematical and Biostatistical Analyses, Institute of Genetics and Genomics, University of Geneva, Geneva, Switzerland

[3] Cantonal Health Service, General Directorate for Health, Geneva, Switzerland

[4] Service of Prevention and Infection Control, Directorate of Medicine and Quality, University Hospital Geneva, HUG, Geneva, Switzerland

[5] Geneva Centre for Emerging Viral Diseases, Geneva University Hospitals, 1205 Geneva, Switzerland

[6] Division of Tropical and Humanitarian Medicine, Geneva University Hospitals, Geneva, Switzerland

[7] Primary Care Division, Geneva University Hospitals, Geneva, Switzerland

[8] Laboratory of Virology, Division of Laboratory Medicine, Geneva University Hospitals & Faculty of Medicine, University of Geneva, 1205 Geneva, Switzerland

[9] Division of Infectious Diseases, Geneva University Hospitals, 1205 Geneva, Switzerland

[10] Centre for Vaccinology, Department of Pathology and Immunology, University of Geneva, Switzerland

[#] *Corresponding authors*

Benjamin Meyer: Benjamin.Meyer@unige.ch

Isabella Eckerle: Isabella.Eckerle@hcuge.ch

Pauline Vetter: Pauline.Vetter@hcuge.ch

* Equally contributed

Abstract

Background

Viral load (VL) is one determinant of secondary transmission of SARS-CoV-2. Emergence of variants of
concerns (VOC) Alpha and Delta was ascribed, at least partly, to higher VL. Furthermore, with parts of
the population vaccinated, knowledge on VL in vaccine breakthrough infections is crucial. As RNA VL
is only a weak proxy for infectiousness, studies on infectious virus presence by cell culture isolation
are of importance.

Methods

We assessed nasopharyngeal swabs of COVID-19 patients for quantitative infectious viral titres (IVT)
by focus-forming assay and compared to overall virus isolation success and RNA genome copies. We
assessed infectious viral titres during the first 5 symptomatic days in a total of 384 patients:
unvaccinated individuals infected with pre-VOC SARS-CoV-2 (n= 118) or Delta (n= 127) and vaccine
breakthrough infections with Delta (n= 121) or Omicron (n=18).

Findings

Correlation between RNA copy number and IVT was low for all groups. No correlation between IVTs
and age or sex was seen. We observed higher RNA genome copies in pre-VOC SARS-CoV-2 compared
to Delta, but significantly higher IVTs in Delta infected individuals. In vaccinated vs. unvaccinated Delta
infected individuals, RNA genome copies were comparable but vaccinated individuals have
significantly lower IVTs, and cleared virus faster. Vaccinated individuals with Omicron infection had
comparable IVTs to Delta breakthrough infections.

Interpretation

Quantitative IVTs can give detailed insights into virus shedding kinetics. Vaccination was associated
with lower infectious titres and faster clearance for Delta, showing that vaccination would also lower
transmission risk. Omicron vaccine breakthrough infections did not show elevated IVTs compared to
Delta, suggesting that other mechanisms than increase VL contribute to the high infectiousness of
Omicron.

Funding

This work was supported by the Swiss National Science Foundation 196644, 196383, NRP (National
Research Program) 78 Covid-19 Grant 198412, the Fondation Ancrage Bienfaisance du Groupe Pictet
and the Fondation Privée des Hôpitaux Universitaires de Genève.

Introduction

As of 2 January 2021, the coronavirus disease 2019 (COVID-19) pandemic led to a total of nearly 289 million cases and just over 5.4 million deaths globally (1). Severe acute respiratory coronavirus 2 (SARS-CoV-2), the causative agent of COVID-19, primarily infects the cells of the upper respiratory tract (URT) where it leads to an increasing viral load (VL) during the course of infection(2).

The two key parameters when assessing VL are either RNA genome copies, often expressed in cycle threshold (Ct) values, or infectious virus that can only be assessed by virus isolation in cell culture. Although the process of human-to-human transmission is complex, VL can serve as a proxy, with higher VL posing a greater risk for onward transmission. In several epidemiological studies, higher viral load expressed as viral RNA was associated with an increased secondary transmission in household settings(3, 4). Shedding of infectious SARS-CoV-2 in the URT starts on average two days prior to the beginning of symptoms and gradually declines up to 8 days post onset of symptoms. Even though viral RNA could be detected afterwards, in most studies infectious virus was not detected in respiratory samples collected from immunocompetent individuals later than 8 days post onset of symptoms (5-9). Moreover, the detection of viral RNA does not correlate with infectiousness in an animal model (10). Instead, success of virus isolation in cell culture was found to be a proxy for detection of infectious virus from respiratory specimens and the ability to shed and transmit fully competent viral particles (8, 11, 12). Virus isolation success can only give information about the presence or absence of infectious virus, but is not able to quantify the infectious viral titre in samples of the URT (13).

Since the start of the pandemic, SARS-CoV-2 is constantly evolving, leading to the emergence of new variants. While most variants vanished quickly, other such as D614G, and the designated variants of concern (VOCs) Alpha, Beta, Gamma, Delta and Omicron harbour an apparent selection advantage and manage to replace other existing variants locally or even globally. These VOCs exhibit various mutations and amino acid alterations in viral proteins (14) that lead to immune evasion and/or higher transmissibility to which increased virus shedding among other factors like environmental stability can significantly contribute (15, 16). For Alpha VOC, an approximately 10-fold higher RNA viral load was observed compared to pre-VOC viral strains, that was also associated with an increase in virus isolation success (17, 18). Similarly, Delta VOC also showed 10- to 15-fold higher viral load on RNA level compared to pre-VOC strains (18, 19). However, little is known about the quantity of emitted infectious viral particles for VOCs including Omicron.

There is extensive evidence that introduction of licensed vaccines against SARS-CoV-2, which were designed against the original virus strain, led to the reduction of cases and disease severity. However, the effect of vaccination on infectious viral shedding and onward transmission from vaccinated patients remains controversial. Considering that all currently approved vaccines are administered intramuscularly, the titre of neutralizing antibodies on the mucosal surfaces lining the URT might be limited, and therefore sterilizing immunity is probably only transient if achieved at all (20, 21). Epidemiological studies on the secondary attack rate in households of vaccinated vs unvaccinated index patients led to contrary results. While some studies show a reduction of transmission rates in vaccinated index cases (22-24), one study found no influence of the vaccination status of index cases (25). However, many additional factors, such as the behaviour of index patients, age, comorbidities, the variant present at the time of study, time elapsed after complete vaccination and the vaccine used, can influence the secondary attack rate in these studies. Therefore, it is difficult to delineate the effect of vaccination on viral load from other factors in purely epidemiological studies. An overall reduction

of RNA viral loads was reported in vaccinated COVID-19 patients (BNT162b2 mRNA vaccine or ChAdOx1 nCoV-19 (AZD1222) adenoviral vector vaccine), but no difference was observed at 6 months post vaccination (26, 27). Another study found reduced RNA viral load early after complete vaccination when the Alpha VOC was prevalent, but no difference at later time points when the Delta VOC was dominant (28). In addition, a study that investigated the kinetics of RNA viral load in COVID-19 patients did not find a difference during the first 5 days post symptom onset, i.e. when most human-to-human transmissions occur, but showed a faster decline of RNA viral load in vaccinated patients (29). Similarly, a lower probability of virus isolation success was found in vaccinated vs unvaccinated COVID-19 patients at the same RNA viral load indicating that vaccines can reduce the infectious viral load (30). However, no study quantified infectious virus titres of different VOCs in URT samples of vaccinated and unvaccinated COVID-19 patients.

Dynamics of infectious viral shedding in vaccinated and unvaccinated patients infected with relevant SARS-CoV-2 variants of concern requires detailed investigation. Understanding of the SARS-CoV-2 viral shedding in patients would help the implication of public health decisions to limit the transmission of the virus in the community (31). Here we compare RNA and infectious viral load between pre-VOC strains and Delta VOC in unvaccinated patients as well as in vaccination breakthrough infections due to Delta and Omicron. Respiratory samples from mildly symptomatic patients of different age and sex, sampled in the first five days post onset of symptoms were used for this study. By quantifying infectious viral titres from URT specimens, we show that patients infected with SARS-CoV-2 Delta variant of concern harbour elevated levels of infectious viral titres, while vaccination leads to a reduction of infectious virus.

Methods

Participants

Sample collection and setting

Nasopharyngeal swabs for diagnostics of SARS-CoV-2 by RT-PCR collected from symptomatic individuals in the outpatient testing center of the Geneva University Hospital were included in this study. Infection with SARS-CoV-2 was diagnosed by RT-PCR assay (Cobas 6800, Roche). All samples originate from the diagnostic unit of the virology laboratory of the hospital and were received for primary diagnosis of SARS-CoV-2. Remaining samples were stored at -80°C, usually on the same day or within 24h. All samples had only one freeze-thaw cycle for the purpose of this study. All specimens from vaccinated individuals were characterized by full genome sequencing for their infecting SARS-CoV-2 variant. Initial identification of the Omicron VOC was done by S gene target failure of the TaqPath COVID19 assay (Thermofisher) and confirmed by partial Sanger sequencing of the Spike (32). Final confirmation was done by next-generation sequencing. No sequence information was obtained for patients infected with pre-VOC SARS-CoV-2, as all samples were collected in 2020, in a time period before first detection of the VOCs in Switzerland. Clinical information of the patients was collected by a standardized questionnaire in our testing Centre and/or through the Cantonal Health Service.

Viral load quantification by qRT-PCR

Viral loads in each sample were determined by quantitative real time PCR (RT-qPCR) using
SuperScript™ III Platinum™ One-Step qRT-PCR Kit (Invitrogen). RT-PCR for SARS-CoV-2 E gene and
quantification of genome copy number was performed as described previously (33).

Quantification of SARS-CoV-2 by focus-forming assay.

Vero E6 and Vero E6-TMPRSS were cultured in complete DMEM GlutaMax I medium supplemented
with 10% fetal bovine serum, 1x Non-essential Amino Acids, and 1% antibiotics
(Penicillin/Streptomycin) (all reagents from Gibco, USA). Vero-TMPRSS were kindly received from
National Institute for Biological Standards and Controls (NIBSC, Cat. Nr. 100978).

Nasopharyngeal swab samples were serially diluted and applied on a monolayer of VeroE6 cells in
duplicates. Following 1 hour of incubation at 37°C, the virus-containing inoculum was removed and
the cells were overlaid with prewarmed medium mixed with 2·4% Avicel (DuPont) at a 1:1 ratio. Plates
were incubated at 37°C for 24 hours and then fixed using 6% paraformaldehyde for 1 hour at room
temperature. Cells were permeabilized with 0·1% Triton X-100 and blocked with 1% BSA (Sigma).
Plates were incubated with an anti-SARS-CoV monoclonal nucleocapsid protein primary antibody
targeting SARS-CoV-2 nucleocapsid protein (Geneva Antibody facility; JS02) for 1 hour at room
temperature and then with peroxidase-conjugated secondary antibody (Jackson ImmunoResearch,
#109-036-09) for 30 minutes at room temperature. Foci were visualized using True Blue HRP substrate
(Avantor) and imaged on an ELISPOT reader (CTL). Focus-forming assay for comparison of infectious
viral loads in Delta vs Omicron was performed in Vero E6-TMPRSS cells.

Virus isolation

Nasopharyngeal samples were applied on the monolayer of Vero E6 cells in 24 well plates. 100 µl of
each sample was added and inoculated for 1 hour at 37°C. Following the incubation, the infectious
supernatant was discarded and virus culture medium was added. 50 µL of the medium was collected
to define the viral load at day 0. 3 to 4 days post inoculation the medium was replaced with fresh one,
and 6 days post infection the infectious medium was collected to define the viral load at day 6. The
change of viral load for at least 1 log of SARS-CoV-2 genome copies from day 0 till day 6 was considered
as successful isolation.

Statistical analysis

All statistical analyses were performed using R Statistical Software version 4.1.1 (Foundation for
Statistical 185 Computing, Austria) and Prism version 8.0.1 (GraphPad, San Diego, CA, USA).

Ethical approval

The study was approved by the Cantonal ethics committee (CCER Nr. 2021-01488). All study
participants and/or their legal guardians provided informed consent.

176 Results

177 In this study, we analysed the viral load characteristics in the URT of unvaccinated pre-VOC- as well as
178 vaccinated and unvaccinated Delta VOC-infected COVID-19 subjects up to 5 days post onset of
179 symptoms (DPOS). We included a total of 384 samples in our cohort of which 118 originated from
180 patients infected with pre-VOC SARS-CoV-2 and 248 from patients infected with the Delta VOC. Of the
181 Delta VOC infected patients, 121 were vaccinated twice prior infection and 127 were unvaccinated. In

addition, we included 18 vaccinated individuals recently infected with Omicron. None of the patients infected with pre-VOC SARS-CoV-2 were vaccinated due to the absence of licensed vaccines at the time of infection. All patients had mild symptoms at the time of sampling. Samples of pre-VOC infected patients were collected between April 7[th] and September 9[th] 2020, before circulation of any VOCs, while samples of Delta VOC infected patients were collected from June 26[th] until December 4[th] 2021 and sample of Omicron VOC infected patients from 14[th] to 17[th] of December 2021. All vaccinated patients included in this study were diagnosed positive at least 14 days after 2[nd] dose, which complies with the definition of vaccination breakthrough given by the Centers for Disease Control and Prevention (34). 132/139 patients were vaccinated with mRNA vaccines, one was vaccinated with a non-replicating viral vector vaccine (CoviVac) and for six patients the vaccine used is not known. The median time between 2[nd] dose and breakthrough infection was 79·5 (IQR 40·5-139 days) for Delta infections and 136 (IQR 85-176) for Omicron infections. All three groups of patients (pre-VOC, Delta VOC unvaccinated and Delta VOC vaccinated) had a similar age and sex distribution (see **Table 1).**

We quantified genome copies and infectious viral titres in SARS-CoV-2-positive nasopharyngeal swabs using qRT-PCR and focus-forming assay. Only specimens with a CT values below 27 for the E gene target of the diagnostic RT-PCR (Cobas, Roche), as determined by the clinical laboratory, were included in our study, since previous studies from us and others have shown that in the pre-Omicron era infectious virus cannot be reliably isolated from samples with higher CT values(12, 35). To validate our focus forming assay, we compared it to the ability to successfully isolate virus in cell culture. Virus isolation success has been used as a correlate of infectious viral shedding for SARS-CoV-2(6, 36-38), but lacks the ability to differentiate between high and low viral load samples. We were able to quantify viral titres using the focus forming assay in 91·9%, 91·7% and 83·8% of culture positive samples in the Pre-VOC, Delta and Delta vaccinated group, respectively, indicating a high sensitivity for our assay **(Figure 1A).** Overall, the Cohens kappa agreement was 0·69, 0·71 and 0·53 for the 3 groups, showing a moderate to substantial agreement **(Figure 1 B).**

Low correlation between genome copies and infectious viral titres

First, we investigated whether RNA genome copies are a good proxy for infectious virus shedding. We observed only a very low correlation (R^2 = 0·119, p=0·0001) between viral genome copies and infectious virus particles for pre-VOC samples **(Figure 2 A),** while the samples from unvaccinated and vaccinated Delta patients showed slightly higher, yet still low correlation (R^2 = 0·312, p <0·0001 and R^2 = 0·3962, p <0·0001, respectively) **(Figure 2B, C).**

No correlation between infectious viral load and age and sex of patients

Next, we tested if infectious viral loads from patient samples are associated with patient age and sex. We did not observe any correlation between the age and infectious viral load for all three groups **(Supplementary figure S1).** Similarly, no significant differences of infectious viral loads between male and female patients were detected for pre-VOC or Delta variant samples from vaccinated or unvaccinated patients **(Supplementary figure S2).**

Delta VOC infected unvaccinated patient have higher infectious viral load

Next, we compared genome copies and infectious viral loads in pre-VOC and Delta VOC samples from unvaccinated patients during the first 5 symptomatic days. Overall, pre-VOC samples had significantly higher genome copies (0·653 log, p<0·0001) compared to Delta VOC, but infectious viral titres were

significantly higher in Delta VOC infected individuals (0·343 log, p=0·0373) **(Figure 3A)**. When we
analysed the kinetics of viral shedding, we found that genome copies for pre-VOC samples were higher
at one and two dpos but similar to Delta VOC at 0, 3, 4, 5 dpos **(Figure 3B)**. Conversely, infectious virus
shedding was higher for Delta VOC at 3-5 dpos, but similar at 0-2 dpos **(Figure 3C)**. In addition, we
observed that genome copies remained largely stable until 5 dpos, with only a minimal decline at 5
dpos, while infectious viral load substantially declined **(Figure 3B and C).**

The association of the levels SARS-CoV-2 infectious shedding with patient age and sex is highly debated
(17). In this study we also did not detect a correlation between patients age or sex with their
infectiousness. However, there is increasing evidences of more severe outcomes of COVID-19 disease
in older male patients **(36, 38, 39).** Thus, to eliminate possible confounders, 84 Delta infected patients
were matched with pre-VOC infected patients in regard to sex, age and dpos. Similarly, significantly
higher infectious viral loads (0·51 log, p=0·001170) were detected in Delta variant samples in
comparison to matched pre-VOC samples **(Supplementary figure 3A).**

Vaccinated patients have lower infectious viral load than unvaccinated patients

To determine the influence of vaccination on virus shedding, we compared genome copies and
infectious viral loads in unvaccinated and vaccinated patients infected with the Delta VOC during the
first 5 dpos. Overall, genome copies did not significantly differ between vaccinated and unvaccinated
patients, while infectious viral load was significantly decreased in vaccinated patients (0·68 log,
****p<0.0001) **(Figure 4A)**. The kinetics of RNA genome copies were largely similar between vaccinated
and unvaccinated patients with a slight trend of faster decline for vaccinated patients starting at 4
dpos **(Figure 4B)**. In contrast, infectious viral titres were substantially lower in vaccinated patients at
all dpos with the biggest effect observed at 3-5 dpos **(Figure 4C)**. Still, at 5 dpos we were able to detect
infectious virus in 7/13 (53·8%) vaccinated and 11/13 (84·6%) unvaccinated patients. Additionally, 67
Delta VOC infected patients were matched with Delta VOC vaccination breakthrough patients in
regard to age, sex and dpos. Infectious viral titres were elevated in unvaccinated patients in
comparison to vaccine breakthroughs (0·97 log, p<0·0001) **(Supplementary figure 3B)** confirming a
significant reduction of infectious viral titres among vaccinated patients. We further analysed if there
is a correlation between infectious viral titres and the time interval since the administration of the last
vaccine dose. We observed a high heterogeneity between patient samples, that resulted in no
significant correlation between the time post vaccination and infectious viral shedding
(Supplementary figure S4).

In previously vaccinated subjects infection with Omicron VOC results in similar infectious viral loads
like Delta

Upon emergence of new VOC Omicron, we analysed the infectious viral shedding in vaccinated
patients infected with this variant. We compared RNA and infectious viral loads in nasopharyngeal
swab samples of 18 Omicron and 17 Delta infected patients. Vaccine breakthrough infection with
Omicron or Delta variant resulted in comparable SARS-CoV-2 genome copies (p= 0·3345). Modestly
lower infectious viral titres were detected in Omicron infected patients in comparison to Delta
infected patients, however this difference was not statistically significant (0·69 log, p= 0.1033) (Figure
5). Same moderate but non-significant reduction of infectious viral titres was observed for Omicron
samples when matching patients for age, sex and dpos (**Supplementary figure 3C**).

Discussion

In this study we analysed virus shedding in COVID-19 patients infected with pre-VOC, Delta VOC and Omicron VOC SARS-CoV-2 and evaluated the impact of vaccination on viral load in the URT during the first 5 dpos. To our knowledge, this is the first study which quantified infectious viral loads in patients infected with different SARS-CoV-2 variants and vaccination breakthrough cases. We could demonstrate a higher infectious viral load in unvaccinated Delta-infected compared to pre-VOC-infected patients and showed a significant reduction of infectious viral loads in vaccinated patients. Furthermore, we found no difference in infectious viral load between Delta and Omicron breakthrough cases.

Determining the magnitude and timing of infectiousness of COVID-19 patients is a key requirement to make informed public health decisions on the duration of isolation of patients and on the need to quarantine contacts. Infectiousness is strongly influenced by viral load in the URT of infected patients (4). However, in most studies viral load is measured as RNA genome copies and not actual infectious virus. In this study we could show that detection of RNA genome copies in URT swab samples is only poorly correlated with infectious virus shedding. This is in line with several other studies that found that RNA is a poor indicator of infectiousness especially in the presence of neutralising antibodies (12, 37).In addition, in a golden hamster animal model it was demonstrated that only infectious virus but not RNA is a good proxy for onward transmission (10).

Virus isolation success in cell culture has been widely used as a proxy for infectiousness (6, 12, 33, 40). Several studies have shown that virus isolation success significantly drops when RNA viral loads are below 6 log10 copies per mL in viral transport medium, or collected later than 8 days post symptom onset. Of note, with only a qualitative result on either successful isolation or not, overall virus isolation success cannot distinguish between high and low infectious viral loads in a patient sample, a key determinant of the size of the transmitted inoculum. Differences in infectious virus load however can have an impact on transmission probability. Therefore, we decided to use a focus forming assay instead that can reliably quantify infectious viral particles from original patient specimens. Focus forming assays have long been used as a standard to quantify excretion of virus in animal infection models for respiratory viruses such as SARS-CoV-2 and influenza virus and are therefore considered one of the best available proxies for infectiousness (41, 42).

Within the first 5 dpos, we found higher RNA genome copies in swabs of unvaccinated patients infected with pre-VOC compared to Delta VOC, but infectious viral titres were higher for Delta VOC. These results are in disagreement with other studies that analysed only nucleic acid detection and found 3-10-fold higher RNA genome copies in Delta VOC infected patients compared to pre-VOC (18, 43). However, these studies did not control for days post symptom onset, age or sex. Other studies found either no difference in RNA genome copies between Delta and pre-VOC swabs (44) or more than 1000-fold higher viral load for Delta VOC (45, 46), documenting the difficulty of comparing RNA viral loads of virus variants during different phases of the pandemic, especially when not taking additional information such as days post symptom onset into account. Conversely, in agreement with our results, a higher virus isolation success rate was observed for Delta compared to pre-VOC SARS-CoV-2 or Alpha VOC (47, 48).

Vaccines have been shown to tremendously reduce symptomatic SARS-CoV-2 infections. However, the impact of vaccination on the infectiousness of breakthrough cases is less clear. We could show that infectious viral load but not RNA genome copies are reduced in vaccinated Delta VOC patients during the first 5 days post symptom onset. In this time period approximately 50% of transmissions occur for pre-VOC SARS-CoV-2 strains (5), indicating that a reduction of viral load could considerably

contribute to a decreased secondary attack rate. Similar observations were made in other studies that
found no difference in RNA genome copies between vaccinated and unvaccinated early after symptom
onset (29, 30), but detected a lower probability of virus isolation success (30). Conversely, another
study detected up to 10-fold reduced RNA genome copies in vaccinated patients but only during the
first 60 days after complete vaccination (27). Similarly, two more studies reported decreased RNA
genome copies for vaccine breakthrough infection with pre-VOC and Alpha VOC SARS-CoV-2 (26), but
the effect vanished around 6 months post vaccination when Delta VOC was present (28). Of note, we
were still able to detect infectious viral particles in 53·8%% of vaccinated subjects at 5 dpos, indicating
that it might be beneficial is not shortened to 5 days as recently recommended by the CDC(49)..
Whether lower infectious viral load translates into lower secondary attack rate remains controversial
and depends on other influencing factors, i.e. environmental stability of virus particles. Several studies
did find a correlation between higher viral load and increased secondary attack rate, with viral load of
the index case being the leading driver of transmission (3, 4). In agreement with these finding,
epidemiological studies also showed reduced transmission rates when index cases were vaccinated,
but the effect size depends on the prevalent variant, the vaccine used and the time elapsed since
complete vaccination (22). In contrast, another study found that the vaccination status of the index
case did not influence the secondary attack rate (25). While VL is a key element of transmission, the
process of human-to-human transmission is complex and other factors, such as recommended
protection measures in a certain country at a certain time, overall incidence, perceived risks and the
context of contacts (household *vs* community transmission) can influence outcomes in the studies
reported.

To date, few data exist on viral load in vaccine breakthrough infections caused by the Omicron VOC
due to the recent emergence in late November 2021. Reduced neutralization of Omicron by infection-
and vaccine-derived antibodies was reported *in vitro* and epidemiological studies show an increased
risk of (re-)infection with Omicron in vaccinated and previously infected individuals (50, 51).
Furthermore, very high transmissibility of Omicron breakthrough infections was observed, with high
secondary attack rates even among vaccinated individuals (52). Higher RNA viral loads as described in
some studies were discussed as one potential contributing factor for the emergence of Alpha and
Delta VOC, although for Delta we could only confirm this for infectious viral load in our data. The
contribution of VL to the high transmissibility of Omicron is not known so far, neither is the mechanism
behind higher transmissibility of Omicron. First in vitro data hint towards alternative entry
mechanisms as well as early replication peak in cell culture (53), but no clinical data for this
phenomenon exist so far. Our findings indicate that with comparable RNA viral load as well as
comparable infectious virus shedding, the higher transmissibility in Omicron doesn't seem to be
related to the shedding of an increased number of infectious viral particles in vaccinated individuals.

Our study has several limitations. We included only samples with Ct values below 27, collected during
the first 5 dpos but not afterwards. Therefore, absolute numbers on RNA copies are biased towards
higher viral loads as patients with low viral load were not included here. However, patients with low
viral load have likely little relevance in terms of transmission and other factors, such as poor swab
quality can be a confounding factor leading to low viral loads. Furthermore, our focus was on
infectious virus shedding and it has been shown that SARS-CoV-2 culture is unlikely to be successfully
from samples with higher Ct values (35) and that the vast majority of secondary transmission occurs
before 5 dpos although this needs to be formally assessed in Omicron cases (5). Due to its recent
emergence, we did not yet have access to samples from Omicron infected unvaccinated individuals.
Last, we also would like to emphasize that almost all patients in this study were vaccinated with mRNA
vaccines that induce high titres of neutralizing antibodies in the blood but relatively low mucosal

antibodies. Therefore, our results cannot be generalized to other vaccines, i.e. those that are used mainly in low- and middle-income countries.

In conclusion, this study provides strong evidence for higher infectiousness of the Delta VOC as well as a significantly lower infectiousness and a faster clearance of infectious virus in vaccinated individuals. In addition, we could show that Omicron has similar infectious viral loads than Delta VOC. Furthermore, we show a more detailed picture of viral load assessment in addition to overall virus isolation success, and that quantifying viral loads can give better insights into viral shedding kinetics in acute SARS-CoV-2 infection.

Funding

This work was supported by the Swiss National Science Foundation 196644, 196383, NRP (National Research Program) 78 Covid-19 Grant 198412, the Fondation Ancrage Bienfaisance du Groupe Pictet and the Fondation Privée des Hôpitaux Universitaires de Genève.

Acknowledgments

We thank all patients for their willingness to participate in our research. We thank the staff of the laboratory of virology from the University Hospitals of Geneva for their support.

Table

	Pre-VOC SARS-CoV-2	SARS-CoV-2 Delta VOC	Vaccine breakthrough infections (Delta VOC)	Vaccine breakthrough infections (Omicron VOC)
Number	118	127	121*	18
Sampling dates	April 7 - September 9 2020	June 26 –August 29 2021	July 8 -December 4 2021	December 14 – December 17
Age				
Median (range)	36 (17-82)	37 (16-83)	40 (16-83)	35 (14-58)
<25	22 (18·6%)	19 (14·9 %)	14 (11·6%)	
25-35	37 (31·4%)	38 (29·9%)	36 (29·8%)	
35-50	30 (25·4%)	41 (32·3%)	44 (36·3%)	
50-65	23 (19·5%)	25 (19·7%)	24 (19·8%)	
>65	6 (5·1%)	4 (3·1%)	3 (2·5%)	
Sex				
Female	50 (42·4%)	65 (51·2%)	62 (51·2%)	9 (50 %)
Male	68 (57·6%)	62 (48·8%)	59 (48·8%)	9 (50 %)
RT-PCR result, CT (E-gene target, Cobas 6800, Roche)	13·9-26·6	13·8-26·3	16·3-26·1	17.2-25.9
Interval vaccination to infection, days, mean (IQR)	na	na	79·5 (IQR 40·5-139 days)	136 (IQR 85-176)
Vaccine				
BNT162b2	na	na	43	8
mRNA-1273	na	na	73	8
CoviVac	na	na	1	-
Vaccine unknown	na	na	4	2

Table 1. Patient characteristics of the specimens used in this study. RT-PCR, reverse transcription
polymerase chain reaction, CT, cycle threshold, IQR, interquartile range, na, not applicable. *Of the all
121 vaccine breakthrough samples, 104 were titrated in parallel with Delta VOC infection and 17 in
parallel with Omicron vaccine break through infections.

Figure 1. Quantitative infectious viral loads versus overall virus isolation success (**A**) Vero E6 cells
were inoculated with 10-fold serial dilutions of nasopharyngeal swabs collected from SARS-CoV-2
infected individuals. Plates were fixed 27 h post-infection and following the staining with SARS-CoV-2
specific antibodies, the number of focus forming units (FFU)/ml was calculated for each sample.
Error bars indicate mean±SD. p-values were calculated with the one-way ANOVA. ***: p<0·002;
****p<0·0001. (**B**) The total number of positive and negative samples defined by titration and virus
isolation for each patient group. Cohens kappa agreement is shown.

Figure 2. Relationship between RNA viral loads and infectious viral titers. Linear regression analysis
of infectious viral titers in FFU/ml and the corresponding RNA viral loads in nasopharyngeal swabs
from the unvaccinated patients infected with pre-VOC (**A**), and Delta SARS-CoV-2(**B**) as well as Delta
vaccination breakthroughs (**C**).

Figure 3. RNA viral load and infectious viral titers for unvaccinated individuals infected with pre-VOC SARS-CoV-2 vs. Delta (A) Genome copies (left panel) and infectious virus (right panel) for pre-VOC and Delta unvaccinated patients. Error bars indicate mean±SD. The t-test was used to determined differences of means. *p=0·0373; **** p<0·0001. Genome copies **(B)** and infectious viral loads **(C)** measured for pre-VOC and Delta VOC infected patients at different dpos. The solid lines represent the fitted curve calculated using (locally estimated scatterplot smoothing) LOESS method.

Figure 4. RNA viral load and infectious viral titers for unvaccinated vs. vaccinated individuals infected Delta (A) Genome copies (left panel) and infectious virus (right panel) for vaccinated and unvaccinated Delta infected patients. Error bars indicate mean±SD. The t-test was used to determined differences of means. ***p=0·0005; ****p<0·0001. Genome copies **(B)** and infectious viral loads (C) measured for vaccinated and unvaccinated Delta infected patients at different dpos. The solid lines represent the fitted curve calculated using (locally estimated scatterplot smoothing) LOESS method.

Figure 5. SARS-CoV-2 infectious viral loads in vaccine break through infections with Omicron or Delta. (A) Genome copies (left panel) and infectious virus (right panel) for vaccinated patients infected with Delta or Omicron VOC Infectious viral loads (were determined by focus-forming assay on Vero-TMPRSS cells. Significance was determined by t-test. ns: nonsignificant

References

1. Weekly epidemiological update on COVID-19 - 6 January 2022. World Health Organization; 2022.
2. Kawasuji H, Takegoshi Y, Kaneda M, Ueno A, Miyajima Y, Kawago K, et al. Transmissibility of COVID-19 depends on the viral load around onset in adult and symptomatic patients. PLoS One. 2020;15(12):e0243597.
3. Marc A, Kerioui M, Blanquart F, Bertrand J, Mitjà O, Corbacho-Monné M, et al. Quantifying the relationship between SARS-CoV-2 viral load and infectiousness. Elife. 2021;10.
4. Marks M, Millat-Martinez P, Ouchi D, Roberts CH, Alemany A, Corbacho-Monné M, et al. Transmission of COVID-19 in 282 clusters in Catalonia, Spain: a cohort study. Lancet Infect Dis. 2021;21(5):629-36.
5. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. Nat Med. 2020;26(5):672-5.
6. Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, et al. Virological assessment of hospitalized patients with COVID-2019. Nature. 2020;581(7809):465-9.
7. Bullard J, Dust K, Funk D, Strong JE, Alexander D, Garnett L, et al. Predicting Infectious Severe Acute Respiratory Syndrome Coronavirus 2 From Diagnostic Samples. Clin Infect Dis. 2020;71(10):2663-6.
8. Glans H, Gredmark-Russ S, Olausson M, Falck-Jones S, Varnaite R, Christ W, et al. Shedding of infectious SARS-CoV-2 by hospitalized COVID-19 patients in relation to serum antibody responses. BMC Infectious Diseases. 2021;21(1):494.
9. Vetter P, Eberhardt CS, Meyer B, Martinez Murillo PA, Torriani G, Pigny F, et al. Daily Viral Kinetics and Innate and Adaptive Immune Response Assessment in COVID-19: a Case Series. mSphere. 2020;5(6).
10. Sia SF, Yan L-M, Chin AWH, Fung K, Choy K-T, Wong AYL, et al. Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. Nature. 2020;583(7818):834-8.
11. Munker D, Osterman A, Stubbe H, Muenchhoff M, Veit T, Weinberger T, et al. Dynamics of SARS-CoV-2 shedding in the respiratory tract depends on the severity of disease in COVID-19 patients. Eur Respir J. 2021;58(1).
12. van Kampen JJA, van de Vijver DAMC, Fraaij PLA, Haagmans BL, Lamers MM, Okba N, et al. Duration and key determinants of infectious virus shedding in hospitalized patients with coronavirus disease-2019 (COVID-19). Nature Communications. 2021;12(1):267.
13. Despres HW, Mills MG, Shirley DJ, Schmidt MM, Huang M-L, Jerome KR, et al. Quantitative measurement of infectious virus in SARS-CoV-2 Alpha, Delta and Epsilon variants reveals higher infectivity (viral titer:RNA ratio) in clinical samples containing the Delta and Epsilon variants. medRxiv. 2021:2021.09.07.21263229.
14. Rapid risk assessment: Assessing SARS-CoV-2 circulation, variants of concern, non-pharmaceutical interventions and vaccine rollout in the EU/EEA, 15th update. European Centre for Disease Prevention and Control; 2021.
15. Xia S, Wen Z, Wang L, Lan Q, Jiao F, Tai L, et al. Structure-based evidence for the enhanced transmissibility of the dominant SARS-CoV-2 B.1.1.7 variant (Alpha). Cell Discov. 2021;7(1):109.
16. Dejnirattisai W, Zhou D, Supasa P, Liu C, Mentzer AJ, Ginn HM, et al. Antibody evasion by the P.1 strain of SARS-CoV-2. Cell. 2021;184(11):2939-54.e9.
17. Jones TC, Biele G, Mühlemann B, Veith T, Schneider J, Beheim-Schwarzbach J, et al. Estimating infectiousness throughout SARS-CoV-2 infection course. Science. 2021;373(6551).
18. Teyssou E, Delagrèverie H, Visseaux B, Lambert-Niclot S, Brichler S, Ferre V, et al. The Delta SARS-CoV-2 variant has a higher viral load than the Beta and the historical variants in nasopharyngeal samples from newly diagnosed COVID-19 patients. J Infect. 2021;83(4):e1-e3.

19.   Imai K, Ikeno R, Tanaka H, Takada N. SARS-CoV-2 Delta variant saliva viral load is 15-fold
higher than wild-type strains. medRxiv. 2021:2021.11.29.21266980.
20.   Hassan AO, Kafai NM, Dmitriev IP, Fox JM, Smith BK, Harvey IB, et al. A Single-Dose
Intranasal ChAd Vaccine Protects Upper and Lower Respiratory Tracts against SARS-CoV-2.
Cell. 2020;183(1):169-84.e13.
21.   Mostaghimi D, Valdez CN, Larson HT, Kalinich CC, Iwasaki A. Prevention of host-to-host
transmission by SARS-CoV-2 vaccines. Lancet Infect Dis. 2021.
22.   Harris RJ, Hall JA, Zaidi A, Andrews NJ, Dunbar JK, Dabrera G. Effect of Vaccination on
Household Transmission of SARS-CoV-2 in England. N Engl J Med. 2021;385(8):759-60.
23.   Layan M, Gilboa M, Gonen T, Goldenfeld M, Meltzer L, Andronico A, et al. Impact of
BNT162b2 vaccination and isolation on SARS-CoV-2 transmission in Israeli households: an
observational study. medRxiv. 2021:2021.07.12.21260377.
24.   Eyre DW, Taylor D, Purver M, Chapman D, Fowler T, Pouwels KB, et al. The impact of SARS-
CoV-2 vaccination on Alpha & Delta variant transmission. medRxiv.
2021:2021.09.28.21264260.
25.   Singanayagam A, Hakki S, Dunning J, Madon KJ, Crone MA, Koycheva A, et al. Community
transmission and viral load kinetics of the SARS-CoV-2 delta (B.1.617.2) variant in vaccinated
and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study. Lancet
Infect Dis. 2021.
26.   Emary KRW, Golubchik T, Aley PK, Ariani CV, Angus B, Bibi S, et al. Efficacy of ChAdOx1 nCoV-
19 (AZD1222) vaccine against SARS-CoV-2 variant of concern 202012/01 (B.1.1.7): an
exploratory analysis of a randomised controlled trial. Lancet. 2021;397(10282):1351-62.
27.   Levine-Tiefenbrun M, Yelin I, Alapi H, Katz R, Herzel E, Kuint J, et al. Viral loads of Delta-
variant SARS-CoV-2 breakthrough infections after vaccination and booster with BNT162b2.
Nature Medicine. 2021;27(12):2108-10.
28.   Pouwels KB, Pritchard E, Matthews PC, Stoesser N, Eyre DW, Vihta K-D, et al. Effect of Delta
variant on viral burden and vaccine effectiveness against new SARS-CoV-2 infections in the
UK. Nature Medicine. 2021;27(12):2127-35.
29.   Chia PY, Ong SWX, Chiew CJ, Ang LW, Chavatte JM, Mak TM, et al. Virological and serological
kinetics of SARS-CoV-2 Delta variant vaccine breakthrough infections: a multicentre cohort
study. Clin Microbiol Infect. 2021.
30.   Shamier MC, Tostmann A, Bogers S, de Wilde J, IJpelaar J, van der Kleij WA, et al. Virological
characteristics of SARS-CoV-2 vaccine breakthrough infections in health care workers.
medRxiv. 2021:2021.08.20.21262158.
31.   Badu K, Oyebola K, Zahouli JZB, Fagbamigbe AF, de Souza DK, Dukhi N, et al. SARS-CoV-2
Viral Shedding and Transmission Dynamics: Implications of WHO COVID-19 Discharge
Guidelines. Front Med (Lausanne). 2021;8:648660.
32.   Sabine Yerly LK, Manuel Schibler, Isabella Eckerle. Protocol for specific RT-PCRs for marker
regions of the Spike indicative of the Omicron variant (B.1.1.529). Geneva, Switzerland:
Centre for Emerging Viral Diseases, Geneva University Hospitals; December 2,2021.
33.   Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of 2019
novel coronavirus (2019-nCoV) by real-time RT-PCR. Euro Surveill. 2020;25(3).
34.   COVID-19 Vaccine Breakthrough Infections Reported to CDC — United States, January 1–
April 30, 2021. 2021 May 28, 2021.
35.   Essaidi-Laziosi M, Perez Rodriguez FJ, Hulo N, Jacquerioz F, Kaiser L, Eckerle I. Estimating
clinical SARS-CoV-2 infectiousness in Vero E6 and primary airway epithelial cells. Lancet
Microbe. 2021;2(11):e571.
36.   Chen PZ, Bobrovitz N, Premji ZA, Koopmans M, Fisman DN, Gu FX. SARS-CoV-2 shedding
dynamics across the respiratory tract, sex, and disease severity for adult and pediatric
COVID-19. Elife. 2021;10.

37.   Jefferson T, Spencer EA, Brassey J, Heneghan C. Viral Cultures for Coronavirus Disease 2019
Infectivity Assessment: A Systematic Review. Clin Infect Dis. 2021;73(11):e3884-e99.
38.   Takahashi T, Ellingson MK, Wong P, Israelow B, Lucas C, Klein J, et al. Sex differences in
immune responses that underlie COVID-19 disease outcomes. Nature. 2020;588(7837):315-
20.
39.   Owusu D, Pomeroy MA, Lewis NM, Wadhwa A, Yousaf AR, Whitaker B, et al. Persistent SARS-
CoV-2 RNA Shedding Without Evidence of Infectiousness: A Cohort Study of Individuals With
COVID-19. J Infect Dis. 2021;224(8):1362-71.
40.   L'Huillier AG, Torriani G, Pigny F, Kaiser L, Eckerle I. Culture-Competent SARS-CoV-2 in
Nasopharynx of Symptomatic Neonates, Children, and Adolescents. Emerg Infect Dis.
2020;26(10):2494-7.
41.   Wong L-YR, Li K, Sun J, Zhuang Z, Zhao J, McCray PB, et al. Sensitization of Non-permissive
Laboratory Mice to SARS-CoV-2 with a Replication-Deficient Adenovirus Expressing Human
ACE2. STAR Protocols. 2020;1(3):100169.
42.   Blazejewska P, Koscinski L, Viegas N, Anhlan D, Ludwig S, Schughart K. Pathogenicity of
different PR8 influenza A virus variants in mice is determined by both viral and host factors.
Virology. 2011;412(1):36-45.
43.   Christian von Wintersdorff JD, Lieke van Alphen, Petra Wolffs, Brian van der Veer, Christian
Hoebe, Paul Savelkoul. Infections caused by the Delta variant (B.1.617.2) of SARS-CoV-2 are
associated with increased viral loads compared to infections with the Alpha variant (B.1.1.7)
or non-Variants of Concern2021.
44.   Tani-Sassa C, Iwasaki Y, Ichimura N, Nagano K, Takatsuki Y, Yuasa S, et al. Viral loads and
profile of the patients infected with SARS-CoV-2 Delta, Alpha, or R.1 variants in Tokyo. J Med
Virol. 2021.
45.   Siedner MJ, Boucau J, Gilbert RF, Uddin R, Luu J, Haneuse S, et al. Duration of viral shedding
and culture positivity with post-vaccination SARS-CoV-2 delta variant infections. JCI Insight.
2021.
46.   Wang Y, Chen R, Hu F, Lan Y, Yang Z, Zhan C, et al. Transmission, viral kinetics and clinical
characteristics of the emergent SARS-CoV-2 Delta VOC in Guangzhou, China.
EClinicalMedicine. 2021;40:101129.
47.   Boschi C, Aherfi S, Houhamdi L, Colson P, Raoult D, Scola B. Isolation of 4000 SARS-CoV-2
shows that contagiousness is associated with viral load, not vaccine or symptomatic status.
Emerg Microbes Infect. 2021;10(1):2276-8.
48.   Luo CH, Morris CP, Sachithanandham J, Amadi A, Gaston DC, Li M, et al. Infection with the
SARS-CoV-2 Delta Variant is Associated with Higher Recovery of Infectious Virus Compared
to the Alpha Variant in both Unvaccinated and Vaccinated Individuals. Clin Infect Dis. 2021.
49.   Prevention CfDCa. CDC Updates and Shortens Recommended Isolation and Quarantine
Period for General Population. December 27, 2021.
50.   Eggink D, Andeweg SP, Vennema H, van Maarseveen N, Vermaas K, Vlaemynck B, et al.
Increased risk of infection with SARS-CoV-2 Omicron compared to Delta in vaccinated and
previously infected individuals, the Netherlands, 22 November to 19 December 2021.
medRxiv. 2021:2021.12.20.21268121.
51.   Pulliam JRC, van Schalkwyk C, Govender N, von Gottberg A, Cohen C, Groome MJ, et al.
Increased risk of SARS-CoV-2 reinfection associated with emergence of the Omicron variant
in South Africa. medRxiv. 2021:2021.11.11.21266068.
52.   Brandal LT, MacDonald E, Veneti L, Ravlo T, Lange H, Naseer U, et al. Outbreak caused by the
SARS-CoV-2 Omicron variant in Norway, November to December 2021. Euro Surveill.
2021;26(50).
53.   Peacock TP, Brown JC, Zhou J, Thakur N, Newman J, Kugathasan R, et al. The SARS-CoV-2
variant, Omicron, shows rapid replication in human primary nasal epithelial cultures and
efficiently uses the endosomal route of entry. bioRxiv. 2022:2021.12.31.474653.

Supplementary figures

Supplementary figure S1. Linear regression analysis of SARS-CoV-2 titers in FFU/ml and the corresponding age of the patient.

Pre-VOC (dpos 0-2)

Pre-VOC (dpos 3-5)

Delta unvaccinated (dpos 0-2)

Delta unvaccinated (dpos 3-5)

Delta vaccinated (dpos 0-2)

Delta vaccinated (dpos 3-5)

Supplementary figure S2. Comparison of infectious viral shedding measured in female and male patients. Error bars indicate mean±SD. The t-test was used to determine differences of means. ns= nonsignificant.

Supplementary figure S3. SARS-CoV-2 infectious viral loads detected in unvaccinated patients infected with pre-VOC or Delta (A), unvaccinated and vaccinated patients infected with Delta (B), vaccinated patients infected with Delta or Omicron (C) matched by age, sex, and dpos. Error bars indicate mean±SD.. **p=0·001170; ****p<0·0001, ns=nonsignificant.

Supplementary Figure S4. Linear regression analysis of infectious viral shedding and time since the completion of vaccination in Delta infected patients.