1 **TITLE**
2 **Whole genome analysis reveals the genomic complexity in metastatic cutaneous squamous**
3 **cell carcinoma**
4
5 <u>**AUTHORS AND AFFILIATIONS**</u>
6

7 Amarinder S. Thind[1,2], Bruce Ashford [1,2,3,9], Dario Strbenac[4], Ruta Gupta[5], Jonathan R Clark[4,6,7],
8 N. Gopalakrishna Iyer[8], Jenny Mitchell[3] , Jenny Lee[9,10], Simon A Mueller [7,11], Elahe Minaei[2,12],
9 Jay Perry[2,12] and Marie Ranson [2,12]
10
[1]School of Medicine, University of Wollongong, Wollongong, NSW, Australia
[2]Illawarra Health and Medical Research Institute, Wollongong, NSW, Australia
[3]Illawarra Shoalhaven Local Health District Wollongong, NSW, Australia,
[4]Sydney Medical School, Faculty of Medicine and Health, The University of Sydney, NSW,
Australia
[5]Anatomical Pathology, Royal Prince Alfred Hospital, Sydney, NSW, Australia.
[6]Royal Prince Alfred Institute of Academic Surgery, Sydney Local Health District, Sydney,
NSW, Australia
[7]Department of Head and Neck Surgery, Chris O'Brien Lifehouse, Sydney, NSW, Australia
[8]National Cancer Center, Singapore
[9]Department of Medical Oncology, Chris O'Brien Lifehouse, Sydney, NSW, Australia
[10]Department of Clinical Medicine, Macquarie University, Sydney, NSW, Australia
[11]Department of Otorhinolaryngology, Head and Neck Surgery, Zurich University Hospital and
University of Zurich, Zurich, Switzerland
[12]School of Chemistry and Molecular Bioscience, University of Wollongong, Wollongong,
NSW, Australia
27
28 <u>**CORRESPONDENCE**</u>
29 A/Prof Bruce Ashford
30 bruceash@uow.edu.au
31 372 Crown St, Wollongong, New South Wales, Australia 2500
32 +61 2 42266111

<u>ABSTRACT</u>

Metastatic cutaneous squamous cell carcinoma (cSCC) is associated with a high risk of recurrence

and poor prognosis. There is limited published data exploring whole genome sequencing (WGS).

The aim of this project was to provide the first comprehensive genomic understanding of the state

of metastatic cSCC. In this study, we used WGS on matched tumor and blood DNA to detect

somatic genetic alterations from 25 patients with regional metastases of head and neck cSCC. Our

computational analyses interrogate clinical impacts of these genetic alterations on metastatic cSCC

across the cohort for both the coding and non-coding genome.

In the non-coding genome, 3'UTR regions of *EVC* (48%), *PPP1R1A* (48%) and *LUM* (16%) were

significantly functionally altered (Q-value < 0.05). Further, significant functional alterations are

observed in the tumor suppressing lncRNA LINC01003 (68% of specimens, Q-value: 0.0158). In

addition, significant recurrent copy number loss in tumor suppressor

genes *KANSL1* and *PTPRD* and gain in *CALR, CCND1* and *FGF3* was observed for coding

regions. SNV's driver analyses predicted *TP53, CDKN2A,* as potential drivers of the metastasis

cSCC (using 3 different tools). Indel signature analysis highlight dominance of ID signature 13

followed by ID8 & ID9. Interestingly, ID 9 has previously been shown to have no association with

skin melanoma, unlike ID 13 and 8, suggesting some point of difference between these two skin-

based diseases. The overall landscape of variation in metastatic cSCC is dominated by cell cycle

and DNA repair disruption.

INTRODUCTION

Cutaneous squamous cell carcinoma (cSCC) is the second most common malignancy, after basal cell carcinoma (BCC), affecting up to 1 000 000 people in the United States (US) annually [1] . Although primary cSCC is common, lymph node metastasis only occurs in 2-5% of cSCC [2, 3]. cSCC arising in the head and neck (HNcSCC) generally show a predictable pattern of spread, predominantly metastasizing to the intraparotid, level II (upper jugular) and perifacial lymph nodes [3]. cSCC that have metastasized to regional lymph nodes are associated with a worse prognosis, with modest progress made in the management of regionally advanced disease over the last 15 years. Most patients with regional metastases from HNcSCC are managed with a multimodality approach, which usually involves a surgery (parotidectomy and neck dissection) and adjuvant external beam radiotherapy depending on the site and stage at the time of diagnosis [4]. More recently immunotherapy has attracted great interest as a potential alternative for unresectable or distant metastatic disease [5].Despite the very high incidence, relatively little is described or understood about the progression from primary cSCC to metastasis.

We, and others, have reported on somatic variation seen in primary cSCC [6, 7] . Three previous publications have included mutational analysis of metastatic cSCC [8-10] . These studies have employed either targeted next-generation sequencing (against 321, 410 or 504 genes) or whole exome sequencing to look principally at coding variants. The extent of non-coding (including regulatory) regions of the genome was limited and varied in each study. Pickering et al [8] was the only study employing whole exome sequencing and this study included only 7 metastatic samples. There was no regulatory or non-exome regions analysed in this study. Both Li et. al [10] and Zehir et. al [9] used targeted next generation sequencing, with limited non-coding analysis. Zehir et al

[9] specifically included the TERT promoter in their targeted panels but otherwise included no

regulatory elements. Li et al [10] similarly reported no regulatory of non-coding variant analysis.

No discussion in these reports addressed variation in either 5' or 3' untranslated regions (UTR).

We have previously analysed mutational burden in a cohort of 15 cSCC lymph node metastases

using whole genome sequencing (WGS) and reported the predominance of C→T transitions

associated with COSMIC UV mutational signatures 7a and 7b in metastatic cSCC, which were

particularly prevalent in non-coding DNA, and distinct from those seen in mucosal squamous cell

carcinoma [11]. The key findings of the study were the preservation of a UV associated signature

through the metastatic process and the concentration of variants within CCCTC-binding factor

regions. In the current report, we have used an expanded cohort of metastatic cSCC samples

processed by WGS for a detailed analysis of the variation in both coding and non-coding regions.

3'UTRs regulate crucial aspects of post-transcriptional gene regulation [12]. Mutations in these

regions can deregulate gene expression by disrupting miRNA-mRNA interactions, which in both

tumor suppressor genes and oncogenes can drive cancer progression [13, 14] . This variation in so

called *cis-elements* can also impact gene expression by altering translation initiation in cancer [15].

Despite the role of lncRNAs in cSCC not being fully understood, in general terms lncRNA are

thought to influence expression of proteins by pre and post translation influences on DNA/RNA

and proteins, chromatin function, miRNA activity and signalling pathways by an array of

mechanisms [16]. Their effect on miRNA might be augmented by soaking up, or sponging free

within lncRNA which impact this function can therefore have a potentiating effect on mRNA

translation hence biological effect [17].

115 The implications for functional consequences associated with both coding and these non-coding

116 genomic changes are also explored and combined to describe functional implications.

118 **MATERIALS AND METHODS**

119 This study had Institutional Human Research Ethics approval (UOW/ISLHD HREC14/397).

120 Thirty-two patients with resectable metastatic cSCC were identified by the treating surgeons

121 preoperatively. In addition to whole blood (for germline DNA), sections of fresh tumor from nodal

122 metastases were collected during surgery and immediately snap frozen. These sections were used

123 for both DNA/RNA extraction (Qiagen All Prep, Qiagen, Hilden, Germany) and for cellularity

124 estimates. Only samples with > 30% tumor (range 35-95%) proceeded to DNA quality control

125 (QC). QC comprised spectrophotography (Nanodrop 2000 ThermoFisher Scientific Inc), gel-

126 electrophoresis and SNP array. Twenty-five samples passed QC. The remaining seven samples

127 had insufficiently clonal tumor content (median variant reads ≤ 5 or median VAF < 0.1) or had an

128 extreme GC bias as determined by PURPLE [18]. Briefly, if more than 220 copy number segments

129 are unsupported by a corresponding structural variants at either end, the sample will be flagged as

130 Fail-Segment. The mean sequencing coverage of the 25 samples was 94.56X (range:64-143) for

131 tumor and 41.08X (range:30-56) for blood.

133 **Variant Calling and Functional significance of the SNVs and Indels**

134 FASTQ reads were aligned to reference genome GRChr38 using BWA-kit version 0.7.17 (BWA-

135 MEM read aligner) (more details refer to https://github.com/Sydney-Informatics-Hub/Fastq-to-

136 BAM). Further Genome Analysis Tool Kit (GATK) 4.1.2.0 and its BaseRecalibrator tool refined

137 the read alignment. Single nucleotide polymorphisms (SNPs) and insertion-deletion (indel)

variants were called by implementing GATK's Best Practices Workflow. These pipelines use

HaplotypeCaller for germline short variant discovery and Mutech2 caller for somatic short variant

discovery for SNVs and Indels (for details refer to https://github.com/Sydney-Informatics-

Hub/Somatic-ShortV). Furthermore, variants effect prediction and annotations completed using

OpenCravat platform [19]. MAF file was generated based with Variant Effect Predictor

annotations. We used three different methods for driver discovery including OncodriveFML [20]

, MutSigCV [21] and dNdScv [22] .

OncodriveFML

Functional significance of both coding and non-coding variants was predicted using the

OncodriveFML tool [20]. This tool is one of few that is designed for both the non-coding genome

and generally for WGS. It first determines the functional impact of the observed variants in

specified genomic region and later compares it with the functional impact of equal number of

random mutations from the same genomic region. The functional impact of the variants may be

computed in different manner in different genomic elements such as impact on protein function,

RNA structure, impact on microRNA Targets. Of particular interest within non-coding genome,

these regions were the 3' and 5' untranslated regions (UTR), promotor regions and long non-

coding RNA. Within 3'UTR regions of each of these genes, OncodriveFML predicted the SNVs

effect in terms of the binding and interactions of microRNA with mRNA. Others have sought to

analyse and interpret 5'UTR and promoters regions mutations in high mutational burden tumours

and reported limitations [23-25]. Increased mutational density at promoters can be linked to

decrease level of nucleotide excision repair (NER)[24, 25], high mutation rate at active

transcription factor binding sites (TFBS) due to E26 transformation-specific (ETS) TFs (which

induced cyclobutane pyrimidine dimers (CPDs) hotspots) [23].Rheinbay et. Al [26] reported

functional significance analysis issues for melanoma in their non-coding somatic driver study. We

encountered similar issues for promoters and 5' UTR regions in cSCC (Supplementary figure S.1).

MutSigCV

166 This tool identifies genes that are mutated more often than one would expect by chance. It reduces

167 the number of false positives in the generated list of significant genes, especially in tumour samples

168 with high mutation rates. Its model does this correction by incorporating various types of

169 information such as patient-specific mutation frequencies and mutation spectra, gene-specific

170 mutation rates, expression levels and replication times.

dNdScv

172 dNdScv is designed to test for positive and negative selection in cancer genomes. For the better

173 accuracy of the dNdScv model on UV originated cancer, we carefully monitored the annotation of

174 CC>TT changes (sometimes reported as C>T changes). Results report significance for missense,

175 truncating, Indel and then one global value. Global significance was considered for the

176 interpretation of outcomes.

Copy Number Variation

Using PURPLE [18], we derived copy number alterations on 25 metastatic specimens. PURPLE

estimates overall ploidy of the samples by using read depth ratio from COBALT and tumor B-

allele frequency (BAF) from AMBER. The pipeline is available at github of HMF Tools

(https://github.com/hartwigmedical/hmftools). Driver genes with significant amplifications and

deletions were then identified from PURPLE based driver gene identification outputs.

To explore the contribution of our analysis employing WGS to the understanding of the degree of

somatic variation in metastatic cSCC, we then compared short variants of the top 6 statistically

significant genes to all previously reported metastatic cSCC [8-10] using processed data available

on the cBioPortal [27, 28] to those within our cohort. Within cBioPortal, there are 63 samples of

nodal metastatic cSCC analyzed by either WES (n = 7) and targeted NGS (n = 56).

GRIDSS2 and its companion interpreter tool LINX were employed for somatic structural variant

analysis and Gene fusion [29]. COSMIC3 based, SNVs and Indels signatures from the whole

genome were built using deconstructSigs [30] and MutationalPatterns [31] software respectively.

RESULTS

Patient characteristics and clinicopathologic data

Twenty-five metastatic cSCC samples from lymph nodes in the head and neck region were

collected between 2015 and 2019. The median age of patients was 69 (range 30-87) and 24/25

(96%) were male. This sex disparity is in keeping with the disease burden we see in our practice,

particularly for advanced and metastatic cSCC (Ashford et al., manuscript in preparation). During

the period of patient recruitment for this study, 88% of patients with nodal metastasis of cSCC

were male. Further, due to histopathologic tumor percentage estimation and other quality control

measures, in this study only 1 female sample was sequenced. Others have highlighted that age,

male sex and immunosuppression are among risk factors for metastasis [32] while another study

reports sex determined immunological mechanisms as a determinant of the development of

metastasis in cSCC [33]. Two patients were immunocompromised; one patient was on long-term

azathioprine for rheumatoid arthritis and the other was on a combination of cyclophosphamide and

tacrolimus following solid organ transplantation.

The location of the index primary lesion was known in 11 patients ;lip (n = 2), ear (n = 2),

forehead/scalp (n = 4), cheek (n = 2) and nose (n = 1) . Nodal metastases were isolated from the

neck in 13 patients and in the parotid in 12 patients. The majority of patients had either moderately

differentiated (n = 8) or poorly differentiated (n = 12) cSCC, with evidence of extranodal extension

found in 20/25 (80%) nodal samples.

Tumor mutational burden (TMB)

The average TMB for SNV across the 25 cases was 238.7 mutations per megabase (range 32.52to

995.66 mutations/Mb) and for Indel it was indel/megabase 2.25 (range 0.63 to 5.9 mutations/Mb).

(Figure.1(F,G))(S.Table1) . The only female in our cohort had the second highest TMB at 499

mutations/Mb. here was no correlation between age, differentiation, nodal stage or extracapsular

spread of the metastasis and TMB. As we have previously reported, a mutation density across the

genome 171 times higher for non-coding variants compared to coding variants [11]. This was

corroborated in our expanded cohort of metastatic cSCC.

Figure1:- shows the overview of tumor mutational burden, signature and mutations types (A-D). Details of the coding mutations of the cSCC genomes are shown in figure 1(A to D), such as variants classification, variant types etc. Whereas Figure 1 (E, F, G) shows the whole genome-based Mutational Signatures, Indel and SNV Tumor mutational burden for each sample.

Mutational signatures

We   performed   mutational   signature   analyses   based   on   COSMIC   V.3.2

(https://cancer.sanger.ac.uk/signatures/). Signatures are designated as single base substitution

(SBS), double base substitution (DBS) or small insertion and deletion (ID) signatures. SNV

signature analysis of our 25 samples shows SBS signatures 7a and 7b are dominant (Figure 1E)

Substantial representation of SBS7c is also seen. SBS32 and SBS7d are observed in one sample.

Indel signature analysis of our samples showed that ID8, 9 and 13 dominate over others (Figure 2)

(S.Table2).

Figure 2: Indel signatures based on cosmic3 database. In the figure X-axis represent the sum of score for signature
among 25 samples. Y axis shows the indel signatures. In our cohort ID13, ID8 and ID9 predominate over other Indel
signatures. Full details are available in supplementary (S.Table2(Tab2-4)).
243

<u>Short variants</u>

Variation in non-coding regulatory regions

Variants in the 3' UTRs of key cancer associated genes with high functional impact on miRNA-

mRNA interactions were evident in our cohort of metastatic cSCC. Among our chort of samples,

within the 3'UTR region of *EVC, LUM* and *PPP1R1A*, non-coding SNVs showed significant

higher observed functional impact than the predicted functional impact (refer to method section)

(Figure 3b). We observed variation within the 3'UTR of *EVC* in 48% of samples with a high

predicted functional impact. Similarly, high impact mutations in the 3'UTR of *PPP1R1A* were

observed in the 48% of samples. We also observed one unique variant i.e *PPP1R1A*

Chr12:54579896 (G to A) variant with cDNA change of c.*491C>T in five samples (S.Table3)

Figure 3 shows various coding and non-coding genes or regions having significant alterations in cSCC metastasis. (a –c) Plots showing the most significantly altered (in the plots above the red line Q-values are below 0.1 and between the green line & the red line Q-values are 0.25 to 0.1.) Q-values are corrected P-values using the Benjamini/Hochberg

correction (Where P-values are defined by times that the observed value is >= the expected value, divided by the number of randomizations) (a) coding region (genes), (b) 3' UTR regions of the genes, and (c) lncRNA regions in the cohort of 25 patient samples. Plots showing the frequency of observed mutations with respect to the expected frequency of the mutations in the corresponding regions. These analyses were done using OncodriveFML (2.2.0). (d) Detailed sample-level information of the SNVs along with types of variants in the top altered genes (refer to panel a). (e) Plots showing variants of significantly altered 3'UTR regions. (f). Oncplot retrieved from cBioportal to compare the variants from panel (d) with other metastatic cSCC studies, which are either whole exome or targeted panel based. Due to the use of targeted panel in some CSCC studies, not all genes from panel (d) were measured.

lncRNA were analysed to determine a potential impact on cancer associated mRNA. An example of this is in lncRNA LINC01003, which is amongst the frequently mutated lncRNA in our cohort (71% of samples). Other recurrently varied lncRNA in our cohort were RP11-151D14.1, RP4-597N16.4 and RP11-61J19.4 (Figure 3c).

Coding Short Variants

The overwhelming majority of coding SNVs were missense mutations, followed by nonsense mutation, which represented less than 5% of variants. Over 80% of SNV mutations were C>T, consistent with the dominant effect of UV radiation on pyrimidine bases. This is in keeping with our recently published findings on mutational signatures according to COSMIC V.2 (Mueller et al., 2019). Genes most frequently subject to high impact SNV included *TP53, CDKN2A and ZNF730* having Q-value <0.1, whereas Q-value of *OR51S1, CCDC68* and *TET2* is between 0.1 – 0.25 (Figure 3a).

The only sample to show no significant mutation in *TP53* was CSCC_0009. The TMB of this sample was 122/Mb, or 51% of the average across the cohort. The five samples without *CDKN2A* mutations average a TMB of 470/Mb, or 201% of the average for the cohort. In nine cases, high impact short variants of *ZNF730* was observed.

Structural variation and gene fusions

Structural variation analysis revealed substantial chromosomal disruption in the metastases

characterized by both translocation and inversion and often accompanied by allele specific copy

number variation. Minor allele loss, large scale deletions and amplifications were observed .

Despite this widespread genomic instability, only a single coding-coding gene fusion was observed

in our cohort – this was between *STRN* and *DLG2* in sample CSCC_0009 (*STRN*: Exon 1

ENST00000263918 *DLG2* Exon 7 ENST00000376104). *STRN* encodes a calcium-dependent

calmodulin-binding protein [34]. *DLG2* plays a role in pain signalling and deletion is seen in both

human and canine osteosarcoma [35]. CSCC_0009 came from a patient who had undergone liver

transplantation and was on immunosuppressive therapy. The primary tumor that gave rise to this

metastasis showed perineural involvement, but this was present in the metastatic deposit.

Copy Number Variation

299 The extent of chromosomal copy number gain and loss was averaged across the genome for all 25

300 tumor samples (Figure 4). Chr5p and 8q were the most frequently amplified regions, with 18q

301 being the region with the most recurrent deletion. Previously reported [36] recurrent 9p loss was

302 also seen.

Figure 4. Chromosomal and recurrent genetic copy number variation in metastatic cSCC (a) Combined chromosomal CNV across 25 metastatic cSCC samples at chromosomal level. X-axis represents the differences of mean minimum copy number (bands) and Mean of overall samples ploidy (after adjustment for purity). For details please refer to S.Table4 (Chromosome_plot_raw_Mean_difference_file). (b) Figure show chromosome Arm loss and Gain at sample level, where red color is for Gain and Blue color is for loss. Where both arm of Chromosome 7 and 5p is gained, and 8p, 18q and 21q shows loss. (A chromosome arm is defined to be deleted if at least half of its bases are one or more

311 copies less than the sample ploidy. A chromosome arm is defined to be amplified if at least half of its bases are one
312 or more copies more than the sample ploidy.)
313

The recurrent amplified and deleted genes across all 25 tumor samples are presented in Figure 2b.

The most frequently deleted genes were *KANSL1* and *PTPRD*. Recurrent amplification was seen

less commonly.

KANSL1 (Ch17q) showed deletion in 32% of samples. Deletion of *PTPRD* (Chr9p) was seen in

20% of samples. Deletion of *PTPRD* (n=6) and *CDKN2A* (Chr9p) (n=1) did not co-occur in our

cohort (Figure 5), although *PTPRD* loss and significant mutation of *CDKN2A* occurred together

in five samples (CSCC_9, 11, 133, 132 and 134).

Figure 5:- Recurrent copy number gain and loss for individual genes, Deletion CN<0.5 and Amplification, minimum
exonic copy number > 3 x sample ploidy >4 adjusted to cellularity and ploidy. These driver genes were derived by
the use of purple-3.1. For details please refer to S.Table5.
325

Amplification was seen most commonly in *CALR*, *CCND1* and *FGF3* (Figure 5). Amplification

of *CCDN1* and *FGF3* co-occurred in two samples (CSCC_0134 and CSCC_0132). These two

cases had extensive nodal involvement being the only samples with >50% of lymph nodes

involved by tumor.

DISCUSSION

This is the largest study to employ WGS to assess the mutational landscape of metastatic cSCC

and provides unique insights into the expansive array of variation. The analysis presented here

demonstrates the breadth of somatic variation in metastatic cSCC across non-coding and coding

regions, including the identification of novel Indel signature patterns, and highlights for the first

time the nature and depth of this variation within regulatory regions. We have previously published

an analysis of WGS of a smaller cohort of metastatic cSCC in which we focussed on the

preservation of UV associated COSMIC signature 7b and the density of mutations found in the

CTCF binding regions of the genome. Herein we have doubled the samples analysed and adopted

a more comprehensive survey of the extent of variation observed using WGS. The utility of WGS

has enabled an understanding of the extent of variation beyond coding regions and has defined the

prominence of non-coding variation, with special attention devoted to 3'UTR, lncRNA.

Additionally, WGS analysis has allowed us to compare the extent of variation to that previously

reported in metastatic cSCC, previously studies relying on WES and targeted NGS, and to update

and expand the understanding of mutational signature patterns in cSCC.

The rate of TMB within metastatic cSCC is greater than any tumour previously described. At 238

mutations/Mb, this is substantially higher than other cancers known to have high mutational

burden, including melanoma (49 mutations/Mb) [37] and NSCLC (10 mutations/Mb) [36]. This

finding is in keeping with Pickering et al [8] who found a TMB of 61.2 mutation/Mb of four times

that of melanoma in their high risk cSCC. Their series included just seven metastatic samples.

According to the formula established by Yarchoan[38] for assessing predicted response rate to PD-

1 inhibition, response rate = $10.8 \times \log_e(X)$-0.7, where X is the mutational burden (mutations/Mb).

The calculated potential response rate to PD-1 inhibition in our cohort was 39.6%, which is

consistent with current literature [5]. Immune checkpoint inhibitors have recently demonstrated

high, durable response rates and superior tolerability when compared to platinum chemotherapy

in unresectable locally advanced and metastatic cSCC [39]. The high TMB was associated with

substantial structural variation, without recurrent gene fusions.

As we have previously reported in a smaller cohort, the metastatic process preserves the dominant

UV associated variant pattern of C>T, or CC>TT in cSCC. This finding is independent of the

degree of differentiation or any other clinicopathologic feature.

Alexandrov et al detailed patterns of mutational signatures in 23829 tumour samples (1965 WGS)

from the Pan Cancer Analysis of Whole Genomes (PCAWG) datasets defining single base

substitution (SBS), double base substitution (DBS) and small insertion and deletion signatures (ID)

(Alexandrov, Kim et al., 2021). No cutaneous SCC (primary or metastatic) are included in this

dataset. SBS7a/SBS7b/SBS7c/SBS7d were all seen in skin melanoma in Alexandrov et al. In our

metastatic cSCC analysis, SBS7a and b were the most dominant signatures, with less presence of

SBS 7c and d. Alexandrov et al identified 17 ID mutational signatures. COSMIC version 3.2

(https://cancer.sanger.ac.uk) has expanded the ID signatures to 18 [40] .

Indel signature analysis on our 25 metastatic cSCC samples identified the predominance of ID

signatures 8, 9 and 13 (100% of samples effected). ID 8 is thought to be both related to double

strand DNA break repair dysfunction and to age related changes. The mechanism of ID 9 is unclear.

In their analysis, Alexandrov et al found ID 9 in only 2/104 melanoma samples. Thus, our finding

represents a departure from the prevailing dominance of ID 9 in mainly soft tissue sarcoma within

the PCAWG data. Alexandrov described only one cancer type with ID 13 as predominant, this

being skin melanoma. Our analysis revealed 100% of samples with ID 13 in metastatic cSCC. Our

data also provides evidence of concomitance of ID 13 with SBS 7a and 7b in keeping with a UV

association.

We have demonstrated that both Indel and somatic variant signatures are different for metastatic

cSCC than for other malignancies of the head and neck. The potential for both Indel and somatic

variation signatures to define the likely aetiology of nodal metastatic disease from an "unknown

primary SCC" would allow for discrimination between skin and mucosal origin. This would have

tremendous treatment impacts and would dictate the benefits of surgery, radiotherapy and systemic

(both chemotherapy and immunotherapy) treatments. Such analysis could be completed on

smaller genomic regions thus allowing for a cheaper and quicker alternative to WGS or WES.

Copy Number Variation

Loss of *KANSL1* and *PTPRD* were the most prominent CN alterations seen within our 25 samples,

with amplification most common in *CALR*, *CCND1* and *FGF3*. *KANSL1* is a chromatin modifier

gene located on the long arm of chr17 with deletion observed in 32% of samples. Germline loss

of this gene is associated with 17q21.31 microdeletion syndrome [41]. In contrast, epigenetic

analysis of epithelial ovarian cancer identified amplification of *KANSL1* as a means for potential

targeted therapy [42].

Deletion of *PTPRD* was seen in 20% of our samples. Lambert et al. [43] described deletions of

PTPRD in 37% of either primary cSCC that had metastasised or metastatic disease. In addition,

some of their cases also displayed a variant in the minor allele concordant with the deletion leading

to a loss of heterozygosity event. *PTPRD* encodes protein tyrosine phosphatase receptor D, which

belongs to a family of receptors whose action oppose that of the tyrosine kinases, which are central

to cell growth and differentiation and oncogenic transformation (genecards.org). Large scale

genomic events impacting *CDKN2A* can also affect *PTPRD* due to their proximity on chr9 [44].

PTPRD dephosphorylates S*TAT3*, deactivating its tumorigenic activity. In head and neck SCC,

PTPRD inactivation significantly increased *STAT3* hyperactivation, which was associated with

decreased survival and resistance to EGFR-targeted therapy [45]. *PTPRD* has been implicated as

a tumor suppressor in several cancers with inactivating somatic variants found in >50% of GBM

and between 10-20% of head and neck mucosal SCC (HNSCC)[46]. It is possible that *PTPRD*

plays a tumor suppressor role in metastatic cSCC.

Amplification was seen most commonly in *CALR*, *CCND1* and *FGF3* (Figure 5). *CALR* encodes

a ubiquitous endoplasmic reticulum bound calcium receptor [47]. Cellular stress can move *CALR*

fragments to the plasma membrane from the ER and influence immune recognition of cancer cells.

Recent analysis of *CALR* fragments in myeloproliferative disease suggest an immunosuppressive

influence of extracellular *CALR* [48]. Cyclin D1 (*CCND1*) amplification is associated with nodal

metastasis and worse survival in oral SCC [49]. In a review of *CCND1* copy number variation in

metastatic non-cutaneous melanoma, amplification of *CCDN1* was prominent in those patients

whose disease did not respond to immune checkpoint inhibition [50]. *FGF3* amplification is more

common in metastatic breast cancer than primary tumors [51]. Targetable *FGF3* amplification was

associated with a poorer prognosis and lung metastasis in hepatocellular carcinoma [52]. This

amplification was seen in only 2% of total HCC but was most common in those cancers showing

rapid response to sorafenib.

Non coding short variants

We identified substantial somatic variation within the 3'UTR region of *EVC, LUM* and *PPP1R1A.*

EVC effects ciliary Hedgehog (Hh) regulation. Aberrant overexpression of *EVC* (and upregulation

of Hh) has been reported in adult T-cell leukaemia as a result of epigenetic modulation [53]. Within

breast cancer, the expression of *EVC* was found to be reduced in nodal deposits of metastatic breast

cancer compared with primary breast cancer suggesting a role in the metastatic process [54].

PPP1R1A is a protein phosphatase inhibitor which appears to have a variable but significant role

in the metastatic process. In Ewing Sarcoma, it is overexpressed and has been proposed as a driver

of metastasis [55]. Conversely, levels of *PPP1R1A* were reduced in breast cancer when compared

to adjacent non-diseased breast tissue [56]. Within our cohort, we observed a recurrent missense

mutation in the 3'UTR of *PPP1R1A* in five samples.

LINC01003 was the most mutated lncRNA in our cohort (71% of samples). In multiple

myeloma, LINC01003 behaves as a tumor suppressor genomic element. Up-regulation

suppresses multiple myeloma by repressing cell viability and adhesion and promoting apoptosis.

This effect is via its sponge effect on miR-33a-5p and its target *PIM1 [57]*.

Coding short variants

The genes with most recurrent variation were *TP53, CDKN2A*. Of note was the absence of

significant or recurrent SNV affecting NOTCH1/2. Inman et al (2018) compared well

differentiated to moderately and poorly differentiated primary cSCC and identified *NOTCH1*,

NOTCH2, TP53 and *CDKN2A* as the most commonly mutated genes, with *ATP1A1, HERC6,*

MAPK1P1L, GRHL2, TRAPPC9, FLNB and *MAP3K9* identified as common early events in

primary cSCC (Inman et al., 2018). Within this group, *GRHL2* was associated with less well

differentiated tumours including those with a worse prognosis, and the authors suggested it's role

needed further assessment in cSCC. We found a single example of splice variant in our 25 samples

in *GRHL2,* suggesting its role in metastatic disease is limited.

Loss of function mutations within *TP53* and *CDKN2A* adversely impact cell cycle pathway control

and DNA repair mechanisms and mutations in these genes has been frequently reported for cSCC

[58]. Kilnakis et al describe a pattern of *TP53* mutation that differed between primary and

metastatic disease in head and neck (mucosal) SCC [59]. They found an overall lower rate of

mutations in metastatic tumors, but a higher concentration of missense mutations in the DNA

binding regions of the gene. Additionally, they found a higher TMB with *TP53* mutation, but a

worse response to immunotherapy. Burtness et al [60] reported that the extent of TMB in HPV-

negative HNSCC is associated with loss of function mutations in both *TP53* and *CDKN2A* .

Loss of function mutations in *ZNF730* are associated with poor prognosis in cervix, lung and

HNSCC and its tumor suppressive role is mediated via *TINCR* and *LAMC2* to effect cell

differentiation, migration and death [61]. Its role in the metastatic phenotype of cSCC has not been

previously reported and herein we showed a high level of deleterious single nucleotide variation

with nine samples showing either a missense or a stop gained mutation. *ZNF730* showed no

mutation in the only study that included its analysis [8].

OR51S1 encodes a gene that regulates olfactory receptor function via a G protein-coupled receptor

pathway (genecards.org). It could be false positive as reported by Lawrence et al. [21] olfactory

receptor genes are long protein coding genes. In our case the Q-value is 0.115 which is not too

low.

High impact short variants of TET2 were seen in 44% of sample. Deletion of *TET2* is understood

to play a role in haemopoietic malignancy [62]. More recently, loss of *TET2* activity was

associated with reduced tumor chemokine production and PD-L1 expression via IFN-

γ/JAK/STAT signalling pathway, conferring an increased resistance to immunotherapy in

melanoma cells [63]. Previous studies [8-10] have noted this variant in metastatic cSCC at a rate

of 20% (Figure 3f). Further interrogation of the TCGA dataset for metastatic cSCC revealed 63

cases [8-10]   with short variant analysis (Figure 3f). Of these only seven cases had undergone

WES with the remainder having undergone targeted NGS. Not all genes identified as high impact

and recurrent variants in our cohort were included in these studies, which underscores the

deficiencies of targeted analyses in discovery projects.

<u>CONCLUSION</u>

WGS provides insight into the unparalleled burden of mutation within metastatic cSCC. In this

study we describe a complex interaction of inactivation of putative tumor suppressor genes and

amplification of oncogenes with biologically significant regulatory region variation. The

combination of checkpoint dysregulation and the high mutation burden underline the response to

immunotherapy in metastatic cSCC. The examination of Indel and somatic variant signatures

provide a path to more accurate determination of the origin of metastatic SCC. This analysis of

genomic variation provides an opportunity for a deeper understanding of the metastatic cSCC

landscape to be the foundation for the identification of biomarkers in primary cSCC predicting

metastasis.

<u>ACKNOWLEDGMENTS</u>

This work was funded by the Illawarra Cancer Carers, The Head and Neck Research Fund, Royal

Prince Alfred Institute of Academic Surgery, The Cancer Institute NSW translational program

grant, Chris O'Brien Lifehouse, and National Health and Medical Research Council Project Grant

APP1181179.

<u>CONFLICT OF INTEREST STATEMENT</u>

The authors state no conflict of interest

1. Rogers, H.W., et al., *Incidence estimate of nonmelanoma skin cancer (keratinocyte carcinomas) in the US population, 2012.* JAMA dermatology, 2015. **151**(10): p. 1081-1086.
2. Venables, Z.C., et al., *Nationwide incidence of metastatic cutaneous squamous cell carcinoma in England.* JAMA dermatology, 2019. **155**(3): p. 298-306.
3. Forest, V.I., et al., *N1S3: a revised staging system for head and neck cutaneous squamous cell carcinoma with lymph node metastases: results of 2 Australian Cancer Centers.* Cancer: Interdisciplinary International Journal of the American Cancer Society, 2010. **116**(5): p. 1298-1304.
4. Veness, M., *Treatment recommendations in patients diagnosed with high-risk cutaneous squamous cell carcinoma.* Australasian Radiology, 2005. **49**(5): p. 365-376.
5. Migden, M.R., et al., *PD-1 blockade with cemiplimab in advanced cutaneous squamous-cell carcinoma.* New England Journal of Medicine, 2018. **379**(4): p. 341-351.
6. Zilberg, C., et al., *Analysis of clinically relevant somatic mutations in high-risk head and neck cutaneous squamous cell carcinoma.* Modern Pathology, 2018. **31**(2): p. 275-287.
7. Inman, G.J., et al., *The genomic landscape of cutaneous SCC reveals drivers and a novel azathioprine associated mutational signature.* Nature communications, 2018. **9**(1): p. 1-14.
8. Pickering, C.R., et al., *Mutational landscape of aggressive cutaneous squamous cell carcinoma.* Clinical cancer research, 2014. **20**(24): p. 6582-6592.
9. Zehir, A., et al., *Mutational landscape of metastatic cancer revealed from prospective clinical sequencing of 10,000 patients.* Nature medicine, 2017. **23**(6): p. 703-713.
10. Li, Y.Y., et al., *Genomic analysis of metastatic cutaneous squamous cell carcinoma.* Clin Cancer Res, 2015. **21**(6): p. 1447-56.
11. Mueller, S.A., et al., *Mutational patterns in metastatic cutaneous squamous cell carcinoma.* Journal of investigative dermatology, 2019. **139**(7): p. 1449-1458. e1.
12. Barrett, L.W., S. Fletcher, and S.D. Wilton, *Regulation of eukaryotic gene expression by the untranslated gene regions and other non-coding elements.* Cellular and molecular life sciences, 2012. **69**(21): p. 3613-3634.
13. Liu, W. and X. Wang, *Prediction of functional microRNA targets by integrative modeling of microRNA binding and target expression data.* Genome biology, 2019. **20**(1): p. 1-10.
14. Schuster, S.L. and A.C. Hsieh, *The untranslated regions of mRNAs in cancer.* Trends in cancer, 2019. **5**(4): p. 245-262.
15. Kobayashi, H. and Y. Tomari, *RISC assembly: coordination between small RNAs and Argonaute proteins.* Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms, 2016. **1859**(1): p. 71-81.
16. Statello, L., et al., *Author Correction: Gene regulation by long non-coding RNAs and its biological functions.* Nat Rev Mol Cell Biol, 2021. **22**(2): p. 159.
17. Cesana, M., et al., *A long noncoding RNA controls muscle differentiation by functioning as a competing endogenous RNA.* Cell, 2011. **147**(2): p. 358-69.
18. Priestley, P., et al., *Pan-cancer whole-genome analyses of metastatic solid tumours.* Nature, 2019. **575**(7781): p. 210-216.

19. Pagel, K.A., et al., *Integrated Informatics Analysis of Cancer-Related Variants*. JCO Clin Cancer Inform, 2020. **4**: p. 310-317.

20. Mularoni, L., et al., *OncodriveFML: a general framework to identify coding and non-coding regions with cancer driver mutations*. Genome Biol, 2016. **17**(1): p. 128.

21. Lawrence, M.S., et al., *Mutational heterogeneity in cancer and the search for new cancer-associated genes*. Nature, 2013. **499**(7457): p. 214-218.

22. Martincorena, I., et al., *Universal Patterns of Selection in Cancer and Somatic Tissues*. Cell, 2017. **171**(5): p. 1029-1041.e21.

23. Mao, P., et al., *ETS transcription factors induce a unique UV damage signature that drives recurrent mutagenesis in melanoma*. Nat Commun, 2018. **9**(1): p. 2626.

24. Perera, D., et al., *Differential DNA repair underlies mutation hotspots at active promoters in cancer genomes*. Nature, 2016. **532**(7598): p. 259-63.

25. Sabarinathan, R., et al., *Nucleotide excision repair is impaired by binding of transcription factors to DNA*. Nature, 2016. **532**(7598): p. 264-7.

26. Rheinbay, E., et al., *Analyses of non-coding somatic drivers in 2,658 cancer whole genomes*. Nature, 2020. **578**(7793): p. 102-111.

27. Cerami, E., et al., *The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data*. Cancer Discov, 2012. **2**(5): p. 401-4.

28. Gao, J., et al., *Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal*. Sci Signal, 2013. **6**(269): p. pl1.

29. Cameron, D.L., et al., *GRIDSS2: comprehensive characterisation of somatic structural variation using single breakend variants and structural variant phasing*. Genome Biol, 2021. **22**(1): p. 202.

30. Rosenthal, R., et al., *DeconstructSigs: delineating mutational processes in single tumors distinguishes DNA repair deficiencies and patterns of carcinoma evolution*. Genome biology, 2016. **17**(1): p. 1-11.

31. Blokzijl, F., et al., *MutationalPatterns: comprehensive genome-wide analysis of mutational processes*. Genome medicine, 2018. **10**(1): p. 1-11.

32. Tokez, S., et al., *Cumulative incidence and disease-specific survival of metastatic cutaneous squamous cell carcinoma: a nationwide cancer registry study*. J Am Acad Dermatol, 2021.

33. Budden, T., et al., *Female Immunity Protects from Cutaneous Squamous Cell Carcinoma*. Clin Cancer Res, 2021. **27**(11): p. 3215-3223.

34. Du, Q.Y., et al., *High STRN Expression Promotes HCC Invasion and Migration but Not Cell Proliferation or Apoptosis through Facilitating Epithelial-Mesenchymal Transition*. Biomed Res Int, 2020. **2020**: p. 6152925.

35. Shao, Y.W., et al., *Cross-species genomics identifies DLG2 as a tumor suppressor in osteosarcoma*. Oncogene, 2019. **38**(2): p. 291-298.

36. Alexandrov, L.B., et al., *Signatures of mutational processes in human cancer*. Nature, 2013. **500**(7463): p. 415-21.

37. Hayward, N.K., et al., *Whole-genome landscapes of major melanoma subtypes*. Nature, 2017. **545**(7653): p. 175-180.

38. Yarchoan, M., A. Hopkins, and E.M. Jaffee, *Tumor Mutational Burden and Response Rate to PD-1 Inhibition*. N Engl J Med, 2017. **377**(25): p. 2500-2501.

39. Migden, M.R., et al., *Cemiplimab in locally advanced cutaneous squamous cell carcinoma: results from an open-label, phase 2, single-arm trial.* Lancet Oncol, 2020. **21**(2): p. 294-305.

40. Tate, J.G., et al., *COSMIC: the Catalogue Of Somatic Mutations In Cancer.* Nucleic Acids Res, 2019. **47**(D1): p. D941-d947.

41. Zollino, M., et al., *Mutations in KANSL1 cause the 17q21.31 microdeletion syndrome phenotype.* Nat Genet, 2012. **44**(6): p. 636-8.

42. Fejzo, M.S., et al., *Analysis in epithelial ovarian cancer identifies KANSL1 as a biomarker and target gene for immune response and HDAC inhibition.* Gynecol Oncol, 2021. **160**(2): p. 539-546.

43. Lambert, S.R., et al., *Metastatic cutaneous squamous cell carcinoma shows frequent deletion in the protein tyrosine phosphatase receptor Type D gene.* Int J Cancer, 2012. **131**(3): p. E216-26.

44. Ortiz, B., et al., *Deletion of Ptprd and Cdkn2a cooperate to accelerate tumorigenesis.* Oncotarget, 2014. **5**(16): p. 6976-82.

45. Veeriah, S., et al., *The tyrosine phosphatase PTPRD is a tumor suppressor that is frequently inactivated and mutated in glioblastoma and other human cancers.* Proc Natl Acad Sci U S A, 2009. **106**(23): p. 9435-40.

46. Peyser, N.D., et al., *Loss-of-Function PTPRD Mutations Lead to Increased STAT3 Activation and Sensitivity to STAT3 Inhibition in Head and Neck Cancer.* PLoS One, 2015. **10**(8): p. e0135750.

47. Holmström, M.O., et al., *The calreticulin (CALR) exon 9 mutations are promising targets for cancer immune therapy.* Leukemia, 2018. **32**(2): p. 429-437.

48. Liu, P., et al., *Immunosuppression by Mutated Calreticulin Released from Malignant Cells.* Mol Cell, 2020. **77**(4): p. 748-760.e9.

49. Miyamoto, R., et al., *Prognostic significance of cyclin D1 amplification and overexpression in oral squamous cell carcinomas.* Oral Oncol, 2003. **39**(6): p. 610-8.

50. Yu, J., et al., *Genetic Aberrations in the CDK4 Pathway Are Associated with Innate Resistance to PD-1 Blockade in Chinese Patients with Non-Cutaneous Melanoma.* Clin Cancer Res, 2019. **25**(21): p. 6511-6523.

51. Rinaldi, J., et al., *The genomic landscape of metastatic breast cancer: Insights from 11,000 tumors.* PLoS One, 2020. **15**(5): p. e0231999.

52. Arao, T., et al., *FGF3/FGF4 amplification and multiple lung metastases in responders to sorafenib in hepatocellular carcinoma.* Hepatology, 2013. **57**(4): p. 1407-15.

53. Takahashi, R., et al., *Epigenetic deregulation of Ellis Van Creveld confers robust Hedgehog signaling in adult T-cell leukemia.* Cancer Sci, 2014. **105**(9): p. 1160-9.

54. Mamoor, S., *EVC is differentially expressed in lymph node metastasis in human breast cancer.* 2021.

55. Luo, W., et al., *Protein phosphatase 1 regulatory subunit 1A in ewing sarcoma tumorigenesis and metastasis.* Oncogene, 2018. **37**(6): p. 798-809.

56. Yuan, C.L., et al., *Identification of differentially expressed lncRNAs and mRNAs in luminal-B breast cancer by RNA-sequencing.* BMC Cancer, 2019. **19**(1): p. 1171.

57. Wu, L., et al., *Long non-coding RNA LINC01003 suppresses the development of multiple myeloma by targeting miR-33a-5p/PIM1 axis.* Leuk Res, 2021. **106**: p. 106565.

58. Ashford, B.G., et al., *Reviewing the genetic alterations in high-risk cutaneous squamous cell carcinoma: A search for prognostic markers and therapeutic targets.* Head Neck, 2017. **39**(7): p. 1462-1469.

59. Klinakis, A. and T. Rampias, *TP53 mutational landscape of metastatic head and neck cancer reveals patterns of mutation selection.* EBioMedicine, 2020. **58**: p. 102905.
60. Burtness, B., et al., *Correlation of tumor mutational burden (TMB) with CDKN2A and TP53 mutation in HPV-negative head and neck squamous cell carcinoma (HNSCC).* 2020, American Society of Clinical Oncology.
61. Hazawa, M., et al., *ZNF750 is a lineage-specific tumour suppressor in squamous cell carcinoma.* Oncogene, 2017. **36**(16): p. 2243-2254.
62. Kosmider, O., et al., *TET2 gene mutation is a frequent and adverse event in chronic myelomonocytic leukemia.* Haematologica, 2009. **94**(12): p. 1676-81.
63. Xu, Y.P., et al., *Tumor suppressor TET2 promotes cancer immunity and immunotherapy efficacy.* J Clin Invest, 2019. **129**(10): p. 4316-4331.