Network Constraints on Longitudinal Grey Matter Changes in First Episode Psychosis

Sidhant Chopra[1,2,3], Stuart Oldham[1,2], Ashlea Segal[1,2], Alexander Holmes[1,2], Kristina

Sabaroedin[1,2,12], Edwina R. Orchard[1,2,4], Shona M. Francey[5,6], Brian O'Donoghue[5,6], Vanessa

Cropley[7], Barnaby Nelson[5,6], Jessica Graham[5,6], Lara Baldwin[5,6], Jeggan Tiego[1,2], Hok Pan

Yuen[5,6], Kelly Allott[5,6], Mario Alvarez-Jimenez[5,6], Susy Harrigan[5,6,8], Ben D. Fulcher[9], Kevin

Aquino[9,10], Christos Pantelis[7,13], Stephen J Wood[5,6,11], Mark Bellgrove[1], Patrick McGorry[5,6],

Alex Fornito[1,2]

1. Turner Institute for Brain and Mental Health, School of Psychological Sciences, Monash

University, Clayton, Australia

2. Monash Biomedical Imaging, Monash University, Clayton, Australia

3. Department of Psychology, Yale University, New Haven, USA

4. Child Study Centre, Yale University, New Haven, USA

5. Orygen, Parkville, Australia

6. Centre for Youth Mental Health, The University of Melbourne, Melbourne, Australia

7. Melbourne Neuropsychiatry Centre, Department of Psychiatry, University of Melbourne

8. Centre for Mental Health, Melbourne School of Global and Population Health, The

University of Melbourne, Parkville, Australian

9. School of Physics, University of Sydney, New South Wales, Australia

10. Centre for Complex Systems, University of Sydney, New South Wales, Australia

11. School of Psychology, University of Birmingham, Edgbaston, UK

12. Departments of Radiology and Paediatrics, Hotchkiss Brain Institute and Alberta

Children's Hospital Research Institute, University of Calgary, Calgary, AB, Canada

13. NorthWestern Mental Health, Royal Melbourne Hospital & Western Hospital Sunshine,

St Albans, Victoria, Australia

Corresponding Author: Sidhant Chopra (sidhant.chopra@yale.edu)

Yale University

1 Prospect St, New Haven

CT, USA

35   **Abstract**

Background: Different regions of the brain's grey matter are connected by a complex structural

network of white-matter fibres, which are responsible for the propagation of action potentials and the

transport of trophic and other molecules. In neurodegenerative disease, these connections constrain

the way in which grey-matter volume change progresses. Here we investigated whether connectome

architecture also shapes the spatial pattern of longitudinal grey matter volume changes attributable to

illness and antipsychotic medication in first episode psychosis (FEP).

Methods: We conducted a triple-blind randomised placebo-control MRI study where 62 young adults

with FEP received either an atypical antipsychotic or placebo over 6-months. Anatomical MRI scans

were acquired at baseline, 3 months, and 12 months. Deformation-based morphometry was used to

estimate illness-related and antipsychotic-related volume changes over time. Representative

functional and structural brain connectivity patterns were derived from an independent healthy control

group using resting-state functional MRI and diffusion-weighted imaging. We used neighbourhood

deformation models to predict the extent of brain change in each of 332 parcellated areas by the

changes observed in areas to which the index region is either structurally connected or functionally

coupled.

Results: At baseline, we found that empirical illness-related regional volume differences were

robustly correlated with predicted differences using a model constrained by structural connectivity

weights ($r = .54$; $p < .001$). At 3 months and 12 months, we also found a robust correlation between

longitudinal regional illness-related ($r > .52$; $p < .001$) and antipsychotic-related volume change ($r >$

$.59$; $p < .001$) with volumetric changes in structurally connected areas. These correlations were

significantly greater than those observed across various null models accounting for lower-order spatial

and network properties of the data. Associations between empirical and predicted volume-change

estimates were much lower for models that only considered binary structural connectivity (all $r < .37$),

or which were constrained by inter-regional functional coupling (all $r < .43$). Finally, we found that

epicentres of volume change emerged posteriorly early in the illness and shifted to the prefrontal

cortex by later illness stages.

Conclusion: Psychosis- and antipsychotic-related grey-matter volume changes are strongly shaped by

structural brain connectivity. This result is consistent with findings in other neurological disorders and

implies that such connections may constrain pathological processes causing brain dysfunction in FE

Introduction

Psychotic disorders such as schizophrenia are characterised by anatomically widespread differences in grey-matter volume (GMV)[1-7], with many of these differences showing evidence of progression as the illness evolves[8-13]. Meta- and mega-analyses indicate that some of the most robust cross-sectional GMV differences are found in frontal, cingulate and temporal cortices, as well as medial temporal lobe, striatum and thalamus[3-7,14-16], with longitudinal changes identified in frontal, temporal and parietal cortices[9,11]. However, despite a large literature describing the location and nature of these brain changes, the specific mechanisms that give rise to their characteristic spatial pattern remain unknown.

The human brain is an intricate network of functionally specialised regions linked by a complex web of axonal fibres, referred to as a connectome[17]. These fibres enable the widespread coordination of neuronal dynamics and the transport of trophic and other biological molecules throughout the brain. They can also act as conduits for the spread of pathology, such that illness processes originating in one area can propagate to affect distributed systems via multiple mechanisms[18,19]. This principle has been powerfully demonstrated in dementia[20-23], with evidence that GMV reductions in different neurodegenerative conditions appear to spread through the brain in a way that is constrained by the underlying architecture of the brain's white-matter pathways.

Recent work suggests that a network-based spreading process may also be involved in psychosis. Cross-sectional grey-matter alterations correlate with increased fractional anisotropy in regionally adjacent white matter[24], and grey-matter differences in patients are often correlated across spatially distributed regions[25-27]. In recent work, Shafiei, et al. [28] developed a simple neighbourhood-deformation model (NDM) in a sample of people with established schizophrenia that predicted the level of cross-sectional GMV reduction in an area based on the average reductions observed in other areas to which it was structurally connected. The analysis further revealed that transmodal frontal and cingulate cortices were putative epicentres of pathology, being regions associated with relatively large GMV reductions that were also connected to other areas showing large reductions.

Together, these findings support the hypothesis that the spatial patterning of GMV differences in psychotic illness is constrained by connectome architecture. However, all studies addressing this question have been cross-sectional, precluding an opportunity to track how coordinated grey-matter changes through time. Moreover, the reliance on samples of patients taking antipsychotic medication makes it difficult to determine whether coupled grey-matter changes are driven by treatments for the illness or the illness process itself. These effects can only be disentangled through a longitudinal, randomised placebo-controlled study of antipsychotic-naive patients. We recently used such a design

to demonstrate that antipsychotic-medicated and non-medicated patients with First Episode Psychosis

(FEP) show different trajectories of GMV change[29] and functional coupling[30] within the first 3 and

12 months of treatment. Here, we use this cohort to investigate whether patterns of illness- and

antipsychotic-related GMV changes are constrained by the structural connectome and whether the

strength of functional and/or structural coupling modulates this relationship. Using the NDM[28], we

show that coordinated patterns of GMV change in FEP are more tightly constrained by the extent of

structural coupling (SC) than functional coupling (FC) between regions. Critically, the same pattern

holds for both illness-related and antipsychotic-related longitudinal brain changes. We also track the

dynamic progression of putative epicentres of illness-related GMV loss, showing that they emerge

posteriorly early in the illness and shift to the prefrontal cortex by later illness stages. This pattern

contrasts epicentres of antipsychotic-related GMV expansion, which are situated in primary visual

and auditory cortices.

Methods

Overview

Our overall analysis strategy is outlined schematically in Fig. 1. Briefly, we used structural MRI

images from a sample of FEP patients recruited to a placebo-controlled clinical trial of second-

generation antipsychotics to estimate brain-wide, voxel-wise illness-related and antipsychotic-related

GMV changes at baseline, 3 months, and 12 months (Fig. 1A). We parcellated the brain into discrete

regions-of-interest[31,32] and estimated volume changes as the mean test-statistic of all voxels within

that area. We used diffusion and functional MRI to generate group-level representative SC and FC

matrices in an independent sample of healthy adults (Fig. 1B). These matrices were then used in three

variants of the Neighbourhood Deformation Model (NDM) to evaluate the extent to which GMV

differences in one area are coupled to the GMV differences observed in its structurally connected

neighbours[28]. In the first variant, all neighbours make an equal contribution to predicting an index

area's GMV. In the second and third variants, the contribution of each neighbour is scaled by either

the weight of its structural or functional coupling to the index area, as determined by the

representative SC and FC matrices, respectively. Model performance was evaluated using the

Spearman correlation (ρ) between region-wise estimates of observed and predicted volume changes

(Fig. 1C). Finally, performance of each of the three NDM models was compared to three benchmark

null models to ensure that our findings are not driven by lower order properties of the spatial maps or

connectome architecture (Fig. 1D).

138

139

140 Fig. 1. - **Analysis workflow for the Network Deformation Model.** (A) We derived voxel-wise GMV

141 estimates using Deformation-based morphometry (DBM). Five separate contrasts were specified using a

142 robust marginal model to infer baseline GMV differences and longitudinal GMV changes associated with

143 illness and antipsychotic medication at 3 months and 12 months. (B) The contrast statistics were mapped

144 to a brain parcellation comprising 332 regions, and diffusion and functional MRI data from an independent

145 healthy sample were used to generate sample-averaged functional coupling (FC) and structural

146 connectivity (SC) matrices. These matrices were used to model average volume changes in structurally

147 connected neighbours. Under the NDM, the predicted deformation of a node, $\hat{d}_i$, is modelled as a weighted

148 sum of the deformation values observed its structurally connected neighbours (shown as light blue nodes

149 in the example graphs). The weights are given by the adjacency matrix, A_{ij}. Three different matrices were

150 used, yielding three NDM variants; (1) A model denoted as NDM_{SC}, in which $A_{ij} = 1$ if two regions share

151 a connection and $A_{ij} = 0$ otherwise; (2) a model denoted as NDM_{SCw} in which the elements of A_{ij}

152 correspond precisely to the weighted SC matrix, such that the contribution of each neighbour is weighted

153 by the strength of its structural connectivity to the index node; and (3) a model denoted NDM_{FCw}, in which

154 the elements of A_{ij} correspond precisely to the weighted FC matrix, such that the contribution of each

155 neighbour is weighted by its level of FC with the index node. (C) Model performance was evaluated using

156 the Spearman correlation between regional estimates of observed and predicted GMV differences. (D) We

157 also compared model performance to three benchmark null models accounting for spatial autocorrelations

158 in the deformation maps (Spatial$_{model}$ and Spatial$_{spin}$) and basic topological properties of the connectome

159 (see *Model evaluation*).

160

161

Sample characteristics

Clinical trial sample

We recruited 62 people aged 15-25 years (46% female) who were experiencing FEP. All patients had

minimal previous exposure to antipsychotic medication (<7 days of use or lifetime 1750 mg

chlorpromazine equivalent exposure) and a duration of untreated psychosis of less than 6 months. At

baseline, patients were randomised to one of two groups: one given antipsychotic medication

(risperidone or paliperidone) plus intensive psychosocial therapy and the other given placebo plus

intensive psychosocial therapy. For both groups, the treatment period spanned 6-months. MRI was

conducted at baseline, 3 months, and 12 months post-intake. The randomisation phase of the study

terminated at 6 months, so patients in either the antipsychotic or placebo group could have received

antipsychotic medication and ongoing psychosocial interventions after 6 months. In practice, four

patients in the placebo group commenced antipsychotic medication in this intervening period,

in addition to the four patients who had commenced at the 3-month timepoint. Thus, between

the 3-month and 12-month scan, a total of eight patients in the placebo group commenced

antipsychotic medication and were removed from the analysis. All patients in the

antipsychotic continued medication with varying degrees of exposure.

A matched healthy control group comprising 27 individuals with no history of psychiatric or

neurological diagnosis was also recruited and scanned alongside the patient groups. Demographic

details of this sample are provided in Table 1. Further sample characteristics and details about

research and safety protocols can be found elsewhere[33,34]. Ethical approval for the study was granted

by the Melbourne Health Human Research Ethics Committee (MHREC:2007.616).

Independent healthy control sample

A total of 119 healthy participants (66% female) were recruited as part of a larger study conducted at

Monash University. The participants were selected from a large cohort of 439 people as those with

high-quality functional and diffusion MRI scans available and with an age that was within the range

of the clinical trial sample. For further details, see Sabaroedin, et al. [35]. The study was conducted in

accordance with the Monash University Human Research Ethics Committee (MUHREC:

2012001562).

197 ***Table 1 - Sample characteristics of clinical trial and healthy control sample.***

	First episode psychosis		Matched healthy control (N=27)	Independent healthy control (N=119)
	Placebo (Baseline N =30; 3-months N = 21; 12-months N=22)	Antipsychotic (Baseline N =29; 3-months N = 20; 12-months N=14)		
Baseline age, years (SD)	18.8 (2.72)	19.5 (2.94)	21.9 (1.93)	20.9 (1.68)
Baseline Females, N (%)	14 (46.6%)	13 (44.8%)	17 (62.9%)	78 (65.5%)
Diagnosis, N Major depression with psychosis	7	5		
Schizophreniform disorder	5	5		
Psychotic disorder NOS	8	7		
Substance-induced psychotic disorder	4	2		
Delusional disorder	1	4		
Schizophrenia	5	5		
Missing diagnosis	0	1		
Baseline BPRS Total, mean (SD)	59.4 (9.64)	55.8 (10.10)		
Baseline SOFAS, mean (SD)	52.9 (14.0)	51.7 (10.6)		

198 Abbreviations: NOS = not otherwise specified; BPRS = Brief Psychiatric Rating Scale version 4;
199 SOFAS = Social and Occupational Functioning Assessment Scale.
200

Structural MRI processing

Acquisition parameters for structural MRI can be found in Supplement. Prior to processing, raw T1w

scans were visually examined for artefacts and then subjected to an automated quality control

procedure[36]. Three patient scans did not pass image quality control and were excluded due to image

artefacts. The remaining scans were processed using the longitudinal deformation-based morphometry

(DBM) pipeline of the Computational Anatomy Toolbox (version r1113)[37] for the Statistical

Parametric Mapping 12[38] software running in MATLAB version 2019a For detailed information

about DBM processing, see the Supplement. We used DBM to quantify volume changes because it

requires less spatial smoothing[39] than voxel-based morphometry (VBM) and to be comparable to

previous work using the NDM[28]. We also replicated our primary findings using VBM, which has been
used more extensively to quantify GMV in the literature (see *Robustness analyses*).

Quantifying baseline and longitudinal grey matter changes in patients
To estimate group-level baseline and longitudinal spatial patterns of volume change, we used a robust
marginal model implemented in the Sandwich Estimator Toolbox[40], which combines ordinary least
squares estimates of parameters of interest with estimates of variance/covariance based on a robust
sandwich estimator, thus accounting for within-subject correlations across time. This method is
asymptotically robust to misspecification of the covariance model and does not depend on the
assumptions of common longitudinal variance structure across the whole brain. All contrasts were
adjusted for age, sex, and handedness.

We conducted three categories of contrasts (Fig. 1A): (1) illness-related differences at baseline, which
compared all patients to matched healthy controls; (2) illness-related change over time, which
compared GMV changes in the placebo group to matched healthy controls; and (3) antipsychotic-
related changes, which compared GMV changes in the medication group to both the placebo group
and matched healthy controls (see also[13]). Longitudinal contrasts were assessed from baseline to 3
months, as well as from baseline to 12 months, with a linear contrast being used for the latter. The
resulting *t*-statistics were converted to *z*-statistics. Contrasts were specified such that positive values
in the resulting voxel-wise *z*-statistic maps indicate lower volume in patients compared to controls at
baseline contrast, and a greater longitudinal GMV decline in patients compared to controls in the
longitudinal contrasts. We apply the NDM to unthresholded *t*-maps encoding regional GMV changes
as we are interested in capturing the spatial patterning of GMV differences across the entire brain, not
only changes which survive a statistical threshold. Renderings of the unthresholded *t*-maps can be
found in Fig 1A-C and Fig 2A-B.

Brain Parcellation
In order to evaluate network-based models of grey-matter changes, we parcellated the brain into 300
discrete cortical regions of approximately equal size[32], in addition to 32 subcortical areas[31], using
previously published atlases. The deformation value for each region was estimated as the mean *t*-
statistic of all voxels corresponding to that region. The regions comprise the nodes of a network,
which can then be directly related to measures of inter-regional SC and FC.

Healthy structural network
We derived a group-level healthy structural connectome from diffusion-weighted imaging (DWI) data
from an independent sample of 119 adults (Fig. 1B). Acquisition parameters and detailed overview of

DWI processing and optimisation can be found in Supplement. This procedure resulted in a single 332

× 332 weighted group-average SC matrix.

Healthy functional network

We derived a group-level healthy functional connectome from resting-state fMRI data from the same

independent sample of adults (Fig. 1B). Acquisition parameters for functional MRI and detailed

information on fMRI processing can be found in Supplement. Given ongoing controversy around the

application of global signal regression[41,42], we evaluated how this step affected our findings (see

Robustness analyses). After processing and denoising, we computed a whole brain 332 × 332 FC

matrix for each subject using pair-wise Pearson correlations between the timeseries from each of the

332 regions and finally took a mean FC matrix across the sample.

Neighbourhood Deformation Model

We evaluated network constraints on baseline and longitudinal GMV changes using the NDM

introduced by Shafiei, et al. [28]. The model is given by

264
$$\hat{d}_i = \frac{1}{N_i} \sum^{N_i} A_{ij} d_j \ , \qquad (2)$$

where $\hat{d}_i$ is the predicted grey matter volume change in node i, N_i is the number of structurally

connected neighbours of i, d_j is the deformation observed in the j-th neighbour of node i, and A_{ij}

defines the connectivity between nodes i and j.

Three different matrices were substituted for A_{ij}, yielding three variants of the NDM. For the first

model, denoted NDM$_{SC}$, A_{ij}=1 if nodes i and j are connected in the group-average SC matrix and

zero otherwise. As such, under this model, all j structurally connected neighbours make an equal

contribution to predicting the extent of deformation observed in node i.

For the second and third models, denoted NDM$_{SCw}$ and NDM$_{FCw,}$ A_{ij} corresponded to the weighted

SC or FC matrices, respectively. Therefore, under these models, the contributions of node i's

neighbours were weighted by either inter-regional SC (NDM$_{SCw}$) or FC (NDM$_{FCw}$) estimates, such

that neighbours of node i with a more strongly weighted connection made a stronger contribution to

predicting node i's volume change (Fig 1B). In all models, only edges which had a structural

connection were included.

Model evaluation

Model performance was evaluated using the Spearman correlation (ρ) between region-wise estimates

of observed and predicted deformation (Fig. 1C). We also compared the performance of the NDM_{SC},

NDM_{SCw} and NDM_{FCw} models to three benchmark null models. The first (Fig. 1D; $Spatial_{model}$) and

second (Fig. 1D; $Spatial_{spin}$) null models evaluated whether the observed findings were specific to the

empirically observed pattern of grey-matter deformations or were a generic property of the intrinsic

spatial correlation structure of the deformation maps. The third null model (Fig. 1D; Connectome)

tests the hypothesis that any network-based prediction of local grey-matter change is specific to the

actual topology of the connectome itself, and cannot be explained by basic network properties, such

as regional variations in node degree or the spatial dependence of inter-regional connectivity. Further

details about benchmark null models can be found in the Supplement.

Mapping epicentres of illness- and antipsychotic-related volume change

We next sought to identify putative epicentres of illness- and antipsychotic-related volume changes.

As per Shafiei, et al. [28], we defined such epicentres as areas showing high deformation that were also

connected to regions showing high deformation (Fig. 5A). A detailed overview of the epicentre

analysis can be found in Supplement.

All code and group-level data used to run analyses can be found online at:

www.github.com/sidchop/NetworkDeformationModels

Results

Structural connectivity shapes illness-related cross-sectional and longitudinal grey matter changes

We initially evaluated the performance of the three NDMs in capturing baseline differences in

regional grey-matter volume. We observed the strongest correlation between predicted and observed

estimates for the NDM_{SCw} model ($\rho = .54$; Fig. 2G), which was substantially stronger than

correlations observed for the NDM_{SC} and NDM_{FCw} models ($\rho = 0.32$ and $\rho = 0.38$, respectively; Fig.

2D). The performance of the NDM_{SCw} was also significantly better than all three benchmark models

(all $p < 0.004$), whereas the NDM_{SC} and NDM_{FCw} models were only superior to the $Spatial_{model}$ ($p <$

0.003), but not $Spatial_{spin}$ or connectome, benchmarks (all $p > 0.05$).

We observed the same pattern of findings for longitudinal illness-related changes at 3 months and 12

months, with the NDM_{SCw} predictions correlating with empirical estimates at $\rho = 0.52$ (Fig. 2H) and

$\rho = 0.55$ (Fig. 2I), respectively. The NDM_{SCw} again showed significantly better performance than all

three benchmark models (all $p < 0.03$). By comparison, correlations for the NDM_{SC} and NDM_{FCw}

did not exceed $\rho = 0.37$ and only showed significantly better performance than the Spatial$_{model}$ (Fig.

2E-F), but not the Spatial$_{spin}$ or connectome benchmarks (all $p > 0.05$).

Fig. 2. – Baseline and longitudinal illness-related GMV changes are constrained by connectome

anatomy. A-C) The contrast statistics for three illness-related contrasts mapped to a brain parcellation

comprising 332 regions. D-F) Performance of the equally weighted (NDM$_{SC}$), structural connectivity-

weighted (NDM$_{SCw}$), and functional coupling-weighted (NDM$_{FCw}$) models relative to the Spatial$_{model}$,

Spatial$_{spin}$ and connectome null. Black circles indicate the observed rank correlations between

predicted and actual regional deformation values for each model at each timepoint, with red borders

indicating statistical significance ($p < 0.05$). Note that the observed value used for comparison

against the Spatial$_{spin}$ models is different because the subcortex was excluded. G–I) Scatterplots of the

association between observed and predicted regional volume deformation values for the best

performing NDM$_{SCw}$ model at each timepoint. Least-squares linear trend lines were included for

illustration purposes.

Structural connectivity shapes longitudinal antipsychotic-related volume changes

As with illness-related changes, we found that the strongest correlation between observed and

predicted antipsychotic-related grey-matter changes was found using the NDM$_{SCw}$ model at both 3

months ($\rho = 0.59$; Fig. 3E) and 12 months ($\rho = 0.60$; Fig. 3F). This association was statistically

significant when compared to all three null models (all $p < 0.007$; Fig. 3C $-$ D). For comparison,

associations at 3 months and 12 months were smaller for the NDM_{SC} ($\rho = .38$ and $\rho = .17$,

respectively) and NDM_{FCw} models ($\rho = .44$ and $\rho = .27$, respectively). At 3 months, the NDM_{SC} and

NDM_{FCw} models only showed significantly better performance than the $Spatial_{model}$ and $Spatial_{spin}$

($p < .007$) benchmarks. At 12 months, the NDM_{FCw} model only showed significantly better

performance than the $Spatial_{model}$ benchmark ($p < .014$).

Fig. 3 - Baseline and longitudinal antipsychotic-related GMV changes are constrained by

connectome anatomy. A-B) The contrast statistics for two antipsychotic-related contrasts mapped to a

brain parcellation comprising 332 regions. C-D) Performance of the equally weighted (NDM_{SC}),

structural connectivity-weighted (NDM_{SCw}) and functional coupling-weighted (NDM_{FCw}) models

relative to the $Spatial_{model}$, $Spatial_{spin}$, and connectome null. Black circles indicate the observed rank

correlations between predicted and actual regional deformation values for each model at each

timepoint, with red borders indicating statistical significance ($p < 0.05$). Note that the observed

value used for comparison against the $Spatial_{spin}$ models is different because the subcortex was

excluded. E-F) Scatterplots of the association between observed and predicted regional deformation

values for the best performing NDM$_{SCw}$ model at each timepoint. Least-squares linear trend lines were

included for illustration purposes.

Illness-related epicentres

We next investigated regional epicentres of grey-matter change, corresponding to areas that are

connected to other areas showing a high degree of volume change in patients that also show large

changes themselves. At baseline, 21 areas with statistically significant atrophy epicentre scores were

identified in bilateral visual and left medial orbito-frontal cortex (Fig. 4D). At 3 months, areas with

statistically significant atrophy epicentre scores were located within the left temporal pole and

bilateral pre-frontal and orbitofrontal cortices (Fig. 4E). Areas with significant expansion epicentre

scores were largely located in the visual cortex (Fig. 4E). At 12 months, the areas with significant

regional epicentres extended across larger areas of anterior frontal cortex (Fig. 4F).

Fig. 4 – Regional epicentres of illness-related and medication-related GMV change. A) Epicentres

were defined as areas showing high deformation that are also connected to regions showing high

deformation. To identify such regions, for each region and each contrast, we computed a region-

specific epicentre score as the mean of that area's deformation and the average of its neighbours'

deformation, weighted by structural coupling strength. Regions showing epicentre scores significantly

greater ($p_{FWE} < 0.05$) than a spatially constrained null model are shown for antipsychotic-related (A-

B) and illness-related (D-F) effects.

Antipsychotic-related epicentres

At 3 months, we identified expansion-related epicentres in bilateral visual cortex (Fig 4B). At 12
months, additional expansion epicentres were located within the right superior temporal cortex and
left temporal pole, along with a significant atrophy epicentre within the left occipital pole (Fig 4C).

Robustness analyses

To ensure that our findings were influenced not by our inclusion of patients with substance-induced
psychotic disorder, we repeated the analysis after excluding these individuals. The magnitude and
pattern of results remained consistent with our original illness and antipsychotic-related (Fig. S1)
findings.

To ensure that our findings were not specific to the structural connectome derived from the large
independent healthy control sample, we replicated our primary NDM analyses using a representative
structural connectome derived from the matched healthy control sample which was part of the clinical
trial. The magnitude and pattern of results remained consistent with our original illness and
antipsychotic-related findings (Fig. S2) .

Next, we assessed whether our findings were specific to volume estimates derived using DBM by
repeating our primary analyses using volume estimates from VBM. We found that the patterns of
results for our primary analyses remained the same (Fig S3).

There can be several orders of magnitude difference between the deformation values and SC/FC
values. We therefore repeated the analyses after *z*-scoring the deformation values and connectivity
weights and again found comparable results to our original findings (Fig. S4).

Finally, we assessed the impact of implementing global signal regression (GSR) on subject-level FC
matrices before computing the group average FC matrix. We found that implementing GSR did not
alter the pattern in the results, with the NDM_{SCw} model still showing the best performance (Fig. S5).

Discussion

Using a simple neighbourhood deformation model (NDM), we showed that the spatial pattern of both
cross-sectional and longitudinal GMV changes in psychosis is shaped by brain-network architecture,
particularly by the strength of structural connectivity between regions. A similar relationship is
apparent for GMV changes that are attributable either to psychotic illness or antipsychotic medication,
suggesting that the connectome represents a generic constraint on diverse processes driving

coordinated grey-matter variations. We also show a dynamic progression of regional epicentres of

GMV change from posterior to anterior areas, suggesting that the pathological burden within

prefrontal systems increases as the illness evolves.

Network constraints on Illness-related volumes changes in FEP

Our baseline findings in FEP align with the results of Shafiei, et al. [28], who also used an NDM to

show that structural connectivity constrains the spatial patterning of cross-sectional GMV differences

in people with established schizophrenia. This earlier result was observed using the NDM_{SC} model

considered here. In our analysis, we found that the strength of structural connectivity between regions

modulates coupled GMV differences within structurally connected neighbourhoods, given that the

NDM_{SCw} showed clearly superior performance to the NDM_{SC} and NDM_{FCw} models. This result

indicates that GMV differences are more tightly coupled between areas with high structural

connectivity. This modulatory effect of the strength of structural connectivity may be especially

pronounced early in the illness, as the performance of the binary NDM_{SC} model in Shafiei, et al. [43]

analysis of patients with established schizophrenia was superior to the performance of the same model

in our sample. Critically, our findings show that network constraints on cross-sectional GMV

differences cannot be explained by antipsychotic medication, as our FEP sample was antipsychotic-

naïve at the baseline scan.

Moving beyond cross-sectional differences, our longitudinal analysis further demonstrates that both

illness-related and antipsychotic-related changes in GMV are also constrained by connectome

architecture. These results are in line with a spreading process, in which some pathological process

propagates across axonal connections. The precise mechanisms driving this process remain unclear,

but given the absence of evidence indicating an accumulation of pathological proteins in psychotic

illness, it seems likely that the dysfunction of one region may trigger abnormal activity in connected

sites which, over time, may trigger structural changes as a result of aberrant neurotransmission or a

loss of trophic support[19]. This process may be exacerbated by a breakdown of white-matter fibre

integrity, which may further disrupt the inter-regional transport of trophic factors. Accordingly,

widespread but subtle alterations in white matter have been repeatedly demonstrated in FEP

populations[44], are anticorrelated with cortical thickness[24], and may predate the transition to psychosis

in high-risk samples[45,46]. Further work may investigate how coordinated GMV changes relate to

white-matter pathology in patients.

Given that we found that antipsychotic-related longitudinal changes in GMV are also constrained by

connectome architecture, it is possible that neuronal signalling and/or neurotransmitter modulation

induced by antipsychotics in one region may propagate to connected regions via structural

connections. Indeed, higher white-matter integrity[47] and higher global efficiency [48] of the structural

connections are both related to better medication response in FEP patients.

An alternate explanation for our findings may be that more strongly anatomically connected regions

share more similar molecular profiles, resulting in a shared vulnerability to illness- or medication-

related processes[49-51]. Future research should examine the associations between the strength of

structural connectivity and shared molecular features such as receptor profiles, gene expression, and

synaptic density.

Regional epicentres of grey matter change dynamically evolve with illness progression

The Shafiei, et al. [28] cross-sectional analysis of patients with established schizophrenia identified

cingulate and prefrontal areas as putative epicentres of GMV reduction. Our analysis indicates that the

involvement of prefrontal areas dynamically emerges as the illness progresses, shifting from an early

posterior focus at the outset of illness. Progressively more prefrontal areas were identified as

significant epicentres at 12-month compared to 3-month follow-up, suggesting that prefrontal circuits

become increasingly compromised with ongoing illness. These findings align with longitudinal

studies in early-onset schizophrenia demonstrating a dynamic wave of volume contraction

progressing from posterior to anterior regions,[52,53] which may reflect an exaggeration of normal

neurodevelopmental processes[54,55]. The findings also align with evidence of pronounced prefrontal

GMV reductions in the earliest stages of illness[8,56-61]. Notably, these regional epicentres of illness-

related GMV reduction localized to areas that were distinct from epicentres of antipsychotic-related

GMV expansion, which were identified in primary visual and superior temporal cortex. The

involvement of these regions may be related to antipsychotic-related improvements of perceptual

abnormalities in patients[62-64].

Limitations and conclusions

Given the complexity and practical challenges of conducting a prospective triple-blind randomised

control MRI study in antipsychotic-naïve patients, the sample sizes within the patient groups are

small. Relatedly, given the ethical and safety requirements of the study, patients who were recruited

posed low risk of harm to self or others, lived in stable accommodation, had adequate social support

and a short duration of untreated illness. While it's possible that these requirements may have resulted

in a patient group with a less severe form of psychotic illness, the symptom severity and functioning

of our patients (see Table 1) are comparable to epidemiologically representative cohorts of FEP

patients[65,66].

The structural and functional connectomes used in this study were derived from healthy populations
to examine how normative network structure constrains the progression of GMV changes in FEP.
Future studies should examine how altered connectivity in patients impacts our findings. Moreover,
our findings are dependent on group-level summary metrics of brain volume and may not be
representative of volume changes at the individual patient level, which show substantial
heterogeneity[67-69]. Subsequent work could look at whether using individual-level measures of brain
volume and connectivity can improve model predictions.
490
491 In summary, we show that longitudinal GMV changes in early psychosis are strongly constrained by
492 the structural connectivity of the brain, but not its inter-regional functional coupling, regardless of
493 whether those changes are driven by illness or antipsychotic medication. Our findings thus highlight a
494 central role for axonal connectivity for shaping dynamic brain changes in the illness and suggest an
495 increasing disruption of prefrontal systems with ongoing illness.
496

Disclosures/Conflict of Interest
SF, KA, MAJ and AF reported receiving grants from the Australian National Health & Medical
Research Council (NHMRC) during the conduct of the study. CP reported receiving grants from the
Australian NHMRC and from the Lundbeck Foundation and personal fees from Lundbeck Australia
Pty Ltd Advisory Board for talks presented at educational meetings organized by Lundbeck. PM
reported receiving grants from the Australian NHMRC, the Colonial Foundation, the National
Alliance for Research on Schizophrenia and Depression, the Stanley Foundation, the National
Institutes of Health, Wellcome Trust, the Australian and Victorian governments, and Janssen-Cilag
(unrestricted investigator-initiated grant) during the conduct of the study; past unrestricted grant
funding from Janssen-Cilag, AstraZeneca, Eli Lilly, Novartis, and Pfizer; honoraria for consultancy
and teaching from Janssen-Cilag, Eli Lilly, Pfizer, AstraZeneca, Roche, Bristol Myers Squibb, and
Lundbeck. BN was supported by an NHMRC Senior Research Fellowship (1137687) and a University
of Melbourne Dame Kate Campbell Fellowship. This work was supported by the computational
infrastructure provided by the MASSIVE HPC facility (www.massive.org.au).

References

1   Gur, R. E., Turetsky, B. I., Bilker, W. B. & Gur, R. C. Reduced Gray Matter Volume
     in Schizophrenia. *Archives of General Psychiatry* **56**, 905-911,
     doi:10.1001/archpsyc.56.10.905 (1999).

2   Haijma, S. V. *et al.* Brain Volumes in Schizophrenia: A Meta-Analysis in Over 18
     000 Subjects. *Schizophrenia Bulletin* **39**, 1129-1138, doi:10.1093/schbul/sbs118
     (2013).

3   Steen, R. G., Mull, C., Mcclure, R., Hamer, R. M. & Lieberman, J. A. Brain volume
     in first-episode schizophrenia. *British Journal of Psychiatry* **188**, 510-518,
     doi:10.1192/bjp.188.6.510 (2006).

4   van Erp, T. G. M. *et al.* Cortical Brain Abnormalities in 4474 Individuals With
     Schizophrenia and 5098 Control Subjects via the Enhancing Neuro Imaging Genetics
     Through Meta Analysis (ENIGMA) Consortium. *Biological Psychiatry* **84**, 644-654,
     doi:10.1016/j.biopsych.2018.04.023 (2018).

5   van Erp, T. G. M. *et al.* Subcortical brain volume abnormalities in 2028 individuals
     with schizophrenia and 2540 healthy controls via the ENIGMA consortium.
     Molecular Psychiatry **21**, 547-553, doi:10.1038/mp.2015.63 (2016).

6   Gupta, C. N. *et al.* Patterns of Gray Matter Abnormalities in Schizophrenia Based on
     an International Mega-analysis. *Schizophrenia Bulletin* **41**, 1133-1142,
     doi:10.1093/schbul/sbu177 (2015).

7   Vieira, S. *et al.* Neuroanatomical abnormalities in first-episode psychosis across
     independent samples: a multi-centre mega-analysis. *Psychological Medicine* **51**, 340-
     350, doi:10.1017/s0033291719003568 (2021).

8   Pantelis, C. *et al.* Neuroanatomical abnormalities before and after onset of psychosis:
     a cross-sectional and longitudinal MRI comparison. *The Lancet* **361**, 281-288,
     doi:10.1016/S0140-6736(03)12323-9 (2003).

9   Vita, A., De Peri, L., Deste, G. & Sacchetti, E. Progressive loss of cortical gray matter
     in schizophrenia: a meta-analysis and meta-regression of longitudinal MRI studies.
     Translational Psychiatry **2**, e190-e190, doi:10.1038/tp.2012.116 (2012).

10   Akudjedu, T. N. *et al.* Progression of neuroanatomical abnormalities after first-
     episode of psychosis: A 3-year longitudinal sMRI study. *Journal of Psychiatric
     Research* **130**, 137-151, doi:10.1016/j.jpsychires.2020.07.034 (2020).

11   Olabi, B. *et al.* Are there progressive brain changes in schizophrenia? A meta-analysis
     of structural magnetic resonance imaging studies. *Biological Psychiatry* **70**, 88-96,
     doi:10.1016/j.biopsych.2011.01.032 (2011).

12   Andreasen, N. C. *et al.* Progressive Brain Change in Schizophrenia: A Prospective
     Longitudinal Study of First-Episode Schizophrenia. *Biological psychiatry* **70**, 672-
     679, doi:10.1016/j.biopsych.2011.05.017 (2011).

13   Chopra, S. *et al.* Differentiating the Effect of Medication and Illness on Brain Volume
     Reductions in First-Episode Psychosis: A Longitudinal, Randomized, Triple-blind,
     Placebo-controlled MRI study. *medRxiv* (2020).

14   Bora, E. *et al.* Neuroanatomical abnormalities in schizophrenia: A multimodal
     voxelwise meta-analysis and meta-regression analysis. *Schizophrenia Research* **127**,
     46-57, doi:10.1016/j.schres.2010.12.020 (2011).

15   Fornito, A., Yücel, M., Patti, J., Wood, S. J. & Pantelis, C. Mapping grey matter
     reductions in schizophrenia: An anatomical likelihood estimation analysis of voxel-
     based morphometry studies. *Schizophrenia Research* **108**, 104-113,
     doi:10.1016/j.schres.2008.12.011 (2009).

16　Del Re, E. C. *et al.* Anterior-posterior axis of hippocampal subfields across
psychoses: A B-SNIP study. *Biomarkers in Neuropsychiatry* **5**, 100037,
doi:https://doi.org/10.1016/j.bionps.2021.100037 (2021).
17　Sporns, O., Tononi, G. & Kötter, R. The Human Connectome: A Structural
Description of the Human Brain. *PLOS Computational Biology* **1**, e42,
doi:10.1371/journal.pcbi.0010042 (2005).
18　Raj, A. & Powell, F. Models of Network Spread and Network Degeneration in Brain
Disorders. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* **3**, 788-
797, doi:10.1016/j.bpsc.2018.07.012 (2018).
19　Fornito, A., Zalesky, A. & Breakspear, M. The connectomics of brain disorders.
doi:10.1038/nrn3901 (2015).
20　Seeley, W. W., Crawford, R. K., Zhou, J., Miller, B. L. & Greicius, M. D.
Neurodegenerative diseases target large-scale human brain networks. *Neuron* **62**, 42-
52 (2009).
21　Zhou, J., Gennatas, E. D., Kramer, J. H., Miller, B. L. & Seeley, W. W. Predicting
regional neurodegeneration from the healthy brain functional connectome. *Neuron* **73**,
1216-1227 (2012).
22　Raj, A., Kuceyeski, A. & Weiner, M. A network diffusion model of disease
progression in dementia. *Neuron* **73**, 1204-1215, doi:10.1016/j.neuron.2011.12.040
(2012).
23　Shafiei, G. *et al.* Network structure and transcriptomic vulnerability shape atrophy in
frontotemporal dementia. *Brain*, doi:10.1093/brain/awac069 (2022).
24　Di Biase, M. A. *et al.* Linking Cortical and Connectional Pathology in Schizophrenia.
Schizophr Bull **45**, 911-923, doi:10.1093/schbul/sby121 (2019).
25　Wannan, C. M. J. *et al.* Evidence for Network-Based Cortical Thickness Reductions
in Schizophrenia. *American Journal of Psychiatry*, appi.ajp.2019.18040380,
doi:10.1176/appi.ajp.2019.18040380 (2019).
26　Cauda, F. *et al.* The morphometric co-atrophy networking of schizophrenia, autistic
and obsessive spectrum disorders. *Hum Brain Mapp* **39**, 1898-1928,
doi:10.1002/hbm.23952 (2018).
27　Jiang, Y. *et al.* Antipsychotics effects on network-level reconfiguration of cortical
morphometry in first-episode schizophrenia. *medRxiv*, 2021.2001.2017.21249965,
doi:10.1101/2021.01.17.21249965 (2021).
28　Shafiei, G. *et al.* Spatial patterning of tissue volume loss in schizophrenia reflects
brain network architecture. *Biological psychiatry* **87**, 727-735 (2020).
29　Chopra, S. *et al.* Differentiating the effect of antipsychotic medication and illness on
brain volume reductions in first-episode psychosis: A Longitudinal, Randomised,
Triple-blind, Placebo-controlled MRI Study. *Neuropsychopharmacology* **46**, 1494-
1501 (2021).
30　Chopra, S. *et al.* Functional connectivity in antipsychotic-treated and antipsychotic-
naive patients with first-episode psychosis and low risk of self-harm or aggression: a
secondary analysis of a randomized clinical trial. *JAMA psychiatry* **78**, 994-1004
(2021).
31　Tian, Y., Margulies, D. S., Breakspear, M. & Zalesky, A. Topographic organization
of the human subcortex unveiled with functional connectivity gradients. *Nature*
Neuroscience **23**, 1421-1432, doi:10.1038/s41593-020-00711-6 (2020).
32　Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from
Intrinsic Functional Connectivity MRI. *Cerebral Cortex (New York, N.Y.: 1991)* **28**,
3095-3114, doi:10.1093/cercor/bhx179 (2018).

33 Francey, S. M. *et al.* Psychosocial Intervention With or Without Antipsychotic
Medication for First-Episode Psychosis: A Randomized Noninferiority Clinical Trial.
Schizophrenia Bulletin Open **1**, doi:10.1093/schizbullopen/sgaa015 (2020).
34 O'Donoghue, B. *et al.* Staged treatment and acceptability guidelines in early psychosis
study (STAGES): A randomized placebo controlled trial of intensive psychosocial
treatment plus or minus antipsychotic medication for first-episode psychosis with
low-risk of self-harm or aggression. Study protocol and baseline characteristics of
participants. *Early Intervention in Psychiatry* **0**, doi:10.1111/eip.12716 (2019).
35 Sabaroedin, K. *et al.* Functional Connectivity of Corticostriatal Circuitry and
Psychosis-like Experiences in the General Community. *Biological Psychiatry* **86**, 16-
24, doi:10.1016/j.biopsych.2019.02.013 (2019).
36 Esteban, O. *et al.* MRIQC: Advancing the automatic prediction of image quality in
MRI from unseen sites. *PLOS ONE* **12**, e0184661, doi:10.1371/journal.pone.0184661
(2017).
37 Gaser, C. & Dahnke, R. CAT - A Computational Anatomy Toolbox for the Analysis
of Structural MRI Data. *HBM*, 336-348 (2016).
38 Ashburner, J. *et al.* SPM12 manual. *Wellcome Trust Centre for Neuroimaging,*
London, UK **2464** (2014).
39 Schwarz, D. & Kašpárek, T. Comparison of Two Methods for Automatic Brain
Morphometry Analysis. *Radioengineering* **20** (2011).
40 Guillaume, B., Hua, X., Thompson, P. M., Waldorp, L. & Nichols, T. E. Fast and
accurate modelling of longitudinal and repeated measures neuroimaging data.
NeuroImage **94**, 287-302, doi:10.1016/j.neuroimage.2014.03.029 (2014).
41 Murphy, K. & Fox, M. D. Towards a consensus regarding global signal regression for
resting state functional connectivity MRI. *Neuroimage* **154**, 169-173,
doi:10.1016/j.neuroimage.2016.11.052 (2017).
42 Aquino, K. M., Fulcher, B. D., Parkes, L., Sabaroedin, K. & Fornito, A. Identifying
and removing widespread signal deflections from fMRI data: Rethinking the global
signal regression problem. *Neuroimage* **212**, 116614 (2020).
43 Shafiei, G. *et al.* Global network structure and local transcriptomic vulnerability
shape atrophy in sporadic and genetic behavioral variant frontotemporal dementia.
bioRxiv (2021).
44 Yao, L. *et al.* White matter deficits in first episode schizophrenia: An activation
likelihood estimation meta-analysis. *Progress in Neuro-Psychopharmacology and*
Biological Psychiatry **45**, 100-106, doi:https://doi.org/10.1016/j.pnpbp.2013.04.019
(2013).
45 Di Biase, M. A. *et al.* White matter changes in psychosis risk relate to development
and are not impacted by the transition to psychosis. *Molecular Psychiatry*,
doi:10.1038/s41380-021-01128-8 (2021).
46 Carletti, F. *et al.* Alterations in White Matter Evident Before the Onset of Psychosis.
Schizophrenia Bulletin **38**, 1170-1179, doi:10.1093/schbul/sbs053 (2012).
47 Kraguljac, N. V. *et al.* White matter integrity, duration of untreated psychosis, and
antipsychotic treatment response in medication-naïve first-episode psychosis patients.
Molecular Psychiatry **26**, 5347-5356, doi:10.1038/s41380-020-0765-x (2021).
48 Crossley, N. A. *et al.* Connectomic correlates of response to treatment in first-episode
psychosis. *Brain* **140**, 487-496 (2017).
49 Arnatkeviciute, A. *et al.* Genetic influences on hub connectivity of the human
connectome. *Nature Communications* **12**, 4237, doi:10.1038/s41467-021-24306-2
(2021).

50 Fornito, A., Arnatkevičiūtė, A. & Fulcher, B. D. Bridging the Gap between
Connectome and Transcriptome. *Trends Cogn Sci* **23**, 34-50,
doi:10.1016/j.tics.2018.10.005 (2019).
51 Hansen, J. Y. *et al.* Mapping neurotransmitter systems to the structural and functional
organization of the human neocortex. *bioRxiv*, 2021.2010.2028.466336,
doi:10.1101/2021.10.28.466336 (2022).
52 Vidal, C. N. *et al.* Dynamically Spreading Frontal and Cingulate Deficits Mapped in
Adolescents With Schizophrenia. *Archives of General Psychiatry* **63**, 25-34,
doi:10.1001/archpsyc.63.1.25 (2006).
53 Thompson, P. M. *et al.* Mapping adolescent brain change reveals dynamic wave of
accelerated gray matter loss in very early-onset schizophrenia. *Proceedings of the*
National Academy of Sciences **98**, 11650-11655, doi:10.1073/pnas.201243998 (2001).
54 Cropley, V. L. *et al.* Accelerated Gray and White Matter Deterioration With Age in
Schizophrenia. *Am J Psychiatry* **174**, 286-295, doi:10.1176/appi.ajp.2016.16050610
(2017).
55 Gogtay, N., Vyas, N. S., Testa, R., Wood, S. J. & Pantelis, C. Age of onset of
schizophrenia: perspectives from structural neuroimaging studies. *Schizophr Bull* **37**,
504-513, doi:10.1093/schbul/sbr030 (2011).
56 Sun, D. *et al.* Progressive Brain Structural Changes Mapped as Psychosis Develops in
'At Risk' Individuals. *Schizophrenia research* **108**, 85-92,
doi:10.1016/j.schres.2008.11.026 (2009).
57 Sun, D. *et al.* Brain surface contraction mapped in first-episode schizophrenia: a
longitudinal magnetic resonance imaging study. *Molecular Psychiatry* **14**, 976-986,
doi:10.1038/mp.2008.34 (2009).
58 Jalbrzikowski, M. Association of Structural Magnetic Resonance Imaging Measures
With Psychosis Onset in Individuals at Clinical High Risk for Developing Psychosis:
An ENIGMA Working Group Mega-analysis. *JAMA Psychiatry* **78**, 753-766,
doi:10.1001/jamapsychiatry.2021.0638 (2021).
59 Cannon, T. D. *et al.* Progressive reduction in cortical thickness as psychosis develops:
a multisite longitudinal neuroimaging study of youth at elevated clinical risk. *Biol*
Psychiatry **77**, 147-157, doi:10.1016/j.biopsych.2014.05.023 (2015).
60 Chung, Y. *et al.* Cortical abnormalities in youth at clinical high-risk for psychosis:
Findings from the NAPLS2 cohort. *NeuroImage: Clinical* **23**, 101862,
doi:https://doi.org/10.1016/j.nicl.2019.101862 (2019).
61 Del Re, E. C. *et al.* Baseline Cortical Thickness Reductions in Clinical High Risk for
Psychosis: Brain Regions Associated with Conversion to Psychosis Versus Non-
Conversion as Assessed at One-Year Follow-Up in the Shanghai-At-Risk-for-
Psychosis (SHARP) Study. *Schizophrenia Bulletin* **47**, 562-574 (2021).
62 Cadenhead, K., Dobkins, K., McGovern, J. & Shafer, K. Schizophrenia spectrum
participants have reduced visual contrast sensitivity to chromatic (red/green) and
luminance (light/dark) stimuli: new insights into information processing, visual
channel function, and antipsychotic effects. *Frontiers in Psychology* **4**,
doi:10.3389/fpsyg.2013.00535 (2013).
63 Grent-'t-Jong, T. *et al.* Association of Magnetoencephalographically Measured High-
Frequency Oscillations in Visual Cortex With Circuit Dysfunctions in Local and
Large-scale Networks During Emerging Psychosis. *JAMA Psychiatry*,
doi:10.1001/jamapsychiatry.2020.0284 (2020).
64 Palaniyappan, L., Balain, V., Radua, J. & Liddle, P. F. Structural correlates of
auditory hallucinations in schizophrenia: A meta-analysis. *Schizophrenia Research*
137, 169-173, doi:https://doi.org/10.1016/j.schres.2012.01.038 (2012).

65   Leucht, S. *et al.* Clinical implications of Brief Psychiatric Rating Scale scores. *British*
Journal of Psychiatry **187**, 366-371, doi:10.1192/bjp.187.4.366 (2005).
66   Henry, L. P. *et al.* The EPPIC Follow-Up Study of First-Episode Psychosis: Longer-
Term Clinical and Functional Outcome 7 Years After Index Admission. *The Journal*
of Clinical Psychiatry **71**, 716-728, doi:10.4088/JCP.08m04846yel (2010).
67   Wolfers, T. *et al.* Mapping the Heterogeneous Phenotype of Schizophrenia and
Bipolar Disorder Using Normative Models. *JAMA Psychiatry*,
doi:10.1001/jamapsychiatry.2018.2467 (2018).
68   Lv, J. *et al.* Individual deviations from normative models of brain structure in a large
cross-sectional schizophrenia cohort. *Molecular Psychiatry* **26**, 3512-3523,
doi:10.1038/s41380-020-00882-5 (2021).
69   Segal, A. *et al.* Regional, circuit, and network heterogeneity of brain abnormalities in
psychiatric disorders. *medRxiv*, 2022.2003.2007.22271986,
doi:10.1101/2022.03.07.22271986 (2022).
731
732