The emergence of two distinct SARS-CoV-2 Gamma related variants during the second

wave of COVID-19 in Santa Catarina, Southern Brazil and the rapid spread of P.1-like-II

SARS-CoV-2 variant transmission and a regionalization in the Western region.

Authors:

Dayane A. Padilha, Vilmar Benetti-Filho, Renato S. Moreira, Tatiany A. T. Soratto, Guilherme

A. Maia, Ana P. Christoff, Fernando H. Barazzetti, Marcos A. Schörner, Fernanda L. Ferrari,

Carolina L. Martins, Eric K. Kawagoe, Julia K. Wachter, Paula Sacchet, Antuani R. Baptistella,

Aline D. Schlindwein, Bruna K. Coelho, Sandra B. Fernandes, Darcita B. Rovaris, Marlei P. D.

Anjos, Fernanda R. Melo, Bianca Bittencourt, Sthefani Cunha, Karine L. Meneghetti, Nestor

Wendt, Tamela Z. Madaloz, Marcus V. D. Rodrigues, Doris S. M. Souza, Milene H. Moraes,

Rodrigo P. Baptista, Guilherme Toledo-Silva, Guilherme R. Maciel, Edmundo C. Grisard,

Patrícia H. Stoco, Luiz F. V. Oliveira, Maria L. Bazzo, Gislaine Fongaro[1] and Glauber Wagner[1,2]

Affiliations:

Universidade Federal de Santa Catarina, Florianópolis, Santa Catarina, Brazil (D. A. Padilha, V.

Benetti, T. A. T. Soratto, G. A. Maia, F. H. Barazzetti, M. A. Schörner, F. L. Ferrari, C. L.

Martins, E. K. Kawagoe, J. K. Wachter, N. Wendt, T. Z. Madaloz, M. V. D. Rodrigues, D. S. M.

Souza, G. Toledo-Silva, G.R. Maciel, E. C. Grisard, P. H. Stoco, M. L. Bazzo, G. Fongaro and

G.Wagner); and Instituto Federal de Santa Catarina, Lages, Santa Catarina, Brazil (R. S.

Moreira); and Biome-Hub Pesquisa e Desenvolvimento, Florianópolis, Santa Catarina, Brazil (A.

P. Christoff, M. H. Moraes, L. F. V. Oliveira); and Universidade de São Paulo, São Paulo, Brazil

(P. Sacchet); and Universidade do Oeste de Santa Catarina, Joaçaba, Santa Catarina, Brazil (A.

R. Baptistella); and Secretaria de Estado da Saúde, Florianópolis, Santa Catarina, Brazil (A. D.

Schlindwein); and Laboratório Central do Estado da Saúde de Santa Catarina, Florianópolis,

Santa Catarina, Brazil (B. K. Coelho, S. B. Fernandes, D. B. Rovaris, M. P. D. Anjos); and

Diretoria de Vigilância Epidemiológica de Santa Catarina, Florianópolis, Santa Catarina, Brazil

(Fernanda R. Melo, Bianca Bittencourt); and Serviço Nacional de Aprendizagem Industrial,

Chapecó, Santa Catarina, Brazil (S. Cunha, K. L. Meneghetti); and University of Georgia,

Athens, Georgia, USA (R. P. Baptista).

ABSTRACT

COVID-19 has assumed significant and lasting proportions worldwide. Following initial cases in

the Western mesoregion, the State of Santa Catarina (SC), southern Brazil, was heavily affected

as a whole by the pandemic in early 2021. This study aimed to evaluate the dynamics of the

SARS-CoV-2 virus spreading patterns in the SC state through March 2020 to April 2021 using

genomic surveillance. During this period, 23 distinct variants, including two VOCs (Beta and

Gamma) were identified, among which, the Gamma and related lineages were predominant in

the second pandemic wave within SC. However, a regionalization of P.1-like-II in the Western

region was observed, concomitant to the increase in cases, mortality, and case fatality rate (CFR)

index. This is the first evidence of the regionalization of the SARS-CoV-2 transmission and

highlights the importance of tracking variants, dispersion and their impact of SARS-CoV-2 on

the public health system in Brazilian states.

Keywords: SARS-CoV-2; Surveillance; Genome; SARS-CoV-2 P.1 variant.

Introduction

The COVID-19 pandemic, caused by the SARS-CoV-2 virus, is the world's most

significant public health challenge of the last 100 years. This virus has infected more than 270

million people, leading to more than 5,3 million deaths worldwide by December 2021(*1*). In the

same period, since the first reported case on 12/01/2019, this virus infected around 22.2 million

people leading to 617 thousand deaths in Brazil and in the Southern State of Santa Catarina (SC),

1,238,056 cases and 20,101 deaths were reported (*2*).

With the virus' rapid accumulation of polymorphisms and the advance of the pandemic

new variants have emerged (*3*). Thus, genomic surveillance has been essential for monitoring

this disease and vital for local health agencies to have information to mitigate the effects of the

pandemic and adopt strategies to increase population engagement in COVID control policies (*4*).

Until December 2021, more than 6,1 million SARS-CoV-2 genomes have been

sequenced, allowing detection of several variants (*5*). Some new variants have distinctive

biological characteristics from their predecessors, such as better transmission, immunological

escape or distinguishing clinical symptoms (6). Some variants of SARS-CoV-2 have been

defined as Variants of Interest (VOIs) or Variants of Concern (VOCs) with potential impact on

public health. The main VOCs described have been: B.1.1.7/VOC 202012 (Alpha), reported in

December 2020; B.1.351/501Y.V2 (Beta), December 2020; P.1/ B.1.1.248/B1.1.28 (Gamma), in

January 2021; and B.1.617.2 (Delta), in May 2021 (*1*). Recently, the B.1.1.529 (Omicron)

variant was reported in South Africa in November 2021 (1), evidencing that new VOC of SARS-

CoV-2 are still emerging and, thus, genomic surveillance of the virus remains crucial.

In Brazil, the VOC Gamma (P.1) variant, derived of B.1.1.28, was first reported in the

city of Manaus, State of Amazonas, being identified in 42% of RT-PCR positive samples

collected between December 15th and 23th, 2020. Such VOC was not detected by genome

surveillance using samples until November 2020 (*7*). This VOC has 1.7-2.6 times more

transmissible than previous strains circulating in Brazil, showing a 1.2-1.9-fold increase in the

mortality risk to adults and 5 - 8-fold for the young population (*8*), broadening the susceptible

population under risk of infection and hospitalization and the needs of hospitalization and

intensive care (*9*).

Among several detected mutations for VOC Gamma the E484K mutation occurs in the

receptor binding domain (RBD) domain of the S protein and is believed to assist the evasion of

the virus from the immune system (*10*) and, therefore, may lead to lower vaccine efficacy (*11*).

Also present in the RBD domain, the N501Y mutation is associated with increased binding

specificity to the host cell angiotensin-converting enzyme 2 (ACE2) receptor. Subvariants of

VOC Gamma and related lineages (P.1-like-I and P.1-like-II), displaying distinct mutations, have

also emerged during 2021 (*12*). Among these, P.1.1, P.1.2, and P.1-like-II have spread massively

across the border States of Rio Grande do Sul (*13*) and Paraná (14) up to April 2021 and, was

later concomitantly detected in SC state (*12*).

In this context, our study evaluated the dispersion of different variants of SARS-CoV-2

through March 1st, 2020 to April 30th, 2021, in the state of Santa Catarina, especially during the

second wave of the pandemic (January to April 2021) when the VOC Gamma variant became

predominant in Brazil.

Material and methods

 Studied regions and sampling

The State of Santa Catarina (SC) is located in the southern region of Brazil, having

borders with the Paraná (North) and Rio Grande do Sul (South) states, and Argentina (West).

With an estimated population of 7,333,473 inhabitants, the SC state is geographically divided

into six mesoregions: Greater Florianópolis, Itajaí River Valley, Mountain range, Northern,

Southern, and Western SC (*15*).

Genome sequences from 203 SARS-CoV-2 positive samples collected from different

mesoregions of the state of Santa Catarina, Brazil through March 1st, 2020 to April 30th, 2021,

were generated (Figure 1A) (Supplementary Table 1). Additionally, 527 SARS-CoV-2 genome

sequences obtained from samples isolated in SC (Figure 1B) and 49 genome sequences from

samples isolated worldwide, including the original SARS-CoV-2 sequence (NC_045512.2), were

retrieved from the EpiCov database GISAID (*5*) and included in this study (only genome

sequences showing <5% ambiguous nucleotide and > 85% genome sequence coverage). This

study was approved by the Federal University of Santa Catarina Ethics Committee (CAAE:

31521920.8.0000.0121).

RNA purification and sequencing

The study used 203 human nasopharyngeal swab samples from SARS-CoV-2 positive

patients diagnosed via RT-qPCR. Inclusion criteria was a positive RT-PCR for gene E with C_t

values < 25. Total viral RNA was extracted using the RNA Viral Kit (QIAGEN, Hilden,

Germany) and the long amplicon pooled protocol was used (17). Briefly, complementary DNA

(cDNA) was synthesized from viral RNA using 50 μM random primers and 200 U/μL

Superscript IV reverse transcriptase (ThermoFisher Scientific, Waltham, MA, USA). The

obtained cDNA was amplified using 10 µM of 14 primer sets in individual reactions to amplify

different regions of the SARS-CoV-2 genome (approximately 2.5 Kb in each fragment) and 40

U/µL Platinum *Taq* DNA polymerase High Fidelity (Invitrogen, Waltham, MA, USA).

Amplicons were visualized on agarose 2% gel electrophoresis.

The amplicons were then pooled, purified using the AMPure XP (Beckman Coulter,

Brea, CA, USA) and quantified by Quant-iT Picogreen dsDNA assays (Invitrogen, Waltham,

MA, USA). Pools were then processed using the Illumina DNA Prep (Illumina, San Diego, CA,

USA), following manufacturer's protocol. Libraries were quantified using the Collibri Library

Quantification kit (ThermoFisher Scientific, Waltham, MA, USA), adjusted to 11.5 pM and

prepared for paired-end sequencing (2x 150 bp) using the MiSeq Reagent Kit V2 (300-cycles)

(Illumina, San Diego, CA, USA). Sequencing was performed on the Illumina MiSeq platform

(Illumina, San Diego, CA, USA).

SARS-CoV-2 genome assembling and variant analysis

The SARS-CoV-2 genome was assembled according to ARTIC protocol (*18*). Briefly,

raw reads were processed with Trimmomatic v. 0.39 (*19*) to trim low-quality bases and Illumina

adapters (Phred quality score > 25). Trimmed sequences were mapped to the human reference

genome (hg19 - GCF_000001405.13) to remove contaminants using BWA v. 0.7.17 (*20*). Next,

human-free reads were aligned to the SARS-CoV-2 reference genome (NC_045512.2) using

BWA with default parameters. Ivar v. 1.3 (*21*) and mpileup function from samtools v. 1.7 (*22*)

were used to perform variant calling and to generate genomes consensus sequences. Assembled

genomes were analyzed in the Nextclade (*23*) and Pangolin (*24*) platforms to classify the SARS-

CoV-2 variants.

Mutations, deletions, and insertions were used to assess the dissimilarity of the genomes

using the vegan R package v. 2.5.6 (*25*) with the Jaccard index method. The multidimensional

scaling (MDS) and the distribution of variants in SC were plotted using the ggplot2 R package v.

3.3.0 (*26*).

Maximum-likelihood (ML) phylogenetic analysis

The ML phylogeny was performed using all 779 SARS-CoV-2 genome sequences

(Supplementary Table 2). Sequences were aligned by MAFFT v. 7.310 (*27*) with default

parameters and were manually curated using AliView v.1.26 (*28*). A ML tree was constructed

using the alignment, with GTR+F+I as the substitution model predicted by ModelFinder (*29*),

1,000 bootstrap runs using UFBoot tree optimization (30) and SH-like approximate likelihood

ratio test (SH-aLRT) (*31*), using IQ-TREE v. 2.1.3 (*32*). ML tree was visualized and annotated

using the ggtree R package v. 2.0.4 (*33*).

Network analysis of SC VOC Gamma and related lineages during the second wave

The network analysis was performed using a total of 589 Brazilian VOC Gamma and

related genomes retrieved from GISAID, of which, 418 were SC SAR-CoV-2 genomes from

samples obtained from January 5[th], 2021, to April 30[th],2021, and 171 genome sequences from

other Brazilian states sequenced up to January 5[th], 2021 (Supplementary Table 3). The ML

consensus tree as constructed like above, except the substitution model was GTR+F+R2. The

network analysis was performed using online service StrainHub with closeness centrality (*34*),

ML consensus tree, and corresponding metadata.

Spike (S) gene sequence polymorphism and protein structure analysis

For structural analysis of S protein of VOC Gamma, 57 presented full-length S protein

coverage of VOC Gamma sequenced in this study with 722 retrieved from GISAID (December

3rd, 2021), including the reference sequence (NC_045512.2) were aligned using MAFFT (27).

The mutations were represented over the 3D S protein structure PDB 6ZGG (resolution 3.8 Å).

The experimental structure presents a representative coverage of the N-terminal domain (NTD)

(Gln14 to Asp1146). The mutations were highlighted and colored according to observed

frequencies and built with PyMOL version 2.3.3 software (35).

Results

Profile of SARS-CoV-2 variants in State of Santa Catarina after one year of COVID-19

pandemic

A total of 23 distinct SARS-CoV-2 variants were identified among the studied genomes

(Figure 2A). Of these 730 complete genomes, 316 (43.2%) were classified as the VOC Gamma,

followed by 175 (23.9%) of the former VOI Zeta variant (P.2) in and 69 (9.4%) of the P1-like II

variant. Only four sequences of the VOC Alpha variant (B.1.1.7) were observed among the

analyzed samples.

The first sequenced sample was classified as a B.39 variant. Then, the B.1.1.28 and the

B.1.1.33 variants prevailed from May - November 2020, when the P.2 variant gradually became

the most prevalent. The P.2 variant peaked in December 2020 and was displaced quickly by the

P.1 variant. By January 2021, the P.1 variant had already taken over most of the samples in Santa

Catarina (Figure 2A). Until December 2020 the profile of variants is similar among the

mesoregions. However, in the beginning of the second wave) (January to April 2021), the

Western mesoregion has a distinct profile, with the P.1-like-II lineage being predominant in that

region. (Figure 2B). The Mountain Range was sampled only in February 2021. The VOC

Gamma was predominantly identified (n=10) and a single P.1-like-II was detected.

VOC Gamma and related lineages in Santa Catarina.

We inferred phylogenetic relationships between the SARS-CoV-2 genomes from SC and

the other known variants (Figure 3). Most of the SC sequences were grouped in the VOC

Gamma clade, which includes the P.1 and related lineages (P.1.1, P.1.2, P.1.10, P.1-like I and

P.1-like II). However, this parent clade generates two distinct sub-clades, one containing the

variant P.1-like-II and P.1-like-I, and the other clustering sequences classified as P.1 and their

subvariants (P.1.1, P.1.2, and P.1.10). It is noteworthy to mention that the clade formed mainly

by P.1-like-I and P.1-like-II variants is composed of samples from the Western SC, pointing out

a possible independent introduction and regionalized transmission of such variants in that region.

A Multidimensional scale (MDS) plot shows the clustering of the SARS-CoV-2 genomes

according to the mutations observed (Figure 4A). There was no clear clustering of the studied

SARS-CoV-2 variants according to the sampled geographic regions along the sampling period

(Figure 4B). However, in a separate analysis of the data from the VOC Gamma and the related

lineages, it was possible to observe a clustering of a set of genomes from Western SC (Figure

4C), mainly composed by the P.1-like-II variant (Figure 4D).

As the ML and the MDS analyses pointed out the evidence of genetic proximity of the

cluster of P.1 related lineage (P.1-like-II) in the Western SC, we compared all SC VOC Gamma

and related lineages sequences obtained from August 7th, 2020 to April 30th, 2021. These

genomes were classified based on the amino acid substitutions described in P.1-like-I (*36*) and

P.1-like-II (*13*) variants. Mutations that characterize each VOC Gamma related lineage profile

were identified and are summarized in Supplementary Figure 1.

The S protein of the VOC Gamma and related lineages were analyzed, higher mutation

frequencies (>90%) were observed as L18F, P26S, D138Y, and R190S positions of the NTD

region (Figure 5). The typical P.1 mutations were found in the RBD domain (K417T, E484K,

and N501Y), SD domain (D614G and H655Y), and overall structure (T1027I, and V1176F). In

20% of sequences conserved the original amino acid T20 in the NTD, being characterized as P.1-

like-II lineages.

The rapid spread and regionalization of P.1-like-II during the second wave of COVID-19

pandemic in Santa Catarina, Brazil

During the second wave of COVID-19 (January 2021 to April 2021) the VOC Gamma

and subvariants were the most representative SC. However, it does not present a homogeneous

variant distribution pattern in the mesoregions (Figure 2B). It is clear by comparative genome

analysis that many sequences from SC were directly connected to those in Manaus (Amazonas,

Brazil). Network analysis revealed that VOC Gamma and related lineage genomes had a genetic

proximity with genomes sequenced in the State of Amazonas (Figure 6A). The VOC Gamma

and related lineages from the Western SC had fewer interactions with other SC mesoregions than

with the State of Amazonas (Figure 6B). The city of Chapecó-SC, the major municipality of the

Western Santa Catarina, acted as a point of concentration and dissemination of lineages imported

from Manaus to other Western cities (Figure 6C).

Also, it was observed that in February and March 2021, 75% and 60% of the sequences

in the Western SC were classified as P.1-like-II, respectively (Figure 7). Furthermore, in the

same period, this mesoregion had the highest Case Fatality Rate (CFR), mortality, and the

number of deaths in the state of Santa Catarina (Figure 8), which indicates that there was rapid

dissemination of P.1-like-II variant in the region followed by an increase in deaths.

Taken together, these data indicate distinct introductions of VOC Gamma and related

lineages in Santa Catarina, followed by a rapid dispersion of the P.1-like-II variant in the

Western region of SC and a regionalization in the transmission of P.1-like-II in the Western

region.

Discussion

In this study carried out between March 2020-April 2021, we sequenced 203 SARS-

CoV-2 genomes from human samples collected from different regions of the SC state,

representing 27.8% of the virus genomic sequences generated for this region. Assessment of

SARS-CoV-2 genetic profiles (Figure 2A) allowed the identification of VOC Alpha

(20I/501Y.V1/ B.1.1.7) and especially the Gamma (20J/501Y.V3/P.1) from January 2021

onwards, corresponding to the second wave of the pandemic in SC, as also observed in other

Brazilian states (*37, 38*), indicating the homogeneity of the spreading pattern within Brazil at that

time. However, between August 2020 - November 2020 we were able to identify the emergence

of the formerly VOI P.2 (*1*) in SC, which rapidly turned to be more prevalent than the other

variants circulating in the region. This variant, a descendant of B1.1.28, was originally identified

in Brazil in Rio de Janeiro by July 2020, although phylogenetic analyses indicate that the variant

may have arisen as early as February, 2020 (*39*). Similar to our findings in SC, P.2 was reported

to have become predominant in several Brazilian states around the same period, especially in

Northeastern and Southeastern Brazil, when variant P.1 turned to be more prevalent by

beginning 2021 (*39*), except for the State of Amazonas were P.2 variant was first detected in

November 2020 but was overshadowed by P.1 since its early detection and thus never became

the main circulating variant (*38*). After the emergence of VOC Gamma in Manaus City, State of

Amazonas (*7*), the first case of P.1 SC was reported in in Florianópolis City (Greater

Florianópolis) (hCoV-19/Brazil/SC-1326/2021) by January 8[th], 2021, and the first case of VOC

Gamma related lineage (P.1-like-II) in Chapecó City (Western Santa Catarina) on January 22[th],

2021, the later also reported by Gräff et al. (*12*).

Maximum likelihood analysis of P.1-like-II samples obtained from patients living in the

Western mesoregion of SC were clustered in a single, separate clade (Figure 3), while viral

samples from the other SC regions due to scarce P.1-like-II were mostly VOC Gamma and

subvariants, pointing out that introduction and initial spreading of P.1-like-II variant in SC

occurred via Western mesoregion (Figures 4C and 4D). Also, as highlighted in Figure 5 and

Supplementary Figure 1, VOC Gamma is characterized by three mutations in the Receptor

Binding Domain - RBD (K417T, E484K, and N501Y) that can modulate ACE2/RBD affinity,

increasing transmissibility or even the affinity of antibody. In addition to mutations L18F, T20N,

P26S, D138Y and R190S in the N-terminus (NTD), D614G and H655Y in the C-terminus of S1,

and T1027I and V1176F in S2 are also report to VOC Gamma (*40*). All these mutations were

identified in SC, however the cluster 3 and 8 contain the P.1-like-II lineage are formed by

sequences with the absence of T20N mutation and are composed specially by samples from

Western region.

Thus, in this study, analysis of such markers allowed identification of two main lineages

that are distinguished by the presence or absence of the T20N mutation in the S protein

(Supplementary Figure 1). This SARS-CoV-2 protein is highly glycosylated, being involved in

the virus evasion mechanisms of the human immune system through mutations that hinder viral

recognition by the immune system. Most of these glycosylation sites are located in Asp residues

that contains N-glycosylation sites, or Ser and Thr residues that contain O-glycosylation sites.

Among these, the most frequent glycosylation sites were found in N+2, where Asp is followed

by a Ser/Thr residues (41). Remarkably, the herein reported T20N mutation at the NTD is

followed by a Thr (N+2) in all analyzed samples (N20X21T22) and introduces a new potential

O-glycosylation site within the S protein that may increase the virus capability to evade the

immune defenses of viral neutralization by means of antibodies, not altering the potential N-

glycosylation site at N17 that is also present in P.1 variants, where a Thr follows an Asp residue

at position 19 (N17X18T19).

As shown in Figure 4D, by February 2021 we identified a high frequency of the

subvariant P.1-like-II in samples from Western SC if comparing to the other regions of the state,

preceding an increase in the CFR index, mortality rate, and deaths (Figure 8A) on that

mesoregion. Taken together, these data indicate that P.1-like-II had high transmissibility and

effect on the lethality in the Western SC. Naveca et al. (38) showed how the second wave of

COVID-19 in the state of Amazonas, with a sudden rise in cases and number of deaths during

late 2020 and early 2021, directly correlates to the rise in Gamma frequency in the state, which

had reached a massive prevalence by January 2021. Faria et al. (7) demonstrated that the

infection by VOC Gamma is 1.1-1.8 times more severe than those previously observed by other

variants, considering that the results of this variant was related to higher viral loads, greater

transmissibility and an increase in the number of deaths.

The relationship between the increase in the number of cases and the appearance of

variants was also observed outside of Brazil, such as in India and Europe. In India, the

appearance of the Delta variant was observed during the second wave, correlating with the

consequent increase in the number of deaths. In Europe, the second wave was also marked by an

increase in the number of cases and deaths, which correlated with the appearance of the Alpha

variant (42-43).

However, transmission rate of SARS-CoV-2 may be affected by a variety of factors and

the emergence of a given variant could not be considered the unique factor to increase such rate

(*44*). Along several virus-inherent factors such as genetic variability, the lack or misuse of non-

pharmaceutical interventions (NPI), social difficulties for diagnosis or late diagnosis,

uncontrolled population movement (emigration, immigration, tourism) (*45*), the collapse of the

public health system (*46*), lower vaccination (*47*), the overtime decrease of the immune

protection (*46*), late or misdecisions taken by authorities (*49*), and the use of scientifically

unproven preventive therapies (*50 - 52*), are also important factors that might contribute to

increase the transmission and case fatality rates. Examples in different countries show that an

increase in the movement of people and low widespread adherence to the NPI was observed

weeks before the collapse of the public health service, and the consequent increase in the number

of cases and deaths (*53-54*). Indeed, the SARS-CoV-2 circulation is evidenced by the detection

of viral RNA copies in sewage months before being reported in people, as occurred in Santa

Catarina state in November 2019 (55). Thus, we cannot rule out that conjunction of both viral

and other biological, environmental and behavioral factors have significant influence on the

occurrence of SARS-CoV-2 variants presenting variable infectivity and pathological effects.

Such may have contributed to the differentially increased number of cases, hospitalization, and

deaths (CRF) in a short period of time in the Western SC as also previously observed in Manaus

(Amazon, Brazil) having totally distinct environmental conditions than SC.

With the following report of the VOC Omicron in South Africa in November 2021 (*1*),

this variant was rapidly detected worldwide (1) and reported in Brazil in São Paulo, late

November 2021, evidencing the ongoing viral adaptations and the continuing pandemic

spreading pattern. Considering these facts, it is of utmost importance the needs for maintenance

of a detailed and standardized genomic surveillance of SARS-CoV-2 variants towards precise

and specific identification of VOCs, avoiding erroneous epidemiological inferences and allowing

evaluation of vaccine efficacy across viral variants.

Genomic surveillance of the SARS-CoV-2 pandemic has been crucial to assess the

accumulation of mutations, mostly due to the enormous quantity of viral replication events, and

the biological outcomes of the genetic variability of each variant on the epidemiology,

transmissibility, infectivity, pathogenesis and vaccine efficacy and effectiveness. Since major

concerns over the genetic evolution of SARS-CoV-2 are related to an increased spreading

capability and to the escape from the immune system, genomic surveillance provides essential

data to prevent the current pandemic from lasting even longer.

Conclusion

Worldwide assessment of circulating variants of SARS-CoV-2 is crucial to predict and

evaluate the impact of the COVID-19 pandemic over time. In SC (Southern Brazil), an attractive

touristic destination, the VOC Gamma was the most prevalent variant detected among the

infected population. Despite such general predominance, our data suggest a primary, distinct

introduction, and rapid spreading of the P.1-like-II in the Western regions of SC in the initial

months of 2021, where it was responsible for a distinct and increased hospitalization and

mortality rates. This study allowed us to observe the distinct patterns of circulation of different

SARS-Cov-2 variants within SC regions. Thus, the continuous real–time genomic surveillance of

SARS-CoV-2 in Santa Catarina will offer precise epidemiological support for implementation

and review of policies and actions carried out by public health authorities towards control of

COVID-19 transmission and assessment of vaccine efficacy.

Acknowledgments

This work is dedicated to all SC citizens who suffered or passed away due to COVID-19.

We also are indebted to Hospital Universitário/HU/EBSERH from Federal University of Santa

Catarina (Florianópolis, SC, Brazil), Hospital Regional do Oeste (Chapecó, SC, Brazil) and

Hospital Regional Santa Terezinha (Joaçaba, SC, Brazil) for granting access to interview patients

and perform sample collection. This work was supported by Santa Catarina Research Foundation

(Fundação de Amparo à Pesquisa e Inovação of Santa Catarina, FAPESC, Santa Catarina, Brazil)

(grant number COV2020051000065), CAPES (Coordination for the Improvement of Higher

Education Personnel, Brazil), CNPq (National Council for Scientific and Technological

Development, Brazil) and UFSC (Federal University of Santa Catarina). We also are thankful to

all institutions that provide access to genome data at EpiCov (GISAID). DAP, VBF, TATP,

GAM, CLM, EKK and TZM were recipients of FAPESC, CAPES or CNPq scholarships.

Disclaimers

The opinions expressed by authors contributing to this journal do not necessarily reflect the

opinions of the Federal University of Santa Catarina, Brazil, or the institutions with which the

authors are affiliated. Funders have no role on study design, data analysis or decision to publish.

Author Bio (first author only, unless there are only 2 authors)

Dayane A. Padilha is a PhD student in Biotechnology and Biosciences at the Federal University

of Santa Catarina (UFSC) and recipient of a CAPES scholarship. She works at both Applied

Virology and Bioinformatics Laboratories at UFSC. Her main interest is on virus genomics

applied to genomic surveillance.

Footnotes (if applicable)

[1] These authors were co–principal investigators.

[2] Corresponding author.

References

1. World Health Organization. WHO Coronavirus (COVID-19) [Internet]. 2019 Dec 30

[cited 2021 Dec 12] https://www.who.int

2. National Council of Health Secretaries - Covid-19 CONASS Panel [Internet]. 2021 May

[cited 2021 Dec 4] https://www.conass.org.br/painelconasscovid19/ .

3. Singh J, Pandit P, McArthur AG, Banerjee A, Mossman K. Evolutionary trajectory of

SARS-CoV-2 and emerging variants. Virol J. 2021 Aug 13;18(1):166.

4. Meredith LW, Hamilton WL, Warne B, Houldcroft CJ, Hosmillo M, Jahun AS, et al.

Rapid implementation of SARS-CoV-2 sequencing to investigate cases of healthcare

associated COVID-19: a prospective genomic surveillance study. Lancet Infect Dis. 2021

Mar;21(3):e36. Corrected and republished from Lancet Infect Dis. 2020

Nov;20(11):1263-1272.

5. Global Initiative on Sharing Avian Influenza Data. [Internet]. 2008 [cited 2021 Nov 27]

https://www.gisaid.org .

6. Tao, K., Tzou, P.L., Nouhin, J. *et al.* The biological and clinical significance of emerging

SARS-CoV-2 variants. Nat Rev Genet. 2021; 22**:** 757–773.

7. Faria NR, Mellan TA, Whittaker C, Claro IM, Candido DDS, Mishra S, et al. Genomics

and epidemiology of P.1 SARS-CoV-2 lineage in Manaus, Brazil. Science. 2021 May 21;

372(6544):815-821.

8. Hemmer CJ, Löbermann M, Reisinger EC. COVID-19: epidemiology and mutations: An

update [in German]. Radiologe. 2021 Oct;61(10):880-887.

9. Nonaka CKV, Gräf T, Barcia CAL, Costa VF, de Oliveira JL, Passos RDH, et al. SARS-

CoV-2 variant of concern P.1 (Gamma) infection in young and middle-aged patients

admitted to the intensive care units of a single hospital in Salvador, Northeast Brazil,

February 2021. Int J Infect Dis. 2021 Oct;111:47-54.

10. Greaney AJ, Starr TN, Gilchuk P, Zost SJ, Binshtein E, Loes AN, et al. Complete

Mapping of Mutations to the SARS-CoV-2 Spike Receptor-Binding Domain that Escape

Antibody Recognition. Cell Host Microbe. 2021 Jan 13;29(1):44-57.

11. Liu Z, VanBlargan LA, Bloyet LM, Rothlauf PW, Chen RE, Stumpf S, Zhao H, Errico

JM, Theel ES, Liebeskind MJ, Alford B, Buchser WJ, et al. Identification of SARS-CoV-

2 spike mutations that attenuate monoclonal and serum antibody neutralization. Cell Host

Microbe. 2021 Mar 10;29(3):477-488.e4. doi: 10.1016/j.chom.2021.01.014.

12. Gräf T, Bello G, Venas TMM, Pereira EC, Paixão ACD, Appolinario LR, et al.

Identification of SARS-CoV-2 P.1-related lineages in Brazil provides new insights about

the mechanisms of emergence of Variants of Concern. [Internet]. 2021 Jun [cited 2021

Nov 30] https://doi.org/10.21203/rs.3.rs-580195/v1 .

13. Varela APM, Prichula J, Mayer FQ, Salvato RS, Sant'Anna FH, Gregianini TS, Martins

LG, Seixas A, Veiga ABGD. SARS-CoV-2 introduction and lineage dynamics across

three epidemic peaks in Southern Brazil: massive spread of P.1. Infect Genet Evol. 2021

Dec;96:105144.

14. Oliveira MM, Schemberger MO, Suzukawa AA, Riediger IN, do Carmo Debur M,

Becker G, et al. Re-emergence of Gamma-like-II and emergence of Gamma-S:E661D

SARS-CoV-2 lineages in the south of Brazil after the 2021 outbreak. Virol J. 2021 Nov

17;18(1):222.

15. Brazilian Institute of Geography and Statistics - Official Territorial Area - Federative

Unit Consultation [Internet]. 2010 [cited 2021 Oct 15] https://www.ibge.gov.br/cidades-

e-estados/sc.html .

16. Santa Catarina Epidemiological Surveillance Directorate [Internet]. 2020 Mar [cited 2021

Nov 7] https://www.dive.sc.gov.br/ .

17. Eden J-S., Sim E. SARS-CoV-2 Genome Sequencing Using Long Pooled Amplicons on

Illumina Platforms. [Internet]. 2020 Apr 04 [cited 2021 Nov 24]

https://www.protocols.io/view/sars-cov-2-genome-sequencing-using-long-pooled-amp-

befyjbpw?step=81 .

18. Young, E., Oakeson, K. Utah DoH ARTIC/Illumina Bioinformatic Workflow [Internet].

2021 Jan [cited 2021 Nov 28] https://github.com/CDCgov/SARS-CoV-

2_Sequencing/tree/master/protocols/BFX-UT_ARTIC_Illumina .

19. Bolger, A. M., Lohse, M., Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014 Aug 1;30(15):2114-20.

20. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009 Jul 15;25(14):1754-60.

21. Grubaugh ND, Gangavarapu K, Quick J, Matteson NL, De Jesus JG, Main BJ, et al. An

amplicon-based sequencing framework for accurately measuring intrahost virus diversity

using PrimalSeq and iVar. Genome Biol. 2019 Jan 8;20(1):8.

22. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. Gigascience. 2021 Feb

16;10(2):giab008.

23. Nextclade Web 1.10.0 [Internet]. 2020 Jun 11 [cited 2021 Nov 9].

https://clades.nextstrain.org/ .

24. Pangolin COVID-19 Lineage Assigner [Internet]. 2021 [cited 2021 Nov 9].

https://pangolin.cog-uk.io/ .

25. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. Vegan:

Community Ecology Package. R package version 2.0-2 [Internet]. 2020 Nov 28 [cited

2021 Dec 5] https://CRAN.R-project.org/package=vegan .

26. Wickham, H. Ggplot2: Elegant Graphics for Data Analysis (ed. Springer-Verlag)

cheatsheet. New York; 2016.

27. Katoh K, Standley DM. A simple method to control over-alignment in the MAFFT

multiple sequence alignment program. Bioinformatics. 2016 Jul 1;32(13):1933-42.

28. Larsson A. AliView: a fast and lightweight alignment viewer and editor for large

datasets. Bioinformatics. 2014 Nov 15;30(22):3276-8.

29. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder:

fast model selection for accurate phylogenetic estimates. Nat Methods. 2017

Jun;14(6):587-589.

30. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the

Ultrafast Bootstrap Approximation, Molecular Biol. Evol. 2018 Feb;35(2):518–522.

31. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New

Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the

Performance of PhyML 3.0, System. Biol. 2010 May;59(3):307–321.

32. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et

al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the

Genomic Era. Mol Biol Evol. 2020 May 1;37(5):1530-1534.

33. Yu G. Using ggtree to Visualize Data on Tree-Like Structures. Curr Protoc

Bioinformatics. 2020 Mar;69(1):e96.

34. Schneider AB, Ford CT, Hostager R, Williams J, Cioce M, Çatalyürek ÜV, et al.

StrainHub: a phylogenetic tool to construct pathogen transmission networks.

Bioinformatics. 2020 Feb 1;36(3):945-947.

35. Schrödinger, LLC. The PyMOL Molecular Graphics System, Version 2.0 [Internet]. 2017

Sep 20 [cited 2021 Nov 17] https://pymol.org/2/ .

36. Naveca F, Nascimento V, Souza V, Corado A, Nascimento F, Silva G, et al. Phylogenetic

relationship of SARS-CoV-2 sequences from Amazonas with emerging Brazilian variants

harboring mutations E484K and N501Y in the Spike protein. Genom. Epidemiol.

[Internet]. 2021 Jan 11 [cited 2021 Nov 12] https://virological.org/t/phylogenetic-

relationship-of-sars-cov-2-sequences-from-amazonas-with-emerging-brazilian-variants-

harboring-mutations-e484k-and-n501y-in-the-spike-protein/585 .

37. Castro, M. C. *et al*. Spatiotemporal pattern of COVID-19 spread in Brazil. Science. 2021

May 21;372(6544):821-826.

38. Naveca FG, Nascimento V, de Souza VC, Corado AL, Nascimento F, Silva G, et al.

COVID-19 in Amazonas, Brazil, was driven by the persistence of endemic lineages and

P.1 emergence. Nat Med. 2021 Jul;27(7):1230-1238.

39. Lamarca AP, de Almeida LGP, Francisco RDS Jr, Lima LFA, Scortecci KC, Perez VP, et

al. Genomic surveillance of SARS-CoV-2 tracks early interstate transmission of P.1

lineage and diversification within P.2 clade in Brazil. PLoS Negl Trop Dis. 2021 Oct

13;15(10):e0009835.

40. Dejnirattisai, W *et al*. Antibody evasion by the P.1 strain of SARS-CoV-2. Cell. 2021

May 27;184(11):2939-2954.e9.

41. Tian W, Li D, Zhang N, Bai G, Yuan K, Xiao H, et al. O-glycosylation pattern of the

SARS-CoV-2 spike protein reveals an "O-Follow-N" rule. Cell Res. 2021

Oct;31(10):1123-1125.

42. Arnab Sarkar, Alok Kumar Chakrabarti, Shanta Dutta. Pathogens. Covid-19. Infection in

India: A Comparative Analysis of the Second Wave with the First Wave. Pathogens.

2021 Sep 21;10(9):1222.

43. Katarzyna J, Samuel A, Pascal A and Mondher T. On the association between SARS-

COV-2 variants and COVID-19 mortality during the second wave of the pandemic in

Europe. Journal of Market Access & Health Policy. 2021 VOL. 9, 2002008.

44. Sabino EC, Buss LF, Carvalho MPS, Prete CA Jr, Crispim MAE, Fraiji NA, et al.

Resurgence of COVID-19 in Manaus, Brazil, despite high seroprevalence. Lancet. 2021

Feb 6;397(10273):452-455.

45. Flaxman S, Mishra S, Gandy A, Unwin HJT, Mellan TA, Coupland H, et al. Estimating

the effects of non-pharmaceutical interventions on COVID-19 in Europe. Nature. 2020

Aug;584(7820):257-261.

46. da Silva SJR, Pena L. Collapse of the public health system and the emergence of new

variants during the second wave of the COVID-19 pandemic in Brazil. One Health. 2021

Dec;13:100287.

47. Kupek E. Low COVID-19 vaccination coverage and high COVID-19 mortality rates in

Brazilian elderly. Rev Bras Epidemiol. 2021 Sep 10;24:e210041.

48. Lumley SF, O'Donnell D, Stoesser NE, Matthews PC, Howarth A, Hatch SB, et al.

Antibody Status and Incidence of SARS-CoV-2 Infection in Health Care Workers. N

Engl J Med. 2021 Feb 11;384(6):533-540.

49. Malta M, Strathdee SA, Garcia PJ. The Brazilian tragedy: Where patients living at the

'Earth's lungs' die of asphyxia, and the fallacy of herd immunity is killing people.

EClinicalMedicine. 2021 Feb 12;32:100757.

50. Furlan L, Caramelli B. The regrettable story of the "Covid Kit" and the "Early Treatment

of Covid-19" in Brazil. Lancet Reg Health Am. 2021 Dec;4:100089.

51. Abella BS, Jolkovsky EL, Biney BT, Uspal JE, Hyman MC, Frank I, et al. Efficacy and

Safety of Hydroxychloroquine vs Placebo for Pre-exposure SARS-CoV-2 Prophylaxis

Among Health Care Workers - A Randomized Clinical Trial. JAMA Intern Med. 2021

Sep 30;181(2):195-202.

52. Popp M, Stegemann M, Metzendorf MI, Gould S, Kranke P, Meybohm P, Skoetz N,

Weibel S. Ivermectin for preventing and treating COVID-19. Cochrane Database Syst

Rev. 2021 Jul 28;7(7):CD015017.

53. Garcia LP, Duarte E. Nonpharmaceutical interventions for tackling the COVID-19

epidemic in Brazil. Epidemiol Serv Saude. 2020 Apr 9;29(2):e2020222.

54. Djin-Ye O, Silke B, Barbara B, Janine R, Frank S, Susanne D, Marianne W, Max VK,

MM, Thorsten W, Ralf D. Trends in respiratory virus circulation following COVID-19-

targeted nonpharmaceutical interventions in Germany, January - September 2020:

Analysis of national surveillance data. Lancet Reg Health Eur. 2021 Jul;6:100112.

55. Fongaro G, Stoco PH, Souza DSM, Grisard EC, Magri ME, Rogovski P, Schörner MA,

Barazzetti FH, Christoff AP, de Oliveira LFV, Bazzo ML, Wagner G, Hernández M,

Rodríguez-Lázaro D. The presence of SARS-CoV-2 RNA in human sewage in Santa

Catarina, Brazil, November 2019. Sci Total Environ. 2021 Jul 15;778:146198.

Address for Correspondence

Address for correspondence: Glauber Wagner, Laboratório de Bioinformática, Departamento de

Microbiologia, Imunologia e Parasitologia (MIP), Centro de Ciências Biológicas (CCB),

Universidade Federal de Santa Catarina (UFSC), Campus João David Ferreira Lima, Bloco E,

Sala 207, Trindade, CEP 88040-900, Florianópolis, Santa Catarina, Brazil. Phone: +55 48 3721

2956. E-mail: glauber.wagner@ufsc.br

(A)

527
(B)

Figure 1. Distribution of sequenced SARS-CoV-2 genomes in Santa Catarina from March 1st,

2020 to April 30th 2021. **(A)** Map showing the Santa Catarina SARS-CoV-2 sequences obtained

from GISAD EpiCov Database (upper map) and samples sequenced in this study (lower map).

(B) Spatial and temporal evolution of COVID-19 cases in the state of Santa Catarina, according

to the Santa Catarina Epidemiological Surveillance Directorate (*16*) during the study period. The

Greater Florianópolis and the Western Santa Catarina mesoregions were the most represented

because they presented the highest cases / 100,000 habitants rates between January 1st 2021 and

April 30th 2021.

(A)

542
543
544 (B)

Figure 2. SARS-CoV-2 variants profile in the state of Santa Catarina from March 2020 to April

2021. **(A)** Analysis of the variant profile considering the whole state of Santa Catarina. **(B)** The

variant profile by Santa Catarina mesoregion. From March to September 2020, the variants

identified were in the highest proportion of B.1.1.28 (Blue) and B.1.1.33 (Light red). From

September 2020 to January 2021, the highest proportion was of P.2 variant, followed by VOC

Gamma and related lineages from February to April 2021.

Figure 3. Phylogenomic reconstruction analysis of Santa Catarina SARS-CoV-2 sequences from

March 1st, 2020 to April 30th 2021. Maximum likelihood (ML) phylogenetic tree was

reconstructed using GTR+F+I substitution model, with 1000 iterations in Ultra-Fast Bootstrap

mode with SH-like approximate likelihood ratio test (SH-aLRT). All 779 sequences were aligned

with 49 reference sequences of distinct SARS-CoV-2 variants available in GISAID. Dots

represent the variant or lineages indicated on the right-bottom legend. According to the right-up

legend, the outside bars indicate the sample mesoregion of Santa Catarina or others. The green

shadow indicates the VOC Gamma and related lineages (P.1.1, P1.2, P.1.10 and P.1.7) clade.

Lilac shadow indicates the P.1-like-II clade.

(A)   (B)

(C)   (D)

Figure 4. Multidimensional scale (MDS) plot of Jaccard dissimilarity matrix between genomes

based on the presence and absence of mutations, deletions, and insertions labeled by **(A)** variants

and **(B)** region. Only VOC Gamma and related lineages MDS analysis labeled by **(C)** region and

(D) variant or lineages.

Figure 5. Overlay of VOC Gamma Spike protein mutations identified in the state of Santa

Catarina during the COVID-19 pandemic. Residues involved in mutations are presented as

spheres, colored from blue to red (low to high mutation frequency). The mutation sites are also

indicated in the two-dimensional representation of the Spike protein, which is colored according

to protein regions. The NTD domain is colored in green and the RBD domain is colored in

purple. From the total of 778 analyzed sequences of the VOC Gamma variant (5' – 3'), red

represents the frequency of 93.6-100%, orange the frequency of 88.8%, yellow the frequency of

3.98%, green the frequency of 1.79%, and blue the frequency of less than 1%. Image generated

using PyMOL version 2.3.3 (*43*).

(A)   (B)

(C)

Figure 6. Network analysis of VOC Gamma and related lineages sequenced in the state of Santa

Catarina, Brazil, from January 3rd, 2020 to April 30th 2021 with the other Brazilian states. The

ML analysis was performed on 13,018 Brazilian VOC Gamma related genomes obtained from

GISAID until April 30th, 2021 and 316 and submitted to StrainHub (*42*) analysis considering the

closeness parameter. The sizes of nodes are scaled by closeness metric and the arrows reflect the

directionality of possible transmissions between the Brazilian states **(A)**, Santa Catarina

mesoregions **(B)** and Santa Catarina cities **(C)**. The thickness of the lines and arrows represents

the frequency of putative transmissions.

Figure 7: Relative frequencies of the SARS-CoV-2 VOC Gamma and related lineages by

mesoregions of the state of Santa Catarina during the second wave of the COVID-19 pandemic,

January to April 2021.

Figure 8. Epidemiological data from each mesoregion of the state of Santa Catarina during the

second COVID-19 pandemic wave (January 2020 to April 2021). Case fatality rate (CFR),

mortality rate, number of deaths and population size. Data source: Santa Catarina

Epidemiological Surveillance Directorate (*16*).

Supplemental Materials

Supplementary Table 1. Synthesis of 203 samples sequenced in the present study. Sequence

name, GISAD accession numbers, city, mesoregion, age, gender, variant and clade assignment.

Supplementary Table 2. List of GISAD accession numbers of sequences used on ML analysis,

by Pangolin lineage.

Supplementary Table 3. GISAD accession numbers of sequences used on Santa Catarina VOC

Gamma and related lineages network analysis.

Supplementary Figure **1**. The non-synonymous amino acid substitution profile of the VOC

Gamma and related lineages genomes in the state of Santa Catarina from January 3rd 2020 to

April 30th 2021.