Antigenic evolution of SARS-CoV-2 in immunocompromised hosts

Cameron A. Smith[1], Ben Ashby[1,2,3,]*

[1]Department of Mathematical Sciences, University of Bath, Bath, BA2 7AY, UK.

[2]Milner Centre for Evolution, University of Bath, Bath, BA2 7AY, UK.

[3]Department of Mathematics, Simon Fraser University, Burnaby, BC, V5A 1S6, Canada.

*Corresponding author. Email: benashbyevo@gmail.com

Keywords: Antigenic evolution, SARS-CoV-2, COVID-19, variant, immunocompromised, epistasis

Abstract

Prolonged infections of immunocompromised individuals have been proposed as a crucial source of new variants of SARS-CoV-2 during the COVID-19 pandemic. In principle, sustained within-host antigenic evolution in immunocompromised hosts could allow novel immune escape variants to emerge more rapidly, but little is known about how and when immunocompromised hosts play a critical role in pathogen evolution. Here, we use a simple mathematical model to understand the effects of immunocompromised hosts on the emergence of immune escape variants in the presence and absence of epistasis. We show that when the pathogen does not have to cross a fitness valley for immune escape to occur (no epistasis), immunocompromised individuals have no qualitative effect on antigenic evolution (although they may accelerate immune escape if within-host evolutionary dynamics are faster in immunocompromised individuals). But if a fitness valley exists between immune escape variants at the between-host level (epistasis), then persistent infections of immunocompromised individuals allow mutations to accumulate, therefore facilitating rather than simply speeding up antigenic evolution. Our results suggest that better genomic surveillance of infected immunocompromised individuals and better global health equality, including improving access to vaccines and treatments for individuals who are immunocompromised (especially in lower- and middle-income countries), may be crucial to preventing the emergence of future immune escape variants of SARS-CoV-2.

Lay Summary

We study the role that immunocompromised individuals may play in the evolution of novel variants of the coronavirus responsible for the COVID-19 pandemic. We show that immunocompromised hosts can be crucial for the evolution of immune escape variants. Targeted treatment and surveillance may therefore prevent the emergence of new variants.

Introduction

Understanding how and when variants of SARS-CoV-2, the causative agent of COVID-19, are likely to evolve is key to managing the future of the pandemic. Multiple variants of concern have evolved since the start of the pandemic, with higher transmissibility evolving on at least two occasions, by the Alpha (B.1.1.7) variant (relative to the wildtype) [1], and by the Delta (B.1.617.2) variant (relative to Alpha) [2,3], with the latter becoming the globally dominant strain in 2021 [4]. Other variants such as Beta (B.1.351) and Omicron (B.1.1.529) have additionally shown evidence of immune escape, indicating antigenic evolution [5–7] (Omicron has also been linked with an increase in transmission [8,9]). With increasing numbers of people acquiring immunity to SARS-CoV-2, either through infection or vaccination, we should expect a shift towards antigenic evolution rather than higher transmissibility or greater virulence as the primary driver of new variants of concern [10]. The extent to which SARS-CoV-2 may evolve antigenically in future, thereby allowing it to evade host immunity partially or fully, is currently unknown. However, the emergence and rapid spread of Omicron towards the end of 2021 has demonstrated that antigenic evolution is both possible and under strong selection. The unusual nature of Omicron (possessing a large number of mutations in the spike protein but only distantly related to the dominant variant at the time, Delta [11]) has led to speculation that it underwent long-term within-host evolution in an immunocompromised individual who was unable to clear the infection [12]. We explore this hypothesis using a simple mathematical model to understand the potential importance of immunocompromised individuals for the antigenic evolution of SARS-CoV-2.

A fundamental tenet of evolutionary epidemiology is that the rate of antigenic evolution depends on a balance between immune pressure and mutation supply [13–15]. The greater the proportion of the population that is immune, the greater the strength of selection for immune escape but mutation supply is constrained as few hosts can be infected. Conversely, if many hosts are susceptible to infection, then mutation supply may be plentiful but selection for immune escape is relatively weak.

Hence, the rate of antigenic evolution should be maximised at an intermediate level of immune

pressure, whereby moderate pathogen prevalence leads to a plentiful supply of mutations for

selection to act upon, and the strength of selection for immune escape is reasonably strong.

Rapid deployment of vaccinations against SARS-CoV-2 combined with the relaxation of non-

pharmaceutical interventions in many countries led to both strong immune pressure and high

numbers of infections in the latter half of 2021. For example, by the end of November 2021 the UK

had fully vaccinated 68% of the population while still experiencing over 620 confirmed cases per

million (approximately 70% of the previous peak in January 2021) [16]. At the time, the Delta variant

was dominant globally and accounted for over 99% of infections in the UK [16]. Yet, despite apparently

favourable evolutionary conditions for immune escape there were no indications of the Delta variant

exhibiting antigenic evolution in the UK or elsewhere. This may indicate that closely related immune

escape variants were suppressed, perhaps by transiently boosted innate, B, and T cell responses, or

due to epistasis (e.g. less transmissible). Instead, the initial BA.1 sublineage of the Omicron variant,

first detected in South Africa and reported to the World Health Organization on November 24, 2021

[11], was able to substantially escape host immunity and evolved from a distant clade. This variant

contains 30 mutations to the spike protein (used for binding to host cell receptors) and has been

shown to evade over 85% of neutralizing antibodies [7]. Relative to Delta, it exhibits substantially

lower vaccine effectiveness [17] and is estimated to be over five times as likely to lead to reinfection

[6]. The BA.1 sublineage of Omicron became the dominant variant in the UK within a month and

replaced Delta in many countries in early 2022 [16], with the BA.2 sublineage later replacing BA.1 [18].

The BA.1 sublineage of Omicron confirms that substantial immune escape is not only possible for

SARS-CoV-2 but also that selection for immune escape towards the end of 2021 was very strong.

According to the conceptual model of antigenic evolution as a balance between immune pressure and

mutation supply [13], this suggests that the lack of adaptation to evade host immunity by the Delta

lineage was simply due to insufficient mutation supply. However, this is difficult to reconcile with the

high number of cases at the time, implying mutation supply was plentiful. Furthermore, if mutation

supply was the key constraint, how did an immune escape variant appear from an obscure clade that

was responsible for few infections?

Several hypotheses have been proposed for the sudden emergence of the Omicron variant from a

distant clade. One possibility is that omicron evolved in an animal host following infection by a human,

and then jumped back into the human population. Alternatively, it could have evolved in a remote

population without being detected until it began to spread more widely in late 2021. However, neither

of these explanations are especially convincing. Evolution in an animal host would have not only

required two jumps across the human-animal species barrier, but also selection in the animal host

would have had to correspond to increased fitness in the human population through immune escape.

It is more plausible that Omicron was able to substantially escape immunity in humans because it had

experienced selection for immune escape in humans. Similarly, evolution in a remote population does

not appear to be plausible as it fails to explain why a similar array of mutations were not seen in

regions where mutation supply was significantly higher (due to more infections) and immune pressure

was strong due to vaccine and naturally-acquired immunity.

A more promising hypothesis is that the Omicron variant arose due to long-term within-host evolution

in an immunocompromised individual, who was most likely infected between March and August 2021

[11]. While an immunocompetent individual would be expected to clear infection after a relatively

short period, an immunocompromised person may fail to fully clear the infection, allowing the virus

to coevolve with the immune system [19]. Indeed, longitudinal sequencing from an

immunocompromised patient who was infected for over 150 days with SARS-CoV-2 revealed rapid

accumulation of mutations [20]. These mutations appeared to be adaptive at the within-host level

due to their concentration in the spike protein, with several common to other variants of concern.

Similar results have been observed in other patients with long-term infections of SARS-CoV-2 [21,22],

including those who have been treated with convalescent plasma, indicating antigenic evolution

within the host [23] (although some immunocompromised individuals show little to no within-host

evolution of SARS-CoV-2 [24]).

It is currently unclear how important immunocompromised individuals are for the antigenic evolution

of SARS-CoV-2, or for pathogen evolution more generally. Do infections of immunocompromised

individuals simply accelerate antigenic evolution or do they play a key role in facilitating immune

escape? In the first case, infection of immunocompromised individuals speeds up antigenic evolution

due to a faster rate of adaptation within these hosts, leading to the emergence of new immune escape

variants on shorter timescales than would be possible in an immunocompetent population. Such a

scenario would suggest that although immunocompromised individuals might speed up antigenic

evolution, they are not essential for it to occur. In the second case, long-term infections of

immunocompromised individuals allow the virus to accumulate mutations that are advantageous (or

neutral) at the within-host level but may be disadvantageous (or neutral) at the between-host level.

If there is epistasis between mutations at the between host level (i.e. fitness depends on the context

of which other mutations are present), as indicated by recent experiments [25], then sustained

adaptation within immunocompromised individuals may allow the virus to traverse valleys in the

fitness landscape, which would otherwise be very difficult to cross, to reach another peak. The second

scenario would therefore suggest that long-term infections in immunocompromised individuals play

a disproportionate role in the antigenic evolution of pathogens such as SARS-CoV-2.

Here, we analyse a simple phenomenological model to explore the potential importance of

immunocompromised hosts for the antigenic evolution of SARS-CoV-2. We show that in the absence

of epistasis, antigenic evolution readily occurs regardless of the frequency of immunocompromised

individuals in the population. If epistasis is present, however, such that the virus must traverse a

fitness valley at the between-host level to escape host immunity, then immunocompromised hosts

are crucial for antigenic evolution to occur. These patterns are robust irrespective of whether within-

host evolutionary dynamics are faster in immunocompromised individuals and for a wide range of

parameters affecting cross-immunity, the strength of epistasis, the proportion of the population that

is immunocompromised and their duration of infection relative to immunocompetent hosts.

Model description

We adapt the model of antigenic evolution presented by Gog and Grenfell [26] to incorporate

immunocompromised individuals and epistasis. The model assumes that there are $n = 30$ variants

equally spaced in a line, with adjacent variants differing by a single mutation. Hosts are classed as

either entirely susceptible to a variant, or entirely immune to it. Cross-immunity between variants is

therefore 'polarising', which means that when an individual is infected by variant i, a proportion σ_{ij}

of those currently susceptible to variant j become fully immune to it for life (no waning immunity) and

a proportion $1 - \sigma_{ij}$ remain fully susceptible to variant j. This assumption greatly reduces the

complexity of the model as it means we do not need to track all infection histories, which would

require at least $(2 + n)2^n \approx 137$ billion classes with $n = 30$. The strength of cross-immunity between

variants i and j is given by

$$\sigma_{ij} = \exp\left\{-\frac{(i-j)^2}{2\eta}\right\}, \tag{1}$$

where $\eta > 0$ controls the breadth of cross-immunity (large values of η give broad cross-immunity

between distant variants, whereas small values of η limit cross-immunity to closely related variants;

Fig. 1a). We assume that the population is large, well-mixed, and of constant size ($N = 10^7$), with a

proportion p of individuals who are immunocompromised (only able to produce a weak immune

response; subscript C) and a proportion $1 - p$ who are immunocompetent (able to produce a normal

or "healthy" immune response; subscript H). For simplicity, we ignore host demographics (births and

deaths) and mortality from infection, as we are only interested in the antigenic evolution of the virus

over a relatively short timescale.

Figure 1: Population-level model. (a) Cross-immunity,σ_{ij}, for variants i and j, with lighter colours corresponding to greater

cross-immunity. (b) Illustration of the normalised transmission rate for each variant, showing a fitness valley. (c) Model

schematic.

Let S_H^i (respectively S_C^i) be the proportion of the population that is immunocompetent (resp.

immunocompromised) and susceptible to variant $i \in \{1, \dots, n\}$, and I_H^i (resp. I_C^i) be the proportion of

the population that is immunocompetent (resp. immunocompromised) and infected with variant i. To

incorporate a fitness valley at the between-host level, we assume that the transmission rate of variant

i is given by

$$\beta_i(\xi) = \tilde{\beta}\left(1 + \frac{\xi}{2}\left(\cos\left(\frac{2\pi(i-1)}{n-1}\right) - 1\right)\right),$$ (2)

where $\tilde{\beta}$ is the maximum transmission rate and ξ controls the strength of epistasis (Fig. 1b).

Preliminary analysis revealed that other functional forms with qualitatively similar properties produce

results consistent with those presented below. When $\xi = 0$, there is no epistasis as $\beta_i(\xi) = \tilde{\beta}$ for all

variants. When $0 < \xi < 1$, epistasis reduces the transmission rate for variants intermediate between

1 and n, reaching a minimum of $\beta_i(\xi) = 1 - \xi$, with $\beta_1(\xi) = \beta_n(\xi) = \tilde{\beta}$ for all ξ (Fig. 1b).

Healthy and immunocompromised individuals are identical in our model except for their infectious

periods and rates of within-host antigenic evolution. The infectious period for immunocompromised

individuals, $1/\gamma_C = 140$ days, is assumed to be 20 times longer than that for healthy individuals,

$1/\gamma_H = 7$ days. These values are chosen to be illustrative and reasonable parameter variation does

not qualitatively affect our results. The rate of within-host antigenic evolution (i.e. the per-capita

transition rate between adjacent variants in the antigenic space) is governed by parameters μ_H and

μ_C in healthy and immunocompromised individuals, respectively. We assume that the virus mutates

at a constant rate, leading to a constant rate of antigenic evolution for a given host type. While our

primary model implicitly captures a simplified version of within-host evolutionary dynamics by

assuming a constant rate of antigenic evolution per host type, we justify this assumption by exploring

a separate within-host only model in the Appendix, which demonstrates that a constant rate of

antigenic evolution is a reasonable approximation. In our primary model, we either set $\mu_H = \mu_C$ so

that the rate of within-host antigenic evolution is the same in healthy and immunocompromised hosts,

or set $\mu_H < \mu_C$ to investigate the impact of a faster rate of within-host antigenic evolution in

immunocompromised individuals. The rate of antigenic evolution may differ between host types due

to differences in the viral population size within a host or the strength of selection for immune escape.

To allow for random mutations, we simulate our model using the stochastic τ-leaping method [27] for

the underlying ordinary differential equations (ODEs)

$$\frac{dS_k^i}{dt} = -\sum_{j=1}^{n} \beta_j(\xi) S_k^i \sigma_{ij} \left(I_H^j + I_C^j \right),\tag{3}$$

$$\frac{dI_k^i}{dt} = \beta_i(\xi) S_k^i \left(I_H^i + I_C^i \right) - \left(\gamma_k + \frac{\mu_k}{1 + \delta_i} \right) I_k^i + \frac{1}{2} \mu_k M_k^i,\tag{4}$$

where M_k^i is the set of variants adjacent to i in the one-dimensional antigenic space ($M_k^i = I_k^{i-1} +$

I_k^{i+1} for $i \in \{2, \dots, n-1\}$, with boundary conditions $M_k^1 = I_k^2$ and $M_k^n = I_k^{n-1}$), and $\delta_i = 1$ if $i \in \{1, n\}$

and is 0 otherwise to control the rate of antigenic evolution at the boundaries. A schematic for this

system can be found in Fig. 1c.

We run 10 simulations for each parameter combination up to $t_{max} = 1460$ time steps (days), as

preliminary analysis revealed that either antigenic evolution reaches the boundary of antigenic space

within this timeframe, or the infection is driven extinct. Note however, that this duration is arbitrary

and varies inversely with μ_C and μ_H. We say that a variant is 'observed' if it exceeds a threshold of

0.01. In each simulation, we summarise the dynamics by measuring the total number of variants

observed and the maximum distance in antigenic space between observed variants.

Results

We wish to establish when antigenic evolution proceeds as a gradual diffusion through antigenic space, or in large jumps. We therefore focus our analysis on the strength of epistasis on transmissibility ξ, the strength of cross-immunity η, the proportion of the population that is immunocompromised p, the relative rate of adaptation (antigenic evolution) in immunocompromised hosts μ_C/μ_H and the relative infectious period γ_H/γ_C. In the absence of epistasis (or when epistasis is sufficiently weak), the virus diffuses gradually through antigenic space (Figs. 2a and 2c). As the host population accumulates immunity to the current dominant variant, selection favours the next variant in line that can substantially escape immunity, leading to successive epidemic waves at regular intervals. This occurs regardless of whether within-host evolution is assumed to be faster in immunocompromised individuals (Fig. 2e).

Figure 2: Antigenic evolution with or without immunocompromised individuals and epistasis. (a) No epistasis in an entirely immunocompetent population ($p = 0, \xi = 0$). (b) Strong epistasis in an entirely immunocompetent population ($p = 0, \xi = 0.8$). (c) No epistasis and a small immunocompromised subpopulation ($p = 0.05, \xi = 0$). (d) Strong epistasis and a small

When epistasis is sufficiently strong, however, the proportion of the population that is

immunocompromised plays a crucial role in antigenic evolution (Figs. 2b, d, f). If very few individuals

are immunocompromised, the epidemic quickly burns out with little antigenic evolution, as the virus

is unable to cross the fitness valley caused by epistasis at the between-host level (Fig. 2b). But if a

sufficient proportion of the population is immunocompromised, then the virus can cross this fitness

valley due to within-host evolution in this subpopulation (Fig. 2d). Immunocompromised hosts

experience longer infections, on average, which allows the virus to accumulate mutations and cross

the fitness valley. When the virus has acquired enough mutations in the immunocompromised such

that between-host transmissibility is restored to a sufficiently high level, it is able to spread in the rest

of the host population. Again, this process is sped up if the within-host evolutionary dynamics are

assumed to be faster in immunocompromised individuals, but the qualitative dynamics are unchanged

(Fig. 2f).

Our results are qualitatively robust to variation in key model parameters, although our sensitivity

analysis reveals two notable interactions (**Figure**). When varying the strength of epistasis and the

extent of cross-immunity between variants, we find that, intuitively, immunocompromised individuals

are especially important for traversing the fitness valley if epistasis is stronger or if cross-immunity is

broader (**Figure** a). This is because stronger epistasis makes the fitness valley deeper and broader

cross-immunity reduces the pool of susceptible hosts across a wider range of variants. However, if

epistasis is sufficiently strong (around $\xi = 0.9$ in **Figure** a) a large jump in antigenic space to a distant

variant occurs regardless of the strength of cross-immunity. Our sensitivity analysis also reveals that

as the proportion of the population that is immunocompromised decreases, a jump in antigenic space

becomes less likely and requires a longer relative infectious period in immunocompromised hosts (Fig.

3b). This suggests that better treatment of immunocompromised hosts (to reduce the average

duration of infection), improved genomic surveillance of these hosts (to identify novel variants of

concern), and better prevention and treatment of pre-existing conditions (to reduce the proportion

of the population that is immunocompromised) may greatly reduce the likelihood of new variants

emerging at distant fitness peaks.

Figure 3: Sensitivity analysis. Top row: maximum distance between observed variants (darker shading indicates larger jumps

in antigenic space); bottom row: total number of variants observed. (a) Varying the strength of cross-immunity (η) and

epistasis (ξ) when 5% of the population is immunocompromised (p=0.05). (b) Varying the percentage of the population that

is immunocompromised and the relative recovery periods (with $\eta = 5$ and $\xi = 0.8$). All other parameters as in Table A3. All

datapoints are averaged over 10 simulations.

Discussion

The presence of immunocompromised individuals has been suggested as an important driver behind

not only the emergence of the Omicron variant of SARS-CoV-2, but also other variants of concern,

including Alpha and Delta [20]. Using a simple model of antigenic evolution, we have shown that

prolonged infections of immunocompromised individuals allow pathogens to accumulate sufficient

mutations to overcome epistasis at the between-host level, facilitating the emergence of novel

immune escape variants. Our model was motivated by the sudden emergence of the Omicron variant

from a distant clade to the dominant variant at the time, coupled with longitudinal sequencing from

a immunocompromised patients that indicate rapid within-host evolution [18–20]. Given relatively

high levels of infection (and hence mutation supply; Fig. 4) combined with rapidly increasing immune

pressure in mid- to late-2021, conditions for the Delta lineage to exhibit antigenic evolution seemed

to be favourable. Mutation supply or lack of immune pressure therefore do not appear to have been

the fundamental constraint for the lack of antigenic evolution by the Delta variant, which suggests

that either epistasis or transiently-boosted innate immunity constrained immune escape. Indeed, our

model suggests that novel immune escape variants readily evolve when epistasis is relatively weak.

When epistasis is stronger, reducing transmissibility for variants between fitness peaks, we find that

immunocompromised individuals play a key role in antigenic evolution, effectively allowing the

pathogen to traverse the fitness valley to reach a new peak. Note that while faster within-host

adaptation in immunocompromised individuals speeds up the rate of antigenic evolution, unlike

epistasis it does not qualitatively affect the outcome. Crucially, our results also suggest that improving

treatment for those who are immunocompromised can greatly reduce the likelihood of new variants

emerging.

Figure 4: Phylogenetic tree for SARS-CoV-2 variants shortly after the emergence of Omicron. Three variants of concern

(Alpha, Delta and Omicron) are highlighted to illustrate that there had been high mutation supply for the Delta variant.

Data downloaded from Nextstrain (nextstrain.org) on 08/02/2022 [18,28] and provided by the Global Initiative for Sharing

All Influenza Data (GISAID, gisaid.org) [29–31]. Data plotted using the ggtree software package in R [32–34].

In real populations, individuals cannot simply be classed as either healthy or immunocompromised;

they vary in the extent to which they are able to mount an immune response due to age,

comorbidities, genetic or other environmental factors. Fitness landscapes for the pathogen may also

differ between individuals and populations and real antigenic space is likely to be far more complex

than our simplified one-dimensional space (note that multi-dimensional antigenic evolution can lead

to more varied behaviour including branching and coalescence of phylogenetic pathways; see for

example [35] and [36]). While our model does not capture the full complexity of antigenic evolution

in real populations, it has important implications for our understanding of future immune escape

variants of SARS-Cov-2, and for pathogen evolution more generally. Crucially, our model suggests that

the lack of antigenic evolution by Delta followed by the emergence of Omicron is consistent with

epistasis constraining immune escape, but epistasis may be overcome when immunocompromised

individuals are infected for sufficiently long periods. Hence, rather than simply accelerating antigenic

evolution, prolonged infections of immunocompromised individuals may be critical for the evolution

of immune escape variants. We tentatively speculate that it may be difficult for SARS-CoV-2 to evolve

antigenically through incremental mutations, and future variants may require multiple (epistatic)

mutations to substantially escape host immunity. If true, a possible implication is that vaccines against

SARS-CoV-2 may not need to be regularly updated in a similar manner to seasonal influenza vaccines

as antigenic evolution is more constrained, but if new variants do emerge, they may substantially

escape prior immunity and be harder to predict. However, it is also possible that Delta was unusual in

being limited in its scope for antigenic evolution and that other variants may not experience similar

constraints, or that Delta may have eventually exhibited antigenic evolution if Omicron had not

emerged.

Our results agree with previous models which suggest that immunocompromised individuals are more

likely to facilitate or accelerate within-host pathogen evolution, for example due to a longer average

duration of infection or higher viral load [37,38]. However, while we find immunocompromised hosts

to play a crucial role in pathogen evolution at the population level, other studies have concluded the

opposite as these individuals only make up a small proportion of infections [37,38]. The reason for this

discrepancy is likely due to contrasting assumptions regarding within-host fitness, immunity, and traits

under selection. For example, van Egeren et al. [38] assume that a fitness valley exists at the within-

host level with two or three mutations required to cross, whereas our model assumes that the fitness

valley only exists at the between-host level (transmission) but may require many more mutations to

traverse. If a fitness valley exists at the within-host level, then intuitively the importance of

immunocompromised individuals for pathogen evolution will be lower. Their model also focused on a

static measure of relative fitness and did not consider antigenic evolution explicitly, whereas in our

model the fitness of a particular variant depends on the level of immunity in the population, and so

will vary over the course of the epidemic. Nevertheless, both models concur that longer duration

infections, especially those of immunocompromised individuals, can play a disproportionate role in

the evolution of novel variants, and are of particular concern for SARS-CoV-2 evolution.

We assumed that the rate of antigenic evolution during an infection was constant (but may vary by

host type), which was motivated by the separate within-host model discussed in the Appendix. For

immunocompetent hosts, who typically clear infection within two weeks [39], this means that there

is relatively little time for new variants to emerge for onwards transmission, which slows down

adaptation and can prevent epistatic mutations accumulating. But for immunocompromised hosts,

who may experience much longer infections (upwards of 150 days [20]), the coevolutionary dynamics

between the virus and the host immune system could allow many (potentially epistatic) mutations to

accumulate. Interestingly, this hypothesis is consistent with previous theoretical [40] and

experimental [41–43] studies showing that coevolution can both accelerate adaptation and allow a

pathogen to cross fitness valleys caused by epistasis.

In this study we have focused on evolution in immunocompromised individuals as a source of immune

escape variants such as Omicron, but two alternative explanations must also be considered [12]. The

first is that Omicron evolved early in the pandemic in a remote population (e.g., in southern Africa)

before eventually taking off globally in late 2021. However, the combination of a large number of

mutations in the spike protein and strong selection for immune escape in the wider population

renders this explanation unsatisfactory, as it does not explain why these mutations did not appear in

places where mutation supply and immune pressure were both high, such as the UK in late 2021. The

second is that Omicron evolved in an animal host following transmission from humans, before crossing

the species barrier into humans once again. This requires crossing the species barrier twice and

selection to favour mutations that are beneficial in both the animal species and humans. We believe

it is more plausible that Omicron evolved in an immunocompromised individual, especially since this

is consistent with longitudinal sequencing of within-host evolution [20,21].

While our model is informative, it does not capture the true complexity of antigenic space, the impact

of vaccinations and non-pharmaceutical interventions, variation in disease outcomes, and the

evolution of other disease characteristics such as transmissibility and virulence. This is by design so

that our model requires as few assumptions as possible and so that the model can be adapted for

other pathogens in future. We did not attempt to capture these effects, as our results are intended to

be illustrative of the key roles that epistasis and immunocompromised individuals may play in the

antigenic evolution of SARS-CoV-2 (and other pathogens). Modelling of immunocompromised

individuals during the COVID-19 pandemic has largely focused on their increased risk of mortality,

rather than their potential importance for pathogen evolution [44–46]. Our study emphasises the

need to consider both aspects.

We stress that while our results suggest that infected immunocompromised individuals may play a

significant role in the antigenic evolution of SARS-CoV-2, we urge caution in how this message is

interpreted and communicated. We urge particular caution with regards to the implications of our

results for people who are immunocompromised. People may be immunocompromised for a variety

of reasons, including uncontrolled HIV, undergoing treatment for cancer, or as a transplant recipient,

and some conditions still wrongly attract stigma. Although Omicron was first detected in South Africa,

which is estimated to have the highest HIV prevalence in the world (7.7 million people, with many

infections uncontrolled [47]), this variant may have evolved in an individual without HIV and may have

evolved elsewhere. Rather than stigmatising people who are immunocompromised, our results

emphasise the need for global health equality and for better genomic surveillance, especially for

immunocompromised people infected with SARS-CoV-2. Improving access to vaccines and

treatments, especially in lower- and middle-income countries, and facilitating wider surveillance for

new variants is crucial for limiting the emergence of new variants in the COVID-19 pandemic.

Appendix

Within-host model

The model in the main text focuses on population-level dynamics and implicitly models within-host

dynamics by assuming that: (1) immunocompetent and immunocompromised hosts differ in terms of

their average infectious period; and (2) antigenic evolution occurs at a constant rate. Here, we

consider the dynamics of a simple within-host model to justify the implicit within-host dynamics in

our population-level model.

Let V_i be the viral abundance of variant $i \in \{1, \dots, n\}$ within a single infected host and let R_i be the

strength of the corresponding immune response. The virus grows exponentially with rate r in the

absence of an immune response and decreases through the immune response at rate $\kappa \sum_{j=1}^{n} \tilde{\sigma}_{ij} R_j$,

where κ is the per-capita rate of virus removal by the host immune system and $\tilde{\sigma}_{ij}$ is the probability

that an immune response for variant j causes cross-immunity to variant i such that

$$\tilde{\sigma}_{ij} = \exp\left\{-\frac{(i-j)^2}{2\tilde{\eta}}\right\} \tag{A1}$$

where $\tilde{\eta}$ controls the breadth of cross-immunity between variants (similar to η in the main text). The

virus also mutates to adjacent variants in the antigenic space with rate $\tilde{\mu}$. The immune response to

variant i increases at per-capita rate $\kappa q V_i$ and decays with rate d. The parameter q controls the

strength of host immune system such that larger values indicate an immune system that can respond

well to infection (immunocompetent) and smaller values indicate a weaker immune response

(immunocompromised).

As with the between-host model, we use the stochastic τ-leaping method [27] to simulate the within-

host dynamics, corresponding to the following set of ODEs

$$\frac{dV_i}{dt} = rV_i - \kappa V_i \sum_{j=1}^{n} \tilde{\sigma}_{ij} R_j - \frac{\tilde{\mu} V_i}{1 + \delta_i} + \frac{1}{2} \tilde{\mu} \widetilde{M}_i \tag{A2}$$

$$\frac{dR_i}{dt} = (\kappa q V_i - d) R_i \tag{A3}$$

where $\widetilde{M}_i$ is the set of variants adjacent to i in the one-dimensional antigenic space ($\widetilde{M}_i = V_{i-1} + V_{i+1}$

for $i \in \{2, \dots, n-1\}$, with boundary conditions $\widetilde{M}_1 = V_2$ and $\widetilde{M}_n = V_{n-1}$), and $\delta_i = 1$ if $i \in \{1, n\}$ and

is 0 otherwise to control the mutation rate at the boundaries.

When the host is immunocompetent (large q), the infection is rapidly cleared, with little within-host

evolution (Fig. A1a). But when the host is immunocompromised (small q), the infection persists over

much longer timescales, with immune pressure leading to successive selective sweeps as the virus

diffuses through the antigenic space at a constant rate (Fig. A1b). If the mutation rate is faster in

immunocompromised hosts (larger $\tilde{\mu}$), the coevolutionary dynamics of the virus and the immune

response are simply accelerated (Fig. A1c compared with Fig. A1b). These results justify the simplifying

assumptions in our population-level model regarding within-host dynamics, where we assume that

there is a constant rate of antigenic evolution, which may differ between host types.

Figure A1: Within-host dynamics. *(a) Immunocompetent host ($q = 1.0$), (b) Immunocompromised host ($q = 0.1$), (c)*

immunocompromised host with faster mutation $q = 0.1, \tilde{\mu} = 0.02$). Values of the viral load are normalised by the maximum

value attained. All other parameters as in Table A4. Dynamics are shown for a single simulation.

Simulation algorithm

We simulate our within-host and population-level models using the τ-leaping method [27], which is

an approximate stochastic simulation algorithm. We define the propensity functions α_E^i in Table A1,

which give the rates of event type E for each variant index i. These propensity functions are then used

to update the system synchronously at a time interval of one day using random numbers from the

Poisson distribution $P(\alpha_E^i)$. Source code for the simulations is available in the Supplementary Material

and at https://github.com/ecoevotheory/Smith_and_Ashby_2022.

Event	Rate
Infection of host type $k \in \{H, C\}$ by variant i	$\alpha^i_{\text{Inf}, k} = \beta_i S^i_k (I^i_H + I^i_S)$
Gain of full immunity to variant i due to infection by variant j (host type $k \in \{H, C\}$)	$\alpha^i_{\text{Imm}, j, k} = \beta_j \sigma_{ij} S^i_k (I^j_H + I^j_S)$
Recovery by host type $k \in \{H, C\}$ from variant i	$\alpha^i_{\text{Rec}, k} = \gamma_k I^i_k$
Mutation from variant i to variant $i \pm 1$ for host type $k \in \{H, C\}$.	$\alpha^i_{\text{Mut}, k} = \dfrac{\mu_k}{1 + \delta_i} I^i_k$

423   *Table A1: The propensity functions for each of the event types in the population-level model.*

Event	Rate
Growth of variant i	$\alpha^i_{\text{Grow}} = r V_i$
Removal of variant i by the corresponding immune response	$\alpha^i_{\text{Rem}} = \kappa V_i R_i$
Removal of variant i by immune response with index j	$\alpha^i_{\text{Cross-rem}, j} = \kappa \tilde{\sigma}_{ij} V_i R_j$
Mutation from variant i to variant $i \pm 1$	$\alpha^i_{\text{Mut},} = \dfrac{\tilde{\mu}}{1 + \delta_i} V_i$
Decay of immune response for variant i	$\alpha^i_{\text{Dec}} = d R_i$

424   *Table A2: The propensity functions for the various event types for the within-host model.*

Parameter	Description	Value
p	Proportion immunocompromised	0.05
ξ	Strength of epistasis (proportional reduction in transmission)	0.8
N	Population size	10^7
n	Number of variants	30
μ_k	Mutation rate for $k \in \{H, C\}$	0.01 per day
R_0	Basic reproductive number	3.0
γ_H	Recovery rate (immunocompetent)	1/7 per day
γ_C	Recovery rate (immunocompromised)	1/140 per day
t_{max}	Final time	1460 days
η	Strength of cross-immunity	10.0

426   *Table A3: Default parameters for the between-host model.*

Parameter	Description	Value
n	Number of variants	30
$\tilde{\mu}$	Mutation rate	5×10^{-3} per day
r	Viral growth rate	1.0 per day
κ	Viral clearance rate by immune response	2×10^{-2} per unit viral load per day
q	Relative strength of host immune system	0.1
t_{max}	Final time	150 days
d	Decay rate of immune response	10^{-4} per day
$\tilde{\eta}$	Strength of cross-immunity	0.1

428   *Table A4: Default parameters for the within-host model.*

Acknowledgements

We thank Angus Buckling for helpful discussions. CAS is funded by the Natural Environment Research Council (NE/V003909/1). BA is funded by the Natural Environment Research Council (NE/N014979/1 and NE/V003909/1).

Author contributions

BA conceived the study, CAS carried out the analysis, and both co-authored the manuscript.

Data accessibility

Source code for the simulations is available in the Supplementary Material and at:

https://github.com/ecoevotheory/Smith_and_Ashby_2022

References

1. Davies NG, Abbott S, Barnard RC *et al.* Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science (1979)* 2021;**372**, DOI: 10.1126/science.abg3055.

2. Hagen A. How Dangerous is the Delta Variant. *American Society for Microbiology* 2021.

3. SPI-M. *SPI-M-O: Consensus Statement on COVID-19.*, 2021.

4. Nexstrain. Genomic epidemiology of novel coronavirus - Global subsampling. 2021.

5. Cele S, Gazy I, Jackson L *et al.* Escape of SARS-CoV-2 501Y.V2 from neutralization by convalescent plasma. *Nature* 2021;**593**:142–6.

6. Ferguson N, Ghani A, Cori A *et al. Report 49: Growth, Population Distribution and Immune Escape of Omicron in England.*, 2021.

7. Cao Y, Wang J, Jian F *et al.* Omicron escapes the majority of existing SARS-CoV-2 neutralizing antibodies. *Nature* 2021, DOI: 10.1038/d41586-021-03796-6.

8. Lyngse FP, Mortensen LH, Denwood MJ *et al.* SARS-CoV-2 Omicron VOC Transmission in Danish Households. *medRxiv* 2021:2021.12.27.21268278.

9. Sofonea MT, Roquebert B, Foulongne V *et al.* From Delta to Omicron: analysing the SARS-CoV-2
epidemic in France using variant-specific screening tests (September 1 to December 18, 2021).
medRxiv 2022:2021.12.31.21268583.

10. Day T, Gandon S, Lion S *et al.* On the evolutionary epidemiology of SARS-CoV-2. *Current Biology*
2020;**30**:R849–57.

11. Viana R, Moyo S, Amoako DG *et al.* Rapid epidemic expansion of the SARS-CoV-2 Omicron variant
in southern Africa. *medRxiv* 2021:2021.12.19.21268028.

12. Kupferschmidt K. Where did "weird" omicron come from? 2021.

13. Grenfell BT, Pybus OG, Gog JR *et al.* Unifying the Epidemiological and Evolutionary Dynamics of
Pathogens. *Science (1979)* 2004;**303**:327–32.

14. Saad-Roy CM, Morris SE, Metcalf CJE *et al.* Epidemiological and evolutionary considerations of
SARS-CoV-2 vaccine dosing regimes. *Science (1979)* 2021;**372**:363–70.

15. Ashby B, Thompson RN. Non-pharmaceutical interventions and the emergence of pathogen
variants. *Medrxiv* 2021, DOI: 10.1101/2021.05.27.21257938.

16. Ritchie H, Mathieu E, Rodés-Guirao L *et al.* Coronovirus Pandemic (COVID-19). *Published online at*
OurWorldInData.org 2020.

17. Andrews N, Stowe J, Kirsebom F *et al.* Effectiveness of COVID-19 vaccines against the Omicron
(B.1.1.529) variant of concern. *medRxiv* 2021:2021.12.14.21267615.

18. Hadfield J, Megill C, Bell SM *et al.* Nextstrain: real-time tracking of pathogen evolution.
Bioinformatics 2018;**34**:4121–3.

19. Marchi J, Lässig M, Walczak AM *et al.* Antigenic waves of virus–immune coevolution. *Proceedings*
of the National Academy of Sciences 2021;**118**:e2103398118.

20. Corey L, Beyrer C, Cohen MS *et al.* SARS-CoV-2 Variants in Patients with Immunosuppression.
New England Journal of Medicine 2021;**385**:562–6.

21. Borges V, Isidro J, Cunha M *et al.* Long-Term Evolution of SARS-CoV-2 in an Immunocompromised
Patient with Non-Hodgkin Lymphoma. *mSphere* 2021;**6**, DOI: 10.1128/msphere.00244-21.

22. Weigang S, Fuchs J, Zimmer G *et al.* Within-host evolution of SARS-CoV-2 in an
immunosuppressed COVID-19 patient as a source of immune escape variants. *Nature*
Communications 2021;**12**:6405.

23. Kemp SA, Collier DA, Datir RP *et al.* SARS-CoV-2 evolution during treatment of chronic infection.
Nature 2021;**592**:277–82.

24. Álvarez H, Ruiz-Mateos E, Juiz-González PM *et al.* SARS-CoV-2 Evolution and Spike-Specific CD4+
T-Cell Response in Persistent COVID-19 with Severe HIV Immune Suppression. *Microorganisms*
2022;**10**:143.

25. Martin DP, Lytras S, Lucaci AG *et al.* Selection analysis identifies unusual clustered mutational
changes in Omicron lineage BA.1 that likely impact Spike function. *bioRxiv* 2022, DOI:
10.1101/2022.01.14.476382.

26. Gog JR, Grenfell BT. Dynamics and selection of many-strain pathogens. *Proceedings of the*
National Academy of Sciences 2002;**99**.

27. Gillespie DT. Approximate accelerated stochastic simulation of chemically reacting systems. *The*
Journal of Chemical Physics 2001;**115**:1716–33.

28. Sagulenko P, Puller V, Neher RA. TreeTime: Maximum-likelihood phylodynamic analysis. *Virus*
Evolution 2018;**4**, DOI: 10.1093/ve/vex042.

29. Khare S, Gurry C, Freitas L *et al.* GISAID's Role in Pandemic Response. *China CDC Weekly*
2021;**3**:1049–51.

30. Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative contribution to
global health. *Global Challenges* 2017;**1**:33–46.

31. Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data – from vision to reality.
Eurosurveillance 2017;**22**, DOI: 10.2807/1560-7917.ES.2017.22.13.30494.

32. Yu G. Using ggtree to Visualize Data on Tree-Like Structures. *Current Protocols in Bioinformatics*
2020;**69**, DOI: 10.1002/cpbi.96.

33. Yu G, Lam TT-Y, Zhu H *et al.* Two Methods for Mapping and Visualizing Associated Data on
Phylogeny Using *Ggtree*. *Molecular Biology and Evolution* 2018;**35**:3041–3.

34. Yu G, Smith DK, Zhu H *et al.* ggtree: an R package for visualization and annotation of phylogenetic
trees with their covariates and other associated data. *Methods in Ecology and Evolution* 2017;**8**:28–
36.

35. Yan L, Neher RA, Shraiman BI. Phylodynamic theory of persistence, extinction and speciation of
rapidly adapting pathogens. *Elife* 2019;**8**, DOI: 10.7554/eLife.44205.

36. Zinder D, Bedford T, Gupta S *et al.* The Roles of Competition and Mutation in Shaping Antigenic
and Genetic Diversity in Influenza. *PLoS Pathogens* 2013;**9**:e1003104.

37. Eden JS, Chisholm RH, Bull RA *et al.* Persistent infections in immunocompromised hosts are
rarely sources of new pathogen variants. *Virus Evolution* 2017;**3**, DOI: 10.1093/ve/vex018.

38. van Egeren D, Novokhodko A, Stoddard M *et al.* Controlling long-term SARS-CoV-2 infections is
important for slowing viral evolution. 2021, DOI: 10.1101/2021.04.10.21255251.

39. Kissler SM, Fauver JR, Mack C *et al.* Viral dynamics of acute SARS-CoV-2 infection and
applications to diagnostic and public health strategies. *PLoS Biology* 2021;**19**, DOI:
10.1371/journal.pbio.3001333.

40. Ashby B, Gupta S, Buckling A. Effects of epistasis on infectivity range during host-parasite
coevolution. *Evolution (N Y)* 2014;**68**:2972–82.

41. Paterson S, Vogwill T, Buckling A *et al.* Antagonistic coevolution accelerates molecular evolution.
Nature 2010;**464**:275–8.

42. Hall AR, Scanlan PD, Buckling A. Bacteria-Phage Coevolution and the Emergence of Generalist
Pathogens. *The American Naturalist* 2011;**177**:44–53.

43. Meyer JR, Dobias DT, Weitz JS *et al.* Repeatability and Contingency in the Evolution of a Key
Innovation in Phage Lambda. *Science (1979)* 2012;**335**:428–32.

44. Smith CA, Yates ChristianA, Ashby B. Critical weaknesses in shielding strategies for COVID-19.
PLOS Global Public Health 2022.

45. Neufeld Z, Khataee H, Czirok A. Targeted adaptive isolation strategy for COVID-19 pandemic.
Infect Dis Model 2020;**5**, DOI: 10.1016/j.idm.2020.04.003.

46. van Bunnik BAD, Morgan ALK, Bessell PR *et al.* Segmentation and shielding of the most
vulnerable members of the population as elements of an exit strategy from COVID-19 lockdown.
Philosophical Transactions of the Royal Society B: Biological Sciences 2021;**376**, DOI:
10.1098/rstb.2020.0275.

47. Mahy M, Marsh K, Sabin K *et al.* HIV estimates through 2018. *AIDS* 2019;**33**:S203–11.

539