Effectiveness of BNT162b2 and mRNA-1273 COVID-19 boosters against SARS-CoV-2 Omicron (B.1.1.529) infection in Qatar

Laith J. Abu-Raddad, PhD[1,2,3,4*†], Hiam Chemaitelly, PhD[1,2,3†], Houssein H. Ayoub, PhD[5], Sawsan AlMukdad, MSc[1,2], Patrick Tang, MD PhD[6], Mohammad R. Hasan, PhD[6], Peter Coyle, MD[7,8,9], Hadi M. Yassine, PhD[8,10], Hebah A. Al-Khatib, PhD[8,10], Maria K. Smatti, MSc[8,10], Zaina Al-Kanaani, PhD[7], Einas Al-Kuwari, MD[7], Andrew Jeremijenko, MD[7], Anvar Hassan Kaleeckal, MSc[7], Ali Nizar Latif, MD[7], Riyazuddin Mohammad Shaik, MSc[7], Hanan F. Abdul-Rahim, PhD[4], Gheyath K. Nasrallah, PhD[8,10], Mohamed Ghaith Al-Kuwari, MD[11], Adeel A. Butt, MBBS MS[3,7,12], Hamad Eid Al-Romaihi, MD[13], Mohamed H. Al-Thani, MD[13], Abdullatif Al-Khal, MD[7], and Roberto Bertollini, MD MPH[13]

[1]Infectious Disease Epidemiology Group, Weill Cornell Medicine-Qatar, Cornell University, Doha, Qatar
[2]World Health Organization Collaborating Centre for Disease Epidemiology Analytics on HIV/AIDS, Sexually Transmitted Infections, and Viral Hepatitis, Weill Cornell Medicine–Qatar, Cornell University, Qatar Foundation – Education City, Doha, Qatar
[3]Department of Population Health Sciences, Weill Cornell Medicine, Cornell University, New York, New York, USA
[4]Department of Public Health, College of Health Sciences, QU Health, Qatar University, Doha, Qatar
[5]Mathematics Program, Department of Mathematics, Statistics, and Physics, College of Arts and Sciences, Qatar University, Doha, Qatar
[6]Department of Pathology, Sidra Medicine, Doha, Qatar
[7]Hamad Medical Corporation, Doha, Qatar
[8]Biomedical Research Center, Member of QU Health, Qatar University, Doha, Qatar
[9]Wellcome-Wolfson Institute for Experimental Medicine, Queens University, Belfast, United Kingdom
[10]Department of Biomedical Science, College of Health Sciences, Member of QU Health, Qatar University, Doha, Qatar
[11]Primary Health Care Corporation, Doha, Qatar
[12]Department of Medicine, Weill Cornell Medicine, Cornell University, New York, New York, USA
[13]Ministry of Public Health, Doha, Qatar

[†]Both authors contributed equally.

Word count: Abstract: 268 words, Main Text: 3,456 words.
Number of tables: 3.
Number of figures: 3.
Running head: Effectiveness of mRNA boosters against Omicron.
Keywords: COVID-19; vaccine; booster; Omicron; breakthrough infection; cohort study; immunity; epidemiology.

[*]Correspondence to Professor Laith J. Abu-Raddad, E-mail: lja2002@qatar-med.cornell.edu.

Abstract

BACKGROUND: Waning of COVID-19 vaccine protection and emergence of SARS-CoV-2 Omicron (B.1.1.529) variant have expedited efforts to scale up booster vaccination. This study compared protection afforded by booster doses of the BNT162b2 (Pfizer-BioNTech) and mRNA-1273 (Moderna) vaccines, compared to the primary series of only two doses in Qatar, during a large, rapidly growing Omicron wave.

METHODS: In a population of 2,232,224 vaccinated persons with at least two doses, two matched, retrospective cohort studies were implemented to investigate effectiveness of booster vaccination against symptomatic SARS-CoV-2 infection and against COVID-19 hospitalization and death, up to January 9, 2022. Association of booster status with infection was estimated using Cox proportional-hazards regression models.

RESULTS: For BNT162b2, cumulative symptomatic infection incidence was 2.9% (95% CI: 2.8-3.1%) in the booster-dose cohort and 5.5% (95% CI: 5.3-5.7%) in the primary-series cohort, after 49 days of follow-up. Adjusted hazard ratio for symptomatic infection was 0.50 (95% CI: 0.47-0.53). Booster effectiveness relative to primary series was 50.1% (95% CI: 47.3-52.8%). For mRNA-1273, cumulative symptomatic infection incidence was 1.9% (95% CI: 1.7-2.2%) in the booster-dose cohort and 3.5% (95% CI: 3.2-3.9%) in the primary-series cohort, after 35 days of follow-up. The adjusted hazard ratio for symptomatic infection was 0.49 (95% CI: 0.43-0.57). Booster effectiveness relative to primary series was 50.8% (95% CI: 43.4-57.3%). There were fewer cases of severe COVID-19 in booster-dose cohorts than in primary-series cohorts, but cases of severe COVID-19 were rare in all cohorts.

CONCLUSIONS: mRNA booster vaccination is associated with modest effectiveness against symptomatic infection with Omicron. The development of a new generation of vaccines targeting a broad range of variants may be warranted.

Introduction

Waning of coronavirus disease 2019 (COVID-19) vaccine protection[1-6] and emergence of the immune-evasive severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) Omicron[7] (B.1.1.529) variant of concern[8-10] have stimulated efforts to scale-up COVID-19 booster vaccination. Qatar launched its third dose/booster vaccination program in mid-September 2021, using both the BNT162b2[11] (Pfizer-BioNTech) and mRNA-1273[12] (Moderna) mRNA vaccines, the same two vaccines that have been used since the launch of the COVID-19 immunization program in this country.[13-15]

Initially, booster eligibility was restricted to the elderly, immunocompromised persons, and persons with severe or multiple chronic conditions, but then expanded by age group to the rest of the population. Eligibility was first restricted to persons who had completed the primary series of two doses at least 8 months earlier, but subsequently this interval was shortened to 6 months. The same vaccine was used in both primary series and booster vaccinations for the majority of the population, but for mRNA-1273, the booster dose was half the dose used in the primary series. As booster vaccination was being scaled up, the Omicron variant was introduced into the population, leading to the largest SARS-CoV-2 epidemic wave that the country has experienced since the start of the pandemic.

Leveraging the integrated national COVID-19 datasets, effectiveness of booster vaccination against SARS-CoV-2 symptomatic infection and against COVID-19 hospitalization and death, relative to that of the primary series of two doses, was investigated in the national cohort of vaccinated individuals in Qatar, during the rapidly growing Omicron wave.[10]

Methods

Data sources and study design

The study analyzed the national, federated databases for COVID-19 vaccination, laboratory testing, hospitalization, and death, retrieved from the integrated nationwide digital-health information platform. Databases include all SARS-CoV-2-related data and associated demographic information, with no missing information, since pandemic onset. These include all polymerase chain reaction (PCR) tests, vaccination records, COVID-19 hospitalizations, infection severity and mortality classifications per World Health Organization (WHO) guidelines,[16,17] in addition to sex, age, and nationality information retrieved from the national registry. Further description of these national databases can be found in previous studies.[1,13,14,18-24]

Effectiveness of booster vaccination relative to the primary series was estimated for both the BNT162b2 and mRNA-1273 vaccines using a matched, retrospective cohort study design that emulated a target trial.[25,26] The study compared incidence of SARS-CoV-2 symptomatic breakthrough infection in the cohort of individuals who completed >7 days after the booster dose (booster-dose cohort) to incidence in the cohort of individuals who have not yet received their booster dose (primary-series cohort). The 7-day cutoff between administration of the booster and start of follow-up was informed by earlier studies[26-29] to ensure sufficient time for build-up of booster protection. A 14-day cutoff was also investigated in a sensitivity analysis.

For BNT162b2, all individuals with at least two doses between January 5, 2021 (date of first second-dose BNT162b2 vaccination in Qatar) and January 9, 2022 (end of study) were eligible for inclusion in the study, provided that they had no record of a prior PCR-confirmed infection before the start of follow-up. The same applied to mRNA-1273, but the corresponding dates were January 24, 2021 and January 9, 2022, respectively.

Individuals in the booster-dose cohort were exact-matched in a 1:1 ratio by sex, 10-year age group, and nationality to individuals in the primary-series cohort, to control for known differences in the risk of exposure to SARS-CoV-2 infection in Qatar.[18,21,30-32] Matching by these factors was shown in an earlier study of similar design to provide adequate control of bias arising from differences in infection exposure in Qatar.[3] Individuals were also exact-matched by the calendar week of second vaccine dose to control for time since vaccination and waning of vaccine immunity over time.[1,3,6,15]

Matching was performed through an iterative process that ensured that each control in the primary-series cohort is alive, infection-free, and has not received the third dose at the start of follow-up, which was defined, for each matched pair, at the 8th day after the individual in the booster-dose cohort received the booster dose. Controls in the primary-series cohort who received the booster dose at a future date were eligible for recruitment into the booster-dose cohort, provided that they were alive and infection-free at the start of follow-up.

Both members of each matched pair were censored at the event of the control receiving the booster dose, to ensure exchangeability.[26] Accordingly, individuals were followed up until the first of one of the following events: a documented SARS-CoV-2 infection (defined as the first PCR-positive test after the start of follow-up regardless of presence of symptoms or reason for PCR testing), or booster-dose vaccination of the control (with matched pair censoring), death, or end of study censoring (January 9, 2022).

Investigated primary outcome was symptomatic infection defined as a PCR-positive nasopharyngeal swab conducted because of clinical suspicion due to presence of symptoms compatible with a respiratory tract infection. Investigated secondary outcome was any severe,[16] critical,[16] or fatal[17] COVID-19. Classification of COVID-19 case severity (acute-care

hospitalizations),[16] criticality (ICU hospitalizations),[16] and fatality[17] followed WHO guidelines, and assessments were made by trained medical personnel using individual chart reviews. Details of the COVID-19 severity, criticality, and fatality classification are found in Section 1 in Supplementary Appendix.

Each person hospitalized for COVID-19 underwent an infection severity assessment every three days until discharge or death. We classified individuals who progressed to severe, critical, or fatal COVID-19 between the time of the PCR-positive test and the end of the study based on their worst outcome, starting with death,[17] followed by critical disease,[16] and then severe disease.[16]

Laboratory methods

Details of laboratory methods for real-time reverse-transcription PCR (RT-qPCR) testing are found in Section 2 in Supplementary Appendix.

Statistical analysis

Frequency distributions and measures of central tendency were used to describe the full and matched cohorts. Group comparison was performed using standardized mean differences (SMDs), with an SMD <0.1 indicating adequate matching.[33] Cumulative incidence of symptomatic infection was defined as the proportion of individuals at risk whose primary endpoint was a documented symptomatic infection during follow-up, and was estimated in each cohort using the Kaplan–Meier estimator method.[34] Equality of failure functions was assessed using the log-rank test. Incidence rate of symptomatic infection in each cohort, which was defined as the number of identified symptomatic infections divided by the number of person-weeks contributed by all individuals in the cohort, was estimated, along with its 95% confidence

interval (CI), using a Poisson log-likelihood regression model with the STATA 17.0[35] *stptime* command.

The hazard ratio comparing incidence of symptomatic infection in both cohorts and corresponding 95% CI were calculated using Cox regression adjusted for the matching factors with the STATA 17.0[35] *stcox* command. Shoenfeld residuals and log-log plots for the survival curves were used to test the proportional-hazards assumption and to investigate its adequacy. 95% CIs were not adjusted for multiplicity. Interactions were not considered. Vaccine effectiveness of booster vaccination relative to the primary series was estimated using the equation: $\text{Vaccine effectiveness} = 1 - \text{adjusted hazrad ratio}$.

In addition to estimating the booster-vaccine effectiveness against symptomatic infection with Omicron, an additional sub-analysis was conducted to estimate effectiveness against symptomatic infection with Delta (B.1.617.2).[7] Here, the end of the study period was December 1, 2021, that is at first detection of Omicron in Qatar.[10,36] During this specific follow-up period, Qatar was experiencing a Delta-dominated low-incidence phase.[1,13,14,23,36-38] This analysis was only possible for the BNT162b2 vaccine as the time of follow-up was limited for the mRNA-1273-vaccine cohorts.

Since optimal effectiveness of the booster dose may require more than 7 days to develop, the main analyses for the BNT162b2 and mRNA-1273 vaccines were repeated, but with the start of follow-up on the 15th day after receiving the booster dose, instead of the 8th day.

Booster-vaccine effectiveness was also calculated against any documented infection regardless of presence of symptoms or reason for PCR testing.

Statistical analyses were conducted in STATA/SE version 17.0.[35]

Oversight

Hamad Medical Corporation and Weill Cornell Medicine-Qatar Institutional Review Boards approved this retrospective study with waiver of informed consent, since analysis was done on routinely collected data. The study was reported following the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines. The STROBE checklist can be found in Table S1 of the Supplementary Appendix.

Results

Study population for BNT162b2 vaccine

Between January 5, 2021 and January 9, 2022, 1,294,138 individuals received at least two doses of BNT162b2 and 230,526 received a third, BNT162b2 booster dose. The median date at the first dose was May 3, 2021, May 24, 2021 at the second dose, and December 6, 2021 at the third dose. The median time elapsed between the first and second doses was 21 days (interquartile range (IQR), 21-22 days). The median time elapsed between the second and booster doses was 249 days (IQR, 234-267 days).

As of January 9, 2022, 88,729 and 13,948 breakthrough infections were recorded among those who received either only two or three BNT162b2 doses, respectively. Among infections after the second dose, but before the booster dose, 210 progressed to severe, 18 to critical, and 19 to fatal COVID-19. Of the infections after the booster dose, 12 progressed to severe, but none to critical or fatal COVID-19.

Figure 1 shows the population selection process for the BNT162b2-booster study. Table 1 shows baseline characteristics of the full and matched cohorts. Median age in the matched cohorts was 43 years (IQR, 35-53). The matched cohorts were balanced on the matching factors.

Effectiveness of BNT162b2 booster against Omicron

The median time of follow-up was 21 days (IQR, 8-36 days) for the booster-dose and primary-series BNT162b2 cohorts. A total of 7,210 infections, 2,035 symptomatic and 5,175 non-symptomatic at diagnosis, were recorded in the booster-dose cohort 8 days or more after receiving the booster dose (Figure 1). None of these infections progressed to severe, critical, or fatal COVID-19 as of end of this study. Although COVID-19 hospitalizations occurred among those who received a booster dose (note section above), all of these cases occurred due to infections within the first 7 days after the booster dose, after censoring for those being followed, or among those not part of the matched cohort—none of these cases were due to infections during time of follow-up of the matched cohort.

A total of 10,994 infections, 3,858 symptomatic and 7,136 non-symptomatic, were recorded in the primary-series cohort. Of these infections, 15 progressed to severe COVID-19, but none to critical or fatal COVID-19.

Cumulative incidence of symptomatic infection was estimated at 2.9% (95% CI: 2.8-3.1%) for the booster-dose cohort and at 5.5% (95% CI: 5.3-5.7%) for the primary-series cohort, 49 days after the start of follow-up (Figure 2). Infection incidence was overwhelmingly predominated by Omicron. The median start date of follow-up was December 9, 2021, after introduction of Omicron in Qatar, but a few days before onset of the large Omicron-wave exponential-growth phase on December 19, 2021.[10,36] Of all PCR-documented infections between December 1, 2021 and January 9, 2022 (end of study), that is the duration over which most time of follow-up occurred, 97.5% of infections occurred after onset of the Omicron-wave exponential-growth phase. Of 98 sequenced random specimens that were collected between December 19 and

December 31, 2021, 86 (88%) were confirmed as Omicron infections and 12 (12%) as Delta infections.[36]

Symptomatic infection incidence rate was estimated at 35.1 (95% CI: 33.6-36.6) per 10,000 person-weeks in the booster-dose cohort and at 67.2 (95% CI: 65.1-69.3) per 10,000 person-weeks in the primary-series cohort (Table 2). The hazard ratio for symptomatic infection, adjusted for sex, 10-year age group, nationality group, and calendar week of dose 2, was estimated at 0.50 (95% CI: 0.47-0.53). Effectiveness of BNT162b2 booster dose relative to the primary series of only two doses was estimated at 50.1% (95% CI: 47.3-52.8%).

The hazard ratio for any severe, critical, or fatal COVID-19 was estimated at 0.00 (95% CI: 0.00-0.29). Effectiveness of the BNT162b2 booster dose against any severe, critical, or fatal COVID-19, relative to the primary series, was estimated at 100.0% (95% CI: 71.4-100.0%).

Effectiveness of BNT162b2 booster against Delta

In the Delta-variant analysis, that is with the end of study on December 1, 2021, cumulative incidence of symptomatic infection was estimated at 0.05% (95% CI: 0.02-0.1%) for the BNT162b2 booster-dose cohort and at 0.2% (95% CI: 0.1-0.3%) for the primary-series cohort, 49 days after the start of follow-up (Figure 3). Symptomatic infection incidence rate was estimated at 0.6 (95% CI: 0.3-1.2) per 10,000 person-weeks in the booster-dose cohort and at 3.6 (95% CI: 2.6-4.8) per 10,000 person-weeks in the primary-series cohort (Table 3). The adjusted hazard ratio for symptomatic infection was estimated at 0.14 (95% CI: 0.06-0.33). Effectiveness of BNT162b2 booster dose relative to the primary series was estimated at 86.1% (95% CI: 67.3-94.1%).

Study population for mRNA-1273 vaccine

Between January 24, 2021 and January 9, 2022, 889,022 individuals received at least two doses of mRNA-1273 and 62,427 received the third, mRNA-1273 booster dose. The median date at the first dose was May 28, 2021, June 27, 2021 at the second dose, and December 22, 2021 at the third dose. The median time elapsed between the first and second doses was 28 days (IQR, 28-30 days). The median time elapsed between the second and booster doses was 228 days (IQR, 211-241 days).

As of January 9, 2022, 35,599 and 2,631 breakthrough infections were recorded among those who received either only two or three mRNA-1273 doses, respectively. Of the infections after the second dose, but before the booster dose, 17 progressed to severe, 5 to critical, and none to fatal COVID-19. Of the infections after the booster dose, 3 progressed to severe, but none to critical or fatal COVID-19.

Figure 1 shows the population selection process for the mRNA-1273-booster study. Table 1 shows the baseline characteristics of the full and matched cohorts. Median age in the matched cohorts was 40 years (IQR, 33-46). The matched cohorts were balanced on the matching factors.

Effectiveness of mRNA-1273 booster against Omicron

The median time of follow-up was 14 days (IQR, 5-26 days) for the booster-dose and primary-series mRNA-1273 cohorts. A total of 1,463 infections, 317 symptomatic and 1,146 non-symptomatic at diagnosis, were recorded in the booster-dose cohort 8 days or more after receiving the booster dose (Figure 1). None of these infections progressed to severe, critical, or fatal COVID-19 as of end of this study. A total of 1,880 infections, 589 symptomatic and 1,291 non-symptomatic, were recorded in the primary-series cohort. None of these infections progressed to severe, critical, or fatal COVID-19.

Cumulative incidence of symptomatic infection was estimated at 1.9% (95% CI: 1.7-2.2%) for the booster-dose cohort and at 3.5% (95% CI: 3.2-3.9%) for the primary-series cohort, 35 days after the start of follow-up (Figure 2). Infection incidence was overwhelmingly predominated by Omicron. The median date of start of follow-up was December 23, 2021, after onset of the Omicron-wave exponential-growth phase on December 19, 2021.[10,36]

Symptomatic infection incidence rate was estimated at 32.8 (95% CI: 29.4-36.6) per 10,000 person-weeks in the booster-dose cohort and at 60.6 (95% CI: 55.9-65.7) per 10,000 person-weeks in the primary-series cohort (Table 2). The adjusted hazard ratio for symptomatic infection was estimated at 0.49 (95% CI: 0.43-0.57). Effectiveness of the mRNA-1273 booster dose relative to the primary series was estimated at 50.8% (95% CI: 43.4-57.3%).

mRNA-1273 vaccine effectiveness against any severe, critical, or fatal COVID-19 could not be calculated because there were no recorded cases in both the booster and primary-series cohorts as of end of this study (Figure 1).

Additional analyses

For the BNT162b2-vaccine analysis, with the start of follow-up on the 15th day after the booster dose, effectiveness of the booster dose against symptomatic (Omicron) infection, relative to the primary series, was estimated at 50.3% (95% CI: 47.5-53.0%) (Figure S1 and Table S2). The corresponding effectiveness for mRNA-1273 was estimated at 50.1% (95% CI: 41.4-57.6%) (Figure S1 and Table S2). Both effectiveness estimates were similar to the main analysis estimates.

Considering any documented (Omicron) infection, regardless of presence of symptoms or reason for PCR testing, as the primary outcome, effectiveness of the booster dose relative to the primary

series was estimated at 38.2% (95% CI: 36.3-40.0%) for BNT162b2 and at 28.8% (95% CI: 23.6-33.6%) for mRNA1273.

During follow-up in the BNT162b2-vaccine analysis, 26.6% of individuals in the booster-dose cohort and 23.1% of individuals in the primary-series cohort had a PCR test done. During follow-up in the mRNA-1273-vaccine analysis, 21.3% of individuals in the booster-dose cohort and 14.6% of individuals in the primary-series cohort had a PCR test done. Those in the booster dose cohorts had higher testing rates due to higher pre-travel testing (Table S3). It appears that many individuals took the booster dose in preparation for planned travel during the year end holidays season.

Adjusting the effectiveness estimates for any documented (Omicron) infection by the ratio of testing rates across the booster and primary-series cohorts (Table S3), effectiveness of the booster dose relative to the primary series was estimated at 47.7% (95% CI: 46.0-49.3%) for BNT162b2 and at 54.0% (95% CI: 50.7-57.2%) for mRNA-1273.

Discussion

BNT162b2 booster vaccination was associated with an 86% reduction in incidence of symptomatic infection for the Delta variant, but with only 50% reduction for the Omicron variant. For the mRNA-1273 booster, the reduction in incidence of Omicron was also similar at 51%. There were fewer cases of severe COVID-19 in the booster-dose cohorts than in the primary-series cohorts, but cases of severe COVID-19 were rare in both booster-dose and primary-series cohorts.

These findings, in the context of a global expansion of Omicron and dwindling incidence of Delta, suggest that a longer term strategy for a global response is development and

administration of a new generation of vaccines targeting a broad range of variants, or pan-coronavirus vaccines, to confront future SARS-CoV-2 variants, rather than continuing with a strategy of repeated booster vaccination employing existing vaccines. To illustrate how Omicron has eroded the effectiveness of current mRNA vaccines, the cumulative incidence of any documented infection in the vaccinated cohorts in Qatar was <1% after 6 months of follow-up during times in which incidence was predominated by Beta and Delta,[3] but reached ten-fold higher in the present study to approximately 10% among recently boosted persons, after only 2-3 weeks of follow-up during the Omicron wave.

The above estimated effectiveness measures for Delta and Omicron are broadly consistent with growing evidence for effectiveness of mRNA vaccines against these variants in other countries,[26-29,39-50] but in our study, effectiveness of boosters was compared to that of the primary series, and not to unvaccinated persons. Effectiveness was also assessed against specifically symptomatic infection, with results being also presented for any documented infection, symptomatic or asymptomatic.

This study has limitations. With the high and durable effectiveness of BNT162b2 and mRNA-1273 primary series against hospitalization and death,[1,6] the lower severity of Omicron,[51] the relatively young population of Qatar,[18,52] and the time lag between infection and severe forms of COVID-19, there were too few confirmed severe, critical, and fatal COVID-19 cases at time of writing of this report to be able to precisely estimate effectiveness of the boosters against COVID-19 hospitalization and death. As an observational study, the vaccinated cohorts were neither blinded nor randomized, so potentially unmeasured or uncontrolled confounding cannot be excluded. While matching was done for sex, age, nationality, and calendar week of the second vaccine dose, it was not possible for other factors, such as comorbidities or occupation, as such

data were not available to investigators. With the large Omicron wave in Qatar, use of rapid antigen testing was expanded to supplement PCR testing starting from January 5, 2022, but rapid antigen testing data were not available for analysis.

Matching was done to control for factors known to affect infection risk of exposure in Qatar,[18,21,30-32] as well as variant exposure.[1,13,14,23,36-38] Matching by age may have (partially) reduced the potential for bias due to co-morbidities. The number of individuals with severe chronic conditions is also small in Qatar's young population.[18,52] Matching by nationality may have (partially) controlled the differences in occupational risk, in consideration of the association between nationality and occupation in Qatar.[18,21,30-32] The prescription for matching was recently shown to provide adequate control of the differences in the risk of exposure to the infection.[3] Use of rapid antigen testing may not have differentially affected the investigated cohorts. A strength of this study is exclusion of those with a documented prior infection, as presence of prior infection may affect the protection afforded by the booster.[15]

In conclusion, mRNA boosters are associated with high effectiveness against Delta infection, but modest effectiveness against Omicron infection. Against future SARS-CoV-2 waves that may be driven by newly emerging variants, these findings suggest that a longer term strategy for a global response is development and administration of a new generation of vaccines targeting a broad range of variants, or pan-coronavirus vaccines, to confront future SARS-CoV-2 variants, rather than continuing with a strategy of repeated booster vaccination employing existing vaccines.

Acknowledgements

We acknowledge the many dedicated individuals at Hamad Medical Corporation, the Ministry of Public Health, the Primary Health Care Corporation, Qatar Biobank, Sidra Medicine, and Weill Cornell Medicine-Qatar for their diligent efforts and contributions to make this study possible. The authors are grateful for institutional salary support from the Biomedical Research Program and the Biostatistics, Epidemiology, and Biomathematics Research Core, both at Weill Cornell Medicine-Qatar, as well as for institutional salary support provided by the Ministry of Public Health, Hamad Medical Corporation, and Sidra Medicine. The authors are also grateful for the Qatar Genome Programme and Qatar University Biomedical Research Center for institutional support for the reagents needed for the viral genome sequencing. The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the article. Statements made herein are solely the responsibility of the authors.

Author contributions

LJA conceived and co-designed the study, led the statistical analyses, and co-wrote the first draft of the article. HC co-designed the study, performed the statistical analyses, and co-wrote the first draft of the article. HY, HAK, and MS conducted viral genome sequencing. PT and MRH conducted the multiplex, RT-qPCR variant screening and viral genome sequencing. All authors contributed to data collection and acquisition, database development, discussion and interpretation of the results, and to the writing of the manuscript. All authors have read and approved the final manuscript.

Competing interests

Dr. Butt has received institutional grant funding from Gilead Sciences unrelated to the work presented in this paper. Otherwise, we declare no competing interests.

References

1. Chemaitelly H, Tang P, Hasan MR, et al. Waning of BNT162b2 Vaccine Protection against SARS-CoV-2 Infection in Qatar. N Engl J Med 2021;385:e83.
2. Goldberg Y, Mandel M, Bar-On YM, et al. Waning Immunity after the BNT162b2 Vaccine in Israel. N Engl J Med 2021.
3. Abu-Raddad LJ, Chemaitelly H, Ayoub HH, et al. Protection offered by mRNA-1273 versus BNT162b2 vaccines against SARS-CoV-2 infection and severe COVID-19 in Qatar. medRxiv In press at the New England Journal of Medicine 2021:2021.11.12.21266250.
4. Andrews N, Tessier E, Stowe J, et al. Duration of Protection against Mild and Severe Disease by Covid-19 Vaccines. N Engl J Med 2022.
5. Tartof SY, Slezak JM, Fischer H, et al. Effectiveness of mRNA BNT162b2 COVID-19 vaccine up to 6 months in a large integrated health system in the USA: a retrospective cohort study. Lancet 2021;398:1407-16.
6. Abu-Raddad LJ, Chemaitelly H, Ayoub HH, et al. Waning of mRNA-1273 vaccine effectiveness against SARS-CoV-2 infection in Qatar. medRxiv In press at the New England Journal of Medicine 2021:2021.12.16.21267902.
7. World Health Organization. Tracking SARS-CoV-2 variants. Available from: https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/. 2021.
8. Schmidt F, Muecksch F, Weisblum Y, et al. Plasma neutralization properties of the SARS-CoV-2 Omicron variant. medRxiv 2021.
9. Planas D, Saunders N, Maes P, et al. Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. Nature 2021.
10. Altarawneh H, Chemaitelly H, Tang P, et al. Protection afforded by prior infection against SARS-CoV-2 reinfection with the Omicron variant. medRxiv 2022:2022.01.05.22268782.
11. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. N Engl J Med 2020;383:2603-15.
12. Baden LR, El Sahly HM, Essink B, et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. N Engl J Med 2021;384:403-16.
13. Abu-Raddad LJ, Chemaitelly H, Butt AA, National Study Group for Covid Vaccination. Effectiveness of the BNT162b2 Covid-19 Vaccine against the B.1.1.7 and B.1.351 Variants. N Engl J Med 2021;385:187-9.
14. Chemaitelly H, Yassine HM, Benslimane FM, et al. mRNA-1273 COVID-19 vaccine effectiveness against the B.1.1.7 and B.1.351 variants and severe COVID-19 disease in Qatar. Nat Med 2021;27:1614-21.
15. Abu-Raddad LJ, Chemaitelly H, Ayoub HH, et al. Association of Prior SARS-CoV-2 Infection With Risk of Breakthrough Infection Following mRNA Vaccination in Qatar. JAMA 2021;326:1930-9.
16. World Health Organization. COVID-19 clinical management: living guidance. Available from: https://www.who.int/publications/i/item/WHO-2019-nCoV-clinical-2021-1. Accessed on: May 15, 2021. 2021.
17. World Health Organization. International guidelines for certification and classification (coding) of COVID-19 as cause of death. Available from: https://www.who.int/classifications/icd/Guidelines_Cause_of_Death_COVID-19-20200420-

EN.pdf?ua=1. Document Number: WHO/HQ/DDI/DNA/CAT. Accessed on May 15, 2021. 2020.

18.   Abu-Raddad LJ, Chemaitelly H, Ayoub HH, et al. Characterizing the Qatar advanced-phase SARS-CoV-2 epidemic. Sci Rep 2021;11:6233.

19.   Abu-Raddad LJ, Chemaitelly H, Coyle P, et al. SARS-CoV-2 antibody-positivity protects against reinfection for at least seven months with 95% efficacy. EClinicalMedicine 2021;35:100861.

20.   Abu-Raddad LJ, Chemaitelly H, Malek JA, et al. Assessment of the Risk of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Reinfection in an Intense Reexposure Setting. Clin Infect Dis 2021;73:e1830-e40.

21.   Ayoub HH, Chemaitelly H, Seedat S, et al. Mathematical modeling of the SARS-CoV-2 epidemic in Qatar and its impact on the national response to COVID-19. J Glob Health 2021;11:05005.

22.   Bertollini R, Chemaitelly H, Yassine HM, Al-Thani MH, Al-Khal A, Abu-Raddad LJ. Associations of Vaccination and of Prior Infection With Positive PCR Test Results for SARS-CoV-2 in Airline Passengers Arriving in Qatar. JAMA 2021;326:185-8.

23.   Tang P, Hasan MR, Chemaitelly H, et al. BNT162b2 and mRNA-1273 COVID-19 vaccine effectiveness against the SARS-CoV-2 Delta variant in Qatar. Nat Med 2021;27:2136-43.

24.   Abu-Raddad LJ, Chemaitelly H, Ayoub HH, et al. Introduction and expansion of the SARS-CoV-2 B.1.1.7 variant and reinfections in Qatar: A nationally representative cohort study. Plos Med 2021;18:e1003879.

25.   Hernan MA, Robins JM. Using Big Data to Emulate a Target Trial When a Randomized Trial Is Not Available. Am J Epidemiol 2016;183:758-64.

26.   Barda N, Dagan N, Cohen C, et al. Effectiveness of a third dose of the BNT162b2 mRNA COVID-19 vaccine for preventing severe outcomes in Israel: an observational study. Lancet 2021;398:2093-100.

27.   Arbel R, Hammerman A, Sergienko R, et al. BNT162b2 Vaccine Booster and Mortality Due to Covid-19. N Engl J Med 2021.

28.   Andrews N, Stowe J, Kirsebom F, Gower C, Ramsay M, Bernal JL. Effectiveness of BNT162b2 (Comirnaty, Pfizer-BioNTech) COVID-19 booster vaccine against covid-19 related symptoms in England: test negative case-control study. medRxiv 2021:2021.11.15.21266341.

29.   Patalon T, Gazit S, Pitzer VE, Prunas O, Warren JL, Weinberger DM. Odds of Testing Positive for SARS-CoV-2 Following Receipt of 3 vs 2 Doses of the BNT162b2 mRNA Vaccine. JAMA Intern Med 2021.

30.   Coyle PV, Chemaitelly H, Ben Hadj Kacem MA, et al. SARS-CoV-2 seroprevalence in the urban population of Qatar: An analysis of antibody testing on a sample of 112,941 individuals. iScience 2021;24:102646.

31.   Al-Thani MH, Farag E, Bertollini R, et al. SARS-CoV-2 Infection Is at Herd Immunity in the Majority Segment of the Population of Qatar. Open Forum Infect Dis 2021;8:ofab221.

32.   Jeremijenko A, Chemaitelly H, Ayoub HH, et al. Herd Immunity against Severe Acute Respiratory Syndrome Coronavirus 2 Infection in 10 Communities, Qatar. Emerg Infect Dis 2021;27:1343-52.

33.   Austin PC. Using the Standardized Difference to Compare the Prevalence of a Binary Variable Between Two Groups in Observational Research. Communications in Statistics - Simulation and Computation 2009;38:1228-34.

34.   Kaplan EL, Meier P. Nonparametric-Estimation from Incomplete Observations. Journal of the American Statistical Association 1958;53:457-81.
35.   StataCorp. Stata Statistical Software: Release 17. College Station, TX: StataCorp LLC. 2021.
36.   Qatar viral genome sequencing data. Data on randomly collected samples. https://www.gisaid.org/phylodynamics/global/nextstrain/. 2021. at https://www.gisaid.org/phylodynamics/global/nextstrain/.)
37.   Hasan MR, Kalikiri MKR, Mirza F, et al. Real-Time SARS-CoV-2 Genotyping by High-Throughput Multiplex PCR Reveals the Epidemiology of the Variants of Concern in Qatar. Int J Infect Dis 2021;112:52-4.
38.   Benslimane FM, Al Khatib HA, Al-Jamal O, et al. One Year of SARS-CoV-2: Genomic Characterization of COVID-19 Outbreak in Qatar. Front Cell Infect Microbiol 2021;11:768883.
39.   Saciuk Y, Kertes J, Shamir Stein N, Ekka Zohar A. Effectiveness of a Third Dose of BNT162b2 mRNA Vaccine. The Journal of Infectious Diseases 2021.
40.   Bar-On YM, Goldberg Y, Mandel M, et al. Protection against Covid-19 by BNT162b2 Booster across Age Groups. N Engl J Med 2021.
41.   Bar-On YM, Goldberg Y, Mandel M, et al. Protection of BNT162b2 Vaccine Booster against Covid-19 in Israel. N Engl J Med 2021;385:1393-400.
42.   Sharma A, Oda G, Holodniy M. Effectiveness of a third dose of BNT162b2 or mRNA-1273 vaccine for preventing post-vaccination COVID-19 infection: an observational study. medRxiv 2021:2021.11.29.21266777.
43.   Andrews N, Stowe J, Kirsebom F, et al. Effectiveness of COVID-19 vaccines against the Omicron (B.1.1.529) variant of concern. medRxiv 2021:2021.12.14.21267615.
44.   Tseng HF, Ackerson BK, Luo Y, et al. Effectiveness of mRNA-1273 against SARS-CoV-2 omicron and delta variants. medRxiv 2022:2022.01.07.22268919.
45.   Buchan SA, Chung H, Brown KA, et al. Effectiveness of COVID-19 vaccines against Omicron or Delta infection. medRxiv 2022:2021.12.30.21268565.
46.   Robles-Fontán MM, Irizarry RA. Effectiveness of different booster regimens for preventing infection and adverse outcomes in Puerto Rico. medRxiv 2021:2021.12.19.21268070.
47.   Hansen CH, Schelde AB, Moustsen-Helm IR, et al. Vaccine effectiveness against SARS-CoV-2 infection with the Omicron or Delta variants following a two-dose or booster BNT162b2 or mRNA-1273 vaccination series: A Danish cohort study. medRxiv 2021:2021.12.20.21267966.
48.   Tartof SY, Slezak JM, Puzniak L, et al. Effectiveness of a Third Dose of BNT162b2 mRNA COVID-19 Vaccine in a Large US Health System: A Retrospective Cohort Study. Available at SSRN: https://ssrn.com/abstract=3989856 or http://dx.doi.org/10.2139/ssrn.3989856 2021.
49.   Sheikh A, Kerr S, Woolhouse M, McMenamin J, Robertson C. Severity of Omicron variant of concern and vaccine effectiveness against symptomatic disease: national cohort with nested test negative design study in Scotland. Availablble at https://www.research.ed.ac.uk/en/publications/severity-of-omicron-variant-of-concern-and-vaccine-effectiveness-. Preprint accessed on January 12, 2021. 2021.
50.   Willett BJ, Grove J, MacLean OA, et al. The hyper-transmissible SARS-CoV-2 Omicron variant exhibits significant antigenic change, vaccine escape and a switch in cell entry mechanism. medRxiv 2022:2022.01.03.21268111.
51.   Wolter N, Jassat W, Walaza S, et al. Early assessment of the clinical severity of the SARS-CoV-2 Omicron variant in South Africa. medRxiv 2021:2021.12.21.21268116.

52.   Seedat S, Chemaitelly H, Ayoub HH, et al. SARS-CoV-2 infection hospitalization, severity, criticality, and fatality rates in Qatar. Sci Rep 2021;11:18182.

53.   Möller S, Ahrenfeldt LJ. Estimating Relative Risk When Observing Zero Events— Frequentist Inference and Bayesian Credibility Intervals. International journal of environmental research and public health 2021;18:5527.

Figure 1. Cohort selection in studying effectiveness of BNT162b2 and mRNA-1273 booster vaccination.

2,232,224 Individuals who received at least 2 COVID-19 vaccine doses between January 5, 2021 and January 9, 2022

48,759 Excluded due to a vaccination record with ChAdOx1 nCoV-19 (AZD1222)

889,372 Excluded due to a vaccination record with mRNA-1273

1,294,713 Excluded due to a vaccination record with BNT162b2

1,294,093 Individuals with 2 BNT162b2 vaccine doses

888,752 Individuals with 2 mRNA-1273 vaccine doses

159,995 Individuals excluded:
- **159,972** Had a PCR-confirmed infection before Dose 2
- **23** Non-COVID-19-related deaths with no ascertained death date

109,208 Individuals excluded:
- **109,188** Had a PCR-confirmed infection before Dose 2
- **20** Non-COVID-19-related deaths with no ascertained death date

960,795 Individuals excluded:
- **931,404** Did not receive Dose 3
- **23,336** Did not complete 7 days after Dose 3 by the end of the study (January 9, 2022)
- **6,053** Had a PCR-confirmed infection before completing 7 days after Dose 3
- **2** Non-COVID-19-related deaths before completing 7 days after Dose 3

738,514 Individuals excluded:
- **726,392** Did not receive Dose 3
- **11,361** Did not complete 7 days after Dose 3 by the end of the study (January 9, 2022)
- **761** Had a PCR-confirmed infection before completing 7 days after Dose 3

173,303 Eligible individuals with 3 vaccine doses

1,134,098 Eligible individuals with 2 or more vaccine doses

41,030 Eligible individuals with 3 vaccine doses

779,544 Eligible individuals with 2 or more vaccine doses

9,343 Not matched

970,138 Not matched

7,214 Not matched

981,930 Not matched

163,960 Individuals who received Dose 3 were matched* to controls who were infection free, alive, and with no Dose 3 vaccination as of the start of follow-up (≥7 days after Dose 3)

163,960 Controls were matched* to individuals who received Dose 3 and followed from ≥7 days after vaccination with Dose 3 of their match

40,504 Individuals who received Dose 3 were matched* to controls who were infection free, alive, and with no Dose 3 vaccination as of the start of follow-up (≥7 days after Dose 3)

40,504 Controls were matched* to individuals who received Dose 3 and followed from ≥7 days after vaccination with Dose 3 of their match

- **7,210** Infected with SARS-CoV-2
  - **2,035** Symptomatic infections†
    - None progressed to severe, critical or fatal COVID-19
  - **5,175** Non-symptomatic infections*
    - None progressed to severe, critical or fatal COVID-19
- **43,522** Censored because their match received Dose 3
- **14** Non-COVID-related deaths during follow-up
- **113,214** Followed until end of study (January 9, 2022)

- **10,994** Infected with SARS-CoV-2
  - **3,858** Symptomatic infections*
    - 8 progressed to severe COVID-19
    - None progressed to critical or fatal COVID-19
  - **7,136** Non-symptomatic infections†
    - 7 progressed to severe COVID-19
    - None progressed to critical or fatal COVID-19
- **43,993** Received Dose 3
- **37** Non-COVID-related deaths during follow-up
- **108,936** Followed until end of study (January 9, 2022)

- **1,463** Infected with SARS-CoV-2
  - **317** Symptomatic infections*
    - None progressed to severe, critical or fatal COVID-19
  - **1,146** Non-symptomatic infections†
    - None progressed to severe, critical or fatal COVID-19
- **4,111** Censored because their match received Dose 3
- **1** Non-COVID-related deaths during follow-up
- **34,929** Followed until end of study (January 9, 2022)

- **1,880** Infected with SARS-CoV-2
  - **589** Symptomatic infections†
    - None progressed to severe, critical or fatal COVID-19
  - **1,291** Non-symptomatic infections*
    - None progressed to severe, critical or fatal COVID-19
- **4,176** Received Dose 3
- **1** Non-COVID-related deaths during follow-up
- **34,447** Followed until end of study (January 9, 2022)

*Individuals with 3 vaccine doses were exact-matched in a 1:1 ratio by sex, 10-year age group, nationality, and calendar week of second vaccine dose to the first eligible individual with at least 2 vaccine doses that had not yet received Dose 3 at the start of the follow-up.
†Symptomatic infection was defined as a PCR-positive test conducted because of clinical suspicion due to presence of symptoms compatible with a respiratory tract infection. Non-symptomatic infection was defined as an infection diagnosed for a reason for PCR testing other than clinical suspicion. Some of the latter infections developed symptoms after diagnosis and some progressed to severe forms of COVID-19.

Table 1. Baseline characteristics of the full and matched primary-series and booster-dose cohorts for the BNT162b2 and mRNA-1273 vaccines.

	BNT162b2						mRNA-1273					
	Full cohorts			Matched cohorts*			Full cohorts			Matched cohorts*		
Characteristics	Individuals who received Dose 3*	Individuals who did not receive Dose 3*	SMD†	Individuals who received Dose 3	Individuals who did not receive Dose 3	SMD†	Individuals who received Dose 3*	Individuals who did not receive Dose 3*	SMD†	Individuals who received Dose 3	Individuals who did not receive Dose 3	SMD†
	N=173,303	N=1,134,098		N=163,960	N=163,960		N=779,544	N=41,030		N=40,504	N=40,504	
Median age (IQR) — years	43 (35-53)	36 (28-44)	0.96‡	43 (35-53)	42 (35-53)	0.01‡	35 (30-42)	40 (33-46)	0.40	40 (33-46)	40 (33-46)	0.01
Age group — no. (%)												
0-19 years	2,132 (1.2)	114,011 (10.1)		2,019 (1.2)	2,019 (1.2)		452 (1.1)	5,727 (0.7)		410 (1.0)	410 (1.0)	
20-29 years	14,322 (8.3)	211,832 (18.7)		13,984 (8.5)	13,984 (8.5)		4,729 (11.5)	186,523 (23.9)		4,681 (11.6)	4,681 (11.6)	
30-39 years	51,461 (29.7)	389,355 (34.3)		50,173 (30.6)	50,173 (30.6)		15,249 (37.2)	332,931 (42.7)		15,149 (37.4)	15,149 (37.4)	
40-49 years	45,467 (26.2)	247,841 (21.9)	0.66	43,789 (26.7)	43,789 (26.7)	0.00	13,843 (33.7)	180,793 (23.2)	0.43	1,3683 (33.8)	13,683 (33.8)	0.00
50-59 years	35,803 (20.7)	114,401 (10.1)		33,162 (20.2)	33,162 (20.2)		5,414 (13.2)	60,101 (7.7)		5,309 (13.1)	5,309 (13.1)	
60-69 years	18,221 (10.5)	42,941 (3.8)		15,978 (9.8)	15,978 (9.8)		1,179 (2.9)	11,351 (1.5)		1,128 (2.8)	1,128 (2.8)	
70+ years	5,897 (3.4)	13,717 (1.2)		4,855 (3.0)	4,855 (3.0)		164 (0.4)	2,118 (0.3)		144 (0.4)	144 (0.4)	
Sex												
Male	107,110 (61.8)	787,914 (69.5)	0.16	102,035 (62.2)	102,035 (62.2)	0.00	27,162 (66.2)	631,681 (81.0)	0.34	26,922 (66.5)	26,922 (66.5)	0.00
Female	66,193 (38.2)	346,184 (30.5)		61,925 (37.8)	61,925 (37.8)		13,868 (33.8)	147,863 (19.0)		13,582 (33.5)	13,582 (33.5)	
Nationality§												
Bangladeshi	7,599 (4.4)	126,300 (11.1)		7,492 (4.6)	7,492 (4.6)		2,805 (6.8)	151,887 (19.5)		2,802 (6.9)	2,802 (6.9)	
Egyptian	11,824 (6.8)	63,381 (5.6)		11,391 (7.0)	11,391 (7.0)		2,062 (5.0)	30,475 (3.9)		2,059 (5.1)	2,059 (5.1)	
Filipino	25,462 (14.7)	102,841 (9.1)		24,388 (14.9)	24,388 (14.9)		7,359 (17.9)	68,803 (8.8)		7,352 (18.2)	7,352 (18.2)	
Indian	49,534 (28.6)	252,232 (22.2)		47,805 (29.2)	47,805 (29.2)		17,145 (41.8)	215,164 (27.6)		17,141 (42.3)	17,141 (42.3)	
Nepalese	3,574 (2.1)	95,937 (8.5)		3,538 (2.2)	3,538 (2.2)		1,638 (4.0)	108,857 (14.0)		1,637 (4.0)	1,637 (4.0)	
Pakistani	5,253 (3.0)	48,554 (4.3)	0.46	5,127 (3.1)	5,127 (3.1)	0.00	1,258 (3.1)	41,912 (5.4)	0.64	1,255 (3.1)	1,255 (3.1)	0.00
Qatari	20,206 (11.7)	158,207 (14.0)		19,809 (12.1)	19,809 (12.1)		627 (1.5)	14,314 (1.8)		626 (1.6)	626 (1.6)	
Sri Lankan	3,718 (2.2)	34,610 (3.1)		3,643 (2.2)	3,643 (2.2)		1,509 (3.7)	32,047 (4.1)		1,504 (3.7)	1,504 (3.7)	
Sudanese	3,422 (2.0)	25,029 (2.2)		3,255 (2.0)	3,255 (2.0)		424 (1.0)	13,216 (1.7)		421 (1.0)	421 (1.0)	
Other nationalities¶	42,711 (24.7)	227,007 (20.0)		37,512 (22.9)	37,512 (22.9)		6,203 (15.1)	102,869 (13.2)		5,707 (14.1)	5,707 (14.1)	
Calendar month of Dose 2**												
January, 2021	9,606 (5.5)	14,168 (1.3)		7,296 (4.5)	7,201 (4.4)		1 (0.0)	3 (0.0)		--	--	
February, 2021	26,614 (15.4)	50,155 (4.4)		23,500 (14.3)	23,315 (14.2)		5 (0.0)	16 (0.0)		2 (0.0)	2 (0.0)	
March, 2021	77,555 (44.8)	212,719 (18.8)		74,269 (45.3)	74,549 (45.5)		222 (0.5)	1,562 (0.2)		196 (0.5)	212 (0.5)	
April, 2021	30,219 (17.4)	140,759 (12.4)		29,857 (18.2)	29,691 (18.1)		17,560 (42.8)	100,750 (12.9)		17,308 (42.7)	17,249 (42.6)	
May, 2021	23,290 (13.4)	242,267 (21.4)		23,074 (14.1)	23,173 (14.1)		20,596 (50.2)	196,771 (25.2)		20,390 (50.3)	20,435 (50.5)	
June, 2021	5,997 (3.5)	163,019 (14.4)		5,942 (3.6)	6,009 (3.7)		2,614 (6.4)	114,129 (14.6)		2,576 (6.4)	2,516 (6.2)	
July, 2021	22 (0.0)	112,949 (10.0)	1.25	22 (0.0)	22 (0.0)	0.01	32 (0.1)	95,502 (12.3)	1.54	--	--	0.04
August, 2021	0 (0)	62,334 (5.5)		--	--		0 (0)	213,870 (27.4)		--	--	
September, 2021	0 (0)	72,422 (6.4)		--	--		0 (0)	49,067 (6.3)		--	--	
October, 2021	0 (0)	28,466 (2.5)		--	--		0 (0)	3,356 (0.4)		--	--	
November, 2021	0 (0)	20,577 (1.8)		--	--		0 (0)	3,342 (0.4)		--	--	
December, 2021	0 (0)	12,255 (1.1)		--	--		0 (0)	551 (0.1)		--	--	
January, 2022	0 (0)	2,008 (0.2)					0 (0)	625 (0.1)				

Abbreviations: IQR, interquartile range; SMD, standardized mean difference.

*Cohorts were matched one-to-one by sex, 10-year age group, nationality, and calendar week of second vaccine dose.

†SMD is the difference in the mean of a covariate between groups divided by the pooled SD. An SMD <0.1 indicates adequate matching.

‡SMD is for the mean difference between groups divided by the pooled SD.

§Nationalities were chosen to represent the most populous groups in Qatar.

¶These comprise 156 other nationalities in Qatar in individuals who received Dose 3 and 191 other nationalities in individuals who did not receive Dose 3 in the unmatched cohorts of those vaccinated with BNT162b2, and 131 other nationalities in individuals who received Dose 3 and 131 other nationalities in individuals who did not receive Dose 3 in the matched cohorts of those vaccinated with BNT162b2. These also comprise 126 other nationalities in Qatar in individuals who received Dose 3 and 171 other nationalities in individuals who did not receive Dose 3 in the unmatched cohorts of those vaccinated with mRNA-1273, and 101 other nationalities in individuals who received Dose 3 and 101 other nationalities in individuals who did not receive Dose 3 in the matched cohorts of those vaccinated with mRNA-1273.

**Cohorts were exact matched using calendar week of Dose 2, but we opted to report the distribution by calendar month for brevity. Accordingly, members of the cohorts who were tested in the same week may appear in different calendar months.

Figure 2. Cumulative incidence of symptomatic infection in the matched booster-dose and primary-series cohorts for the BNT162b2 (A) and mRNA-1273 (B) vaccines.

Time (days)	0	7	14	21	28	35	42	49
No. at risk								
Vaccinated with 3 doses	163,960	120,776	102,113	83,979	64,250	44,759	27,383	15,170
Vaccinated with 2 doses	163,960	123,266	102,365	83,169	63,166	43,732	26,648	14,891

Time (days)	0	7	14	21	28	35
No. at risk						
Vaccinated with 3 doses	40,504	28,326	21,093	14,841	8,888	4,257
Vaccinated with 2 doses	40,504	28,355	20,948	14,689	8,721	4,166

Table 2. Effectiveness of BNT162b2 and mRNA-1273 booster vaccination.

Epidemiological measure	BNT162b2		mRNA-1273	
	Estimate (95% CI)	Effectiveness in % (95% CI)	Estimate (95% CI)	Effectiveness in % (95% CI)
Total follow-up time—booster-dose cohort (person-weeks)	580,361	--	96,659	--
Total follow-up time—primary-series cohort (person-weeks)	574,302	--	97,253	--
Incidence rate of symptomatic infection—booster-dose cohort (per 10,000 person-weeks)	35.1 (33.6-36.6)	--	32.8 (29.4-36.6)	--
Incidence rate of symptomatic infection—primary-series cohort (per 10,000 person-weeks)	67.2 (65.1-69.3)	--	60.6 (55.9-65.7)	--
Unadjusted hazard ratio for symptomatic infection	0.50 (0.48-0.53)	49.8 (47.0-52.5)	0.49 (0.43-0.57)	50.7 (43.3-57.2)
Adjusted hazard ratio for symptomatic infection[*]	0.50 (0.47-0.53)	50.1 (47.3-52.8)	0.49 (0.43-0.57)	50.8 (43.4-57.3)
Unadjusted odds ratio for hospitalization and death[†]	0.00 (0.00-0.29)[‡]	100.0% (71.4-100.0%)[‡]	--	--

Abbreviations: CI, confidence interval.

[*]Cox regression analysis adjusted for sex, 10 age-groups, 10 nationality groups, and calendar week of second vaccine dose.

[†]Cox regression could not be implemented because of zero events in the booster-dose cohort for BNT162b2 and because of zero events in both the booster-dose cohort and the primary-series cohort for mRNA-1273.

[‡]Because of zero events in the booster-dose cohort and rarity of events in the primary-series cohort, the hazard ratio comparing hospitalizations and deaths in both cohorts was approximated using the odds ratio with exact 95% CIs calculated using the Baptista and Pike method.[53]

Figure 3. Cumulative incidence of symptomatic infection in the matched booster-dose and primary-series cohorts for the BNT162b2 vaccine during the Delta-predominated low-incidence phase.

Time (days)	0	7	14	21	28	35	42	49
No. at risk								
Vaccinated with 3 doses	58,821	30,928	20,079	15,016	10,826	7,415	5,174	3,155
Vaccinated with 2 doses	58,821	31,138	20,087	14,941	10,768	7,375	5,141	3,129

Table 3. Effectiveness of BNT162b2 booster vaccination during the Delta-predominated low-incidence phase.

Epidemiological measure	Estimate (95% CI)	Effectiveness in % (95% CI)
Total follow-up time—booster-dose cohort (person-weeks)	120,914	--
Total follow-up time—primary-series cohort (person-weeks)	120,599	--
Incidence rate for symptomatic infection—booster-dose cohort (per 10,000 person-weeks)	0.6 (0.3-1.2)	--
Incidence rate for symptomatic infection—primary-series cohort (per 10,000 person-weeks)	3.6 (2.6-4.8)	--
Unadjusted hazard ratio for symptomatic infection	0.14 (0.06-0.33)	86.0 (67.0-94.0)
Adjusted hazard ratio for symptomatic infection[*]	0.14 (0.06-0.33)	86.1 (67.3-94.1)

Abbreviations: CI, confidence interval.

[*]Cox regression analysis adjusted for sex, 10 age-groups, 10 nationality groups, and calendar week of second vaccine dose.

Supplementary Appendix

Table of Contents

Section 1. COVID-19 severity, criticality, and fatality classification

Severe Coronavirus Disease 2019 (COVID-19) disease was defined per the World health Organization (WHO) classification as a severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infected person with "oxygen saturation of <90% on room air, and/or respiratory rate of >30 breaths/minute in adults and children >5 years old (or ≥60 breaths/minute in children <2 months old or ≥50 breaths/minute in children 2-11 months old or ≥40 breaths/minute in children 1–5 years old), and/or signs of severe respiratory distress (accessory muscle use and inability to complete full sentences, and, in children, very severe chest wall indrawing, grunting, central cyanosis, or presence of any other general danger signs)".[1] Detailed WHO criteria for classifying SARS-CoV-2 infection severity can be found in the WHO technical report.[1]

Critical COVID-19 disease was defined per WHO classification as a SARS-CoV-2 infected person with "acute respiratory distress syndrome, sepsis, septic shock, or other conditions that would normally require the provision of life sustaining therapies such as mechanical ventilation (invasive or non-invasive) or vasopressor therapy".[1] Detailed WHO criteria for classifying SARS-CoV-2 infection criticality can be found in the WHO technical report.[1]

COVID-19 death was defined per WHO classification as "a death resulting from a clinically compatible illness, in a probable or confirmed COVID-19 case, unless there is a clear alternative cause of death that cannot be related to COVID-19 disease (e.g. trauma). There should be no period of complete recovery from COVID-19 between illness and death. A death due to COVID-19 may not be attributed to another disease (e.g. cancer) and should be counted independently of preexisting conditions that are suspected of triggering a severe course of COVID-19". Detailed WHO criteria for classifying COVID-19 death can be found in the WHO technical report.[2]

Section 2. Laboratory methods

Nasopharyngeal and/or oropharyngeal swabs were collected for PCR testing and placed in Universal Transport Medium (UTM). Aliquots of UTM were: 1) extracted on a QIAsymphony platform (QIAGEN, USA), KingFisher Flex (Thermo Fisher Scientific, USA), MGISP-960 (MGI, China), or ExiPrep 96 Lite (Bioneer, South Korea) followed by testing with real-time reverse-transcription PCR (RT-qPCR) using TaqPath COVID-19 Combo Kits (Thermo Fisher Scientific, USA) on an ABI 7500 FAST (Thermo Fisher Scientific, USA); 2) tested directly on the Cepheid GeneXpert system using the Xpert Xpress SARS-CoV-2 (Cepheid, USA); or 3) loaded directly into a Roche cobas 6800 system and assayed with the cobas SARS-CoV-2 Test (Roche, Switzerland). The first assay targets the viral S, N, and ORF1ab gene regions. The second targets the viral N and E-gene regions, and the third targets the ORF1ab and E-gene regions.

All PCR testing was conducted at the Hamad Medical Corporation Central Laboratory or Sidra Medicine Laboratory, following standardized protocols.

Table S1. STROBE checklist for cohort studies.

	Item No	Recommendation	Main Text page
Title and abstract	1	(*a*) Indicate the study's design with a commonly used term in the title or the abstract	Abstract
		(*b*) Provide in the abstract an informative and balanced summary of what was done and what was found	Abstract
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Introduction
Objectives	3	State specific objectives, including any prespecified hypotheses	Introduction
Methods			
Study design	4	Present key elements of study design early in the paper	Methods ('Data sources and study design', paragraph 2)
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Methods ('Data sources and study design') & Figure 1
Participants	6	(*a*) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (*b*) For matched studies, give matching criteria and number of exposed and unexposed	Methods ('Data sources and study design', paragraphs 1-6) & Figure 1
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Methods ('Data sources and study design', paragraphs 2-8) & Table 1
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Methods ('Data sources and study design' & 'Laboratory methods'), Table 1 & Sections 1-2 in Appendix
Bias	9	Describe any efforts to address potential sources of bias	Methods ('Data sources and study design', paragraphs 4-5 & 'Statistical analysis', paragraph 4)
Study size	10	Explain how the study size was arrived at	Figure 1
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Methods ('Data sources and study design', paragraphs 4 and 8) & Table 1
Statistical methods	12	(*a*) Describe all statistical methods, including those used to control for confounding	Methods ('Statistical analysis')
		(*b*) Describe any methods used to examine subgroups and interactions	NA
		(*c*) Explain how missing data were addressed	NA, see Methods ('Data sources and study design', paragraph 1)
		(*d*) If applicable, explain how loss to follow-up was addressed	NA, see Methods ('Data sources and study design', paragraph 1)
		(*e*) Describe any sensitivity analyses	Methods ('Data sources and study design', paragraph 2 & 'Statistical analysis', paragraphs 3-4)
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	Results ('Study population for BNT162b2 vaccine' & 'Study population for mRNA-1273 vaccine), Figure 1, & Table 1
Descriptive data	14	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Results ('Study population for BNT162b2 vaccine' & 'Study population for mRNA-1273 vaccine) & Table 1
		(b) Indicate number of participants with missing data for each variable of interest	NA, see Methods ('Data sources and study design', paragraph 1)
		(c) Summarise follow-up time (eg, average and total amount)	Results ('Study population for BNT162b2 vaccine' & 'Study population for mRNA-1273 vaccine), Figure 2, & Table 2
Outcome data	15	Report numbers of outcome events or summary measures over time	Results ('Effectiveness of BNT162b2 booster against Omicron' & 'Effectiveness of mRNA-1273 booster against Omicron), Figures 1-2, & Table 2

Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Results ('Effectiveness of BNT162b2 booster against Omicron' & 'Effectiveness of mRNA-1273 booster against Omicron), Figure 2, & Table 2
		(b) Report category boundaries when continuous variables were categorized	Table 1
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Results ('Effectiveness of BNT162b2 booster against Delta' & 'Additional analyses), Figure 3, Table 3, & Figure S1 & Tables S2-S3 in Appendix
Discussion			
Key results	18	Summarise key results with reference to study objectives	Discussion, paragraphs 1-3
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Discussion, paragraphs 4-5
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Discussion, paragraph 6
Generalisability	21	Discuss the generalisability (external validity) of the study results	Discussion, paragraphs 4-5
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Acknowledgements

Figure S1. Cumulative incidence of symptomatic infection in the matched booster-dose and primary-series cohorts for each of the BNT162b2 (A) and mRNA-1273 (B) vaccines, with start of follow-up on the 15th day after booster dose.

Time (days)	0	7	14	21	28	35	42	49
No. at risk								
Vaccinated with 3 doses	138,629	108,972	89,530	68,679	47,953	28,976	15,915	11,718
Vaccinated with 2 doses	138,629	109,909	88,863	67,610	46,900	28,290	15,609	11,464

Time (days)	0	7	14	21	28	35
No. at risk						
Vaccinated with 3 doses	29,774	21,827	15,375	9,199	4,392	1,060
Vaccinated with 2 doses	29,774	21,747	15,271	9,053	4,314	1,039

Table S2. Effectiveness of BNT162b2 and mRNA-1273 booster vaccination, with start of follow-up on the 15th day after booster dose.

Epidemiological measure	BNT162b2		mRNA-1273	
	Estimate (95% CI)	Effectiveness in % (95% CI)	Estimate (95% CI)	Effectiveness in % (95% CI)
Total follow-up time—booster-dose cohort (person-weeks)	465,476	--	65,523	--
Total follow-up time—primary-series cohort (person-weeks)	460,012	--	65,023	--
Incidence rate for symptomatic infection—booster-dose cohort (per 10,000 person-weeks)	39.9 (38.1-41.8)	--	35.7 (31.4-40.6)	--
Incidence rate for symptomatic infection—primary-series cohort (per 10,000 person-weeks)	78.5 (76.0-81.1)	--	67.8 (61.8-74.5)	--
Unadjusted hazard ratio for symptomatic infection	0.50 (0.47-0.53)	50.0 (47.1-52.7)	0.50 (0.43-0.59)	50.0 (41.2-57.4)
Adjusted hazard ratio for symptomatic infection[*]	0.50 (0.47-0.53)	50.3 (47.5-53.0)	0.50 (0.42-0.59)	50.1 (41.4-57.6)

Abbreviations: CI, confidence interval.

[*]Cox regression analysis adjusted for sex, 10 age-groups, 10 nationality groups, and calendar week of second vaccine dose.

Table S3. Distribution of the reason for PCR testing during follow-up for A) all PCR tests during follow-up and B) identified infections, among the matched booster-dose and primary-series cohorts for the BNT162b2 and mRNA-1273 vaccines.

| | BNT162b2 | | | | | | mRNA-1273 | | | | | |
| | A) All PCR tests | | | B) PCR-confirmed infections | | | A) All PCR tests | | | B) PCR-confirmed infections | | |
	Individuals who received Dose 3*	Individuals who did not receive Dose 3*	SMD†	Individuals who received Dose 3	Individuals who did not receive Dose 3	SMD†	Individuals who received Dose 3*	Individuals who did not receive Dose 3*	SMD†	Individuals who received Dose 3	Individuals who did not receive Dose 3	SMD†
Number of tests	**N=65,589**	**N=55,369**		**N=7,210**	**N=10,994**		**N=10,742**	**N=6,962**		**N=1,463**	**N=1,880**	
Reason for PCR testing												
Clinical suspicion	5,677 (8.7)	7,169 (13.0)		2,035 (28.2)	3,858 (35.1)		824 (7.7)	926 (13.3)		317 (21.7)	589 (31.3)	
Contact tracing	2,930 (4.5)	2,791 (5.0)		867 (12.0)	1,080 (9.8)		359 (3.3)	409 (5.9)		124 (8.5)	196 (10.4)	
Survey	6,847 (10.4)	7,377 (13.3)	0.28	951 (13.2)	1,570 (14.3)	0.20	821 (7.6)	856 (12.3)	0.43	180 (12.3)	233 (12.4)	0.30
Individual request	2,268 (3.5)	2,005 (3.6)		421 (5.8)	644 (5.9)		509 (4.7)	322 (4.6)		100 (6.8)	125 (6.7)	
Pre-travel	34,361 (52.4)	21,683 (39.2)		2,288 (31.7)	2,811 (25.6)		6,462 (60.2)	2,775 (39.9)		667 (45.6)	619 (32.9)	
Port of entry	12,349 (18.8)	12,893 (23.3)		455 (6.3)	680 (6.2)		1,674 (15.6)	1,554 (22.3)		60 (4.1)	82 (4.4)	
Healthcare routine testing	959 (1.5)	1,335 (2.4)		150 (2.1)	316 (2.9)		86 (0.8)	103 (1.5)		13 (0.9)	31 (1.7)	
Other	198 (0.3)	116 (0.2)		43 (0.6)	35 (0.3)		7 (0.1)	17 (0.2)		2 (0.1)	5 (0.3)	

Abbreviations: SMD, standardized mean difference.

*Cohorts were matched one-to-one by sex, 10-year age group, nationality, and calendar week of second vaccine dose.

†SMD is the difference in the mean of a covariate between groups divided by the pooled SD.

References

1.   World Health Organization. COVID-19 clinical management: living guidance. Available from: https://www.who.int/publications/i/item/WHO-2019-nCoV-clinical-2021-1. Accessed on: May 15 2021. 2021.
2.   World Health Organization. International guidelines for certification and classification (coding) of COVID-19 as cause of death. Available from: https://www.who.int/classifications/icd/Guidelines_Cause_of_Death_COVID-19-20200420-EN.pdf?ua=1. Document Number: WHO/HQ/DDI/DNA/CAT. Accessed on May 31, 2021. 2021.