PBPK-led guidance for cystic fibrosis patients taking elexacaftor-tezacaftor-ivacaftor with

nirmatrelvir-ritonavir for the treatment of COVID-19

Hong, E.[1]; Almond, L.M.[2]; Chung P.S.[3,4]; Rao, A.P.[3,4]; Beringer, P.M.[1,4,*]

[1]Department of Clinical Pharmacy, School of Pharmacy, University of Southern California, 1985

Zonal Ave, Los Angeles, CA 90033, USA

[2] Certara UK Ltd, Simcyp Division, 1 Concourse Way, Level 2-Acero, Sheffield, S1 2BJ, United

Kingdom

[3]Division of Pulmonary and Critical Care Medicine, Department of Medicine, Keck School of

Medicine, University of Southern California, 1975 Zonal Ave, Los Angeles, CA 90033, USA

[4]USC Anton Yelchin CF Clinic, 1510 San Pablo St, Los Angeles, CA 90033, USA.

[*]Corresponding author: 1985 Zonal Avenue, Los Angeles, CA 90033, USA, Tel: 323-442-1402,

Fax: 323-442-1395, Email: beringer@usc.edu

Conflict of interest statement: LMA is an employee of Certara UK Limited (Simcyp Division). EH,

PSC, APR, PMB declared no competing interests for this work.

Funding source: This work was supported by the Cystic Fibrosis Foundation [BERING21AO] and

the Anton Yelchin Foundation.

Keywords: Physiologically based pharmacokinetic (PBPK), Drug-drug interactions, Cystic fibrosis

transmembrane conductance regulator (CFTR) modulator therapy, Nirmatrelvir-ritonavir, COVID-

19, Infectious disease

Abstract

Background: Cystic fibrosis transmembrane conductance regulator (CFTR) modulating therapies

including elexacaftor, tezacaftor, and ivacaftor (ETI) are primarily eliminated through cytochrome

P450 (CYP) 3A-mediated metabolism. This creates a therapeutic challenge to the treatment of

COVID-19 with nirmatrelvir-ritonavir in people with cystic fibrosis (pwCF) due to the potential for

significant drug-drug interactions (DDI). However, pwCF are more at risk of serious illness

following COVID-19 infection and hence it is important to manage the DDI risk and provide

treatment options.

Methods: CYP3A-mediated DDI of ETI was evaluated using a physiologically based

pharmacokinetic (PBPK) modeling approach. Modeling was performed incorporating physiological

information and drug dependent parameters of ETI to predict the effect of ritonavir (the CYP3A4

inhibiting component of the combination) on pharmacokinetics of ETI. The ETI models were

verified using independent clinical pharmacokinetic and DDI data of ETI with a range of CYP3A

modulators.

Results: When ritonavir was administered on day 1 through 5, the predicted AUC ratio of ivacaftor

(the most sensitive CYP3A substrate) on day 6 was 9.31, indicating that its metabolism was strongly

inhibited. Based on the predicted DDI, the dose of ETI should be reduced when co-administered with

nirmatrelvir-ritonavir to elexacaftor 200mg-tezacaftor 100mg-ivacaftor 150mg on days 1 and 5, with

resumption of full dose ETI on day 9, considering the residual inhibitory effect of ritonavir as a

mechanism-based inhibitor.

Conclusions: Coadministration of nirmatrelvir-ritonavir requires a significant reduction in the ETI

dosing frequency with delayed resumption of full dose due to the mechanism-based inhibition with

ritonavir.

1. Introduction

The introduction of the Cystic Fibrosis Transmembrane Conductance Regular (CFTR) modulator, a triple combination of elexacaftor, tezacaftor, and ivacaftor (ETI, TRIKAFTA®) has resulted in significant improvements in lung function and nutritional status in people with cystic fibrosis (pwCF)(1). While ETI is indicated in up to 90% of the CF population(1), all 3 components are eliminated mainly through cytochrome P450 (CYP) 3A-mediated hepatic metabolism(2), and therefore present a therapeutic challenge in pwCF due to the potential for significant drug-drug interactions (DDI). The use of strong CYP3A inducers will increase the metabolism of ETI resulting in reduced exposure and a potential lack of efficacy, while concomitant therapy with agents that inhibit CYP3A will increase ETI levels placing the patient at increased risk of dose-related adverse effects. Therefore, the safe and effective use of CFTR modulators requires appropriate DDI management with concomitant CF medications.

One notable therapeutic challenge is in the treatment of COVID-19 (SARS-CoV-2). In pwCF, viral respiratory tract infections can lead to acute pulmonary exacerbations with a negative impact on lung function(3). COVID-19 infection triggers a cytokine storm which can lead to the life-threatening respiratory distress syndrome, potentially putting CF population infected with COVID-19 be at high risk of serious illness(4). The U.S. Food and Drug Administration (FDA) has recently issued an emergency use authorization (EUA) for the use of the nirmatrelvir-ritonavir for the treatment of mild-to-moderate COVID-19. Nirmatrelvir-ritonavir treatment significantly reduces hospital admissions and deaths among people with COVID-19 who are at high risk of severe illness(5). Nirmatrelvir is co-administered with ritonavir, a CYP3A inhibitor, to boost nirmatrelvir concentrations to achieve therapeutic levels(5). However, due to the potent inhibition effect of ritonavir, it may increase plasma concentrations of drugs that are primarily metabolized by CYP3A. Therefore, co-administration of nirmatrelvir-ritonavir is contraindicated with drugs highly dependent on CYP3A for clearance and for which elevated concentrations are associated with serious and/or

life-threatening reactions. Since all 3 components of ETI are eliminated mainly through CYP3A,

nirmatrelvir-ritonavir is expected to exhibit a significant drug interaction with ETI. Thus, the use of

nirmatrelvir-ritonavir in pwCF would require an adjusted dosing regimen of ETI to prevent increased

plasma concentrations and potential adverse drug reactions. However, there is currently no clinical

data available regarding the interactions of ETI with nirmatrelvir-ritonavir and no specific dosing

guidelines have been established. Therefore, there is an urgent need for the proper guidance

regarding the use of nirmatrelvir-ritonavir for pwCF to prevent progression of COVID-19 to severe

disease.

This study aimed to investigate the magnitude of the drug interactions of ritonavir-ETI, to

simulate possible treatment scenarios and provide dosing recommendations to overcome the

interaction. The CYP3A inhibition-mediated drug interaction of ETI was evaluated using a

physiologically based pharmacokinetic (PBPK) simulation-based approach. PBPK simulation is a

tool to predict the pharmacokinetic behavior of drugs in humans by integrating the information from

multiple *in vitro* and clinical studies, exploring the effects of drug (e.g., physicochemical properties)

and system (e.g., physiological) information on drug exposure. The predictive performance of PBPK

simulations for CYP enzyme-based DDIs has been well established[6, 7], and this strategy is

increasingly included during regulatory review by the FDA as an alternative for exploring DDI

potential to provide dosing recommendations in product labeling[8]. The present study contributes to

improved treatment for COVID-19 in pwCF, by providing tools to evaluate and potentially overcome

clinically important drug interactions involving highly active CFTR modulator therapy.

2. Methods

The workflow adopted for PBPK model development, verification, and application are

illustrated in **Figure 1**. The models were implemented within the Simcyp Simulator (version 19;

Certara, Sheffield, UK).

Figure 1. PBPK modeling framework detailing the processes of model development and verification

that were performed in this study. Successful model verification must precede the application of the

PBPK model of ETI for predictions of drug interactions with nirmatrelvir-ritonavir.

2.1. Model Development

PBPK Models of ETI. In the default healthy population library file (Sim-Healthy volunteers)

provided in Simcyp®, the distribution of ages and proportion of female were corrected to reflect the

demographics of CF population based on patient registry 2020 annual report published by cystic

fibrosis foundation(9). Specifically, the frequency of population aged 18-21 years, was adjusted from

4.5% in healthy population to 13.1% in CF. Also, the proportion of females was adjusted from 0.32

in healthy population to 0.48 in CF. The mean body mass index (BMI) of healthy population (23.5

kg/m2) was similar to that observed in CF (21.2 kg/m2 in 2005 to 23.1 kg/m2 in 2020) and no

further adjustment was needed, reflecting how BMI in pwCF has been increased over the years with

continued improvements in CF care (9, 10). All other system parameters were kept as the default

healthy, and this assumption is in line with the PK parameters of ETI not differing between healthy

adults and patients with CF(11-13). This population library file was used for all simulations. For trial

design, we used a total size of 100 population (10 trials and 10 subjects in each trial). The PBPK

model input parameters for ETI are summarized in **Table S1-S3** and described in detail below.

The ivacaftor model consists of the advanced dissolution, absorption, and metabolism (ADAM)

model and minimal PBPK model. The ADAM model divides the gastrointestinal tract into nine

segments, where the drug absorption from each segment is predicted by a function of drug

dissolution, luminal degradation, metabolism, transport, etc(14). The minimal PBPK model consists

of no more than five compartments, that include the gastrointestinal tract, blood, liver, and up to two

additional compartments, reducing the complexity of the model while allowing for mechanistic

simulations in the compartment of interest(15). The ivacaftor model was constructed based on

available physicochemical properties and clinical data from published PK studies(11, 16-19).

Ivacaftor is a diprotic acid with an acid dissociation constant (pKa) of 9.4 and 11.6. Ivacaftor has a

logP of 5.68, and predominantly binds to albumin with fraction of unbound drug in plasma of 0.001.

The blood/plasma partition ratio was 0.55. The absorption parameter $P_{\text{eff,man}}$ was predicted by the

Simcyp calculator based on its physicochemical properties. The distribution parameters including

V_{sac} and V_{ss} were obtained from published data. The *in vitro* studies and clinical DDI data suggest

that ivacaftor is predominantly eliminated through CYP3A4 mediated hepatic metabolism(13).

Therefore, the excretion was set to enzyme kinetics to quantify its metabolism by CYP3A. The

intrinsic clearance by CYP3A4 was back calculated from the oral clearance observed in healthy

subjects (19.0 L/h)(11). The fraction of ivacaftor being metabolized by CYP3A4 (fmCYP3A4) was

set to 98% in order to capture the observed drug interactions of ivacaftor with the strong CYP3A4

modulators, ketoconazole or rifampin. The F_{g} was also optimized to 0.50 using observed DDI data.

The additional human liver microsomes (HLM) clearance representing non-CYP3A pathways was

also estimated using the retrograde calculator and entered into the model to capture the observed

pharmacokinetics profiles.

The tezacaftor and elexacaftor PBPK models were constructed based on the data from the PBPK

review section within the NDA documents and the publication of Tsai et al(12, 13, 17). Briefly, the

elexacaftor and tezacaftor models consist of the first-order absorption and a minimal PBPK model.

In the NDA document, it was described that the absorption and distribution parameters were

obtained from the observed PK profiles following clinical phase 1-3 studies, and the fmCYP3A4 of

elexacaftor and tezacaftor were set to 67% and 73.2% based on human ADME studies. Using the

retrograde model, the rCYP3A4 CL_{int} values of elexacaftor and tezacaftor attributed to CYP3A4

were calculated to be 0.233 and 0.175 µL/min per picomole of isoform. The M1-tezacaftor, active

metabolite of tezacaftor, was also incorporated into the tezacaftor PBPK model based on data in the

NDA document(12). The other active metabolites, M23-elexacaftor and M1-ivacaftor, were not

incorporated into the PBPK analyses due to insufficient information to build the model.

PBPK Models of CYP3A Modulators. Rifampin, ketoconazole, fluconazole, itraconazole, and its

primary metabolite, hydroxy itraconazole, are prototypical CYP3A modulators that have been

implicated in clinical drug interaction studies with elexacaftor, tezacaftor, or ivacaftor. Ritonavir is

also a CYP3A modulator that we aimed to predict interactions with ETI. For simulation of DDIs, the

validated compound files of these CYP3A modulators provided in Simcyp® (version 19) were used.

2.2. Model Verification: PK Simulations

The PK profiles of ETI following a single oral dose administration and multiple administrations of

clinically relevant doses (elexacaftor 200mg qd, tezacaftor 100mg qd, and ivacaftor 150mg q12h)

were first simulated to verify the performance of the PBPK models. ETI was orally administered

under fed conditions to mimic the clinical setting, where the fat-containing food is required for

optimal absorption of ETI. The simulated data were qualified using the observed PK data in a CF

population aged older than 17 years old. The prediction accuracy for the area under the curve (AUC)

and maximum plasma concentration (C_{max}) values were calculated as a ratio of mean observed values

over mean predicted values. Successful model performance was defined by mean ratios of AUC and

C_{max} within a two-fold range as previously described(20, 21).

2.3. Model Verification: DDI Simulations

Upon accurate recapitulation of ETI's PK, the models were further assessed against the clinical DDI

data to verify fmCYP3A4 and establish if the models were adequate for the assessment of victim

DDI liability. For verification simulations, the dose and schedule of drugs were matched to the

design of the corresponding clinical DDI study in healthy subjects. For elexacaftor, itraconazole

solution (200 mg) was administered daily from Day 1 to Day 10 and a single 20 mg dose of

elexacaftor was administered orally on Day 5(13). For tezacaftor, the itraconazole solution (200 mg)

was administered BID on Day 1 and QD from Day 2 to Day 14 and tezacaftor 25 mg was

administered daily from Day 1 to Day 14(12). For ivacaftor, four clinical DDI studies were

conducted with ritonavir, ketoconazole, fluconazole, or rifampin(11, 22). For ritonavir, 50mg was

administered daily from Day 1 to Day 18 and a single 150 mg dose of ivacaftor was administered on

Day 15. The ketoconazole 400 mg was administered daily from Day 1 to Day 10 and a single 150 mg

dose of ivacaftor was administered on Day 4. For fluconazole, 400mg was administered on Day 1

and 200mg was administered daily from Day 2 to Day 8, and ivacaftor 150 mg was administered

BID from Day 1 to Day 8. For rifampin, 600mg was administered daily from Day 1 to Day 10 and a

single 150 mg dose of ivacaftor was administered on Day 6. To quantify the DDIs, the geometric

mean ratios of AUC or C_{max} with or without the presence of CYP3A4 modulators were calculated.

The assessment of DDI prediction success was based on whether predictions fall within a two-fold

range of the observed data.

2.4. Model Application: DDI Predictions of ETI with Ritonavir

Although ritonavir-nirmatrelvir (PAXLOVID) is a fixed dose combination of 2 drugs, nirmatrelvir

has not been included in simulations as there is no clinical evidence that it modulates CYP3A4

activity, and we have focused on interactions with ritonavir which is a potent CYP3A4 inhibitor. The

verified PBPK-DDI model was applied to (1) predict the effect of ritonavir on the PK of ETI and (2)

determine a potential dose alteration of ETI to overcome the CYP3A inhibition mediated by

ritonavir. We first simulated the steady-state PK of standard dose ETI alone and when co-

administered with 100mg ritonavir twice daily for 5 days based on the instruction for dosage and

administration in the FDA-approved fact sheet of nirmatrelvir-ritonavir (5). In addition, since

ritonavir acts as a mechanism-based CYP3A4 inhibitor by covalently binding to the CYP3A4(23),

simulations were run until CYP3A4 and the PK of ETI had returned to baseline (10 days after

ritonavir discontinuation). We then simulated several adjusted dosing regimens of ETI when co-

administered with ritonavir to find the regimen that could provide the closest PK profiles of standard

dosing of ETI alone.

3. Results

3.1. Development and Verification of the Models of ETI

1) Parameter Sensitivity Analysis of the Fractional Metabolism of Ivacaftor by CYP3A4 on the

Predicted AUC Ratio with CYP3A Modulators

For the PBPK model of ivacaftor, in the absence of an *in vitro* estimate, clinical interaction data with

strong modulators of CYP3A4 were used to assign fmCYP3A4. First, we predicted DDI with

ketoconazole by varying the fmCYP3A4 value of ivacaftor from 95% to 100% (**Figure S1A**), since

it has been reported that the fractional metabolism of ivacaftor assigned to CYP3A4 is greater than

95%(17). An fmCYP3A4 of 98 % predicted the AUC ratio (GMR 6.95) of ivacaftor within the

bioequivalence limit (80-125%) of the observed AUC ratio (GMR 8.45). Repeating this analysis for

rifampicin, indicated the magnitude of DDI was well captured with fmCYP3A4 between 98 and

100% (simulated GMR 0.11 vs. observed 0.11) (**Figure S2B**). Taken together, the fmCYP3A4 value

of 98% was chosen as it describes observed DDIs between ivacaftor and ketoconazole or rifampin

within the bioequivalence limit.

2) PBPK Models of ETI Recapitulated Clinically Observed PK Profiles.

Model predictive performance of ETI was assessed using observed pharmacokinetic data sets from

clinical trials(12, 13). The observed and simulated plasma concentration-time profiles of ETI

following a single oral dose administration of elexacaftor 200mg, tezacaftor 100mg, and ivacaftor

100mg in healthy subjects are shown in **Figure S2**. For elexacaftor and tezacaftor, the mean plasma

concentrations were used in the graph while the median plasma concentrations were used for

ivacaftor as median value was reported from ivacaftor single dose PK study. The pharmacokinetic

profile of ETI after single oral dose administration was captured by the PBPK model.

Further verification of the model was performed by simulating the steady-state PK of ETI when

administered with multiple oral doses of elexacaftor 200mg qd, tezacaftor 100mg qd, and ivacaftor

150mg q12h. The predicted steady-state AUC and C_{max} of ETI were in the range of 0.9 to 1.2 of the

observed values demonstrating the excellent performance of the model. The observed and simulated

PK parameters of ETI are summarized in **Table 1**.

Table 1. Comparison of PK parameters between simulated and observed data for model verification of ETI

PK study			Steady-state PK parameters			
			Simulated		Observed	
Drug	Regimen		Cmax (mg/L)	AUC* (mg·h/L)	Cmax (mg/L)	AUC* (mg·h/L)
elexacaftor	200mg qd	Mean	8.1	158.0	8.8	167.0
		CV(%)	40.0	45.7	24.6	30.2
		Simulated/ observed	0.9	0.9		
tezacaftor	100mg qd	Mean	8.3	114.0	6.7	92.4
		CV(%)	38.8	49.9	20.8	25.8
		Simulated/ observed	1.2	1.2		
ivacaftor	150 mg q12h	Mean	1.6	13.4	1.3	12.1
		CV(%)	51.4	61.2	27.8	34.5
		Simulated/ observed	1.2	1.1		

* AUC(0-24h) for elexacaftor and tezacaftor, and AUC(0-12h) for ivacaftor.

3) PBPK-DDI Models of ETI Recapitulated Clinically Observed Drug Interactions

Although preliminary PK simulations verified the predicted PK ETI, given that the PBPK models of ETI are intended to be applied for the characterization of DDIs involving CYP3A modulation, it is essential to verify the victim properties defined in the models by simulating independent clinical DDI studies with a range of perpetrator drugs. The robustness of the model was assessed by comparing

the magnitude of simulated DDIs of ETI with that observed from the clinical trials. The PBPK-DDI

models accurately recapitulated the observed DDI magnitude (**Table 2**).

Table 2. Summary of the simulated vs. observed Geometric Mean Ratio (GMR) of PK parameters in
the presence and absence of CYP3A modulators

DDI study	PK Parameters	Simulated GMR (90% CI)	Observed GMR (90% CI)	Ratio (Simulated/ Observed)
Ivacaftor +/- Ritonavir	Cmax Ratio	2.61 (2.47, 2.77)	2.28 (1.84, 2.83)	1.14
	AUC Ratio	3.64 (3.37, 3.94)	3.06 (2.36, 3.97)	1.19
Ivacaftor +/- Ketoconazole	Cmax Ratio	2.04 (1.96, 2.12)	2.65 (2.21, 3.18)	0.77
	AUC Ratio	6.95 (6.44, 7.49)	8.45 (7.14, 10.01)	0.82
Ivacaftor +/- Fluconazole	Cmax Ratio	2.72 (2.64, 2.81)	2.47 (1.93, 3.17)	1.10
	AUC Ratio	3.33 (3.21, 3.46)	2.95 (2.27, 3.82)	1.13
Ivacaftor +/- Rifampin	Cmax Ratio	0.29 (0.27, 0.31)	0.20 (0.17, 0.24)	1.45
	AUC Ratio	0.11 (0.10, 0.13)	0.11 (0.10, 0.14)	1.00
Tezacaftor +/- Itraconazole	Cmax Ratio	2.62 (2.53, 2.72)	2.83 (2.62, 3.07)	0.93
	AUC Ratio	3.85 (3.65, 4.07)	4.02 (3.71, 4.63)	0.96
Elexacaftor +/- Itraconazole	Cmax Ratio	1.08 (1.08, 1.09)	1.05 (0.98, 1.13)	1.03
	AUC Ratio	2.00 (1.94, 2.06)	2.83 (2.59, 3.10)	0.71

3.2. DDI Simulation of ETI with Ritonavir

1) Simulated DDI of ETI and Ritonavir Suggests Significant DDI with ETI

The verified PBPK-DDI models of ETI were used to simulate the standard dose of ETI when co-administered with nirmatrelvir/ritonavir to determine the magnitude of the DDI for its intended use for treatment of COVID-19. To mimic the clinical setting of nirmatrelvir/ritonavir administration, we simulated steady state PK of ETI, then ritonavir q12h for 5 days while continuing ETI standard dosing during and after ritonavir administrations. We calculated the C_{max} and AUC ratio of ETI in the presence and absence of ritonavir on day 6 of co-administration (**Table S4**). The magnitude of DDI achieves its maximum level on day 6, as ETI haven't gotten to the steady-state after ritonavir administration, so drugs are still accumulating after the last dose of ritonavir (**Figure 2A, B, and C**). Further, the maximum CYP3A4 inhibition effect is maintained through day 6 (**Figure 2E**). The simulated geometric mean AUC ratio was highest for ivacaftor (9.31, 90% CI: 8.28, 10.47), followed by tezacaftor (3.11, 90% CI: 2.96, 3.27) and elexacaftor (2.31, 90% CI: 2.20, 2.42). Plasma concentrations of ETI in the presence and absence of ritonavir is shown in **Figure 2**. Although ritonavir itself is eliminated the day after discontinuation (**Figure 2D**), the CYP3A4 inhibition is time-dependent(24), so the inhibition is prolonged and the recovery to baseline time is reliant on the turnover of the CYP3A4 itself (**Figure 2E**). Thus, baseline steady state of all ETI drugs is predicted to be re-established on day 15. Crucially, this indicates that dose adjustment of ETI in case of co-administration with nirmatrelvir/ritonavir would be required to extend beyond the 5 days of co-administration.

Figure 2. Plasma Concentration Profile of Elexacaftor (A), Tezacaftor (B), Ivacaftor (C), and Ritonavir (D), and the % of Active CYP3A4 Enzyme (E) Over Time. Green: without ritonavir, Red: with ritonavir administered day 1 through day 5.

2) Altered Dose of ETI to Recapitulate the PK profile of Standard Dose ETI Alone

We next utilized the models to simulate ETI dose adjustments when these agents are co-administered

with ritonavir and determine how long the adjusted dosage needed to be maintained, to overcome the

enzyme inhibition effect mediated by ritonavir. Based on the simulated effects of ritonavir,

elexacaftor 200mg, tezacaftor 100mg, ivacaftor 150mg in the morning (2 orange tablets) every 4

days (administered on day 1 and day 5 and resumed full dose on day 9) provided similar steady-state

PK profile of the conventional regimen of ETI alone (**Figure 3**). The trough concentrations of ETI

were all above the EC50 targets, which are 0.99 mg/L, 0.5 mg/L, and 0.048 mg/L for elexacaftor,

tezacaftor, and ivacaftor, respectively(11-13). Since CYP3A4 inhibition dynamics mediated by

ritonavir changes over time, we measured the mean C_{max} and AUC of reduced dosing of ETI

regarding the first dose on day 1 and the second dose on day 5 and calculated the percentage of ETI

standard regimen alone (**Table 3**). The AUC(0-96h) of reduced dosing regimen ranged from 83.0-

142.5% of ETI alone. Resumption of the full dose of ETI on day 9 is based on simulations to

optimize the concentration profiles of all components of ETI, where the level of elexacaftor and

tezacaftor do not become lower than 80% of the standard regimen before resuming the full dose,

while striving to maintain levels of ivacaftor below 125% of the standard regimen after resuming the

full dose. At day 9, the CYP3A4 enzyme activities were recovered to 60% of the steady-state values.

Figure 3. Plasma Concentration Profile of ETI. Green: standard dose without ritonavir, Red: reduced

dose with ritonavir 150mg q12h administered day 1 through day 5. (EC50 for tezacaftor and

ivacaftor: obtained from exposure-response analysis in clinical trials regarding the reduction of sweat

chloride, EC50 for elexacaftor: obtained from in vitro study of chloride transport in

phe508del/phe508del human bronchial epithelial cells as no in vivo data are available.)

Table 3. Predicted mean Cmax and AUC of reduced dose of ETI (two orange tablets q96h instead of

two orange tablets in the morning and one blue tablet in the evening) with ritonavir 150mg q12h

administered day 1 through day 5

Drug regimen with ritonavir administered day 1 through 5			Cmax and % of standard dose ETI alone		AUC and % of standard dose ETI alone	
Drug	Regimen	Days	Cmax (mg/L)	% of ETI alone	AUC* (mg·h/L)	% of ETI alone
Elexacaftor	200mg on day 1	Day 1-2	8.7	107.4	185.9	117.7
		Day 1-5			605.3	95.8
	200mg on day 5	Day 5-6	7.9	97.5	168.9	106.9
		Day 5-9			524.4	83.0
Tezacaftor	100mg on day 1	Day 1-2	8.7	104.8	158.2	138.8
		Day 1-5			451.0	98.9

	100mg on day 5	Day 5-6	9.2	110.8	165.6	145.3
		Day 5-9			426.8	93.6
Ivacaftor	150mg on day 1	Day 1-2	1.8	112.5	35.9	134.0
		Day 1-5			125.2	116.8
	150mg on day 5	Day 5-6	2.7	168.8	54.4	203.0
		Day 5-9			152.8	142.5

*AUC(0-24h) for day 1-2 and day 5-6, AUC(0-96h) for day 1-5 and day 5-9.

In addition, we simulated an alternate dosing regimen, which is elexacaftor 100mg, tezacaftor 50mg,

ivacaftor 75mg in the morning (1 orange tablet) administered every 2 days. This regimen provided

concentration profiles closer to the standard regimen of ETI alone with less fluctuations of

peak/trough concentrations (**Figure S3 and Table S5**). Especially for ivacaftor on day 5 which

showed higher C_{max} (2.7 mg/L, 168.8% of standard regimen) in case of 150mg q96h, the C_{max} of

ivacaftor was 2.1 mg/L (131.3 % of standard regimen) with the 75mg q48h. However, the dosing

regimen of two orange tablets every 3-4 days is consistent with recommendations for other strong

CYP3A inhibitors and the C_{max} and AUC values between the two regimens were not demonstrably

different.

For the dosing recommendation of tezacaftor/ivacaftor (SYMDEKO), since it is provided as a fixed

dose yellow tablet consisting of tezacaftor 100mg and ivacaftor 150mg, the same dosing

recommendation above (tezacaftor-ivacaftor 100-150mg q96h) can be applied. Also, for ivacaftor

150mg tablet (KALYDECO), the same dosing recommendation (one tablet q96h) can be applied, but

alternatively, the dosing interval of ivacaftor could be further increased to 5 days rather than 4 days,

to recapitulate similar PK profile of standard regimen. When ivacaftor 150mg was administered on

day 6 instead of day 5, the AUC(0-24h) was decreased to 48.31 mg·h/L (180.2% of standard

regimen) from 54.4 mg·h/L (203.0% of standard regimen) (**Table S6**). Taken together, the suggested

dosing schedule of CFTR modulators co-administered with nirmatrelvir/ritonavir is described in

Figure 4.

Figure 4. Suggested Dosing Schedule of CFTR modulators co-administered with nirmatrelvir-

ritonavir

4. Discussion

All three components of ETI are eliminated predominantly through hepatic metabolism along with

limited renal excretion. The clinical DDI study with strong CYP3A4 inhibitors (Ketoconazole and

Itraconazole) or inducer (Rifampin) showed that ETI are the sensitive CYP3A4 substrates.

Therefore, the safe and effective use of CFTR modulators is complicated by DDI management with

concomitant CF medications, as CYP3A4 modulation by inducers or inhibitors can lead to altered

systemic exposure, resulting in variability in drug response. CF patients often take multiple

antibiotics including rifamycins, macrolides, and azole antifungals, which potentially inhibit or

induce CYP3A4-mediated metabolism of ETI. Recently Tsai et. al. published a ETI PBPK model to

evaluate exposures during the transition from mono or dual combination of CFTR modulators to

ETI(17). We extended the models by refining the ivacaftor model and further validating the ETI

PBPK-DDI model with published clinical DDI data. Since the ETI PBPK-DDI model we employed

could robustly predict PK parameters and the observed drug interactions of ETI, it can provide an

approach to the evaluation and management of other potential DDIs involving CFTR modulators.

In particular, we aimed to provide guidance for ETI dose adjustment with ritonavir, the CYP3A

inhibitor and the component of nirmatrelvir/ritonavir for the treatment of COVID-19. From the ETI-

ritonavir DDI simulations, we found that when ritonavir 100mg q12h was administered for 5 days, it

led to the AUC ratio of ivacaftor as 9.31 (90% CI: 8.28, 10.47) which far exceeded the observed and

simulated AUC ratio (3.06) when ivacaftor was administered with ritonavir 50 mg q24h(22). The

increase in interaction with the therapeutic regimen shows that dose adjustments should not be

estimated purely on the basis of the available clinical study. Further, through the simulations we

found that the elevated concentrations of ETI were continued even after ritonavir is eliminated due to

the irreversible CYP3A4 inhibition effect. The suggested reduced dosing regimen that is maintained

until 4 days after ritonavir is discontinued provided a similar PK profile of the standard regimen of

ETI alone. However, dose-dependent adverse reactions of ivacaftor should be more closely

monitored due to its high peak concentrations on day 5. This study has important clinical

implications, bridging the gap between the available clinical DDI study and that of the therapeutic

regimen and providing proper dosing guidelines for treatment of COVID-19 with

nirmatrelvir/ritonavir in pwCF receiving concomitant CFTR modulator therapy.

A limitation of this study is that in the absence of data, population system parameters, such as

plasma protein levels in pwCF was not incorporated into the modelling. However, changes in

demography reflecting CF population were incorporated. Furthermore, a prior study evaluating the

hepatic clearance of drugs showing that CYP3A enzyme activity is unaffected in pwCF(25) and the

current weight of evidence based on comparisons of ETI PK in healthy volunteers compared to

patients suggests they are comparable(11-13). Previous studies indicate that differences in

pharmacokinetics of drugs in CF is attributed to differences in body composition and plasma protein

concentrations secondary to nutritional deficiencies(26). However, the BMI of pwCF has increased

over the years with continued improvements in CF-care, including highly effective CF modulators

and nutritional support(9), to the extent it is now similar to that of healthy volunteers(10).

   A potential limitation of this work is that we did not include models of all active metabolites of

ETI. M1-tezacaftor is an important metabolite due to its similar potency with parent drug as well as

its high metabolite to parent AUC ratio (157.8%)(12). Since M1-tezacaftor is also metabolized by

CYP3A4(12), its formation and elimination may be altered with ritonavir co-administration. Using

the PBPK model of M1-tezacaftor, we were able to determine that the reduced dose of ETI provided

mean AUC(0-96h) for M1-tezacaftor during the co-administration period with ritonavir that was

80.9% of the standard regimen of ETI alone. We did not include models of active metabolites of

ivacaftor (M1-ivacaftor) and elexacaftor (M23-elexacaftor), since there was insufficient information

to build the models incorporating them. While it is currently unknown whether ritonavir would alter

the levels of M23-elexacaftor, the exposure of this metabolite is significantly reduced when

compared with the parent compound and is not considered to contribute significantly to the overall

efficacy(13). For M1-ivacaftor, there is evidence that its plasma concentration of is decreased by

35% in the presence of rifampin, indicating that M1-ivacaftor may also metabolized by

CYP3A4(27). This perhaps suggests that the reduced dose of ivacaftor may not significantly reduce

M1-ivacaftor exposure when co-administered with ritonavir, since ritonavir will inhibit the

metabolism of both ivacaftor and M1-ivacaftor.

   In conclusion, using a PBPK modeling approach, we demonstrated that nirmatrelvir/ritonavir can

be administered concomitantly with ETI in pwCF with proper dose adjustment. The outcome of this

study ensures the use of nirmatrelvir/ritonavir for the treatment of COVID-19 in patients with CF

while continuing to receive highly active CFTR modulators. In addition, this work provides tools to

evaluate and potentially overcome clinically important DDIs involving highly active CFTR

modulator therapy.

References

(1) Heijerman, H.G.M. *et al.* Efficacy and safety of the elexacaftor plus tezacaftor plus ivacaftor combination regimen in people with cystic fibrosis homozygous for the F508del mutation: a double-blind, randomised, phase 3 trial. *Lancet* **394**, 1940-8 (2019).

(2) Garg, V. *et al.* Pharmacokinetic and Drug-Drug Interaction Profiles of the Combination of Tezacaftor/Ivacaftor. *Clinical and translational science* **12**, (2019).

(3) Colombo, C. *et al.* Impact of COVID-19 on people with cystic fibrosis. *The Lancet Respiratory medicine* **8**, (2020).

(4) Ye, Q., Wang, B. & Mao, J. The pathogenesis and treatment of the `Cytokine Storm' in COVID-19. *The Journal of infection* **80**, (2020).

(5) PAXLOVID [FACT SHEET FOR HEALTHCARE PROVIDERS: EMERGENCY USE AUTHORIZATION FOR PAXLOVID], N.Y., NY, Pfizer, DEC 2021.

(6) Wagner, C., Pan, Y., Hsu, V., Sinha, V. & Zhao, P. Predicting the Effect of CYP3A Inducers on the Pharmacokinetics of Substrate Drugs Using Physiologically Based Pharmacokinetic (PBPK) Modeling: An Analysis of PBPK Submissions to the US FDA. *Clinical pharmacokinetics* **55**, (2016).

(7) Zhao, P. *et al.* Applications of physiologically based pharmacokinetic (PBPK) modeling and simulation during regulatory review. *Clinical pharmacology and therapeutics* **89**, (2011).

(8) *US Food and Drug Administration. Physiologically Based Pharmacokinetic Analyses — Format and Content Guidance for Industry. FDA.* <https://www.fda.gov/media/101469/download> (2018).

(9) Cystic Fibrosis Foundation Patient Registry 2020 Annual Data Report, Bethesda, Maryland ©2021 Cystic Fibrosis Foundation.

(10) Courville, A. *et al.* Increase in body mass index from normal weight to overweight in a cross-sectional sample of healthy research volunteers. *Nutrition research (New York, NY)* **34**, (2014).

(11) *U.S. Food and Drug Administration. Center for Drug Evaluation and Research. Clinical Pharmacology and Biopharmaceutics review(s), Ivacaftor. Available online: https://www.accessdata.fda.gov/drugsatfda_docs/nda/2012/203188Orig1s000ClinPharmR.pdf.*

(12) *U.S. Food and Drug Administration. Center for Drug Evaluation and Research. Clinical Pharmacology and Biopharmaceutics review(s), Tezacaftor/Ivacaftor. Available online: https://www.accessdata.fda.gov/drugsatfda_docs/nda/2018/210491Orig1s000ClinPharmR.pdf.*

(13) *U.S. Food and Drug Administration. Center for Drug Evaluation and Research. Multi-Discipline review, Elexacaftor/ Tezacaftor/Ivacaftor. Available online: https://www.accessdata.fda.gov/drugsatfda_docs/nda/2019/212273Orig1s000MultidisciplineR.pdf.*

(14) Jamei, M. *et al.* Population-based mechanistic prediction of oral drug absorption. *The AAPS journal* **11**, (2009).

(15) Sager, J., Yu, J., Ragueneau-Majlessi, I. & Isoherranen, N. Physiologically Based Pharmacokinetic (PBPK) Modeling and Simulation Approaches: A Systematic Review of Published Models, Applications, and Model Verification. *Drug metabolism and disposition: the biological fate of chemicals* **43**, (2015).

(16) *Australian Register of Therapeutic Goods, AusPAR Attachment 1: Product Information for lumacaftor ivacaftor (Orkambi 200/125).* <https://www.tga.gov.au/sites/default/files/auspar-lumacaftor-ivacaftor-160908-pi.pdf> (2016).

(17) Tsai, A. *et al.* Physiologically Based Pharmacokinetic Modeling of CFTR Modulation in People with Cystic Fibrosis Transitioning from Mono or Dual Regimens to Triple-

435   Combination Elexacaftor/Tezacaftor/Ivacaftor. *Pulmonary therapy*, 1-12 (2020).

436 (18) Matthes, E. *et al.* Low free drug concentration prevents inhibition of F508del CFTR
437   functional expression by the potentiator VX-770 (ivacaftor). *British journal of pharmacology*
438   **173**, (2016).

439 (19) *Kalydeco [package insert]. Cambridge, MA: Vertex Pharmaceuticals Inc.;.*
440   <https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/203188s022l_207925s003lbl.p
441   df> (2012).

442 (20) Galetin, A., Ito, K., Hallifax, D. & Houston, J.B. CYP3A4 substrate selection and substitution
443   in the prediction of potential drug-drug interactions. *The Journal of pharmacology and*
444   *experimental therapeutics*  **314**, (2005).

445 (21) Wang, Y.H. Confidence assessment of the Simcyp time-based approach and a static
446   mathematical model in predicting clinical drug-drug interactions for mechanism-based
447   CYP3A inhibitors. *Drug metabolism and disposition: the biological fate of chemicals*  **38**,
448   (2010).

449 (22) Liddy, A.M., McLaughlin, G., Schmitz, S., D'Arcy, D.M. & Barry, M.G. The
450   pharmacokinetic interaction between ivacaftor and ritonavir in healthy volunteers. In: *Br J*
451   *Clin Pharmacol*, Vol. 83 2235-41 (2017).

452 (23) Rock, B., Hengel, S., Rock, D., Wienkers, L. & Kunze, K. Characterization of ritonavir-
453   mediated inactivation of cytochrome P450 3A4. *Molecular pharmacology*  **86**, (2014).

454 (24) Katzenmaier, S., Markert, C., Riedel, K., Burhenne, J., Haefeli, W. & Mikus, G. Determining
455   the time course of CYP3A inhibition by potent reversible and irreversible CYP3A inhibitors
456   using A limited sampling strategy. *Clinical pharmacology and therapeutics*  **90**, (2011).

457 (25) De Sutter, P.J. *et al.* Pharmacokinetics in Patients with Cystic Fibrosis: A Systematic Review
458   of Data Published Between 1999 and 2019. *Clinical pharmacokinetics*, 1-23 (2020).

459 (26) Shah, N. *et al.* Novel Population Pharmacokinetic Approach to Explain the Differences
460   between Cystic Fibrosis Patients and Healthy Volunteers via Protein Binding. *Pharmaceutics*
461   **11**, (2019).

462 (27) *AusPAR Attachment 2 Extract from the Clinical Evaluation Report for Lumacaftor / Ivacaftor,*
463   *2015*.