**Long title: Childhood vaccination coverage by socioeconomic, geographic,**

**maternal, child, and healthcare characteristics in Nigeria: equity impact**

**analysis**

**Short title: Inequities in childhood vaccination coverage by socioeconomic**

**characteristics in Nigeria**

Sarah V Williams[1,3*], Tanimola Akande[2], Kaja Abbas[3]

[1] Royal Free London, NHS Foundation Trust, NHS Foundation Trust, Pond Street, London, NW3 2QG,

United Kingdom

[2] Department of Epidemiology & Community Health, University of Ilorin, Nigeria

[3] Department of Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine,

Keppel Street, London, WC1E 7HT United Kingdom

*Corresponding author: sarah.williams122@nhs.net

## Abstract

**Background:** Nigeria has a high proportion of the world's underimmunised children. We estimated the inequities in childhood immunisation coverage associated with socioeconomic, geographic, maternal, child, and healthcare characteristics among children aged 12-23 months in Nigeria using a social determinants of health perspective.

**Methods:** We first conducted a systematic review focused on inequities in childhood immunisation in low- and middle-income countries to identify the social determinants of childhood immunisation. To identify inequities in basic vaccination coverage in Nigeria (1-dose BCG, 3-dose DTP-HepB-Hib (diphtheria, tetanus, pertussis, hepatitis B and Haemophilus influenzae type B), 3-dose polio, and 1-dose measles) of 6,059 children aged 12-23 months using the 2018 Nigeria Demographic and Health Survey, we conducted multiple logistic regression to analyse the association of basic vaccination with these determinants.

**Results:** We identified social determinants of immunisation: household wealth, religion, and ethnicity (socioeconomic characteristics); region and place of residence (geographic characteristics); maternal age at birth, maternal education, and maternal household head status (maternal characteristics); sex of child, and birth order (child characteristics); antenatal care and birth setting (healthcare characteristics). While basic vaccination coverage was 31% (95% CI: 29–33), 19% (95% CI:18–21) of children aged 12-23 months had received none of the basic vaccines. After controlling for background characteristics, there was a significant increase in the odds of basic vaccination by household wealth, antenatal care, delivery in a health facility, maternal age and maternal education. Children of Fulani ethnicity in comparison to children of Igbo ethnicity had lower odds of receiving basic vaccinations.

**Conclusions:** Children from the poorest households, of Fulani ethnicity, who were born in home settings, and with young mothers with no formal education nor antenatal care, were associated with

43    lower odds of basic vaccination in Nigeria. Basic vaccination coverage is below target levels for all

44    groups and we recommend a proportionate universalism approach for addressing the immunisation

45    barriers.

46    **Keywords:** Childhood immunisation, vaccine equity, social determinants of immunisation,

47    demographic and health survey, Nigeria

## Introduction

Nigeria is the most populous country in Africa with around 202 million people in 2020 and its

population is predicted to double by 2050 [1]. It is a multi-ethnic country with 36 autonomous states

and the Federal Capital Territory. Around 83 million people (40% of total population) live below the

poverty line while an additional 53 million people (25% of total population) are vulnerable to falling

below the poverty line [2]. Economic growth has been slow with challenges including ongoing

conflict in parts of the country, inconsistent regulatory environment, poor power supply and

infrastructure  [2].

Vaccination is a highly cost-effective public health intervention and beyond the direct benefits to

population health, vaccines provide additional economic and social benefits to individuals and

society [3,4]. Infectious diseases remain a leading cause of death among under-5-year-old children,

and an additional 1.5 million deaths could be avoided every year with improvements in global

vaccination coverage [5,6]. Model-based estimates, not including COVID-19, project 51 million

deaths to be prevented by vaccination during 2021-2030 [7]. There have been substantial

improvements in vaccine introductions and vaccination coverage in low- and middle-income

countries since the inception of Gavi, the Vaccine Alliance in 2000 [8]. However, the prevalence of

zero-dose children, that is children aged 12-23 months who had not received any of the routine

childhood vaccines, was 7.7% in low- and middle-income countries during 2010-2019 [9]. The

importance of improving vaccination coverage was recognised in the Sustainable Development Goals

(SDGs), with immunisation contributing to 14 of the 17 SDGs and includes reduction on poverty and

hunger and improving social equity [10]. Vaccination coverage and equity are a strategic goal of the

global Immunisation Agenda 2030, with the aim to reach equitable coverage at national and district

levels by addressing immunisation barriers posed by location, age, socioeconomic status, and gender

[11].

The Expanded Programme on Immunisation (EPI) was established by the World Health Organization

(WHO) in 1974 to improve vaccination services globally [12], and Nigeria began nationwide

implementation of EPI in 1979, [13] which was later changed to the National Programme on

Immunization. Although the vaccines in the routine immunisation programme (Table A1) for under

5-year-old children are available with no out-of-pocket charges [14], Nigeria has the most under-

immunised children in the world with 4.5 million in 2018 [15]. The immunisation system challenges

in Nigeria include weak institutions, service delivery, funding, infrastructure, poor coordination

between the National Programme on Immunization and non-governmental organisations delivering

vaccination services, and a lack of political commitment in some regions [14,16]. There are fewer

adequately skilled healthcare personnel in rural areas and northern states, and poor retention and

frequent transfers of workers. Security is also an issue, with attacks on healthcare workers in recent

years. Attitudes of communities and caregivers are important too, with a lack of knowledge about

vaccination and mistrust of services hindering vaccination uptake [16,17].

In the context of wider immunisation system challenges in Nigeria, we focused on factors associated

with inequities in basic vaccination coverage through the social determinants of health model. This

model framework has been explicitly linked to health equity by the WHO Commission on Social

Determinants of Health [18] and considers the social, cultural, political, economic, commercial and

environmental factors that shape the conditions in which people are born, grow, live, work and age,

and these factors are determined by wealth, power and resources. In this study, we use the social

determinants of health model framework, which encompasses the individual, parental, household,

environment, and national policy levels that influence inequities in basic vaccination coverage

among children in Nigeria (see Figure 1). We refer to vaccine inequity as unfair and avoidable or

remediable differences in health among population groups defined socially, economically,

demographically, or geographically [19]. This is related to but distinct from health inequality, which

indicates the status of imbalances or differences in health among population groups without any

moral judgement on whether the imbalances or differences are fair or not [20,21].

Our aim is to analyse the 2018 Nigeria Demographic and Health Survey (DHS) and estimate the

inequities in basic vaccination coverage (1-dose BCG, 3-dose DTP-HepB-Hib, 3-dose polio, and 1-dose

measles vaccines) associated with socioeconomic, geographic, maternal, child, and healthcare

characteristics among children aged 12-23 months in Nigeria.

*Figure 1 placement*

**Figure 1. Social determinants of childhood immunisation.** Social determinants of health model

framework encompassing the individual, parental, household, environment, and national policy

levels and influencing inequities in basic vaccination coverage among children in Nigeria.

## Methods

### Characteristics selection and systematic review

The WHO had analysed inequality in immunisation using the following characteristics determined

from a literature review: child (gender, birth order), maternal (age at birth, education, ethnicity or

caste), household (sex of household head, household economic status), geographic (rural or urban

place of residence, subnational region) [22]. Using the WHO inequality work as a starting point, we

conducted a systematic review to select the pertinent DHS variables for our inequity analysis. We

conducted a systematic review to analyse qualitative and quantitative peer-reviewed literature of

studies focused on inequities in childhood immunisation in low- and middle-income countries

(LMICs) to identify the social determinants of childhood immunisation. Figure A2 (in appendix)

illustrates the process flow diagram of identification, screening, eligibility, and inclusion of articles

for the systematic review, using the PRISMA (Preferred Reporting Items for Systematic Reviews and

Meta-Analyses) framework [23]. We conducted our search using MEDLINE and additional studies

were identified through hand-searches of reference lists for articles written in the English language,

published between 01/01/2010 to 04/10/2021, for which the full text was available, and contained

the following terms in English or American spellings: (Vaccination coverage or Immunisation

coverage) AND (Determinant, characteristic, predictor) AND (Equity, equality, disparity, inequality,

inequity). We did not search prior to 2010 as papers and the subsequently identified social

determinants may be less relevant to the current context. We identified 160 publications, screened

the title and abstract, assessed full articles for eligibility, and included 49 publications in our

systematic review (see Table 1). In addition to the characteristics analysed by the WHO inequality

report on immunisation [22], we identified that antenatal care and birth setting had evidence of

association with vaccination coverage.

**Table 1. Systematic review of social determinants of childhood immunisation.**

| Location | Key inferences | Source | |
|---|---|---|---|
| Afghanistan | Approximately 60% of children aged 1-4 years were under vaccinated or not vaccinated, with large disparities among provinces and higher coverage in urban areas. Increased odds of full immunisation were associated with giving birth in government institution, | Shenton et al., 2018 | [54] |

| | | | |
|---|---|---|---|
| | having a higher number of antenatal care visits, and visiting a health facility in the past 12 months. Maternal involvement in household decision making was also significantly associated with vaccination status. | | |
| Angola | In Bom Jesus in Angola, 37% of under 5-year-old children had completed the vaccination schedule, but coverage was higher for under 1-year-old children. Coverage was higher in rural areas. The percentage of children completing the vaccination schedule varied according to child age, mother's education, family size, ownership of household appliances, and means of disposal of domestic waste. | Oliveira et al., 2014 | [55] |
| Bangladesh | There was no significant disparity of vaccination coverage by ethnicity. There was significant variation of vaccination coverage by the child's gender (lower for females), household ownership of mobile phones and by household geographical location. | Rahman et al., 2018 | [56] |
| Bangladesh | Maternal age, education and wealth have a significant effect on coverage of most vaccines. The number of microfinancing | Vyas et al., 2019 | [57] |

| | organisations in the community also increased the odds of children being vaccinated, whilst poor accessibility or being far from a health centre, specifically an Upazilla Health Complex, decreased the odds of a child being vaccinated. There are differences between regions, but even in regions with relatively higher coverage, some communities had poor coverage. | | |
|---|---|---|---|
| Cameroon | Analysis of five rounds of DHS surveys from 1991 to 2001 showed that children in the latter rounds were more likely to be fully immunised. The likelihood of children being fully immunised was affected by having a vaccination card, birth order, maternal age and maternal education. | Nda'chi Deffo et al., 2020 | [58] |
| Democratic Republic of the Congo | In urban areas, children of educated mothers had higher measles vaccination coverage in comparison to children of mothers with no formal education. In rural areas, children of wealthier households had higher measles vaccination coverage. Vaccination card usage rates were higher in urban areas than rural areas. | Ashbaugh et al., 2018 | [59] |
| Ethiopia | Around 25% of children aged 12-23 months | Lakew et al., 2015 | [60] |

| | | | |
|---|---|---|---|
| | were fully vaccinated and coverage varied with urbanicity. Predictors of full coverage were sources of information from the vaccination card, received postnatal check-up within two months after birth, region, women's awareness of community conversation program, and wealth index. | | |
| Ethiopia | Children from poorer households, rural regions of Afar and Somali, no maternal education, and female-headed households had lower full vaccination coverage. | Geweniger et al., 2020 | [61] |
| Ghana | Vaccination coverage at the end of the first year of life was high except for polio vaccine given at birth. However, many vaccines were given late and there was substantial health inequity across socioeconomic indicators for timeliness of vaccination. | Gram et al., 2014 | [62] |
| India | Slightly more than half of children aged 12-36 months were fully vaccinated in 2008. The analysis adjusted for state of residence, age, gender, household wealth, and maternal education, and other predictors of vaccination were religion, caste, place of delivery, number of antenatal care visits, and maternal tetanus vaccination. Children in urban areas had | Shrivastwa et al., 2015 | [63] |

| | higher odds of being unvaccinated or under-vaccinated than those in rural areas. | | |
|---|---|---|---|
| India | Vaccine coverage in an urban poor area of south east Delhi was lower than estimates from other studies for overall regional urban coverage, indicating a discrepancy in coverage between the urban poor and urban non-poor. Coverage was determined by gender of child, religion, maternal literacy, household's socioeconomic position, being born in a facility and being born outside of Delhi and having a birth certificate. | Devasenapathy et al., 2016 | [64] |
| India | There were regional disparities in vaccination coverage with a north south divide. All doses of vaccination coverage were significantly associated with postnatal care, institutional births, neonatal tetanus protection of the last birth, women's education, and health insurance coverage | Khan et al., 2018 | [65] |
| India | Approximately 50% of children aged 12-23 months were fully vaccinated in a pooled dataset of three time periods (1998–99, 2002–04 and 2007–08), and children recorded in the 2007-08 dataset were more likely to be vaccinated. Vaccination was inversely | Francis et al., 2018 | [66] |

| | | | |
|---|---|---|---|
| | associated with female gender, Muslim religion, lower caste, urban residence, and maternal characteristics (lower educational attainment, non-institutional delivery, fewer antenatal care visits and non-receipt of maternal tetanus vaccination). The most common reason for non-vaccination was that mothers were unaware of the need. | | |
| India | Immunisation was highly associated with maternal education and household wealth. Inequality in immunisation was highest in lower socioeconomic groups among children of mothers with no formal education. | Kannankeril Joseph VJ et al., 2021 | [67] |
| India | Full immunisation was associated with living in urban areas and richer household wealth, and was highest in Manipur and lowest in Nagaland. | Srivastava et al., 2021 | [68] |
| Indonesia | Immunisation was significantly associated with birth order, maternal age, maternal education, paternal occupation, antenatal care, and living in certain regions. Being resident in communities with a higher proportion of public health centres were more likely to be fully immunised. | Siramaneerat et al., 2021 | [69] |

| Kenya | Measles vaccination coverage differed according to household wealth, parents' education, skilled antenatal care visits, birth order and father's occupation. Rural residence reduced inequality in measles vaccination, which may reflect efforts to provide vaccination sites in remote and rural areas of the country. | Van Malderen et al., 2013 | [70] |
|---|---|---|---|
| Kenya | Birth setting, ethnicity, and wealth index were significant predictors of vaccination coverage. Somalis had greater odds of being under or non-vaccinated than the Kikuyu ethnic group, and wealth and birth setting were associated with immunisation status for both Somalis and non-Somalis. | Masters et al., 2019 | [71] |
| Kenya | Children of mothers with no education, born in home settings, in regions with limited health infrastructure, living in poorer households, and of higher birth order are associated with lower rates of full immunisation. | Allan et al., 2021 | [72] |
| Kenya | Immunisation coverage and timeliness of immunisation improved during 2003-2017. In two urban informal settlements, the hazard for being fully immunised varied by household | Mutua et al., 2020 | [73] |

| | wealth and ethnicity. | | |
|---|---|---|---|
| Laos | Community demand for immunisation was lowered by health system barriers of multiple providers, inconsistent record keeping, and inadequate health information system. Lack of understanding of the value of vaccination and immunisation services was demonstrated at the individual and household levels. | Sychareun et al., 2019 | [74] |
| Malawi | There were regional differences in vaccination coverage and regions with higher coverage had a high percentage of deliveries attended by a health professional. Coverage was higher for women who gave birth at a hospital or maternity clinic or had a midwife or nurse at birth. Coverage was significantly correlated with some socio-demographic characteristics: child's age, illiteracy, income, water, and sanitary conditions. | Abebe et al., 2012 | [75] |
| Malawi | In children aged 12-23 months, individual factors had a stronger effect than community factors on vaccination coverage.  Mother's education, frequency of antenatal care visits, use of immunisation cards, household wealth, and geographical region were the most significant factors associated with vaccination | Ntenda et al., 2017 | [76] |

| | coverage. | | |
|---|---|---|---|
| Mozambique | Analysis of data from the 2015 Immunization, AIDS and Malaria Indicators Survey (IMASIDA) indicated that living in the Southern region was a protective factor for full immunisation whilst being in the poorest quintile increased the risk of a child not being fully immunised. | Daca et al., 2020 | [77] |
| Nepal | A quarter of children were not fully immunised. Full immunisation was associated with having an immunisation card, delivery in a public or private institution, maternal education and maternal employment. | Patel et al., 2021 | [78] |
| Nigeria | Socio-economic characteristics explained the disparities in full vaccination and recommended community-level interventions focused on improving full vaccination coverage. | Antai, 2009 | [79] |
| Nigeria | Women who were sole providers of household earnings were associated with higher likelihood of fully immunising their children, while women who lack decision-making autonomy were associated with lower likelihood of fully immunising their children. | Antai, 2012 | [80] |

| Nigeria | Routine immunisation data shows lower coverage in states of northern Nigeria compared with states of southern Nigeria. When comparing determinants of supply side access in two northern and two southern states, regional supply-side disparities were not apparent. However, there was a general sub-optimal supply of services and residents in the northern states were more likely to live within 5km of an immunisation service. This supports socio-cultural explanations of disparities in immunisation coverage. | Eboreime et al., 2015 | [38] |
|---|---|---|---|
| Nigeria | Fully immunised children are concentrated amongst the rich and partially immunised children are concentrated amongst the poor, no significant difference between socioeconomic status for unimmunised. Concentration of fully immunised children among the rich were determined by mother's literacy, living in the rural area, socioeconomic status and geopolitical location. Significant concentration of partially immunised children among the poor were determined by the same factors. | Ataguba et al., 2016 | [81] |
| Nigeria | In a cross-sectional study in south-eastern Nigeria of under 5-year-old children, | Uzochukwu et al., 2017 | [82] |

| | vaccination coverage was poor and below national targets. There were differences in coverage between wealth quartiles with the wealthiest having higher coverage than the poorest. Coverage was highest for children in the first year of life and decreased with age. | | |
|---|---|---|---|
| Nigeria | Children unvaccinated against polio was significantly associated with household wealth index, maternal educational level, maternal employment, geopolitical and neighbourhood illiteracy level. | Uthman et al., 2017 | [83] |
| Nigeria | Coverage varied by place of residence (rural, urban formal, and urban slum) and was significantly associated with place of delivery, antenatal care, maternal education, maternal age at childbirth, religion, place of residence, media exposure and distance to the health facility. Population attributable risk analysis demonstrated the biggest increase in coverage for maternal education and biggest reduction for non-attendance at antenatal care. | Obanewa et al., 2020 | [84] |
| Pakistan | Vaccination coverage of children aged 12-23 months was strongly associated with maternal education, paternal education, extended | Siddiqui et al., 2014 | [85] |

| | | | |
|---|---|---|---|
| | family structure and family wealth. Coverage differed by ethnicity, with Bengali children having the lowest coverage. | | |
| East African countries | Most children in each country had received one of the recommended vaccinations, with regional variation within countries. There were no consistent predictors across countries of complete vaccination status. However, being delivered in a public institution rather than at home was associated with increased odds, except in Burundi. Whether a child had received a check-up within two months of birth was also associated with complete vaccination status in Burundi, Kenya, and Uganda. | Canavan et al., 2014 | [86] |
| West Africa | Vaccination coverage increased between 2000 and 2017. Inequalities in coverage were mainly related to poverty, maternal education and living in certain regions. | Wariri et al., 2019 | [87] |
| Sub-Saharan Africa | Poor households were more likely to have missed opportunities for vaccination. In some countries, literacy, maternal education, media access and the number of under 5-year-old children contributed to missed opportunities for vaccination. | Ndwandwe et al., 2018 | [88] |

| | | | |
|---|---|---|---|
| Sub-Saharan Africa | Education inequality in missed opportunities for vaccination was mainly explained by differential effects such as neighbourhood socioeconomic status, presence of under 5-year-old children, media access and household wealth index. | Sambala et al., 2018 | [89] |
| Sub-Saharan Africa | Vaccination coverage was lower in rural areas and household wealth, birth order and distance to health facilities were important contributors to the rural-urban gap. | Ameyaw et al., 2021 | [90] |
| Conflict-impacted countries | In conflict affected countries, low vaccination coverage and outbreaks of vaccine preventable diseases are a concern. Many outbreaks were in conflict-affected areas or in displaced populations. In 14 of the 16 conflict affected countries, DTP3 coverage was below the global average of 85% in 2014. | Grundy et al., 2019 | [50] |
| Gavi-supported countries | Following a systematic analysis of inequalities in vaccination coverage in 45 Gavi supported countries, recommendations were made for Gavi equity monitoring. In addition to the wealth index, other measures of vulnerability should be monitored: maternal education, place of residence, child sex and the multidimensional poverty index. Both | Arsenault et al., 2017 | [91] |

| | absolute and relative measures should be tracked. | | |
|---|---|---|---|
| Gavi-supported countries | DHS data from 45 Gavi supported countries demonstrated that vaccination coverage and inequalities varied between countries. There were wealth, education, and multidimensional poverty index (MPI) inequalities in vaccination coverage. Country level predictors of vaccination coverage included political stability, government expenditure on health, government effectiveness and control of corruption, greater land area, linguistic fractionalisation, and gender inequality. There were similar associations across the three indicators of inequality (wealth, maternal education and multidimensional poverty) for country level predictors of inequality. | Arsenault et al., 2017 | [92] |
| Low- and middle-income countries | By analysing data from 241 nationally representative household surveys in 96 countries, being unvaccinated was strongly associated with education of the caregiver, education of caregiver's partner, caregiver's tetanus toxoid status, wealth index and type of family member participation in decision-making when the child is ill. Tetanus toxoid | Bosch-Capblanch et al., 2012 | [93] |

| | | | |
|---|---|---|---|
| | status was the strongest predictor of unvaccinated status. | | |
| Low- and middle-income countries | Twenty-five studies were reviewed to assess the relevance of gender inequality on the structural, health system, community, and individual levels. Vaccination programmes often target mothers and can therefore reinforce a community's gender and social dynamics, with women facing barriers at every level to accessing vaccinations: access to education, income, as well as autonomous decision-making about time and resource allocation were evident barriers. | Merten et al., 2015 | [94] |
| Low- and middle-income countries | The greatest disparities were among children born to women with no formal education compared to children born to women with secondary or higher education. Coverage was lower in rural areas compared to urban areas and in the lowest wealth quintile compared to the richest quintile. No gender differences were observed. | Hinman et al., 2015 | [95] |
| Low- and middle-income countries | Rural-urban migrant children were less likely to be immunised than urban non-migrants and the general population. Coverage estimates were lower for most vaccines for | Awoh et al., 2016 | [96] |

| | rural-urban migrants than for the general population. | | |
|---|---|---|---|
| Low- and middle-income countries | Vaccination coverage varied between and within countries, and coverage was lowest in children from poorer households with pro-rich inequality in most countries. | Hosseinpoor et al., 2016 | [97] |
| Low- and middle-income countries | In most countries, vaccination coverage was pro-rich, and coverage was higher in urban areas than rural areas. Antenatal care was concentrated among wealthier mothers and was significantly associated with the concentration of vaccination coverage among wealthier children. | Hajizadeh et al., 2018 | [98] |
| Low- and middle-income countries | The WHO analysed inequality in immunisation using the following characteristics determined from a literature review: child (gender, birth order), maternal (age at birth, education, ethnicity or caste), household (sex of household head, household economic status), geographic (place of residence - rural or urban, subnational region). The analysis focused on ten countries that account for more than 70% of children globally who did not receive basic vaccination. | WHO, 2018 | [22] |

| Low- and middle-income countries | DHS data was examined for socioeconomic inequalities in the completion rates of vaccines in the Gambia, the Kyrgyz Republic and Namibia. Higher completion of vaccination was observed in rural areas. Completion rate for BCG, DTP3, polio (3-doses), and measles vaccines had a pro-poor distribution with children of lower socioeconomic status more likely to be immunised than children from higher socioeconomic status. In the Gambia and Namibia, the difference in completion rates for childhood immunisation between rural and urban areas was the primary contributor for the concentration of child vaccination among the poor. In the Kyrgyz Republic, household wealth was a key determinant for child vaccination. Birth order, maternal age, maternal education, and skilled birth attendance were also factors in some of the countries. | Hajizadeh,. 2019 | [99] |
|---|---|---|---|

Systematic review of studies focused on inequities in childhood immunisation in low- and middle-

income countries.

**Demographic and health survey**

We analysed the 2018 Nigeria DHS which was conducted between August to December 2018 [24].

The DHS are nationally representative household surveys focusing on population, health, and

nutrition in LMICs [25]. The DHS sample is a two-stage stratified cluster sample with sampling

weights applied to ensure that results are representative. There are four questionnaires: household

questionnaire, woman's questionnaire, man's questionnaire, and biomarker questionnaire. The

country is divided into clusters with 30 households selected from each cluster. The woman's

questionnaire was asked to women aged 15-49 years and provides the data for our study. All women

aged 15-49 years in the sampled households were included and the survey was successfully

conducted in 1,389 clusters after 11 clusters were dropped following deteriorating security in those

areas during data collection. In addition, in the state of Borno, 11 of the 27 Local Government Areas

(LGAs) in the state were dropped due to insecurity. Clusters selected from these dropped LGAs were

replaced with clusters from the remaining LGAs and so may not be representative of the entire state

[24]. The study population for our analysis was women aged 15-49 years with a child aged 12 to 23

months old. Of the 41,821 women interviewed, 33,924 women had a child aged 59 months or

younger and immunisation data was collected for 6,059 living children aged between 12-23 months.

We applied sampling weights to the survey dataset to adjust for disproportionate sampling and non-

response, thereby ensuring that the sample was representative of the population.

**Vaccination coverage**

The information of whether a child has received a vaccination is gathered from the child's

vaccination card. If that is not available or if a vaccine has not been recorded, then the mother is

asked which vaccines have been given to her child [24]. Our primary outcome and dependent

variable of interest was basic vaccination coverage, that is the proportion of children receiving 1-

dose BCG, 3-dose DTP-HepB-Hib, 3-dose polio, and 1-dose measles vaccines. DHS gives the

vaccination status of "received" and "not received" for individual vaccines. Binary variables of

"received all three doses" and "not received all three doses"were generated for the three dose

vaccines. The basic vaccination variable was generated as a binary variable of "received all basic

vaccinations" and "not received all basic vaccinations" by combining the variables for 1-dose BCG, 3-

dose DTP-HepB-Hib, 3-dose polio, and 1-dose measles vaccines.

**Inequity analysis**

We conducted simple logistic regression to estimate crude odds ratios and assess basic vaccination

coverage disaggregated by socioeconomic, geographic, maternal, child, and healthcare

characteristics as measured by the DHS and identified through the systematic review and the WHO

inequality report on immunisation [22]. Due to the survey design, a p-value for all categories within a

characteristic was determined using the Adjusted Wald's test through the "test" function in Stata

[26]. These p-values were used to assess the association between basic vaccination coverage and the

characteristics [27] and if a p-value was <0.05 then the characteristic was included in the model. We

checked for collinearity between variables and avoided multicollinearity in the development of a

parsimonious model [27] using the "regress" in Stata  and calculation of the variance inflation factor.

However, no variables required removal from the model.

Interaction was tested using the "contrast" function in Stata. This gives F statistics which are

adjusted for the survey design, with the p-value demonstrating the statistical significance of the

interaction. As in the WHO report [22,27], interactions were tested between mother's education and

household wealth, mother's age and household wealth, mother's age and education, and place of

residence and household wealth. We conducted multivariable logistic regression to estimate

adjusted odds ratios (AORs) for socioeconomic, geographic, maternal, child, and healthcare

characteristics associated with basic vaccination coverage (see Table 3 for the characteristics

included in the multivariable logistic regression and Tables A4 in the appendix for the strata specific

AORs between place of residence and household wealth).

We analysed inequity further by estimating the Erreygers concentration indices for maternal

education, antenatal care, and household wealth to assess if basic vaccination coverage and

vaccination card usage had progressive, regressive, or equal distribution based on each of these

characteristics. The concentration index value shows how much of a health measure is concentrated

in an advantaged or disadvantaged group. Values range from +1 to -1, with a value of zero meaning

there is no inequity and positive values indicating that a health measure is concentrated in the more

advantaged groups [28].

**Reproducibility of analysis**

We conducted the survey analysis using the Stata statistical software [26], and visualisations were

generated using the R statistical software [29]. The analysis code is publicly accessible on GitHub at

https://github.com/svwilliams122/vaccine-equity-Nigeria and the 2018 Nigeria DHS data set is

accessible upon registration on the DHS website at

https://dhsprogram.com/methodology/survey/survey-display-528.cfm.

# Results

Through the systematic review for characteristics selection (see Table 1), we streamlined the social

determinants of childhood immunisation to: household wealth, religion, and ethnicity for

socioeconomic characteristics; region and place of residence for geographic characteristics; maternal

age at birth, maternal education, and maternal household head status for maternal characteristics;

sex of child and birth order for child characteristics; and antenatal care and birth setting for

healthcare characteristics.

**Vaccination coverage**

Among the 6,059 children aged 12-23 months in the 2018 Nigeria DHS following the application of

sample weights, 2,100 (35%) of them lived in urban areas. The coverage for the individual vaccines

of BCG, DTP-HepB-Hib, polio, and measles was higher in urban areas compared to rural areas (Figure

2 and Table A3). At the national level, the mean basic vaccination coverage was 31% (95% CI: 29-33)

with coverage of single vaccinations ranging from 48% (46-50%) for the third dose of the polio

vaccine to 73% (71-75%) for the first dose of the polio vaccine.

*Figure 2 placement*

**Figure 2. Vaccination coverage and vaccination card usage rates in Nigeria.** Vaccination coverage

among children aged 12-23 months in Nigeria and disaggregated by urban and rural areas of

residence. Vaccination card coverage is relatively higher in urban areas in comparison to rural areas,

and is associated with higher vaccination coverage. Basic vaccination includes 1-dose BCG, 3-dose

DTP-HepB-Hib (diphtheria, tetanus, pertussis, hepatitis B and Haemophilus influenzae type B), 3-

dose polio, and 1-dose measles vaccines.

Vaccination cards were available for 49% (46-51%) of children, among which 57% (55-60%) had

received all basic vaccinations. Among the 51% (49-54%) of children without vaccination cards, only

6.4% (5.3-7.7%) of them had received all basic vaccinations. Almost one fifth of children (19% (18-

21%)) had not received any of the basic vaccinations and in rural areas almost a quarter had received

none. For polio vaccinations, 26% (24-28%) of children had received none of the three doses and for

DTP vaccinations, 35% (33-37%) of children had received none. There was a higher proportion of

zero-dose children in rural areas compared to urban areas.

**Inequity analysis**

Figure 3 shows the basic vaccination coverage among children aged 12-23 months in Nigeria

disaggregated by socioeconomic (household wealth, religion, ethnicity), geographic (region, place of

residence), maternal (maternal age at birth, maternal education, maternal household head status),

child (sex of child, birth order), and healthcare (antenatal care, birth setting) characteristics. For

socioeconomic characteristics, children living in wealthier households had higher basic vaccination

coverage ranging from 17% (15-20%) to 49% (45-53%) from the poorest to the richest wealth

quintiles. For religion, basic vaccination coverage was higher among children of the Catholic faith at

49% (43-54%) while most children (58%) were of Islamic faith with relatively lower coverage of 23%

(21-25%). Regarding ethnicity, basic vaccination coverage ranged from 12% (9-16%) to 56% (51-61%)

among children of Fulani and Igbo ethnicities respectively. For geographic characteristics, 65% of

children lived in rural areas but basic vaccination coverage was higher for children in urban areas at

44% (41-47%) in comparison to 23% (21-25%) in rural areas. At the regional level, basic vaccination

coverage among children ranged from 20% (17-23%) in the North West region to 57% (51-62%) in

the South East region (Figure 4). For maternal characteristics, children of mothers aged 35-49 years

had higher basic vaccination coverage at 34% (31-39%) in comparison to 16% (12-21%) for children

of younger mothers aged 15-19 years. Basic vaccination coverage increased with higher levels of

maternal education, with basic vaccination coverage among children of mothers with no education,

primary education, and secondary education or higher at 15% (13-17%), 33% (28-38%), and 47% (45-

50%) respectively. Children living in female-headed households had relatively higher basic

vaccination coverage of 39% (35-44%) in comparison to 30% (28-32%) in male-headed households.

For child characteristics, basic vaccination coverage was similar among female and male children at

31% (29-33%) and 31% (29-34%) respectively, while coverage decreased by birth order with 36% (32-

40%) and 23% (20-26%) among first-born and sixth-born respectively. For healthcare characteristics,

children of women who had a higher number of antenatal care visits during their pregnancy had

higher basic vaccination coverage at 41% (39-44%) for four or more visits and 11% (8.7-13%) for no

or unknown number of visits. More than half of all women gave birth at home, and basic vaccination

coverage among these children was relatively lower at 20% (18-22%) in comparison to 47% (44-49%)

among children born in a public or private clinical facility.

*Figure 3 placement*

**Figure 3. Basic vaccination coverage in Nigeria.** Basic vaccination coverage among children aged 12-

23 months in Nigeria by socioeconomic (household wealth, religion, ethnicity), geographic (region,

place of residence), maternal (maternal age at birth, maternal education, maternal household head

status), child (sex of child, birth order), and healthcare (birth setting, antenatal care) characteristics.

*Figure 4 placement*

**Figure 4. Basic vaccination coverage in Nigeria at the regional level.** Basic vaccination coverage

among children aged 12-23 months in Nigeria at the regional level.

Figure 5 shows the concentration curve for household wealth-related inequity in basic vaccination

coverage, while Table 2 presents the inequities in vaccination card usage rates and vaccination

coverage among children aged 12-23 months disaggregated by maternal education, antenatal care,

and household wealth in Nigeria, based on Erreygers concentration indices. With respect to

maternal education, antenatal care, and household wealth, each of these characteristics had a

regressive pro-advantage distribution, with higher vaccination card usage rates and coverage among

children from wealthier households, higher maternal education and more antenatal care.

*Figure 5 placement*

**Figure 5. Wealth-related inequity in basic vaccination coverage in Nigeria.** Concentration curve for

household wealth-related inequity in basic vaccination coverage among children aged 12-23 months

in Nigeria.

**Table 2. Inequities in vaccination card usage rates and vaccination coverage in Nigeria.**

|  | Maternal education | | Antenatal care | | Household wealth | |
|---|---|---|---|---|---|---|
|  | ECI | Probability | ECI | Probability | ECI | Probability |
| Vaccination card | 0.40 | <0.0001 | 0.36 | <0.0001 | 0.30 | <0.0001 |
|  | | | | | | |
| Vaccine | | | | | | |
| BCG | 0.48 | <0.0001 | 0.44 | <0.0001 | 0.32 | <0.0001 |
| Measles – first dose | 0.41 | <0.0001 | 0.36 | <0.0001 | 0.30 | <0.0001 |
| DTP-HepB-Hib – third dose | 0.48 | <0.0001 | 0.39 | <0.0001 | 0.31 | <0.0001 |
| Polio – third dose | 0.22 | <0.0001 | 0.22 | <0.0001 | 0.17 | <0.0001 |
| Basic vaccination | 0.32 | <0.0001 | 0.27 | <0.0001 | 0.24 | <0.0001 |

Inequities in vaccination card usage rates and vaccination coverage among children aged 12-23

months disaggregated by maternal education, antenatal care, and household wealth in Nigeria,

based on Erreygers concentration indices (ECI).

Table 3 and Figure 6 presents the inequities in basic vaccination coverage in Nigeria among children

aged 12-23 months associated with socioeconomic (household wealth, religion, ethnicity),

geographic (region, place of residence), maternal (maternal age at birth, maternal education,

maternal household head status), child (sex of child, birth order), and healthcare (antenatal care,

birth setting) characteristics. After controlling for other background characteristics through multiple

logistic regression, the AORs were significant for the associations between basic vaccination

coverage and household wealth, religion, ethnicity, maternal age at birth, maternal education,

antenatal care, and birth setting. This model and the AORs include the interaction between

household wealth and place of residence.

**Table 3. Inequities in basic vaccination coverage in Nigeria associated with socioeconomic,**

**geographic, maternal, child, and healthcare characteristics.**

| Characteristics | Population (n = 6059) | Mean basic vaccination coverage (% and 95% confidence interval) | Crude odds ratio (OR and 95% confidence interval) | p-value | Adjusted odds ratio (AOR and 95% confidence interval) | p-value |
|---|---|---|---|---|---|---|
| **Socioeconomic** | | | | | | |
| *Household wealth* | | | | | | |
| Poorest | 1301 | 17.2 (14.5, 20.3) | 1 | <0.0001 | 1 | |
| Poorer | 1278 | 24.6 (21.6, 27.8) | **1.57** **(1.22, 2.01)** | | 1.38 (0.96, 2.00) | 0.086 |
| Middle | 1239 | 29.2 | **1.98** | | **1.73** | **0.005** |

| | N | % (CI) | OR (CI) | p | aOR (CI) | p |
|---|---|---|---|---|---|---|
| | | (25.9, 32.7) | **(1.53, 2.56)** | | **(1.17, 2.53)** | |
| Richer | 1156 | 38.2 (34.1, 42.5) | **2.97 (2.27, 3.90)** | | **3.10 (2.03, 4.74)** | **<0.001** |
| Richest | 1085 | 49.1 (45.1, 53.1) | **4.64 (3.59, 6.00)** | | **3.21 (2.06, 5.00)** | **<0.001** |
| *Religion* | | | | | | |
| Catholic | 586 | 48.8 (43.3, 54.3) | 1 | <0.0001 | 1 | |
| Other Christian | 1883 | 44.0 (41.0, 47.1) | 0.82 (0.65, 1.04) | | 0.98 (0.76, 1.26) | 0.852 |
| Islam | 3538 | 22.6 (20.5, 24.8) | **0.31 (0.24, 0.39)** | | 0.96 (0.67, 1.38) | 0.823 |
| Traditional | 15 | 22.2 (5.9, 56.3) | 0.30 (0.07, 1.36) | | 0.87 (0.23, 3.25) | 0.830 |

| | N | % (CI) | OR (CI) | p-value | Adjusted OR (CI) | p-value |
|---|---|---|---|---|---|---|
| Other | 37 | 14.0 (6.6, 27.3) | **0.17 (0.07, 0.40)** | | **0.25 (0.10, 0.60)** | **0.002** |
| *Ethnicity* (n = 6057) | | | | | | |
| Igbo | 847 | 56.3 (51.3, 61.1) | 1 | <0.0001 | 1 | |
| Ekoi | 25 | 51.5 (31.6, 70.9) | 0.82 (0.35, 1.94) | | 1.96 (0.74, 5.20) | 0.178 |
| Fulani | 568 | 12.2 (9.0, 16.2) | **0.11 (0.07, 0.16)** | | **0.51 (0.26, 0.97)** | **0.039** |
| Hausa | 1796 | 19.9 (17.3, 22.8) | **0.19 (0.15, 0.25)** | | 0.73 (0.39, 1.36) | 0.324 |
| Ibibio | 105 | 42.1 (31.9, 53.1) | **0.57 (0.35, 0.90)** | | 1.45 (0.77, 2.75) | 0.251 |
| Igala | 55 | 27.6 | **0.30** | | 0.61 | 0.283 |

| | | | | | | |
|---|---|---|---|---|---|---|
| | | (14.4, 46.5) | **(0.13, 0.69)** | | (0.24, 1.51) | |
| Ijaw/Izon | 171 | 33.5 (24.7, 43.7) | **0.39 (0.24, 0.63)** | | 1.05 (0.54, 2.04) | 0.891 |
| Kanuri/Beriberi | 148 | 14.4 (8.8, 22.7) | **0.13 (0.07, 0.24)** | | 0.42 (0.15, 1.19) | 0.102 |
| Tiv | 148 | 24.6 (16.8, 34.5) | **0.25 (0.15, 0.43)** | | 0.56 (0.27, 1.16) | 0.117 |
| Yoruba | 622 | 40.8 (35.6, 46.1) | **0.53 (0.40, 0.72)** | | 0.70 (0.40, 1.23) | 0.215 |
| Other/Don't know | 1574 | 35.9 (32.4, 39.5) | **0.43 (0.34, 0.56)** | | 1.03 (0.63, 1.67) | 0.912 |
| **Geographic** | | | | | | |
| *Region* | | | | | | |

| | | | | | | |
|---|---|---|---|---|---|---|
| North Central | 1061 | 30.9 (26.3, 35.8) | 1 | <0.0001 | 1 | |
| North East | 1303 | 22.9 (19.6, 26.5) | **0.66** **(0.50, 0.89)** | | 1.06 (0.75, 1.51) | 0.724 |
| North West | 1697 | 19.9 (17.1, 23.1) | **0.56** **(0.42, 0.74)** | | 1.00 (0.67, 1.49) | 1.000 |
| South East | 698 | 56.6 (51.0, 62.1) | **2.93** **(2.13, 4.01)** | | 1.35 (0.78, 2.35) | 0.284 |
| South South | 637 | 41.8 (37.2, 46.6) | **1.61** **(1.20, 2.16)** | | 0.86 (0.57, 1.27) | 0.441 |
| South West | 663 | 42.6 (37.3, 48.0) | **1.66** **(1.22, 2.27)** | | 0.83 (0.56, 1.24) | 0.357 |
| *Place of residence* | | | | | | |
| Urban | 2100 | 44.2 | 1 | <0.0001 | 1 | |

|  |  | (41.0, 47.5) |  |  |  |  |
| --- | --- | --- | --- | --- | --- | --- |
| Rural | 3959 | 22.7<br><br>(20.7, 24.8) | **0.37**<br><br>**(0.31, 0.44)** |  | 0.88<br><br>(0.56, 1.37) | 0.560 |
| **Maternal** |  |  |  |  |  |  |
| *Maternal age at birth (years)* |  |  |  |  |  |  |
| $15 - 19$ | 366 | 15.8<br><br>(11.9, 20.9) | 1 | <0.0001 | 1 |  |
| $20 - 34$ | 4330 | 31.4<br><br>(29.3, 33.5) | **2.43**<br><br>**(1.72, 3.43)** |  | **1.69**<br><br>**(1.15, 2.47)** | **0.007** |
| $35 - 49$ | 1363 | 34.5<br><br>(30.6, 38.6) | **2.80**<br><br>**(1.93, 4.05)** |  | **2.25**<br><br>**(1.46, 3.49)** | **<0.001** |
| *Maternal education* |  |  |  |  |  |  |
| None | 2614 | 14.8 | 1 | <0.0001 | 1 |  |

| | | | | | | |
|---|---|---|---|---|---|---|
| | | (12.9, 16.9) | | | | |
| Primary | 881 | 32.7 (28.1, 37.6) | **2.80 (2.15, 3.63)** | | **1.51 (1.14, 1.99)** | **0.004** |
| Secondary or higher | 2564 | 47.3 (44.5, 50.0) | **5.17 (4.27, 6.26)** | | **1.79 (1.39, 2.31)** | **<0.001** |
| *Sex of household head* | | | | | | |
| Male | 5434 | 30.3 (28.4, 32.3) | 1 | <0.0001 | 1 | |
| Female | 625 | 39.1 (34.7, 43.8) | **1.48 (1.21, 1.81)** | | 1.09 (0.86, 1.37) | 0.474 |
| **Child** | | | | | | |
| *Sex* | | | | | | |
| Male | 3148 | 31.4 | 1 | 0.81 | 1 | |

| | | | | | | |
|---|---|---|---|---|---|---|
| | | (28.9, 34.0) | | | | |
| Female | 2911 | 31.0 (28.7, 33.3) | 0.98 (0.85, 1.14) | | 1.02 (0.88, 1.19) | 0.766 |
| *Birth order* | | | | | | |
| 1st | 1157 | 35.8 (32.1, 39.7) | 1 | <0.0001 | 1 | |
| 2nd – 3rd | 2081 | 33.9 (31.0, 36.9) | 0.92 (0.75, 1.13) | | 0.91 (0.73, 1.14) | 0.417 |
| 4th – 5th | 1412 | 31.8 (28.6, 35.2) | 0.84 (0.66, 1.06) | | 1.01 (0.79, 1.30) | 0.909 |
| 6th or higher | 1409 | 22.7 (20.0, 25.6) | **0.53 (0.42, 0.65)** | | 0.86 (0.65, 1.14) | 0.292 |
| **Healthcare** | | | | | | |

*Antenatal care* (n = 5824)

| | | | | | | |
|---|---|---|---|---|---|---|
| None/Don't know | 1529 | 10.6 (8.7, 12.9) | 1 | <0.0001 | 1 | |
| 1 – 3 visits | 976 | 24.3 (21.3, 27.6) | **2.70** **(2.07, 3.53)** | | **2.15** **(1.62, 2.85)** | **<0.001** |
| 4 or more visits | 3319 | 41.5 (39.2, 43.8) | **5.96** **(4.71, 7.53)** | | **2.87** **(2.21, 3.72)** | **<0.001** |

*Birth setting*

| | | | | | | |
|---|---|---|---|---|---|---|
| Home setting | 3415 | 19.5 (17.5, 21.7) | 1 | <0.0001 | 1 | |
| Public or private facility | 2549 | 46.7 (44.0, 49.4) | **3.61** **(3.06, 4.26)** | | **1.32** **(1.08, 1.61)** | **0.006** |
| Other | 95 | 31.2 (19.8, 45.5) | 1.87 (0.98, 3.56) | | 0.77 (0.40, 1.50) | 0.444 |

Inequities in basic vaccination coverage (1-dose BCG, 3-dose DTP-HepB-Hib, 3-dose polio, and 1-dose

measles) in Nigeria among children aged 12-23 months associated with socioeconomic (household

wealth, religion, ethnicity), geographic (region, place of residence), maternal (maternal age at birth,

maternal education, maternal household head status), child (sex of child, birth order), and

healthcare (antenatal care, birth setting) characteristics. Crude and adjusted odds ratios were

estimated using simple and multiple logistic regression respectively.

*Figure 6 placement*

**Figure 6. Inequities in basic vaccination coverage in Nigeria.** Inequities in basic vaccination coverage

among children aged 12-23 months in Nigeria associated with socioeconomic (household wealth,

religion, ethnicity), geographic (region, place of residence), maternal (maternal age at birth,

maternal education, maternal household head status), child (sex of child, birth order), and

healthcare (birth setting, antenatal care) characteristics.

Children living in households of richest wealth quintiles had 221% higher odds (AOR: 3.21 (2.06,

5.00), p < 0.001) of receiving basic vaccinations in comparison to children of the poorest wealth

quintiles. However, children living in rural areas in the richer quintile had 64% lower odds (AOR: 0.36

(0.24, 0.53), p < 0.001) and children in rural areas in the richest quintile 50% lower odds (AOR: 0.50

(0.35, 0.72), p <0.001) than children in urban areas. Children whose religion was classified as 'other'

had 75% lower odds  (AOR: 0.25 (0.10, 0.60), p < 0.002) of receiving basic vaccinations than children

of Catholic faith. In comparison to children of Igbo ethnicity, children of Fulani ethnicity had 49%

lower odds of receiving basic vaccinations (AOR: 0.51 (0.26, 0.97), p = 0.039). Children of mothers

aged 35-49 years had 125% higher odds (AOR: 2.25 (1.46, 3.49), p < 0.001) of receiving basic

vaccination than children of mothers aged 15-19 years . Children of mothers with secondary or

higher education had 79% higher odds (AOR: 1.79 (1.39, 2.31), p < 0.001) of receiving basic

vaccination in comparison to children of mothers with no formal education. Children of mothers

who had four or more antenatal care visits had 187% higher odds  (AOR: 2.87 (2.21, 3.72), p < 0.001)

of  receiving basic vaccinations than children of mothers who had no antenatal care. Children born in

clinical facilities had 32% higher odds (AOR 1.32 (1.08, 1.61), p = 0.006) than children born in home

settings of receiving basic vaccinations.

**Discussion**

We conducted a systematic review to identify the social determinants of childhood immunisation in

low- and middle-income countries. We selected household wealth, religion, and ethnicity for

socioeconomic characteristics; region and place of residence for geographic characteristics; maternal

age at birth, maternal education, and maternal household head status for maternal characteristics;

sex of child and birth order for child characteristics; and antenatal care and birth setting for

healthcare characteristics.

Based on the characteristics identified from the systematic review, we applied a social determinants

framework to assess basic vaccination coverage (1-dose BCG, 3-dose DTP-HepB-Hib, 3-dose polio,

and 1-dose measles) among children aged 12-23 months in Nigeria using the 2018 Nigeria DHS

survey dataset.

The associations identified in this study between basic vaccination coverage and socioeconomic,

geographic, maternal, child, and healthcare characteristics identified are supported by other studies.

Basic vaccination coverage was associated with household wealth, and families in the richest quintile

are more likely to live in urban areas with better access to functional private and public health

facilities that provide immunisation services [24,30]. Mothers in urban areas are more likely to use

preventive healthcare services including childhood immunisation, due to their proximity to

healthcare facilities in urban settings and higher educational status in comparison to mothers in

rural areas with higher travel costs [31–33]. Children in the richer households of rural areas had

reduced odds of basic vaccination coverage in comparison to children in the poorest households of

urban areas. Children of Fulani ethnicity had lower odds of basic vaccination compared to Igbo

children, and there is evidence that awareness of immunisation is low amongst Fulani mothers

[34,35]. Further, Fulani ethnic groups reside in mostly rural settings and are nomadic, limiting their

access to health services including immunisation services.

We found that basic vaccination coverage was associated with maternal education. Educated

mothers have better awareness and knowledge on childhood immunisation and are more able to

overcome cultural barriers to vaccination [36]. Maternal age was associated with basic vaccination

coverage and older mothers may have more experience with antenatal clinics and have greater

awareness of immunisation services from previous children [22]. They are also more likely to have

financial access to immunisation services and live in urban settings with improved access to

immunisation services.

Delivery in a health facility and antenatal care were associated with basic vaccination, and to give

birth in a health facility indicates that mothers have overcome barriers to accessing health services,

and mothers will also receive information on childhood immunisation from healthcare workers

there. Hence, utilisation of health services by mothers leads to improved immunisation status of

their children [37].

At the regional level, basic vaccination among children ranged from the lowest coverage in the North

West region to nearly three times higher coverage in the South East region, although residents in

Northern Nigeria are more likely to have immunisation services within 5 km [38]. The large degree of

autonomy of different states and the impact of ongoing conflict in parts of the country, in addition

to socio-cultural reasons, can explain in part the geographical disparity in immunisation services

[14]. However, regional differences in coverage were not significant in the multivariable logistic

regression model.

Despite vaccination being provided at no cost to individuals, coverage was still below target levels

for even the most advantaged groups and therefore we recommend a proportionate universalism

approach with actions proportionate to the level of disadvantage [39,40]. There is some evidence of

payment being required for vaccination, even in public health facilities, and this may threaten

vaccination uptake [32]. Both the oversight of immunisation services and public awareness of

vaccinations' no cost status should be strengthened.

Higher coverage with more antenatal care indicates greater engagement with health services and

thereby providing more opportunities for vaccine education. The integration of immunisation

services to nutrition programmes and paediatric outpatient departments of primary healthcare

centres has been shown to improve coverage and decrease drop-out rates in South Sudan [41].

Hence, we recommend efforts to engage with families during health care visits outside of routine

immunisation services and to take advantage of these missed opportunities to reach under-

immunised children [42].

Our estimates of vaccination coverage in Nigeria for 2018 were lower than national estimates based

on administrative reporting from health service providers, though closer to WHO and UNICEF

Estimates of National Immunization Coverage (WEUNIC) (see Appendix A5) [43]. The coverage gaps

have been shown to be systematically underestimated by administrative reporting in Nigeria [44]. In

80% of Nigerian states, the basic vaccination coverage was below the national target of 95% or

higher coverage and below the Gavi target of 90% DTP3 coverage [45]. The below target coverage

for first doses indicates challenges with access to immunisation services while the decrease in

coverage for subsequent doses indicates drop-out due to insufficient knowledge on dose completion

[9,46,47]. The COVID-19 pandemic has disrupted vaccination globally and coverage was lower than

expected in Nigeria in 2020 [48], highlighting the urgent need for catch-up vaccination to close the

immunity gaps and prevent vaccine-preventable disease outbreaks [49]. The role of conflict on these

disparities is also important to understand. A literature review of conflict and vaccination inferred

that conflict-affected countries had vaccination coverage below global levels and conflict affected

vaccination services and human resources, including attacks on healthcare workers, which has

happened in Nigeria [14,50]. Our study inferences complement the findings in related studies in

Nigeria and other countries (see Table 1).

The proportion of zero-dose children was high and raises the risk of vaccine-preventable disease

outbreaks, with nearly 25% of children in rural areas not receiving a single dose of any of the basic

vaccinations. Zero-dose children are also likely to lack access to health and welfare services and to

suffer from multiple sources of deprivation [51,52]. More than 50% of children do not have

vaccination cards and nearly 60% of children do not have birth registrations [53] - improving uptake

of vaccination cards and birth registrations would help in part to address the barriers for vaccine

access for all children, including zero-dose children.

Our study has limitations, and we cannot estimate causal-effect relationships nor temporal

inferences due to the cross-sectional study design of 2018 Nigeria DHS. Our study has similar biases

that are associated with DHS surveys, including recall bias, measurement bias, and social desirability

bias which tend to overestimate vaccination coverage. In particular, only 49% of children had a

vaccination card and when a vaccination card was not available for a child, their mother was asked

to recall their vaccinations, which may lead to an overestimation of coverage.

For future work, we recommend qualitative research to understand the barriers and enablers of

childhood vaccination and their associations with socioeconomic, geographic, maternal, child, and

healthcare characteristics which would be valuable to adapt vaccination programmes to improve

coverage equitably in Nigeria.

**Conclusions**

We identified the inequities in basic vaccination coverage by socioeconomic, geographic, maternal,

child, and healthcare characteristics among children aged 12-23 months in Nigeria using a social

determinants of health perspective. In conclusion, we infer that inequities in basic vaccination were

associated with lower coverage among children living in poorer households, belonging to Fulani

ethnicity, born in home settings in Nigeria, with younger mothers at birth, with mothers with no

formal education and with mothers who had no antenatal care visits. We recommend a

proportionate universalism approach with targeted vaccination programmes proportionate to the

level of disadvantage for addressing the immunisation barriers faced by these underserved

subpopulations. This will improve coverage and reduce inequities in childhood immunisation

associated with socioeconomic, geographic, maternal, child, and healthcare characteristics in

Nigeria.

**Abbreviations**

AOR          Adjusted odds ratio

BCG          Bacille Calmette-Guérin vaccine

DHS          Demographic Health Survey

DTP          Diphtheria, tetanus and pertussis containing vaccine

DTP-HepB-Hib Diphtheria, tetanus, pertussis, hepatitis B and Haemophilus influenzae type B

ECI          Erreygers concentration indices

EPI          Expanded Programme of Immunization

Gavi         Global Alliance for Vaccines and Immunisations

| 436 | LGA | Local Government Area |
| 437 | LMIC | Low- and Middle-Income Countries |
| 438 | PRISMA | Preferred Reporting Items for Systematic Reviews and Meta-Analyses |
| 439 | SDG | Sustainable Development Goals |
| 440 | UNICEF | United Nations Children's Fund |
| 441 | WHO | World Health Organisation |
| 442 | WUENIC | WHO and UNICEF estimates of national immunization coverage |

## Declarations

**(a) Ethics Approval and Consent to Participate:**

This study was approved by the ethics committee (Reference: 21600) of the London School of

Hygiene & Tropical Medicine. For DHS surveys in general, the procedures and questionnaires are

approved by the ICF Institutional Review Board (IRB), and the country-specific DHS survey protocols

are reviewed by the ICF IRB and an IRB in the host country. Before each interview is conducted for

the DHS survey, an informed consent statement is read to the respondent, who may accept or

decline to participate. Further information on informed voluntary participation in the DHS survey is

available here: https://dhsprogram.com/Methodology/Protecting-the-Privacy-of-DHS-Survey-

Respondents.cfm All methods in this study were carried out following the relevant rules and

regulations.

**(b) Consent for Publication**

Not applicable.

**(c) Availability of data and materials:**

The analysis code is publicly accessible on GitHub at https://github.com/svwilliams122/vaccine-

equity-Nigeria. The 2018 Nigeria DHS data set is publicly available and accessible upon registration

for legitimate research purposes on the DHS website at

https://dhsprogram.com/methodology/survey/survey-display-528.cfm. To download DHS datasets,

researchers must register as a DHS data user at https://dhsprogram.com/data/new-user-

registration.cfm

**(d) Competing interests**

The authors declare that they have no known competing financial interests or personal relationships

that could have appeared to influence the work reported in this paper.

**(e) Funding**

KA is supported by the Vaccine Impact Modelling Consortium (OPP1157270).

**(f) Acknowledgment**

We thank the DHS Program for access to the 2018 Nigeria Demographic and Health Survey dataset.

**(g) Author's Contribution**

SVW and KA conceptualised the study. SVW undertook the analysis and wrote the initial draft. TA

provided contextual expertise of the national immunisation programme in Nigeria to explain the

findings. SVW, TA, and KA contributed to the analysis, interpretation of results, and the final drafting

of the manuscript. All authors have read and approved the final manuscript.

**Bibliography**

1.   UNDP. World Population Prospects - Population Division - United Nations. 2019 [cited 1 Nov
2019]. Available: https://population.un.org/wpp/

2.   World Bank. Nigeria Overview: Development news, research, data | World Bank. 2020 [cited
21 Sep 2021]. Available: https://www.worldbank.org/en/country/nigeria/overview

3.   Piot P, Larson HJ, O'Brien KL, N'kengasong J, Ng E, Sow S, et al. Immunization: vital progress,
unfinished agenda. Nature. 2019;575: 119–129. doi:10.1038/s41586-019-1656-7

4.   Rodrigues CMC, Plotkin SA. Impact of vaccines; health, economic and social perspectives.
Front Microbiol. 2020;11: 1526. doi:10.3389/fmicb.2020.01526

5.   UNICEF. IGME: Levels and Trends in Child Mortality - UNICEF DATA. 2020 [cited 21 Sep 2021].
Available: https://data.unicef.org/resources/levels-and-trends-in-child-mortality/

6.   WHO. Immunization. 2019 [cited 21 Sep 2021]. Available: https://www.who.int/news-
room/facts-in-pictures/detail/immunization

7.   Carter A, Msemburi W, Sim SY, A.M. Gaythorpe K, Lindstrand A, Hutubessy RCW. Modeling
the Impact of Vaccination for the Immunization Agenda 2030: Deaths Averted Due to
Vaccination Against 14 Pathogens in 194 Countries from 2021-2030. SSRN Journal. 2021.
doi:10.2139/ssrn.3830781

8.   Gavi. Facts and figures. 2021 [cited 21 Sep 2021]. Available:
https://www.gavi.org/programmes-impact/our-impact/facts-and-figures

9.   Cata-Preta BO, Santos TM, Mengistu T, Hogan DR, Barros AJD, Victora CG. Zero-dose children
and the immunisation cascade: Understanding immunisation pathways in low and middle-
income countries. Vaccine. 2021;39: 4564–4570. doi:10.1016/j.vaccine.2021.02.072

10.  Decouttere C, De Boeck K, Vandaele N. Advancing sustainable development goals through
immunization: a literature review. Global Health. 2021;17: 95. doi:10.1186/s12992-021-

00745-w

11.   WHO. Immunization Agenda 2030: A Global Strategy to Leave No One Behind. 2020 [cited 11

Jan 2021]. Available: https://www.who.int/teams/immunization-vaccines-and-

biologicals/strategies/ia2030

12.   Keja K, Chan C, Hayden G, Henderson RH. Expanded programme on immunization. World

Health Stat Q. 1988;41: 59–63.

13.   Sorungbe AO. Expanded programme on immunization in Nigeria. Rev Infect Dis. 1989;11 Suppl

3: S509-11. doi:10.1093/clinids/11.supplement_3.s509

14.   National Primary Health Care Development Agency. Comprehensive EPI Multi-Year Plan 2016 -

2020. Federal Ministry of Health, Nigeria; 2015.

15.   Gavi. Successfully transitioning Nigeria from Gavi support. 2018 [cited 21 Sep 2021]. Available:

https://www.gavi.org/sites/default/files/board_meetings/2018/6_Jun/05%20-

%20Successfully%20transitioning%20Nigeria%20from%20Gavi%20support.pdf

16.   WHO Africa. NERICC-Nigeria's panacea to routine immunization and primary health care

strengthening. | WHO | Regional Office for Africa. 2019 [cited 21 Sep 2021]. Available:

https://www.afro.who.int/news/nericc-nigerias-panacea-routine-immunization-and-primary-

health-care-strengthening

17.   Oku A, Oyo-Ita A, Glenton C, Fretheim A, Eteng G, Ames H, et al. Factors affecting the

implementation of childhood vaccination communication strategies in Nigeria: a qualitative

study. BMC Public Health. 2017;17: 200. doi:10.1186/s12889-017-4020-6

18.   Marmot M, Friel S, Bell R, Houweling TAJ, Taylor S, Commission on Social Determinants of

Health. Closing the gap in a generation: health equity through action on the social

determinants of health. Lancet. 2008;372: 1661–1669. doi:10.1016/S0140-6736(08)61690-6

19.   WHO. Social determinants of health. 2021 [cited 18 Oct 2022]. Available:

https://www.who.int/health-topics/social-determinants-of-health#tab=tab_3

20.    Mulholland E, Smith L, Carneiro I, Becher H, Lehmann D. Equity and child-survival strategies.

Bull World Health Organ. 2008;86: 399–407. doi:10.2471/blt.07.044545

21.    Arcaya MC, Arcaya AL, Subramanian SV. Inequalities in health: definitions, concepts, and

theories. Glob Health Action. 2015;8: 27106. doi:10.3402/gha.v8.27106

22.    WHO. WHO | Explorations of inequality: Childhood immunization. 2018 [cited 18 Sep 2019].

Available: https://apps.who.int/iris/handle/10665/272864

23.    Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for

systematic reviews and meta-analyses: The PRISMA statement. PLoS Med. 2009;6: e1000097.

doi:10.1371/journal.pmed.1000097

24.    NPC/Nigeria, ICF.  Nigeria Demographic and Health Survey 2018. National Population

Commission Nigeria and ICF; 2019.

25.    Croft TN, Marshall AMJ, Allen CK. Guide to DHS Statistics. Rockville, Maryland, USA: ICF; 2018.

26.    StataCorp. Stata: software for statistics and data science. College Station, Texas: StataCorp LP;

2019.

27.    Kirkwood BR, Sterne JAC. Essential Medical Statistics. 2nd ed. Wiley; 2003.

28.    WHO. Health Inequality Monitoring with a special focus on low- and middle-income countries.

World Health Organization; 2013.

29.    R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R

Foundation for Statistical Computing; 2019.

30.    Adebowale A, Obembe T, Bamgboye E. Relationship between household wealth and

childhood immunization in core-North Nigeria. Afr Health Sci. 2019;19: 1582–1593.

doi:10.4314/ahs.v19i1.33

31.    Adenike O-B, Adejumoke J, Olufunmi O, Ridwan O. Maternal characteristics and immunization

status of children in North Central of Nigeria. Pan Afr Med J. 2017;26: 159.

doi:10.11604/pamj.2017.26.159.11530

32.  Sibeudu FT, Uzochukwu BS, Onwujekwe OE. Rural-urban comparison of routine immunization

utilization and its determinants in communities in Anambra States, Nigeria. SAGE Open Med.

2019;7: 2050312118823893. doi:10.1177/2050312118823893

33.  Okafor IP, Dolapo DC, Onigbogi MO, Iloabuchi IG. Rural-urban disparities in maternal

immunization knowledge and childhood health-seeking behavior in Nigeria: a mixed method

study. Afr Health Sci. 2014;14: 339–347. doi:10.4314/ahs.v14i2.8

34.  Dao MY, Brieger WR. Immunization for the migrant fulani: identifying an under-served

population in southwestern Nigeria. Int Q Community Health Educ. 1994;15: 21–32.

doi:10.2190/VKHL-JXC5-WK91-8QGQ

35.  Afolabi RF, Salawu MM, Gbadebo BM, Salawu AT, Fagbamigbe AF, Adebowale AS. Ethnicity as

a cultural factor influencing complete vaccination among children aged 12-23 months in

Nigeria. Hum Vaccin Immunother. 2021;17: 2008–2017. doi:10.1080/21645515.2020.1870394

36.  Balogun SA, Yusuff HA, Yusuf KQ, Al-Shenqiti AM, Balogun MT, Tettey P. Maternal education

and child immunization: the mediating roles of maternal literacy and socioeconomic status.

Pan Afr Med J. 2017;26: 217. doi:10.11604/pamj.2017.26.217.11856

37.  Adedokun ST, Uthman OA, Adekanmbi VT, Wiysonge CS. Incomplete childhood immunization

in Nigeria: a multilevel analysis of individual and contextual factors. BMC Public Health.

2017;17: 236. doi:10.1186/s12889-017-4137-7

38.  Eboreime E, Abimbola S, Bozzani F. Access to Routine Immunization: A Comparative Analysis

of Supply-Side Disparities between Northern and Southern Nigeria. PLoS ONE. 2015;10:

e0144876. doi:10.1371/journal.pone.0144876

39.  Carey G, Crammond B, De Leeuw E. Towards health equity: a framework for the application of

574           proportionate universalism. Int J Equity Health. 2015;14: 81. doi:10.1186/s12939-015-0207-6

575    40.    Marmot M, Bell R. Fair society, healthy lives. Public Health. 2012;126 Suppl 1: S4-10.

576           doi:10.1016/j.puhe.2012.05.014

577    41.    Idris IO, Obwoya JG, Tapkigen J, Lamidi SA, Ochagu VA, Abbas K. Impact evaluation of

578           immunisation service integration to nutrition programmes and paediatric outpatient

579           departments of primary healthcare centres in Rumbek East and Rumbek Centre counties of

580           South Sudan. Family Med Commun Hlth. 2021;9. doi:10.1136/fmch-2021-001034

581    42.    Restrepo-Méndez MC, Barros AJD, Wong KLM, Johnson HL, Pariyo G, Wehrmeister FC, et al.

582           Missed opportunities in full immunization coverage: findings from low- and lower-middle-

583           income countries. Glob Health Action. 2016;9: 30963. doi:10.3402/gha.v9.30963

584    43.    WHO, UNICEF. WUENIC (WHO/UNICEF Estimates of National Immunization Coverage)

585           estimates - Vaccines monitoring system 2019 Global Summary Reference Time Series. 2019

586           [cited 19 Apr 2020]. Available:

587           https://apps.who.int/immunization_monitoring/globalsummary/timeseries/tswucoveragedtp

588           3.html

589    44.    Gunnala R, Ogbuanu IU, Adegoke OJ, Scobie HM, Uba BV, Wannemuehler KA, et al. Routine

590           Vaccination Coverage in Northern Nigeria: Results from 40 District-Level Cluster Surveys,

591           2014-2015. PLoS ONE. 2016;11: e0167835. doi:10.1371/journal.pone.0167835

592    45.    Gavi. 2019 Annual Progress Report. 2020 [cited 21 Sep 2021]. Available:

593           https://www.gavi.org/sites/default/files/programmes-impact/our-impact/apr/Gavi-Progress-

594           Report-2019_1.pdf

595    46.    Adedire EB, Ajayi I, Fawole OI, Ajumobi O, Kasasa S, Wasswa P, et al. Immunisation coverage

596           and its determinants among children aged 12-23 months in Atakumosa-west district, Osun

597           State Nigeria: a cross-sectional study. BMC Public Health. 2016;16: 905. doi:10.1186/s12889-

598           016-3531-x

599    47.    Odusanya OO, Alufohai EF, Meurice FP, Ahonkhai VI. Determinants of vaccination coverage in
600           rural Nigeria. BMC Public Health. 2008;8: 381. doi:10.1186/1471-2458-8-381

601    48.    Causey K, Fullman N, Sorensen RJD, Galles NC, Zheng P, Aravkin A, et al. Estimating global and
602           regional disruptions to routine childhood vaccine coverage during the COVID-19 pandemic in
603           2020: a modelling study. Lancet. 2021.

604    49.    WHO. Leave No One Behind: Guidance for Planning and Implementing Catch-up Vaccination.
605           2020 [cited 1 Feb 2021]. Available: https://www.who.int/teams/immunization-vaccines-and-
606           biologicals/essential-programme-on-immunization/implementation/catch-up-vaccination/

607    50.    Grundy J, Biggs B-A. The impact of conflict on immunisation coverage in 16 countries. Int J
608           Health Policy Manag. 2019;8: 211–221. doi:10.15171/ijhpm.2018.127

609    51.    WHO. Progress and Challenges with Achieving Universal Immunization Coverage. 2020 [cited
610           22 Dec 2022]. Available: https://www.who.int/publications/m/item/progress-and-challenges-
611           with-achievinguniversal-immunization-coverage

612    52.    Bhatia A, Krieger N, Beckfield J, Barros AJD, Victora C. Are inequities decreasing? Birth
613           registration for children under five in low-income and middle-income countries, 1999-2016.
614           BMJ Glob Health. 2019;4: e001926. doi:10.1136/bmjgh-2019-001926

615    53.    United Nations Children's Fund,. Birth Registration for Every Child by 2030: Are we on track?
616           UNICEF; 2019.

617    54.    Shenton LM, Wagner AL, Carlson BF, Mubarak MY, Boulton ML. Vaccination status of children
618           aged 1-4 years in Afghanistan and associated factors, 2015. Vaccine. 2018;36: 5141–5149.
619           doi:10.1016/j.vaccine.2018.07.020

620    55.    Oliveira MFS de, Martinez EZ, Rocha JSY. Factors associated with vaccination coverage in
621           children < 5 years in Angola. Rev Saude Publica. 2014;48: 906–915. doi:10.1590/S0034-
622           8910.2014048005284

56. Rahman A, Reza AAS, Bhuiyan BA, Alam N, Dasgupta SK, Mostari S, et al. Equity and

determinants of routine child immunisation programme among tribal and non-tribal

populations in rural Tangail subdistrict, Bangladesh: a cohort study. BMJ Open. 2018;8:

e022634. doi:10.1136/bmjopen-2018-022634

57. Vyas P, Kim D, Adams A. Understanding spatial and contextual factors influencing

intraregional differences in child vaccination coverage in Bangladesh. Asia Pac J Public Health.

2019;31: 51–60. doi:10.1177/1010539518813604

58. Nda'chi Deffo R, Fomba Kamga B. Do the dynamics of vaccine programs improve the full

immunization of children under the age of five in Cameroon? BMC Health Serv Res. 2020;20:

953. doi:10.1186/s12913-020-05745-x

59. Ashbaugh HR, Hoff NA, Doshi RH, Alfonso VH, Gadoth A, Mukadi P, et al. Predictors of measles

vaccination coverage among children 6-59 months of age in the Democratic Republic of the

Congo. Vaccine. 2018;36: 587–593. doi:10.1016/j.vaccine.2017.11.049

60. Lakew Y, Bekele A, Biadgilign S. Factors influencing full immunization coverage among 12-23

months of age children in Ethiopia: evidence from the national demographic and health

survey in 2011. BMC Public Health. 2015;15: 728. doi:10.1186/s12889-015-2078-6

61. Geweniger A, Abbas KM. Childhood vaccination coverage and equity impact in Ethiopia by

socioeconomic, geographic, maternal, and child characteristics. Vaccine. 2020;38: 3627–3638.

doi:10.1016/j.vaccine.2020.03.040

62. Gram L, Soremekun S, ten Asbroek A, Manu A, O'Leary M, Hill Z, et al. Socio-economic

determinants and inequities in coverage and timeliness of early childhood immunisation in

rural Ghana. Trop Med Int Health. 2014;19: 802–811. doi:10.1111/tmi.12324

63. Shrivastwa N, Gillespie BW, Kolenic GE, Lepkowski JM, Boulton ML. Predictors of vaccination

in India for children aged 12-36 months. Vaccine. 2015;33 Suppl 4: D99-105.

doi:10.1016/j.vaccine.2015.09.034

64.    Devasenapathy N, Ghosh Jerath S, Sharma S, Allen E, Shankar AH, Zodpey S. Determinants of

childhood immunisation coverage in urban poor settlements of Delhi, India: a cross-sectional

study. BMJ Open. 2016;6: e013015. doi:10.1136/bmjopen-2016-013015

65.    Khan J, Shil A, Prakash R. Exploring the spatial heterogeneity in different doses of vaccination

coverage in India. PLoS ONE. 2018;13: e0207209. doi:10.1371/journal.pone.0207209

66.    Francis MR, Nohynek H, Larson H, Balraj V, Mohan VR, Kang G, et al. Factors associated with

routine childhood vaccine uptake and reasons for non-vaccination in India: 1998-2008.

Vaccine. 2018;36: 6559–6566. doi:10.1016/j.vaccine.2017.08.026

67.    Kannankeril Joseph VJ. Understanding inequalities in child immunization in India: a

decomposition approach. J Biosoc Sci. 2021; 1–13. doi:10.1017/S0021932021000110

68.    Srivastava S, Kumar P, Chauhan S, Banerjee A. Household expenditure for immunization

among children in India: a two-part model approach. BMC Health Serv Res. 2021;21: 1001.

doi:10.1186/s12913-021-07011-0

69.    Siramaneerat I, Agushybana F. Inequalities in immunization coverage in Indonesia: a

multilevel analysis. Rural Remote Health. 2021;21: 6348. doi:10.22605/RRH6348

70.    Van Malderen C, Ogali I, Khasakhala A, Muchiri SN, Sparks C, Van Oyen H, et al. Decomposing

Kenyan socio-economic inequalities in skilled birth attendance and measles immunization. Int

J Equity Health. 2013;12: 3. doi:10.1186/1475-9276-12-3

71.    Masters NB, Wagner AL, Carlson BF, Muuo SW, Mutua MK, Boulton ML. Childhood vaccination

in Kenya: socioeconomic determinants and disparities among the Somali ethnic community.

Int J Public Health. 2019;64: 313–322. doi:10.1007/s00038-018-1187-2

72.    Allan S, Adetifa IMO, Abbas K. Inequities in childhood immunisation coverage associated with

socioeconomic, geographic, maternal, child, and place of birth characteristics in Kenya. BMC

Infect Dis. 2021;21: 553. doi:10.1186/s12879-021-06271-9

73.   Mutua MK, Mohamed SF, Iddi S, Muyingo S, Mwangi B, Kadengye D, et al. Do inequalities exist

in the disadvantaged populations? Levels and trends of full and on-time vaccination coverage

in two Nairobi urban informal settlements. Global Epidemiology. 2020;2: 100044.

doi:10.1016/j.gloepi.2020.100044

74.   Sychareun V, Rowlands L, Vilay P, Durham J, Morgan A. The determinants of vaccination in a

semi-rural area of Vientiane City, Lao People's Democratic Republic: a qualitative study.

Health Res Policy Syst. 2019;17: 2. doi:10.1186/s12961-018-0407-9

75.   Abebe DS, Nielsen VO, Finnvold JE. Regional inequality and vaccine uptake: a multilevel

analysis of the 2007 Welfare Monitoring Survey in Malawi. BMC Public Health. 2012;12: 1075.

doi:10.1186/1471-2458-12-1075

76.   Ntenda PAM, Chuang K-Y, Tiruneh FN, Chuang Y-C. Analysis of the effects of individual and

community level factors on childhood immunization in Malawi. Vaccine. 2017;35: 1907–1917.

doi:10.1016/j.vaccine.2017.02.036

77.   Daca C, Sebastian MS, Arnaldo C, Schumann B. Socio-economic and demographic factors

associated with reproductive and child health preventive care in Mozambique: a cross-

sectional study. Int J Equity Health. 2020;19: 200. doi:10.1186/s12939-020-01303-3

78.   Patel PN, Hada M, Carlson BF, Boulton ML. Immunization status of children in Nepal and

associated factors, 2016. Vaccine. 2021;39: 5831–5838. doi:10.1016/j.vaccine.2021.08.059

79.   Antai D. Inequitable childhood immunization uptake in Nigeria: a multilevel analysis of

individual and contextual determinants. BMC Infect Dis. 2009;9: 181. doi:10.1186/1471-2334-

9-181

80.   Antai D. Gender inequities, relationship power, and childhood immunization uptake in

Nigeria: a population-based cross-sectional study. Int J Infect Dis. 2012;16: e136-45.

doi:10.1016/j.ijid.2011.11.004

81.    Ataguba JE, Ojo KO, Ichoku HE. Explaining socio-economic inequalities in immunization

697           coverage in Nigeria. Health Policy Plan. 2016;31: 1212–1224. doi:10.1093/heapol/czw053

698    82.    Uzochukwu BS, Okeke CC, Envuladu E, Mbachu C, Okwuosa C, Onwujekwe OE. Inequity in

699           access to childhood immunization in Enugu urban, Southeast Nigeria. Niger J Clin Pract.

700           2017;20: 971–977. doi:10.4103/njcp.njcp_375_16

701    83.    Uthman OA, Adedokun ST, Olukade T, Watson S, Adetokunboh O, Adeniran A, et al. Children

702           who have received no routine polio vaccines in Nigeria: Who are they and where do they live?

703           Hum Vaccin Immunother. 2017;13: 2111–2122. doi:10.1080/21645515.2017.1336590

704    84.    Obanewa OA, Newell ML. The role of place of residency in childhood immunisation coverage

705           in Nigeria: analysis of data from three DHS rounds 2003-2013. BMC Public Health. 2020;20:

706           123. doi:10.1186/s12889-020-8170-6

707    85.    Siddiqui NT, Owais A, Agha A, Karim MS, Zaidi AKM. Ethnic disparities in routine immunization

708           coverage: a reason for persistent poliovirus circulation in Karachi, Pakistan? Asia Pac J Public

709           Health. 2014;26: 67–76. doi:10.1177/1010539513475648

710    86.    Canavan ME, Sipsma HL, Kassie GM, Bradley EH. Correlates of complete childhood vaccination

711           in East African countries. PLoS ONE. 2014;9: e95709. doi:10.1371/journal.pone.0095709

712    87.    Wariri O, Edem B, Nkereuwem E, Nkereuwem OO, Umeh G, Clark E, et al. Tracking coverage,

713           dropout and multidimensional equity gaps in immunisation systems in West Africa, 2000-

714           2017. BMJ Glob Health. 2019;4: e001713. doi:10.1136/bmjgh-2019-001713

715    88.    Ndwandwe D, Uthman OA, Adamu AA, Sambala EZ, Wiyeh AB, Olukade T, et al. Decomposing

716           the gap in missed opportunities for vaccination between poor and non-poor in sub-Saharan

717           Africa: A Multicountry Analyses. Hum Vaccin Immunother. 2018;14: 2358–2364.

718           doi:10.1080/21645515.2018.1467685

89.    Sambala EZ, Uthman OA, Adamu AA, Ndwandwe D, Wiyeh AB, Olukade T, et al. Mind the Gap:

What explains the education-related inequality in missed opportunities for vaccination in sub-

Saharan Africa? Compositional and structural characteristics. Hum Vaccin Immunother.

2018;14: 2365–2372. doi:10.1080/21645515.2018.1460985

90.     Ameyaw EK, Kareem YO, Ahinkorah BO, Seidu A-A, Yaya S. Decomposing the rural-urban gap

in factors associated with childhood immunisation in sub-Saharan Africa: evidence from

surveys in 23 countries. BMJ Glob Health. 2021;6. doi:10.1136/bmjgh-2020-003773

91.     Arsenault C, Harper S, Nandi A, Rodríguez JMM, Hansen PM, Johri M. An equity dashboard to

monitor vaccination coverage. Bull World Health Organ. 2017;95: 128–134.

doi:10.2471/BLT.16.178079

92.     Arsenault C, Johri M, Nandi A, Mendoza Rodríguez JM, Hansen PM, Harper S. Country-level

predictors of vaccination coverage and inequalities in Gavi-supported countries. Vaccine.

2017;35: 2479–2488. doi:10.1016/j.vaccine.2017.03.029

93.     Bosch-Capblanch X, Banerjee K, Burton A. Unvaccinated children in years of increasing

coverage: how many and who are they? Evidence from 96 low- and middle-income countries.

Trop Med Int Health. 2012;17: 697–710. doi:10.1111/j.1365-3156.2012.02989.x

94.     Merten S, Martin Hilber A, Biaggi C, Secula F, Bosch-Capblanch X, Namgyal P, et al. Gender

Determinants of Vaccination Status in Children: Evidence from a Meta-Ethnographic

Systematic Review. PLoS ONE. 2015;10: e0135222. doi:10.1371/journal.pone.0135222

95.     Hinman AR, McKinlay MA. Immunization Equity. Am J Prev Med. 2015;49: S399-405.

doi:10.1016/j.amepre.2015.04.018

96.     Awoh AB, Plugge E. Immunisation coverage in rural-urban migrant children in low and middle-

income countries (LMICs): a systematic review and meta-analysis. J Epidemiol Community

Health. 2016;70: 305–311. doi:10.1136/jech-2015-205652

97.     Hosseinpoor AR, Bergen N, Schlotheuber A, Gacic-Dobo M, Hansen PM, Senouci K, et al. State

744           of inequality in diphtheria-tetanus-pertussis immunisation coverage in low-income and

745           middle-income countries: a multicountry study of household health surveys. Lancet Glob

746           Health. 2016;4: e617-26. doi:10.1016/S2214-109X(16)30141-3

747    98.    Hajizadeh M. Socioeconomic inequalities in child vaccination in low/middle-income countries:

748           what accounts for the differences? J Epidemiol Community Health. 2018;72: 719–725.

749           doi:10.1136/jech-2017-210296

750    99.    Hajizadeh M. Decomposing socioeconomic inequality in child vaccination in the Gambia, the

751           Kyrgyz Republic and Namibia. Vaccine. 2019;37: 6609–6616.

752           doi:10.1016/j.vaccine.2019.09.054

**National Policies**

**Geographical:** region, place of residence

**Household:** household wealth, religion, ethnicity

**Maternal:** age at birth, education, household head

**Individual (child and healthcare):** sex of child, birth order, antenatal care, birth setting