Original research

Acquired neutralizing breadth against SARS-CoV-2 variants including Omicron

after three doses of mRNA COVID-19 vaccination and the vaccine efficacy

Koichi Furukawa, MD, Lidya Handayani Tjan, MD, PhD, Yukiya Kurahashi, MD,

Silvia Sutandhio, MD, Mitsuhiro Nishimura, PhD, Jun Arii, DVM, PhD, Yasuko Mori,

MD, PhD

Division of Clinical Virology, Center for Infectious Diseases, Kobe University Graduate

School of Medicine, Kobe, Hyogo 650-0017, Japan

Corresponding author:

Prof Yasuko Mori

Division of Clinical Virology, Center for Infectious Diseases, Kobe University Graduate

School of Medicine.

7-5-1 Kusunoki-cho, Chuo-ku, Kobe, Hyogo Japan, 650-0017

Tel: +81-78-382-6272   Fax: +81-78-382-6879

E-mail: ymori@med.kobe-u.ac.jp

ABSTRACT

To investigate the induction of neutralizing antibodies against Omicron after two and

three vaccine doses in recipients of different ages. Physicians at Kobe University

Hospital who had received the second dose of the BNT162b2 mRNA vaccine. At 2

months after the second vaccinations, the positive rate of neutralizing antibody against

Omicron was 28%, and the titer was significantly lower than those against other

variants, 11.8-fold and 3.6-fold lower than those against D614G and Delta, respectively.

Unlike Delta, that positive rates of neutralizing antibody against Omicron were low in

all age groups, and there was no significant difference in titers among age groups. Seven

months after the 2nd dose, the positive rate of neutralizing antibody against Omicron

decreased to 6%, but after the booster,3rd vaccination, it increased to 100%, and the

titer was much higher than those at 2 and 7 months post-vaccination, 32-fold and

39-fold respectively. The booster vaccination effect was also observed in the younger at

41-fold, middle-aged at 43-fold, and older at 27-fold groups compared to the 7-month

titers. Surprisingly, higher-than-predicted titers of the neutralizing antibodies against

Omicron were induced after the booster vaccination regardless of recipient age, while

this effect was not observed after two doses, indicating the induction of antibodies

36   against common epitopes by the booster vaccination. Three doses can be confidently

37   recommended to suppress the pandemic.

Introduction

The coronavirus disease 2019 (COVID-19) pandemic declared by the World

Health Organization (WHO) in March 2020 continues to affect all countries around the

world. Based on the WHO COVID-19 dashboard as of 20 January 2022, there have

been more than 332 million confirmed cases with more than 5.5 million deaths

worldwide.[1] Currently, the Omicron variant (B.1.1.529) of the original severe acute

respiratory syndrome coronavirus 2 (SARS-CoV-2) is rapidly spreading and has become

the dominant variant all over the world, including Japan.[2-6] In efforts to control the

pandemic, several vaccine platforms have been developed using the original

SARS-CoV-2 virus (Wuhan-1) as the template, and the BNT162b2 mRNA vaccine

(Comirnaty®, BioNTech-Pfizer, Mainz, Germany/New York, U.S.) was granted an

emergency use authorization by the U.S. Food and Drug Administration (FDA) in

December 2020, after showing high effectiveness in preventing infection.[7]

However, since April 2020, a variant bearing a D614G mutation of the spike

protein and several SARS-CoV-2 variants classified as variants of concern (VOC) or

variants of interest (VOI) by the WHO have emerged and spurred the spread of

infection worldwide. As of this manuscript's writing, SARS-CoV-2 variants classified as

VOCs include Alpha (B.1.1.7) first detected in the UK,[8-10] Beta (B.1.351) from South

Africa,[11-14] Gamma (P.1) from Brazil,[15,16] Delta (B.1.617.2) from India,[17] and Omicron

(B.1.1.529) from South Africa and Bostwana, which is the latest VOC (detected in early

November 2021).[18,19] The emergence of these variants poses a tremendous challenge

because the vaccine efficacy is negatively impacted by these variants.

In particular, the Beta and Gamma variants carry the E484K mutation in the

receptor binding domain (RBD) of the spike protein, which allows evasion of

neutralizing antibodies against the original SARS-CoV-2, further compromising the

currently available therapy.[20-22] The Delta variant has L452R, T478K, and P681R

mutations in the RBD, which increases the virus transmissibility□—□leading to a

higher viral load in infected individuals.[23,24] However, the novel Omicron variant is the

most mutated variant, with over 30 mutations in the spike protein and 15 mutations in

the RBD.[25] There is thus great concern about the Omicron variant's higher

transmissibility and immune escapes from COVID-19 vaccines and antibody-based

therapies.[26-29] It has also been demonstrated that vaccine-induced neutralizing antibody

responses decrease with increased time post-vaccination.[30,31]

To counteract the decline of the neutralizing antibody response over time and the

emergence of new variants including Omicron, a third dose of vaccination, which has

been called a "booster vaccination," has been approved and administered to individuals

who were vaccinated >6 months ago; the booster vaccination has been shown to

effectively induce a high titer of neutralizing antibody.[32] However, it is unclear whether

the effective neutralizing antibody against Omicron can be obtained by two or three

doses of vaccination compared to the other variants. It is also not yet clear whether the

characteristics of the vaccine's recipients or side effects are related to the acquired

immunity.

Here, we investigated the neutralizing breadth of sera against D614G, Alpha,

Beta, Gamma, Delta, Kappa, and Omicron variants which were obtained from 82

vaccinated physicians without a history of SARS-CoV-2 infection, and we analyzed the

correlations between the neutralizing antibody titers and the recipients' background and

side effects. We also followed and analyzed the neutralizing antibody titers against

D614G, Delta, and Omicron in individuals after they received the booster (third dose of)

vaccination.

Participants and Methods

Participant recruitment

From June 2021 to July 2021, blood samples from 82 physicians at Kobe University

Hospital (Kobe, Japan) who had already received the second vaccination of the

BNT162b2 mRNA COVID-19 vaccine and had never tested positive for SARS-CoV-2

infection were collected and analyzed (1st trial). As the 2nd trial, blood samples were

collected again from the participants during the period from October to November 2021.

In a 3rd trial, blood samples from the 72 participants who received booster vaccination

were collected in January 2022 and analyzed. The sera of 24 unvaccinated and healthy

adults were also collected and confirmed to have no antibody against SARS-CoV-2;[33]

these sera were used as the negative control group. This study was carried out after

written consent was obtained from all participants. No statistical methods were used to

predetermine the sample size.

Questionnaire about side effects

When blood samples were collected, a questionnaire was used to obtain the participants'

background characteristics and information about any side effects that occurred within

the first week after the first, second, and booster vaccinations. The contents of the

questionnaire included age, sex, symptoms after vaccination such as fever (>37.5°),

general fatigue, injection site pain (0: none, 1: mild, 2: moderate, 3: severe), the side of

the injected arm, and others. The degree of injection site pain was subjectively

determined by each participant.

Measurement of neutralizing activity against SARS-CoV-2

The neutralization assay was performed using authentic viruses as recently

described.[34,35]

Preparation of SARS-CoV-2 variants

We used the SARS-CoV-2 Biken-2 (B2) strain with a D614G mutation as a

conventional variant (accession no. LC644163); it was provided by the BIKEN

Innovative Vaccine Research Alliance Laboratories (Osaka, Japan). The other variants,

i.e., Alpha (GISAID ID: EPI_ISL_804007), Beta (GISAID ID: EPI_ISL_1123289),

Gamma (GISAID ID: EPI_ISL_833366), Delta (GISAID ID: EPI_ISL_2158617),

Kappa (GISAID ID: EPI_ISL_2158613), and Omicron (GISAID ID:

EPI_ISL_7418017) had been isolated and were provided by the National Institute of

Infectious Disease, Japan (Tokyo). Each variant was confirmed by the cDNA sequence

of its spike gene.

Statistical analysis

GraphPad Prism software (ver. 8.4.3) and STATA software (ver. 14.2) were used for the

statistical analysis and preparation of figures. The Friedman test was used to compare

the neutralizing antibody titer among the seven variants. The Mann-Whitney U-test or

Kruskal-Wallis test was used to compare the neutralizing antibody titer or age among

the different groups of participants. If a significant difference was found by the

Friedman test or Kruskal-Wallis test, Dunn's multiple comparison test was then

performed. Results were considered significant when p-values were <0.05 (two-tailed).

Ethical approval

This study was approved by the ethical committee of Kobe University Graduate School

of Medicine (approval code: B200200).

Results

Participant characteristics and side effects after the first, second, and third vaccine

doses

The demographic characteristics of the participants are summarized in Table 1. The

median (interquartile [IQR]) numbers of days between the second dose of vaccination

and the collection of serum samples in the 1st and 2nd trials were 53 [46–59] and 199

[192–203], respectively (hereinafter referred to as "2 months" and "7 months" after the

vaccination). The median number of days between the booster vaccination and the

collection of serum samples in the 3rd trial was 16 days. Because most of the physicians

working in the hospital were men, most (71 of the 82) participants were males (86.6%).

The side effects reported by the participants after the first, second, and booster

vaccination are also listed in Table 1. Side effects at the time of the booster vaccination

were as follows: fever in 24 participants (33%), general fatigue in 45 (63%), and

injection site pain in 65 (89%) (15 mild, 35 moderate, and 15 severe). The frequency of

systemic reactions including fever and fatigue increased in order, with the booster

vaccination being the most frequent, but that of injection site pain did not change. No

participant had serious adverse events.

Neutralizing activity against Omicron compared with the other variants at 2 months

post-vaccination

Although the neutralizing antibody titer varied, all 82 sera had neutralizing activity

against D614G and Alpha. In contrast, most (but not all) of the sera had neutralizing

activity against the Beta, Gamma, Delta, and Kappa variants; neutralizing antibody

positive rates against the respective variants were 87.8%, 95.1%, 92.7%, and 90.2% of

the participants. Surprisingly however, only 28% of the participants had the neutralizing

activity against Omicron (Fig. 1A). The comparison of the antibody titers for each

variant revealed that the antibody titers against Omicron were significantly lower than

those against the six other variants (Fig. 1B). The average neutralizing antibody titers

against Omicron were 11.8-fold (95%CI: 9.9–13.9) lower that that against D614G,

7.3-fold (95%CI: 6.3–8.5)) lower than that against Alpha, 3.2-fold (95%CI: 2.6–3.8))

lower than that against Beta, 4.9-fold (95%CI: 4.1–5.9)) lower than that against

Gamma, 3.6-fold (95%CI: 3.1–4.2)) lower than that against Delta, and 2.6-fold

(95%CI: 2.2–3.0) lower than that against Kappa.

The neutralizing antibody titer against D614G was significantly higher than those

against Beta (3.6-fold; 95%CI: 3.0–4.3), Gamma (2.3-fold; 95%CI: 1.9–2.8), Delta

(3.2-fold; 95%CI: 2.7–3.7), and Kappa (4.5-fold; 95%CI: 3.8–5.2).

Neutralizing activity against Omicron and the other variants at 2 months after

vaccination in each age group

We had divided the participants into three groups by age: the 31 participants (37.8%)

who were ≤38 years old were classified as the younger group; the 32 participants

(39.0%) aged 39–58 years old comprised the middle-aged group; and the 19 participants

(23.2%) who were ≥59 years old constituted the older group (Table 1). Although all of

the participants' sera showed neutralizing activity against D614G and Alpha regardless

of age, sera with negative neutralizing activity against other variants were observed in

all age groups, and the negative proportion tended to be higher in the older group (Fig.

2A). The positive rate of sera with neutralizing antibody against Omicron was low in all

three age groups; only 32% had the neutralizing antibody against Omicron even in the

younger age (≤38 yrs) group. The comparison of neutralizing antibody titers among the

three age groups showed results similar to the positive rate: the median [range] of

neutralizing antibody titer against Delta was significantly lower in the older age group

(3 [<2–8]) compared to the other two groups: (4[<2–32] vs. the younger group, p=0.03)

and (4 [<2–32] vs. the middle-aged group, p=0.04). Similar results were also obtained

for the median [range] of neutralizing titer against Kappa: (2 [<2–4] in the older group)

versus (4 [<2–16] in the younger group, p=0.02) and (4 [<2–16] in the middle-aged

group , p=0.05) (Fig. 2B).

Side effect-related differences in neutralizing activity and recipient characteristics

Our comparison of neutralizing antibody titers against any of the SARS-CoV-2 variants

showed no significant difference in the percentages of participants with and without side

effects, e.g., fever, injection site pain, and fatigue (eFig. 1A in Supplement). However,

the comparison of participant ages and side effects revealed that the median age of the

participants with fever was significantly younger than that of the participants without

fever: 37 years (range 28–64 yrs) versus 46 (29–68) years, p=0.02. There were no

significant between-group differences in the median age of the participants with or

without fatigue (44 years [range 28–68 yrs] vs. 39 [28–64], p=0.97) or regarding the

degrees of injection site pain (none: 41 [31–68] vs. mild: 50 [30–64] vs. moderate: 42

[28–64] vs. severe: 39 [36–51]) (eFig. 1B in Supplement).

Neutralizing activity against three variants at 7 months after two vaccine doses and

after the booster

We determined the positive rates of neutralizing antibody against D614G, Delta, and

Omicron at 2 and 7 months after the two vaccine doses. At 7 months after the

vaccination, 93% of the 82 participants had neutralizing antibody against D614G, 67%

against Delta, and only 6% against Omicron (Fig. 3A). However, surprisingly, after the

booster vaccination, all 72 participants who had received the booster had the

neutralizing antibody against these three variants. Figure 3B illustrates the changes in

the neutralizing antibody titers against these variants during the same period. The

neutralizing antibody titers against D614G and Delta were significantly decreased at 7

months compared to those at 2 months after the second vaccination: D614G, 2.2-fold

(95%CI: 1.9–2.5) lower; Delta, 2.1-fold (95%CI: 1.8–2.4) lower. Although the

neutralizing antibody titers against Omicron also tended to decrease during this period,

no significant difference was observed: median [range] neutralizing antibody titer was

<2 [<2–2] at 2 months versus <2 [<2–4] at 7 months (p=0.54).

Notable, the neutralizing antibody titers after the booster vaccination were much

higher than those at 2 and 7 months post-vaccination against all variants, as follows.

D614G: 13-fold (95%CI: 11–16) and 31-fold (95%CI: 25–38) higher than that at 2 and

7 months, respectively; Delta: 16-fold (13–20) and 35-fold (95%CI: 29–43); Omicron:

32-fold (27–37) and 39-fold (95%CI: 32–46).

Comparison of neutralizing antibody titers after booster vaccination in the three age

groups

We compared the effects of booster vaccination on the neutralizing antibody titers in

each age group (Fig. 4A–C). Seven months after the 2nd vaccination, the neutralizing

antibody titers against D614G and Delta tended to be low in the older age group, and

those against Omicron were much lower in all three age groups. However, the booster

vaccination effectively increased the neutralizing antibody titers for all three variants

including Omicron, regardless of the participants' ages, as follows. The younger,

middle-aged, and older groups values after the booster compared to the values at 7

months were D614G: 28-fold (22–36), 28-fold (95%CI: 18–42), and 47-fold (95%CI:

29–75) greater; those for Delta were 32-fold (95%CI: 23–44), 38-fold (95%CI: 26–55),

and 39-fold (95%CI: 29–52) greater; and those for Omicron were 41-fold (95%CI:

30–56), 43-fold (95%CI: 32–58), and 27-fold (95%CI: 20–36) greater, respectively.

Discussion

We evaluated neutralizing activities against authentic SARS-CoV-2 variants in the sera

of adults vaccinated with two or three doses of the BNT162b2 mRNA vaccine. We also

analyzed the changes in the neutralizing antibody titers after the booster vaccination in

different age groups, and we evaluated the relationships between side effects, the

participants' ages, and the acquired neutralizing antibody titer. This is the first report on

the detailed efficacy of two doses and the booster vaccination for a Japanese population

using authentic viruses including Delta and Omicron.

Our results demonstrated that only 28% of the vaccine recipients had neutralizing

antibodies against Omicron at 2 months after their second vaccination, and the

neutralizing antibody titer against Omicron was significantly lower than those against

all six of the other variants tested. This low positive rate of neutralizing antibody against

Omicron was distinctly different from what we observed for other variants, in which

88%–100% of recipients had the neutralizing antibody against other variants at 2

months after the vaccination, suggesting that the neutralizing antibody against Omicron

cannot be sufficiently induced by only two doses of vaccination, as has been noted in

other reports.[28,29,36,37]

At 7 months after the second vaccination, that tendency was even more

pronounced, with only 6% of the participants showing neutralizing antibodies against

Omicron, suggesting a high risk of breakthrough infection or reinfection. Regarding the

potential cause of this finding, it has been reported that mutations on the spike protein

of Omicron, especially K417N, N440K, G446S, S477N, T478K, E484A, Q493R,

G496S, Q498R, N501Y, and Y505H, were responsible for the variant's broad evasion

from antibodies.[38] In case of the Delta variant, L452R, T478K, and P681R have been

reported to promote the interaction between the spike and angiotensin-converting

enzyme 2 (ACE2) receptor by inducing structural changes in the binding domain,

resulting in immune evasion and increased transmissibility.[39-41] In our present

investigation, 66.7% of the participants had neutralizing antibodies against Delta at 7

months after their second vaccination, suggesting that neutralizing antibodies against

Delta could be more easily induced by two doses of vaccination compared to Omicron.

We also observed that the booster vaccination induced much higher levels of

neutralizing antibodies against Omicron compared to those after two vaccine doses.

This may indicate that the booster vaccination can induce sufficient immunity against

Omicron and the other variants that may appear in the future, by inducing neutralizing

antibodies that recognize common epitopes, and by promoting affinity maturation of the

neutralizing antibody.[42-44]

Our comparison of neutralizing antibody titers among the three age groups at 2

months after the second vaccination showed that neutralizing activity against all seven

tested variants tended to be lower in the older participants (and the activity was

significantly lower only for Delta and Kappa). However, our findings of no significant

difference in the neutralizing antibody titers against Omicron among the present age

groups and that only 32% of the participants in the youngest group had neutralizing

antibodies reflects the universally low neutralizing activity for Omicron, unlike that for

Delta.

However, the increase in the neutralizing antibody level induced by the booster

vaccination did not show a large difference among the three age groups, and the

neutralizing antibody against Omicron was sufficiently induced in all age groups,

suggesting that the effect of the booster vaccination is guaranteed even in older

individuals and that the booster should be recommended for all individuals regardless of

age.

The proportion of participants who had fever or general fatigue after vaccination

in this study increased from the first dose to the third dose of vaccination, whereas the

distribution of injection site pain was almost the same across the doses, indicating that

the vaccinations' systemic side effects such as fever and general fatigue may increase

mainly as a consequence of a stronger immunological response for the second and third

doses.[45-47] However, we did not detect a correlation between the neutralizing antibody

titer induced by two doses of vaccination and vaccination-related side effects, indicating

that fever and other side effects are not specific immune reactions to the viral spike

protein, but rather may be specific to other materials contained in the vaccine. This

relationship remains controversial, and further studies including a booster vaccination

are necessary to evaluate it.[48]

Overall, our findings showed that a sufficient neutralizing antibody titer against

Omicron could not be induced by only two vaccine doses, suggesting a high risk of

breakthrough infection or reinfection. Moreover, the elderly may not be equally

protected from Delta and other variants. Although systemic side effects such as fever

and general fatigue may increase, our study revealed that the third dose of the

vaccination effectively induced neutralizing antibodies against Omicron regardless of

the recipients' age. The booster should therefore be recommended to help prevent the

further spread of the COVID-19 pandemic.

Acknowledgments

We thank Kazuro Sugimura MD, PhD (Superintendent, Hyogo Prefectural Hospital

Agency and Professor, Kobe University) for his full support to promote this study. We

thank BIKEN Innovative Vaccine Research Alliance Laboratories for providing

SARS-CoV-2 B2 strain. We thank the National Institute of Infectious Disease Japan for

providing SARS-CoV-2 Alpha, Beta, Gamma, Delta, Kappa and Omicron variants.

Authors' Statement

All authors attest that they meet the ICMJE criteria for authorship.

Conflicts of Interest

All authors declare no conflicts of interest with respect to this.

Funding

This work was supported by Hyogo Prefectural Government. The funders had no role in

study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

Presentation at meetings

This study has never been presented anywhere.

Contributions

References :

1. World Health Organization. WHO Coronavirus (COVID-19) Dashboard. Accessed 20 January, 2022, https://covid19.who.int

2. Maruki T, Iwamoto N, Kanda K, et al. Two cases of breakthrough SARS-CoV-2 infections caused by the Omicron variant (B.1.1.529 lineage) in international travelers to Japan. *Clin Infect Dis*. Jan 03 2022;doi:10.1093/cid/ciab1072

3. Abbasi J. Omicron Has Reached the US-Here's What Infectious Disease Experts Know About the Variant. *JAMA*. Dec 28 2021;326(24):2460-2462. doi:10.1001/jama.2021.22619

4. Graham F. Daily briefing: Omicron was already spreading in Europe. *Nature*. Dec 01 2021;doi:10.1038/d41586-021-03610-3

5. Gu H, Krishnan P, Ng DYM, et al. Probable Transmission of SARS-CoV-2 Omicron Variant in Quarantine Hotel, Hong Kong, China, November 2021. *Emerg Infect Dis*. Dec 03 2021;28(2)doi:10.3201/eid2802.212422

6. Torjesen I. Covid restrictions tighten as omicron cases double every two to three days. *BMJ*. Dec 09 2021;375:n3051. doi:10.1136/bmj.n3051

7. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. 12 2020;383(27):2603-2615. doi:10.1056/NEJMoa2034577

8. Challen R, Brooks-Pollock E, Read JM, Dyson L, Tsaneva-Atanasova K, Danon L. Risk of mortality in patients infected with SARS-CoV-2 variant of concern 202012/1: matched cohort study. *BMJ*. 03 2021;372:n579. doi:10.1136/bmj.n579

9. Davies NG, Jarvis CI, Edmunds WJ, et al. Increased mortality in community-tested cases of SARS-CoV-2 lineage B.1.1.7. *Nature*. 05 2021;593(7858):270-274. doi:10.1038/s41586-021-03426-1

10. Leung K, Shum MH, Leung GM, Lam TT, Wu JT. Early transmissibility assessment of the N501Y mutant strains of SARS-CoV-2 in the United Kingdom, October to November 2020. *Euro Surveill*. 01 2021;26(1)doi:10.2807/1560-7917.ES.2020.26.1.2002106

11. Cele S, Gazy I, Jackson L, et al. Escape of SARS-CoV-2 501Y.V2 from neutralization by convalescent plasma. *Nature*. 05 2021;593(7857):142-146. doi:10.1038/s41586-021-03471-w

12. Zhou D, Dejnirattisai W, Supasa P, et al. Evidence of escape of SARS-CoV-2 variant B.1.351 from natural and vaccine-induced sera. *Cell*. 04 2021;184(9):2348-2361.e6. doi:10.1016/j.cell.2021.02.037

13. Tegally H, Wilkinson E, Giovanetti M, et al. Detection of a SARS-CoV-2 variant of
concern in South Africa. *Nature*. 04 2021;592(7854):438-443.
doi:10.1038/s41586-021-03402-9
14. Bian L, Gao F, Zhang J, et al. Effects of SARS-CoV-2 variants on vaccine efficacy
and response strategies. *Expert Rev Vaccines*. 04 2021;20(4):365-373.
doi:10.1080/14760584.2021.1903879
15. Faria NR, Mellan TA, Whittaker C, et al. Genomics and epidemiology of the P.1
SARS-CoV-2 lineage in Manaus, Brazil. *Science*. 05 21 2021;372(6544):815-821.
doi:10.1126/science.abh2644
16. Dejnirattisai W, Zhou D, Supasa P, et al. Antibody evasion by the P.1 strain of
SARS-CoV-2. *Cell*. 05 27 2021;184(11):2939-2954.e9.
doi:10.1016/j.cell.2021.03.055
17. Singh J, Rahman SA, Ehtesham NZ, Hira S, Hasnain SE. SARS-CoV-2 variants of
concern are emerging in India. *Nat Med*. Jul 2021;27(7):1131-1133.
doi:10.1038/s41591-021-01397-4
18. Karim SSA, Karim QA. Omicron SARS-CoV-2 variant: a new chapter in the
COVID-19 pandemic. *Lancet*. 12 11 2021;398(10317):2126-2128.
doi:10.1016/S0140-6736(21)02758-6
19. Saxena SK, Kumar S, Ansari S, et al. Characterization of the novel SARS-CoV-2
Omicron (B.1.1.529) Variant of Concern and its global perspective. *J Med Virol*.
Dec 14 2021;doi:10.1002/jmv.27524
20. Chen RE, Zhang X, Case JB, et al. Resistance of SARS-CoV-2 variants to
neutralization by monoclonal and serum-derived polyclonal antibodies. *Nat Med*.
04 2021;27(4):717-726. doi:10.1038/s41591-021-01294-w
21. Wang P, Nair MS, Liu L, et al. Antibody resistance of SARS-CoV-2 variants
B.1.351 and B.1.1.7. *Nature*. 05 2021;593(7857):130-135.
doi:10.1038/s41586-021-03398-2
22. Jangra S, Ye C, Rathnasinghe R, et al. SARS-CoV-2 spike E484K mutation reduces
antibody neutralisation. *Lancet Microbe*. Jul 2021;2(7):e283-e284.
doi:10.1016/S2666-5247(21)00068-9
23. Li Q, Wu J, Nie J, et al. The Impact of Mutations in SARS-CoV-2 Spike on Viral
Infectivity and Antigenicity. *Cell*. 09 03 2020;182(5):1284-1294.e9.
doi:10.1016/j.cell.2020.07.012
24. Johnson BA, Xie X, Bailey AL, et al. Loss of furin cleavage site attenuates
SARS-CoV-2 pathogenesis. *Nature*. 03 2021;591(7849):293-299.
doi:10.1038/s41586-021-03237-4

25. Science Brief: Omicron (B.1.1.529) Variant | CDC, (n.d.). Accessed January 21, 2022, https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/scientific-brief-omicron-variant.html

26. Wang Y, Zhang L, Li Q, et al. The significant immune escape of pseudotyped SARS-CoV-2 variant Omicron. *Emerg Microbes Infect*. Dec 2022;11(1):1-5. doi:10.1080/22221751.2021.2017757

27. Cele S, Jackson L, Khan K, et al. SARS-CoV-2 Omicron has extensive but incomplete escape of Pfizer BNT162b2 elicited neutralization and requires ACE2 for infection. *medRxiv*. Dec 11 2021;doi:10.1101/2021.12.08.21267417

28. Muik A, Lui BG, Wallisch AK, et al. Neutralization of SARS-CoV-2 Omicron by BNT162b2 mRNA vaccine-elicited human sera. *Science*. Jan 18 2022:eabn7591. doi:10.1126/science.abn7591

29. Hoffmann M, Krüger N, Schulz S, et al. The Omicron variant is highly resistant against antibody-mediated neutralization: Implications for control of the COVID-19 pandemic. *Cell*. Dec 24 2021;doi:10.1016/j.cell.2021.12.032

30. Altawalah H. Antibody Responses to Natural SARS-CoV-2 Infection or after COVID-19 Vaccination. *Vaccines (Basel)*. Aug 16 2021;9(8)doi:10.3390/vaccines9080910

31. Tré-Hardy M, Cupaiolo R, Wilmet A, et al. Immunogenicity of mRNA-1273 COVID vaccine after 6 months surveillance in health care workers; a third dose is necessary. *J Infect*. 11 2021;83(5):559-564. doi:10.1016/j.jinf.2021.08.031

32. Bar-On YM, Goldberg Y, Mandel M, et al. Protection of BNT162b2 Vaccine Booster against Covid-19 in Israel. *N Engl J Med*. 10 07 2021;385(15):1393-1400. doi:10.1056/NEJMoa2114255

33. Nagano T, Arii J, Nishimura M, et al. Diligent Medical Activities of a Publicly Designated Medical Institution for Infectious Diseases Pave the Way for Overcoming COVID-19: A Positive Message to People Working at the Cutting Edge. *Clin Infect Dis*. 02 2021;72(4):723-724. doi:10.1093/cid/ciaa694

34. Furukawa K, Arii J, Nishimura M, et al. Seroepidemiological Survey of the Antibody for Severe Acute Respiratory Syndrome Coronavirus 2 with Neutralizing Activity at Hospitals: A Cross-sectional Study in Hyogo Prefecture, Japan. *JMA J*. Jan 2021;4(1):41-49. doi:10.31662/jmaj.2020-0094

35. Furukawa K, Tjan LH, Sutandhio S, et al. Cross-Neutralizing Activity Against SARS-CoV-2 Variants in COVID-19 Patients: Comparison of 4 Waves of the Pandemic in Japan. *Open Forum Infect Dis*. Oct 2021;8(10):ofab430.

448   doi:10.1093/ofid/ofab430

36. Peiris M, Cheng S, Mok CKP, et al. Neutralizing antibody titres to SARS-CoV-2 Omicron variant and wild-type virus in those with past infection or vaccinated or boosted with mRNA BNT162b2 or inactivated CoronaVac vaccines. *Res Sq*. Jan 05 2022;doi:10.21203/rs.3.rs-1207071/v1

37. Yang Y, Gong X, Yang L, et al. Regular and booster vaccination with inactivated vaccines enhance the neutralizing activity against Omicron variant both in the breakthrough infections or vaccinees. *J Infect*. Jan 08 2022;doi:10.1016/j.jinf.2022.01.004

38. Cao Y, Wang J, Jian F, et al. Omicron escapes the majority of existing SARS-CoV-2 neutralizing antibodies. *Nature*. Dec 23 2021;doi:10.1038/s41586-021-04385-3

39. Deng X, Garcia-Knight MA, Khalid MM, et al. Transmission, infectivity, and neutralization of a spike L452R SARS-CoV-2 variant. *Cell*. 06 24 2021;184(13):3426-3437.e8. doi:10.1016/j.cell.2021.04.025

40. Muecksch F, Weisblum Y, Barnes CO, et al. Affinity maturation of SARS-CoV-2 neutralizing antibodies confers potency, breadth, and resilience to viral escape mutations. *Immunity*. 08 10 2021;54(8):1853-1868.e7. doi:10.1016/j.immuni.2021.07.008

41. Saito A, Irie T, Suzuki R, et al. Enhanced fusogenicity and pathogenicity of SARS-CoV-2 Delta P681R mutation. *Nature*. Nov 25 2021;doi:10.1038/s41586-021-04266-9

42. Gaebler C, Wang Z, Lorenzi JCC, et al. Evolution of antibody immunity to SARS-CoV-2. *Nature*. 03 2021;591(7851):639-644. doi:10.1038/s41586-021-03207-w

43. Wang Z, Muecksch F, Schaefer-Babajew D, et al. Naturally enhanced neutralizing breadth against SARS-CoV-2 one year after infection. *Nature*. 07 2021;595(7867):426-431. doi:10.1038/s41586-021-03696-9

44. Schmidt F, Muecksch F, Weisblum Y, et al. Plasma Neutralization of the SARS-CoV-2 Omicron Variant. *N Engl J Med*. Dec 30 2021;doi:10.1056/NEJMc2119641

45. Tré-Hardy M, Cupaiolo R, Papleux E, et al. Reactogenicity, safety and antibody response, after one and two doses of mRNA-1273 in seronegative and seropositive healthcare workers. *J Infect*. 08 2021;83(2):237-279. doi:10.1016/j.jinf.2021.03.025

46. Chapin-Bardales J, Gee J, Myers T. Reactogenicity Following Receipt of mRNA-Based COVID-19 Vaccines. *JAMA*. 06 01 2021;325(21):2201-2202. doi:10.1001/jama.2021.5374

47. Naaber P, Tserel L, Kangro K, et al. Dynamics of antibody response to BNT162b2
vaccine after six months: a longitudinal prospective study. *Lancet Reg Health Eur*.
Nov 2021;10:100208. doi:10.1016/j.lanepe.2021.100208
48. Modenese A, Paduano S, Bargellini A, et al. Neutralizing Anti-SARS-CoV-2
Antibody Titer and Reported Adverse Effects, in a Sample of Italian Nursing Home
Personnel after Two Doses of the BNT162b2 Vaccine Administered Four Weeks
Apart. *Vaccines (Basel)*. Jun 15 2021;9(6)doi:10.3390/vaccines9060652
491

Table 1: Characteristics of participants and side effects of each dose of vaccination.

Characteristics	First trial[a] (n = 82)	Second trial[b] (n = 82)	Third trial[c] (n = 72)
Sex, No. (%)			
Male	71 (87)	71 (87)	61 (85)
Female	11 (13)	11 (13)	11 (15)
Age, years			
Median (IQR)	44 (33-58)	44 (33-58)	44 (33-53)
Age group, No. (%)			
≤ 38 years	31 (38)	31 (38)	31 (43)
39-58 years	32 (39)	32 (39)	27 (38)
≥ 59 years	19 (23)	19 (23)	14 (19)
Days from 2nd dose of vaccination, days (IQR)	53 (46-59)	199 (192-203)	274 (269-278)
Days from booster vaccination, days (IQR)	NA[d]	NA[d]	16 (14-21)

Side effects	1st dose of vaccination (n = 82)	2nd dose of vaccination (n = 82)	Booster vaccination (n = 72)
Fever, No. (%)	0 (0)	16 (20)	24 (33)
Fatigue, No. (%)	4 (5)	14 (17)	45 (63)
Injection site pain, No. (%)	73 (89)	71 (87)	65 (90)
mild	30 (37)	27 (33)	15 (21)

moderate	39 (48)	38 (46)	35 (49)
severe	4 (5)	6 (7)	15 (21)

Table 1 Footnotes:

[a] Samples were collected from June to July 2021

[b] Samples were collected from October to November 2021

[c] Samples were collected in January 2022

[d] NA, not applicable

Figure legends

Fig. 1. Neutralizing antibody against SARS-CoV-2 variants at 2 months post-vaccination. Sera of 82 recipients were tested for neutralizing activity against the SARS-CoV-2 variants D614G, Alpha, Beta, Gamma, Delta, Kappa, and Omicron. The neutralizing antibody titer is represented by the highest serum dilution that did not show any cytopathic effects. **(A)** The positive rate of neutralizing antibody against each variant. Sera with neutralizing activity at twofold dilution (the detection limit) or higher were considered positive. Numbers above the bar indicate the actual positive rate. **(B)** Box plot of the neutralizing antibody titers. The ends of the box represent the upper and lower quartiles, and the median is marked by the *horizontal line* inside the box. The ends of the 'whiskers' represent the minimum and maximum of all data. The titers against the seven variants were compared by the Friedman test and Dunn's multiple comparisons test; two-tailed p-values were calculated.

Fig. 2. Comparison of neutralizing antibodies against SARS-CoV-2 variants among three age groups. The 82 participants were divided into three age groups: 31 participants (37.8%) who were ≤38 years old, 32 participants who were 39–58 years old,

and 19 participants who were ≥59 years old, and the recipients' sera were tested for

neutralizing activities against the seven SARS-CoV-2 variants. (**A**) The positive rates of

neutralizing antibodies against each variant in the three age groups shown. Sera with

neutralizing activity at twofold dilution (detection limit) or higher were considered

positive. Numbers above the bar are the actual positive rate. (**B**) Comparison of

neutralizing antibody titers against each variant among the age groups. Ends of the box

indicate the upper and lower quartiles, and the median is marked by the *horizontal line*

inside the box. Ends of the 'whiskers' = the min. and max. of all data. The

Kruskal-Wallis test and Dunn's multiple comparison test were performed, and two-tailed

p-values were calculated.

Fig. 3. Neutralizing antibody after booster vaccination. Sera of the 72 participants

who received the booster vaccination were tested for neutralizing activity against

D614G, Delta, and Omicron at 2 and 7 months after the two doses of vaccination and

after the booster (third) vaccination. (**A**) Changes in the neutralizing antibody positive

rate against each variant. Sera with neutralizing activity at twofold dilution (detection

limit) or higher were considered positive. Numbers above the bar: the actual positive

rate. (**B**) Changes in the neutralizing antibody titers against each variant. Box ends: the

upper and lower quartiles. The *horizontal line* inside the box: median. Whisker ends: the

min. and max. of all data. The titers were compared by the Friedman test and Dunn's

multiple comparisons test; two-tailed p-values were calculated.

Fig. 4. Changes in the neutralizing antibody titer after booster vaccination in the

age groups. The 72 recipients who received the booster vaccination were divided into

three age groups: 31 participants who were ≤38 years, 27 participants who were 39–58

years, and 14 participants who were ≥59 years. The neutralizing antibody titers against

D614G **(A)**, Delta **(B)**, and Omicron **(C)** at 2 and 7 months after two doses of

vaccination and after the booster vaccination were compared in each age group. Box

ends: the upper and lower quartiles. The *horizontal line* inside the box: median. Whisker

ends: the min. and max. of all data. The Friedman test and Dunn's multiple comparisons

test were performed; two-tailed p-values were calculated.

A Positive rate of neutralizing antibody at 2 months after the vaccination

B Neutralizing antibody titer at 2 months after the vaccination

A Positive rate of neutralizing antibody at 2 months after the vaccination

B Neutralizing antibody titer at 2 months after the vaccination among three different age groups

A Changes of the neutralizing antibody positive rates

B Changes of the neutralizing antibody titers

Changes of the neutralizing antibody titers in three different age groups

A D614G

C Omicron

A   **Neutralizing antibody titer and side effects after the second dose of vaccination**

B   **Age and side effects after the second dose of vaccination**

eFigure 1. Comparison of side effects and neutralizing antibody titers or recipient's age.

Eighty-two recipients were divided according to the appearance of fever, fatigue, and the degree of injection site pain after the second dose of vaccination. The neutralizing antibody titers **(A)** and age of recipients **(B)** were compared by each group. The ends of the box show the upper and lower quartiles, and the median is marked by the horizontal line inside the box. The ends of the "whiskers" represent the minimum and maximum of all data. Lines in eFigure 1B indicate the mean age of recipients in each group. Mann-Whitney U test was performed for the comparison by the presence or absence of fever and fatigue, and Kruskal-Wallis test was performed for the comparison by the degree of injection site pain. Two-tailed p-values were calculated.