Using quantitative magnetic resonance imaging to track cerebral damage in multiple sclerosis: a longitudinal study

Nora Vandeleene[a], Camille Guillemin[a,b], Solène Dauby[a,c], Florence Requier[a,b], Maëlle Charonitis[a,b], Daphne Chylinski[a,b], Evelyne Balteau[a], Pierre Maquet[a,c], Emilie Lommers[a,c*], Christophe Phillips[a,d*]

[a] GIGA CRC in vivo Imaging, University of Liège, Liège, Belgium
[b] Psychology and Cognitive Neuroscience Research Unit, University of Liège, Liège, Belgium
[c] Clinical Neuroimmunology Unit, Neurology Department, CHU Liège, Belgium
[d] GIGA in silico Medicine, University of Liège, Liège, Belgium

∗   These authors equally contributed to the work.

Corresponding author: Christophe PHILLIPS
                                 GIGA – CRC in vivo imaging B30
                                 Allée du 6 août
                                 4000 Liège
                                 Belgium

Acknowledgements
The authors are particularly thankful for the patients who took part in this study.
N.V., E.L. and C.P. are supported by the Fonds de la Recherche Scientifique (F.R.S-FNRS Belgium).

Data availability
The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

Conflict interests
None.

Abbreviations:
ARoC: Annual Rate of Change, **BPF**: Brain Parenchymal Fraction, **CDP**: Confirmed Disability Progression, **cMRI**: Conventional Magnetic Resonance Imaging, **CNS**: Central Nervous System, **GMF**: Grey Matter Fraction, **GLMM**: General Linear Mixed Model, **LF**: Lesion Fraction, **MPM**: Multi-Parameter Mapping, **MRI**: Magnetic Resonance Imaging, **MS**: Multiple Sclerosis, **MT**: Magnetization Transfer, **MTR**:

Magnetization Transfer Ratio, **MTsat**: Saturated Magnetization Transfer, **NABT**:
Normal Appearing Brain Tissue, **NACGM**: Normal Appearing Cortical Grey Matter,
NADGM: Normal Appearing Deep Grey matter, **NAWM**: Normal Appearing White
Matter, **NEDA-3**: No Evidence of Disease Activity, **RRMS**: Relapsing-Remitting
Multiple Sclerosis, **PD**: Proton Density, **PMS**: Progressive Multiple Sclerosis, **qMRI**:
Quantitative Magnetic Resonance Imaging, **R1**: Longitudinal Relaxation Rate (1/T1),
R2*: Transverse Relaxation Rate (1/T2*), **TIV**: Total Intracranial Volume, **TPM**: Tissue
Probability Map, **US:** Unified Segmentation, **USwL**: Unified Segmentation with Lesion

Abstract

Quantitative MRI quantifies tissue microstructural properties and improves the characterization of cerebral tissue damages. With an MPM protocol, 4 parameter maps are constructed: MTsat, PD, R1 and R2*, reflecting tissue physical properties associated with iron and myelin contents. Here, we used qMRI to investigate the microstructural longitudinal changes in MS.

Seventeen MS patients (age 25-65, 11 RRMS) were scanned on a 3T MRI, in two sessions separated with a median of 30 months, and the parameters evolution was evaluated within several tissue classes: NAWM, NACGM and NADGM, as well as focal WM lesions. Tissue segmentation was performed using US-with-Lesion, an adapted version of the Unified Segmentation (US) algorithm accounting for lesions. An individual annual rate of change for each qMRI parameter was computed, and its correlation to clinical status was evaluated. For WM plaques, three areas were defined, and a GLMM tested the effect of area, time points, and their interaction on each median qMRI parameter value.

Patients with a better clinical evolution, i.e., clinically stable or improving state, showed positive annual rate of change in MT and R2* within NAWM and NACGM, suggesting repair mechanisms in terms of increased myelin content and/or axonal density as well as edema/inflammation resorption. When examining WM lesions, qMRI parameters within surrounding NAWM showed microstructural modifications, even before any focal lesion is visible on conventional FLAIR MRI.

The results illustrate the benefit of multiple qMRI data in monitoring subtle changes within normal appearing brain tissues and plaque dynamics in relation with tissue repair or disease progression.

74   Keywords:

75   Quantitative MRI, relaxometry, longitudinal analysis, multiple sclerosis.

77   Key points:

78   •   Patients with a better clinical evolution showed microstructural improvement

79   in term of MT and R2* increase in their normal appearing tissue, suggesting

80   repair mechanisms.

81   •   Using qMRI allows to detect modifications in tissue microstructure in normal

82   appearing tissues surrounding lesions several months before they are visible

83   on conventional MRI.

1. Introduction

Multiple sclerosis (MS) is a chronic autoimmune disease of the central nervous system (CNS). Plaques are the pathological hallmark of MS. They are spread in acute, focal, disseminated and recurrent way throughout the CNS and harbor variable degrees of inflammation, demyelination, gliosis and axonal injury [1, 2]. Plaques are not restricted to the white matter (WM), but are also present in the cortex and deep grey matter (GM) [3-5]. The progressive accumulation of disability principally correlates with the early, diffuse and chronic inflammation within the normal appearing white matter (NAWM) and grey matter (NAGM) that is ultimately responsible for diffuse neuro-axonal loss and neurodegeneration [3, 4, 6]. By contrast, effective repair mechanisms can occur within focal lesions but probably also in normal appearing brain tissue (NABT) [7]. However, our understanding of these complex processes is still fragmentary. The difficulty of acquiring histopathological data on MS patients at various stages of the disease makes it challenging to describe the time course of injury and potential repair mechanisms in MS. Consequently, there is an important need for new imaging techniques to improve in-vivo monitoring of lesion formation, progression and repair in MS [8].

Conventional MRI (cMRI) readily depicts focal WM lesions on T2/FLAIR sequences and is able to distinguish between acute and allegedly chronic lesions. T2-hyperintensities in cMRI constitute the keystone of McDonald diagnostic criteria [9] and also make an important contribution to the monitoring of WM lesion burden. Unfortunately, cMRI sequences are not able to efficiently assess cortical lesions or detect diffuse changes in NABT. Quantitative MRI (qMRI) potentially overcomes these limitations by quantifying physical microstructural properties of cerebral tissue in standardized units. QMRI is more sensitive but also more specific to microstructural properties of CNS tissues. Magnetization transfer ratio (MTR) was

regularly linked to cerebral macromolecular content detected by a greater percentage loss

of magnetization in voxels with a higher myelin content and axons density [10-12]. Post-

mortem studies comparing the relative contribution of these two factors indicate that

myelin has a stronger and more direct influence on MTR than the axonal density, which is

considered as a T1-dependent effect. Tissue water content (inflammation, edema…),

another T1-dependent effect, also accounts for MTR variability [11, 13, 14]. However, the

MT saturation (MTsat) map offers a measure which, unlike MTR, is minimally affected by

longitudinal relaxation and B1 mapping inhomogeneities [15], increasing its sensitivity to

myelin content. Moreover, the brain contrast to noise ratio is larger for the MTsat map than

for MTR, thus improving brain tissue segmentation in healthy subjects [11, 16]. R2* was

usually linked to iron and myelin contents, as paramagnetic iron and diamagnetic myelin

generate microscopic field gradients in the CNS, thus shortening T2* and increasing R2*

(1/T2*). Orientation and density of myelin fibers are also a determining factor of R2* values

[17-19]. Iron is probably a key factor in MS monitoring as it was shown that aberrant iron

metabolism occurs in the course of the disease [20]. Particularly increased iron

concentration within chronic active lesions (i.e iron rim lesion) or deep grey matter

structures was observed [20]. Regarding the longitudinal relaxation rate R1 (1/T1) in the

CNS, its three major determinants are tissue myelination and associated axons, iron, and

extracellular water contents [19, 21, 22]. Finally, proton density (PD) mostly reflects the free

water content of the brain [23].

We have previously shown that a multivariate qMRI approach is useful to assess NABT

microstructural alteration in a cross-sectional study comparing MS patients to healthy

controls [24, 25]. Because each qMRI parameter is differently sensitive to histologically

measured iron and myelin contents, this approach constitutes a fundamental tool for

longitudinal *in vivo* monitoring of MS lesions and NABT evolution at the tissue

microstructural level.

In this longitudinal study, we investigate the evolution of four simultaneously acquired qMRI

parameters (MTsat, PD, R1, R2*) within NABT and WM lesions of 17 MS patients - relapsing

remitting (RRMS) and progressive MS (PMS) - who were scanned two times with at least a

one-year interval, following the same multi-parameter mapping (MPM) protocol at 3 Tesla

[10, 26]. Segmentation of different cerebral tissue classes was computed using an advanced

segmentation technique called Unified Segmentation with Lesion (USwL), an updated

version of the traditional Unified Segmentation (US) algorithm from SPM. This pipeline

accounts for lesions and relies on quantitative parameter maps rather than the standard

weighted images.

Here, we assessed the time course of parameter values in several tissue classes: NAWM,

normal appearing cortical and deep GM (NACGM and NADGM) as well as focal WM lesions.

In addition, we related the changes in NABT to clinical course. We also looked at the tissue

directly surrounding WM lesions, that changed status over time, i.e., switching from NAWM

to lesion.

2. Materials and methods

2.1 Population

Seventeen patients, recruited at the specialized MS outpatient clinic of the CHU Liège,

Belgium, with a diagnosis of MS according to the McDonald criteria 2010 [27], were gathered

from two studies: ten of them were part of the work reported by Lommers et al. 2019 [24],

the other seven were recruited from another MS study taking place at the GIGA Cyclotron

Research Centre – In Vivo Imaging (Liège, Belgium) [28]. For the first study (10 subjects), the

inclusion criteria were (1) age between 18 and 65 years ; (2) Expanded Disability Status Scale

(EDSS) inferior or equal to 6.5 ; (3) absence of relapse within the previous four weeks ; Both

RRMS and PMS patients were recruited. The second study (7 subjects) differs a bit as it

comprises only RRMS patients, and the inclusion criteria were (1) age between 18 and 45, (2)

EDSS between 0 and 4, (3) absence of relapse for at least 6 months prior to the study, (4)

disease duration was below or equal to 5 years. For both studies, compatibility with MRI

and absence of other neurological/psychiatric diseases were required. These studies were

approved by the local ethics committee (approval numbers B707201213806 and

B707201835630, respectively). All patients were followed up and scanned twice on the same

3T MRI scanner, every 1 to 3 years. For each of the 17 MS patients, data from two MRI

sessions were available, at T0 and T1. This cohort included 11 RRMS and 6 (primary and

secondary) PMS patients. Thirteen were receiving disease-modifying treatments (DMTs). The

patients' median age was 36 years (range: 25-65) and the median time interval between two

scans was 30 months (range: 14-61). Demographic data appears in Table 1. Extended

individual information appears in Supplementary data.

172
2.2 MR image acquisition

MRI data were acquired on a 3T whole-body MRI-scanner (Magnetom Prisma, Siemens

Medical Solutions, Erlangen, Germany). The whole-brain MRI acquisitions included a multi-

parameter mapping protocol (MPM), from which one can simultaneously estimate

(semi)quantitative maps of magnetization transfer saturation (MTsat), proton density (PD),

transverse relaxation (R1) and effective longitudinal relaxation (R2*). This protocol arising

from an international collaborative effort [10, 26], has already been used to study brain

microstructure in various conditions including normal aging [26, 9, 30], brain tumor [31],

Parkinson's disease [32-34] as well as MS. It consists of three co-localized 3D multi-echo fast

low angle shot (FLASH) acquisitions at 1mm³ resolution and two additional calibration

sequences to correct for inhomogeneities in the RF transmit field [35, 36]. The FLASH

datasets were acquired with predominantly PD, T1 and MT weighting, referred to in the

following as PDw, T1w and MTw, at multiple echo times. All three had high bandwidth to

minimize off-resonance and chemical shift artifacts. Volumes were acquired in 176 sagittal

slices using a 256x224 voxel matrix. GRAPPA parallel imaging was combined with partial

Fourier acquisition to speed up acquisition time to approximately 20 min. An additional

FLAIR sequence was recorded with spatial resolution 1mm³ and

TR/TE/TI=5000ms/516ms/1800ms. Extra B1 field mapping images (transmit B1+ and receive

B1- fields) were also acquired to reduce spatial inhomogeneities related to B1 effect. This

was essential for proper quantification of T1 (or R1=1/T1) in particular. Finally, B0 field

mapping images, corresponding to both magnitude images and pre-subtracted phase image,

were acquired for image distortions corrections. A summary of the acquisition parameters

appears in Supplementary data.

2.3 MR image processing

All data processing was performed in Matlab (The MathWorks Inc., Natick, MA, USA) using

SPM12 (www.fil.ion.ucl.ac.uk/spm) and three additional dedicated SPM extensions: the

Lesion Segmentation Tool (LST) version 1.2.3 (www.statisticalmodelling.de/lst.html) [37], the

"quantitative MRI and in vivo histology using MRI" toolbox (hMRI, http://hmri.info) [10], and

"US-with-Lesion" tool (USwL, https://github.com/CyclotronResearchCentre/USwLesion).

Quantitative maps - MTsat, PD, R1 and R2* - were estimated using the hMRI toolbox. T1w,

PDw and MTw images acquired at multiple echo times (TE) were extrapolated to TE=0 to

increase signal-to-noise ratio and remove the otherwise remaining R2* bias [10, 24, 38]. The

TE=0 extrapolated MTw, PDw and T1w images were used to calculate MT saturation, R1 and

apparent signal amplitude A* maps. PD map was derived from A* map, which is proportional

to proton density. All quantitative maps were corrected for inhomogeneities from local RF

transmit field (B1+), and R1 quantitative maps were further corrected for imperfect RF

spoiling using the strategy of Preibisch and Deichmann [39]. The receive bias field map (B1-)

was used to correct PD maps for instrumental biases. The R2* map was estimated from all

three multi-echo series (MTw, PDw and R1w) using the ESTATICS model [38].

After generating quantitative maps using the hMRI toolbox for all sessions, spatial

preprocessing involved the following steps (Figure 1): within-patient registration brought the

two serial MR data sets into the individual T0 space, using the longitudinal registration tool

from SPM [40]. For each individual patient, a preliminary WM lesion mask was generated

based on FLAIR and T1w images by the lesion growth algorithm implemented in the LST

toolbox [37], followed by manual corrections by an MS expert (EL) to remove

aberrant/artefactual lesion detections [24]. The images were then segmented using the

USwL toolbox, which consists of an extended version of the traditional Unified Segmentation

(US) algorithm [41] and includes an additional tissue class representing the WM lesion(s).

The USwL method internally generates a subject-specific extended set of tissue probability

maps (TPM) [42]: an extra tissue class, based on the smoothed preliminary lesion mask

warped into template space (using cost function masking during normalization [43]), is

added to account for the lesion, and the original white matter prior map is updated

accordingly [44]. The grey matter TPM was not updated due to a very low number of lesions

present in the cortical ribbon. Multi-channel segmentation was conducted, using MTsat, PD,

R1 and FLAIR images. This pipeline did not use the PD-, T1- and MT-weighted images

acquired for the MPM maps construction, but the parametric maps themselves instead. In

this way, voxels do not depict MR intensities but rather physical quantitative parameters.

The method generated the segmented tissue classes (*a posteriori* tissue, including lesion,

probability maps), as well as spatial warping into standard template space. The preliminary

lesion mask was used as input for the first session data (at T0) then the *a posteriori* lesion

map generated at this initial step served as prior to the subsequent session (at T1).

Segmentation teased out the different tissue classes of interest: NAWM, NACGM and

NADGM, as well as WM lesions. To analyze the microstructure within those tissue classes, *a*

posteriori tissue maps were binarized and tissue-specific independent masks were

constructed: each voxel is assigned to one single tissue class with the highest probability for

that voxel (provided that this probability was above 0.2). The lesion binary mask was further

cleaned for lesions <10mm³ which likely resulted from segmentation errors. Finally, binarized

tissue class masks were in turn applied on the MPM maps to extract voxel values inside

them.

2.4 Brain volume change

Volumetric changes were investigated using the USwL *a posteriori* tissue probability maps.

The following measures of brain volume were computed for each session of each

participant: (1) Total intra-cranial volume (TIV) = volume (NAWM + GM + CSF + lesions), (2)

brain parenchymal fraction (BPF) = volume (NAWM + GM + lesions)/TIV, (3) GM fraction

(GMF) = volume (GM)/TIV, and (4) lesion fraction (LF) = volume (lesion)/TIV. The percentage

of change between both scanning sessions was evaluated for each volumetric measurement,

then annualized changes were computed by dividing these measures by scan intervals (in

years). Results were directly analysed with a t-test (testing if significantly different from 0 at

$p < .05$), but also in the same way as the normal appearing tissues MR parameters in

relation to the patients' clinical status (see next section).

2.5 Analysis of normal appearing tissues

The median value of quantitative MRI parameters was extracted from the three normal

appearing tissues (NAWM, NACGM and NADGM), and an individual annual rate of change

(ARoC) was computed for each parameter in each tissue class, based on the initial and final

values and accounting for the time interval (in years) between scans. This rate of change in

qMRI parameters served as dependent variable in a general linear model testing the effect

of clinical status:

$$Y = \beta_0 + \beta_1 X_{status} + \epsilon$$

Y is the ARoC for a qMRI parameter and tissue class, β's are the regression parameters

corresponding to the associated regressor (with β_0 the intercept), and ϵ the residuals.

X_{status} is a binary categorical variable representing the patient's disease activity status: a

status score of 1 was assigned to patients stable or improving from T0 to T1.

This patient status X_{status} was derived from one score of disease activity: NEDA-3 (No

Evidence of Disease Activity [45]), a composite of three related measures of disease activity.

A score of 0 was assigned in the presence of new clinical relapses (only concerning RRMS

patients) and/or MRI activity (new or enlarged lesions visible on FLAIR T2 or Gadolinium-

enhanced images) and/or sustained disability progression over six months based on

Expanded Disability Status Scale (EDSS). For both RRMS and PMS patients, disability

progression was defined as a 1.0-point increase if the EDSS score was ≤ 4.0 at baseline and

as a 0.5 point increased if the baseline EDSS score was > 4.0. The threshold of 4.0 was

proposed in this study because it is considered as a milestone regarding ambulatory

performance.

NEDA-3 and was evaluated at mid- and end-scanning interval, and a final status score of 0

was given only to patients for which disease activity or progression was noted in both cases,

indicating a clear progression of the disease over the whole interscan interval.

The influence of several clinical measurements such as 25 FWT, 9HPT and SDMT was also

considered to refine the evaluation of disease activity. However complete data were lacking

for several patients. Moreover, when available, these additional clinical parameters did not

modify the final X_{status}. Longitudinal clinical information allowing to derive the disease

activity status for each subject appears in Supplementary data.

Permutation tests were employed for inferences [46]. R-squared value was tested against

computed statistics after permutation of the data. For a number n of permutations, the

X_{status} values were randomly shuffled (constructing a new regressor written X_{status}^{π}), tested

against the unchanged response Y, and generating each time a permuted R-squared value

(noted R_{π}, R_{obs} being the true R-squared value computed without permutation of the data).

The condition $X_{status} \neq X_{status}^{\pi}$ is verified at each permutation. After n permutations (with

$n = 5000$ in this study), a p-value was computed based on the following formula:

$$p = \frac{\#(R_{\pi} > R_{obs})}{n + 1}$$

which estimates the probability of obtaining R_{obs} under the null hypothesis that Y is not

correlated to X_{status}. The null hypothesis is rejected if $p < .05$ FDR-corrected for multiple

comparisons [47], for the 12 tests performed (3 tissue classes with 4 qMRI parameters).

Two-tailed t-tests were applied *post-hoc* on the significant results of permutation tests to

compare the ARoC distribution between disease status, i.e., $X_{status} = 0$ against $X_{status} = 1$.

Inferences were conducted at a significance level of .05.

The same pipeline was applied to the brain volumetric changes (BPF, GMF and LF) to test

their correlation to the disease activity status.

2.6 Analysis of lesions and peripheral tissues

For white matter lesions analysis, we did not use ARoC but exploited directly the qMRI

parameters voxel values. Importantly, with USwL, the prior lesion mask is only used in a

probabilistic way and the estimated posterior lesion map, obtained using MTsat, PD, R1 and

FLAIR images, typically showed more extended lesion than clinically visible on the FLAIR

image alone.

Therefore, we separated focal lesions detected on FLAIR images, with LST and visual

inspection, from their peripheral regions detected on qMRI maps. Two different peripheral

regions were considered: one for each time point (T0 and T1). Therefore, at T0, three distinct

lesion-related regions were isolated:

• The lesions, as clinically defined, corresponding to hyperintensity on conventional FLAIR

MR image acquired at T0 (referred to as 'focal FLAIR lesion').

• The peripheral region detected on qMRI maps at T0, not including the FLAIR lesion

(referred to as 'initial peripheral lesion').

• The peripheral region, detected on qMRI maps at follow up (appearing hyperintense on

FLAIR images or not); this was computed by masking out the T1 lesion mask with the T0

lesion mask (referred to as 'later peripheral lesion'). This region allows us to determine

whether its microstructure at T0 forebodes a full-blown plaque, detectable during

follow up.

The three areas were compared between each other and with NAWM, in order to

characterize them on a microstructural basis (Figure 2). For an accurate lesion-by-lesion

analysis, only enlarging lesions, i.e., present in the three masks, were considered for these

comparisons.

NAWM region consisted of all white matter voxels which did not belong to any of the three

lesion-related regions. The four areas are not overlapping as no voxel could belong to more

than one class at the same time.

For all participants, MTsat, PD, R1 and R2* median values were extracted from each lesion

area, considering lesions individually (between 2 and 66 measurements per subject).

Similarly, the median qMRI values within NAWM were also extracted (one measurement per

subject). These values were extracted from T0 and T1 scans separately. Statistical analyses

were performed in SAS 9.4 (SAS Institute, Cary, NC). None of the qMRI parameter was

normally distributed, therefore we applied a log transformation on each of them prior to

statistical analysis. For each qMRI parameter, a separate Generalized Linear Mixed Model

(GLMM) tested the effect of areas (NAWM and the three lesion-related areas), and time

points (T0 and T1), as well as their interaction (i.e., area*time), on the median qMRI

parameter value, with a first-order autoregressive variance/covariance model and

participants as a random factor (intercept). The degrees of freedom were estimated using

Kenward-Roger's method. Statistical significance was estimated at $p < .05$ after adjustment

for multiple comparison using Tukey's procedure.

3. Results

338   **3.1 Volume changes**

339   BPF annually decreased between T0 and T1 by -0.67 ± 1.12% (significantly different from

340   zero; paired-sample t-tests; $t(16) = 2.57; p = .0204$) whereas LF increased by 22.88 ±

341   26.13% ($t(16) = -3.70; p = .0019$). GMF non-significantly decreased by -0.30 ± 1.44%.

342   **3.2 Analysis of normal appearing tissues**

As expected, changes in MTsat and R2* within NABT between T0 and T1 varied across

subjects (Figure 3). PD and R1 exhibited similar behaviors, see Supplementary data.

At the group level, with the GLM analysis and permutation inference, we observed that the

ARoC of MTsat and R2* positively regressed with disease status as follows (Table 2): MTsat

in NAWM and NACGM and R2* in NAWM significantly increased in patients who fare well

$(X_{status} = 1)$.

Post-hoc t-tests applied on these significant results for a clearer illustration of the difference

in disease status (Figure 4) were all significant at a level of .05.

Regarding BPF and LF, their correlation to the disease activity status was not significant

(Table 2), suggesting that qMRI parameters are more sensitive to subtle microstructural

changes in NABT over time than global morphological measurements

3.3 Analysis of lesion microstructure

The number of enlarging WM lesions between T0 and T1 varied from 2 to 66 across patients, for a

total of 741 identified enlarging lesions among all subject, corresponding on average among patients

to 63% (±31%) of the amount of initial focal lesions. The number of enlarging lesions did not

significantly differ between patients' disease status groups $(t(15) = .244, p = .811)$.

GLMMs found a significant effect of areas (3 lesional regions and NAWM) for MT, R1, R2* and PD

median (MT: $F_3 = 35.34$, $p < .0001$, PD: $F_3 = 68.03$, $p < .0001$, R1: $F_3 = 40.26$, $p < .0001$,

R2*: $F_3 = 32.32$, $p < .0001$). By contrast, neither time effect (T0 vs T1; MT: $F_3 = 0.36$, $p = .5481$,

PD: $F_3 = 1.20$, $p = .2735$, R1: $F_3 = 2.05$, $p = .1520$, R2*: $F_3 = 2.86$, $p = .0911$), nor the

area*time interaction (MT: $F_3 = 0.09$, $p = .9671$, PD: $F_3 = 0.14$, $p = .9346$, R1: $F_3 = 0.14$, $p = $

.9331, R2*: $F_3 = 0.40$, $p = .7565$) were significant, suggesting the microstructural stability of the

initial lesion core. *Post-hoc* tests confirmed significant differences between the four tissue areas.

At T0 and T1, MTsat, R1 and R2* values were significantly larger in the initial peripheral

lesion than FLAIR lesion, in the later peripheral lesion than the initial one, and in the NAWM

than later peripheral lesion. The reverse was observed for PD. The significant difference in

parameters between initial and later peripheral lesion at T0 suggests that subtle

microstructural changes appear in the periphery of the initial lesion, months before their

detection as focal FLAIR lesions at T1. Adjusted *p*-values appear in Figure 5. Detailed

statistical results of the GLMM's appear in Supplementary data.

4. Discussion

This longitudinal study followed up volumetric data and qMRI brain metrics (MTsat, PD, R1,

R2*) in 17 MS patients for a median time interval of 30 months. The main results are

threefold. First, the microstructure of normal appearing brain tissues changes over time and

these modifications concur with, and potentially drive, clinical evolution. This critical finding

suggests that repair mechanism and edema resorption can be monitored *in vivo*. Second, the

microstructure within WM plaques is remarkably heterogeneous. Importantly, at their

periphery, microstructural alterations foreshadow their expansion, as detected by

conventional MRI. Third, as expected, we observed a small but significant brain atrophy and

lesion load increase with time.

Quantitative MRI parameter time course within NABT

In this study, we used a multiparameter mapping protocol that was gradually optimized and

validated for multi-centric studies [48]. It provides high-resolution maps of multiple qMRI

parameters from data acquired during a single scanning session of acceptable duration. A

number of cross-sectional studies using a combination of MT, R1, R2* or PD parameters

reported significant changes in the microstructure of NABT in MS [24, 49-55]. By contrast,

longitudinal analyses of multiparameter qMRI data are scarce. A progressive shortening of

T2/T2* [56] or increase in R2* [57-59] was reported within the basal ganglia, suggesting

increased of myelin and/or iron contents as well as edema resorption. Likewise, PD and T1

increased within a year, suggesting a demyelination and/or axonal loss [60]. MTR

progressively decreases in NAWM of MS patients over one [61] or two years [62]. These

abnormalities tend to be more pronounced in progressive phenotypes [63] and were

associated to a slow, diffuse and global myelin pathology.

Here, we showed that MTsat within NAWM and NACGM and R2* values within NAWM

increase in clinically stable or improving patients. Because MTsat and R2* both correlate

with myelin content [11, 29, 30, 64-66], our results suggest repair mechanisms within NABT

of patients who are responding to disease modifying treatments, despite the initial

myelin/axonal loss and independently from WM focal lesion evolution. Such increases could

also be explained by an edema/inflammation resorption, but less likely than myelin/axonal

density changes since MTsat is the least dependent to water content among the four qMRI

parameters. These results echo cross-sectional analyses showing that healthy controls (HC)

have higher MTsat and R2* values within the same tissue classes compared to MS patients

[24]. Annual rates of change of R1 and PD within NABT were not significantly associated

with the individual clinical status in this study, although R1 reduction within NABT has

already been reported in cross sectional [24, 52, 53] and longitudinal [60] studies comparing

MS subjects to HC.

Lesion microstructure

Focal inflammatory demyelinating lesions have been extensively characterized and are

traditionally classified as active, chronic active (smoldering) or inactive plaques according to

the presence and distribution of plaque-infiltrating macrophages/microglia [67-69]. Focal

WM pathology is a constantly evolving process including episodes of demyelination and

remyelination but also accumulation of irreversible axonal damage. Age, disease duration,

clinical phenotype as well as disease modifying treatment all contribute to the dynamic

nature of focal WM pathology [68, 70]. This accounts for the large inter- and intra-individual

heterogeneity of MS, which conventional MRI is largely unable to capture. By contrast,

quantitative MRI parameters are sensitive to myelin, axonal as well as iron contents and

appear as promising markers of plaque dynamics. For instance, MTR was shown to sharply

decrease within gadolinium enhancing lesions before recovering during the subsequent

months [71-73]. Likewise, reduction of MTR within NAWM, days to weeks before the

formation of a new active lesion was also demonstrated [74, 75], and long-term MTR

changes in WM plaques were observed in relation with disease progression [63, 76]. The

present study broadens the quantitative characterization of plaque dynamics, in keeping

with previous longitudinal studies [56, 77]. Two important findings emerge from the results.

First, qMRI refines lesion segmentation, as compared to the processing based on the sole

FLAIR image. In consequence, the initial lesion revealed by qMRI is typically wider that the

plaque detected in FLAIR. Its periphery is characterized by a decrease in MTsat and R2* as

compared to NAWM, suggesting an incipient demyelination, reminiscent of the so-called

'periplaques' [78]. Moreover, MTsat, R2* and R1 values progressively decrease from NAWM

to plaque core, suggesting a centripetal loss of myelin content. Second, plaque

microstructure is altered in plaque periphery before any observable change in conventional

MRI signals. This finding suggests, in keeping with neuropathological observations [68, 70,

79, 80] that subclinical ongoing inflammation and/or demyelination takes place in the

periphery of an active plaque, well before it is detectable on FLAIR or T1 post-gadolinium

sequences. If confirmed on larger population samples, this finding might significantly modify

treatment management in MS patients.

Oddly enough, plaque qMRI parameters did not significantly change across time. Because

iron concentration increases within chronic active or smoldering lesions [81, 82], we were

expecting a progressive increase in R2* value. This negative result might be due to the small

sample size, the short period of follow up or the limited sensitivity of R2* to local iron

concentration as compared to the combined use of R2* and quantitative susceptibility

mapping (QSM) [18].

Volumetric Data

CNS atrophy occurs in all stages of MS, since the preclinical phase of the disease and

progresses throughout its course, at a much higher rate than one reported in normal aging

[83-86]. In this study, the annual brain percentage volume loss at the group level was 0.67%,

which is in line with previous publications [87]. We also showed a significant increase in

lesion fraction. Volumetric data (ARoC's) were highly variable across subjects: changes in BPF

range from -2.52 to 1.17% and LF from -0.78 to 103.06%. This variability arises from a large

number of factors which do not necessarily relate to MS: age, disease duration, disease

phenotype, disease modifying treatment, circadian rhythm, hydration… [85, 86].

Moreover, annual changes in brain parenchymal fraction as well as lesion fraction only

partially correlated to patients' disease status, in accordance with a large amount of

publications [60, 88]. This highlights the lack of specificity and sensitivity of volumetric

measurements, at least at the individual level.

Study limitation

As mentioned earlier, the small size and heterogeneous aspect of the present dataset

constitute major limitations of this study. Indeed, it is composed of only 17 patients, with a

rather broad range of characteristics such as age, disease duration, disease phenotype,

disease modifying treatment, etc., which are known to influence the disability state of the

patient and thus their ability to put together repair mechanisms within cerebral tissues [1,

68-70, 89, 90]. In addition, the time interval between two scanning sessions varied rather

widely across patients (between 14 and 61 months), although it was brought back to an

annual rate where possible. All of these parameters were imposed by standard clinical follow

up. Therefore, these results should not be over-interpreted but are nevertheless promising

and call for a replication with a larger and more homogeneous or controlled set of MS

patients. Larger longitudinal studies are currently being held and will probably confirm these

preliminary results.

The second significative limitation is the absence of longitudinal MRI data acquired in a

control group of healthy subjects. However, we considered that literature of longitudinal

studies of healthy subjects that analysed tissue microstructure could constitute a solution for

comparison with MS patients. For example, in Bonnier et al. (2017) [56], the control group

did not show any significant differences regarding T1, T2* or MTR measurements over two

years, and the median age of their group is quite similar to ours (34,3 vs 36 years). Also, in

Elkady et al. (2018) [57], they found no longitudinal R2* effect in their control groups, even

with an age range superior to ours. Moreover, the median age of our population (< 60

years), as well as the short period between two scanning sessions (median of 14 months),

suggests that microstructural alterations would not be noticeable in a healthy participants

group, as many quantitative ageing studies detected differences over much larger time

periods [26, 29, 91].

5. Conclusion

These preliminary results highlight the relevance of multiple qMRI data in the monitoring of

MS disease, highlighting subtle changes within NABT and plaque dynamics in relation with

repair or disease progression. Of course, large scale longitudinal study would be needed to

reproduce these findings and better exploit the full potential of qMRI parameters.

References

[1] Lassmann, Hans. "Pathology and Disease Mechanisms in Different Stages of Multiple Sclerosis." *Journal of the Neurological Sciences* 333, no. 1–2 (October 2013): 1–4. https://doi.org/10.1016/j.jns.2013.05.010.

[2] Trapp, Bruce D., John Peterson, Richard M. Ransohoff, Richard Rudick, Sverre Mörk, and Lars Bö. "Axonal Transection in the Lesions of Multiple Sclerosis." *New England Journal of Medicine* 338, no. 5 (January 29, 1998): 278–85. https://doi.org/10.1056/NEJM199801293380502.

[3] Haider, Lukas, Tobias Zrzavy, Simon Hametner, Romana Höftberger, Francesca Bagnato, Günther Grabner, Siegfried Trattnig, Sabine Pfeifenbring, Wolfgang Brück, and Hans Lassmann. "The Topograpy of Demyelination and Neurodegeneration in the Multiple Sclerosis Brain." *Brain* 139, no. 3 (March 2016): 807–15. https://doi.org/10.1093/brain/awv398.

[4] Kutzelnigg, Alexandra, Claudia F. Lucchinetti, Christine Stadelmann, Wolfgang Brück, Helmut Rauschka, Markus Bergmann, Manfred Schmidbauer, Joseph E. Parisi, and Hans Lassmann. "Cortical Demyelination and Diffuse White Matter Injury in Multiple Sclerosis." *Brain* 128, no. 11 (November 1, 2005): 2705–12. https://doi.org/10.1093/brain/awh641.

[5] Gh Popescu, Bogdan F, and Claudia F Lucchinetti. "Meningeal and Cortical Grey Matter Pathology in Multiple Sclerosis." *BMC Neurology* 12, no. 1 (December 2012): 11. https://doi.org/10.1186/1471-2377-12-11.

[6] Frischer, Josa M., Stephan Bramow, Assunta Dal-Bianco, Claudia F. Lucchinetti, Helmut

Rauschka, Manfred Schmidbauer, Henning Laursen, Per Soelberg Sorensen, and Hans

Lassmann. "The Relation between Inflammation and Neurodegeneration in Multiple

Sclerosis Brains." *Brain* 132, no. 5 (May 2009): 1175–89.

https://doi.org/10.1093/brain/awp070.

[7] Brown, Robert A., Sridar Narayanan, and Douglas L. Arnold. "Imaging of Repeated

Episodes of Demyelination and Remyelination in Multiple Sclerosis." *NeuroImage: Clinical* 6

(2014): 20–25. https://doi.org/10.1016/j.nicl.2014.06.009.

[8] Wang, Chenyu Tim, Michael Barnett, and Yael Barnett. "Imaging the Multiple Sclerosis

Lesion: Insights into Pathogenesis, Progression and Repair." *Current Opinion in Neurology* 32,

no. 3 (June 2019): 338–45. https://doi.org/10.1097/WCO.0000000000000698.

[9] Thompson, Alan J, Brenda L Banwell, Frederik Barkhof, William M Carroll, Timothy

Coetzee, Giancarlo Comi, Jorge Correale, et al. "Diagnosis of Multiple Sclerosis: 2017

Revisions of the McDonald Criteria." *The Lancet Neurology* 17, no. 2 (February 2018): 162–

73. https://doi.org/10.1016/S1474-4422(17)30470-2.

[10] Tabelow, Karsten, Evelyne Balteau, John Ashburner, Martina F. Callaghan, Bogdan

Draganski, Gunther Helms, Ferath Kherif, et al. "HMRI – A Toolbox for Quantitative MRI in

Neuroscience   and   Clinical   Research."   *NeuroImage*   194   (July   2019):   191–210.

https://doi.org/10.1016/j.neuroimage.2019.01.029.

[11] Schmierer, Klaus, Francesco Scaravilli, Daniel R. Altmann, Gareth J. Barker, and David H.

Miller. "Magnetization Transfer Ratio and Myelin in Postmortem Multiple Sclerosis Brain."

Annals   of   Neurology   56,   no.   3   (September   2004):   407–15.

https://doi.org/10.1002/ana.20202.

[12] Callaghan, Martina F., Gunther Helms, Antoine Lutti, Siawoosh Mohammadi, and

Nikolaus Weiskopf. "A General Linear Relaxometry Model of R $_1$ Using Imaging Data: General

Linear Relaxometry Model of R1." *Magnetic Resonance in Medicine* 73, no. 3 (March 2015):

1309–14. https://doi.org/10.1002/mrm.25210.

[13] Van Waesberghe, J. H. T. M., W. Kamphorst, C. J. A. De Groot, M. A. A. Van Walderveen,

J. A. Castelijns, R. Ravid, G. J. Lycklama a Nijeholt, et al. "Axonal Loss in Multiple Sclerosis

Lesions: Magnetic Resonance Imaging Insights into Substrates of Disability." *Annals   of*

Neurology   46,   no.   5   (November   1999):   747–54.   https://doi.org/10.1002/1531-

8249(199911)46:5<747::AID-ANA10>3.0.CO;2-4.

[14] Mottershead, J. P., K. Schmierer, M. Clemence, J. S. Thornton, F. Scaravilli, G. J. Barker,

P. S. Tofts, et al. "High Field MRI Correlates of Myelin Content and Axonal Density in Multiple

Sclerosis."   *Journal   of   Neurology*   250,   no.   11   (November   1,   2003):   1293–1301.

https://doi.org/10.1007/s00415-003-0192-3.

[15] Lema, Alfonso, Courtney Bishop, Omar Malik, Miriam Mattoscio, Rehiana Ali, Richard

Nicholas, Paolo A. Muraro, Paul M. Matthews, Adam D. Waldman, and Rexford D. Newbould.

"A Comparison of Magnetization Transfer Methods to Assess Brain and Cervical Cord

Microstructure in Multiple Sclerosis: MT of Brain and c-Spine in MS." *Journal of*

Neuroimaging 27, no. 2 (March 2017): 221–26. https://doi.org/10.1111/jon.12377.

[16] Helms, Gunther, Henning Dathe, and Peter Dechent. "Modeling the Influence of TR and

Excitation Flip Angle on the Magnetization Transfer Ratio (MTR) in Human Brain Obtained

from 3D Spoiled Gradient Echo MRI." *Magnetic Resonance in Medicine* 64, no. 1 (July 2010):

177–85. https://doi.org/10.1002/mrm.22379.

[17] Bagnato, Francesca, Simon Hametner, Emma Boyd, Verena Endmayr, Yaping Shi, Vasiliki

Ikonomidou, Guanhua Chen, et al. "Untangling the R2* Contrast in Multiple Sclerosis: A

Combined MRI-Histology Study at 7.0 Tesla." Edited by Quan Jiang. *PLOS ONE* 13, no. 3

(March 21, 2018): e0193839. https://doi.org/10.1371/journal.pone.0193839.

[18] Hametner, Simon, Verena Endmayr, Andreas Deistung, Pilar Palmrich, Max Prihoda,

Evelin Haimburger, Christian Menard, et al. "The Influence of Brain Iron and Myelin on

Magnetic Susceptibility and Effective Transverse Relaxation - A Biochemical and Histological

Validation Study." *NeuroImage* 179 (October 2018): 117–33.

https://doi.org/10.1016/j.neuroimage.2018.06.007.

[19] Stüber, Carsten, Markus Morawski, Andreas Schäfer, Christian Labadie, Miriam

Wähnert, Christoph Leuze, Markus Streicher, et al. "Myelin and Iron Concentration in the

Human Brain: A Quantitative Study of MRI Contrast." *NeuroImage* 93 (June 2014): 95–106.

https://doi.org/10.1016/j.neuroimage.2014.02.026.

[20] Stankiewicz, James M., Mohit Neema, et Antonia Ceccarelli. « Iron and Multiple

Sclerosis ». *Neurobiology of Aging* 35 (September 2014): S51-58.

https://doi.org/10.1016/j.neurobiolaging.2014.03.039.

[21] Granziera, Cristina, Jens Wuerfel, Frederik Barkhof, Massimiliano Calabrese, Nicola De

Stefano, Christian Enzinger, Nikos Evangelou, et al. "Quantitative Magnetic Resonance

Imaging towards Clinical Application in Multiple Sclerosis." *Brain* 144, no. 5 (June 22, 2021):

1296–1311. https://doi.org/10.1093/brain/awab029.

[22] Kolb, Hadar, Martina Absinta, Erin S. Beck, Seung-Kwon Ha, Yeajin Song, Gina Norato,

Irene Cortese, Pascal Sati, Govind Nair, and Daniel S. Reich. "7T MRI Differentiates

Remyelinated from Demyelinated Multiple Sclerosis Lesions." *Annals of Neurology* 90, no. 4

(October 2021): 612–26. https://doi.org/10.1002/ana.26194.

[23] Edwards, Luke J., Evgeniya Kirilina, Siawoosh Mohammadi, and Nikolaus Weiskopf.

"Microstructural Imaging of Human Neocortex in Vivo." *NeuroImage* 182 (November 2018):

184–206. https://doi.org/10.1016/j.neuroimage.2018.02.055.

[24] Lommers, Emilie, Jessica Simon, Gilles Reuter, Gaël Delrue, Dominique Dive, Christian

Degueldre, Evelyne Balteau, Christophe Phillips, and Pierre Maquet. "Multiparameter MRI

Quantification of Microstructural Tissue Alterations in Multiple Sclerosis." *NeuroImage:*

Clinical 23 (2019): 101879. https://doi.org/10.1016/j.nicl.2019.101879.

[25] Lommers, Emilie, Camille Guillemin, Gilles Reuter, Eve Fouarge, Gaël Delrue, Fabienne

Collette, Christian Degueldre, Evelyne Balteau, Pierre Maquet, and Christophe Phillips.

"VOXEL-BASED Quantitative MRI Reveals Spatial Patterns of Grey Matter Alteration in Multiple

Sclerosis." *Human Brain Mapping* 42, no. 4 (March 2021): 1003–12.

https://doi.org/10.1002/hbm.25274.

[26] Draganski, B., J. Ashburner, C. Hutton, F. Kherif, R.S.J. Frackowiak, G. Helms, and N.

Weiskopf. "Regional Specificity of MRI Contrast Parameter Changes in Normal Ageing

Revealed by Voxel-Based Quantification (VBQ)." *NeuroImage* 55, no. 4 (April 2011): 1423–34.

https://doi.org/10.1016/j.neuroimage.2011.01.052.

[27] Polman, Chris H., Stephen C. Reingold, Brenda Banwell, Michel Clanet, Jeffrey A. Cohen,

Massimo Filippi, Kazuo Fujihara, et al. "Diagnostic Criteria for Multiple Sclerosis: 2010

Revisions to the McDonald Criteria." *Annals of Neurology* 69, no. 2 (February 2011): 292–

302. https://doi.org/10.1002/ana.22366.

[28] Guillemin, C., E. Lommers, G. Delrue, E. Gester, P. Maquet, et F. Collette. « The Complex

Interplay Between Trait Fatigue and Cognition in Multiple Sclerosis ». *Psychologica Belgica*

62, n° 1 (March 2022): 108. https://doi.org/10.5334/pb.1125.

[29] Callaghan, Martina F., Patrick Freund, Bogdan Draganski, Elaine Anderson, Marinella

Cappelletti, Rumana Chowdhury, Joern Diedrichsen, et al. "Widespread Age-Related

Differences in the Human Brain Microstructure Revealed by Quantitative Magnetic

Resonance Imaging." *Neurobiology of Aging* 35, no. 8 (August 2014): 1862–72.

https://doi.org/10.1016/j.neurobiolaging.2014.02.008.

[30] Carey, Daniel, Francesco Caprini, Micah Allen, Antoine Lutti, Nikolaus Weiskopf, Geraint

Rees, Martina F. Callaghan, and Frederic Dick. "Quantitative MRI Provides Markers of Intra-,

Inter-Regional, and Age-Related Differences in Young Adult Cortical Microstructure."

NeuroImage 182 (November 2018): 429–40.

https://doi.org/10.1016/j.neuroimage.2017.11.066.

[31] Reuter, Gilles, Emilie Lommers, Evelyne Balteau, Jessica Simon, Christophe Phillips, Felix

Scholtes, Didier Martin, Arnaud Lombard, and Pierre Maquet. "Multiparameter Quantitative

Histological MRI Values in High-Grade Gliomas: A Potential Biomarker of Tumor

Progression." *Neuro-Oncology Practice* 7, no. 6 (December 4, 2020): 646–55.

https://doi.org/10.1093/nop/npaa047.

[32] Depierreux, Frédérique, Eric Parmentier, Laurane Mackels, Katherine Baquero, Christian

Degueldre, Evelyne Balteau, Eric Salmon, et al. "Parkinson's Disease Multimodal Imaging: F-

DOPA PET, Neuromelanin-Sensitive and Quantitative Iron-Sensitive MRI." *Npj Parkinson's*

Disease 7, no. 1 (December 2021): 57. https://doi.org/10.1038/s41531-021-00199-2.

[33] Nürnberger, Lucas, René-Maxime Gracien, Pavel Hok, Stephanie-Michelle Hof, Udo Rüb,

Helmuth Steinmetz, Rüdiger Hilker, Johannes C. Klein, Ralf Deichmann, and Simon Baudrexel.

"Longitudinal Changes of Cortical Microstructure in Parkinson's Disease Assessed with T1

Relaxometry." *NeuroImage: Clinical* 13 (2017): 405–14.

https://doi.org/10.1016/j.nicl.2016.12.025.

[34] Klein, J.C., M. Rolinski, L. Griffanti, K. Szewczyk-Krolikowski, F. Baig, C. Ruffmann, A.R.

Groves, R.A.L. Menke, M.T. Hu, and C. Mackay. "Cortical Structural Involvement and

Cognitive Dysfunction in Early Parkinson's Disease." *NMR in Biomedicine* 31, no. 4 (April

2018): e3900. https://doi.org/10.1002/nbm.3900.

[35] Lutti, Antoine, Chloe Hutton, Jürgen Finsterbusch, Gunther Helms, and Nikolaus

Weiskopf. "Optimization and Validation of Methods for Mapping of the Radiofrequency

Transmit Field at 3T: Optimized RF Transmit Field Mapping at 3T." *Magnetic Resonance in*

Medicine 64, no. 1 (July 2010): 229–38. https://doi.org/10.1002/mrm.22421.

[36] Lutti, Antoine, Joerg Stadler, Oliver Josephs, Christian Windischberger, Oliver Speck,

Johannes Bernarding, Chloe Hutton, and Nikolaus Weiskopf. "Robust and Fast Whole Brain

Mapping of the RF Transmit Field B1 at 7T." Edited by Wang Zhan. *PLoS ONE* 7, no. 3 (March

12, 2012): e32379. https://doi.org/10.1371/journal.pone.0032379.

[37] Schmidt, Paul, Christian Gaser, Milan Arsic, Dorothea Buck, Annette Förschler, Achim

Berthele, Muna Hoshi, et al. "An Automated Tool for Detection of FLAIR-Hyperintense

White-Matter Lesions in Multiple Sclerosis." *NeuroImage* 59, no. 4 (February 2012): 3774–

83. https://doi.org/10.1016/j.neuroimage.2011.11.032.

[38] Weiskopf, Nikolaus, Martina F. Callaghan, Oliver Josephs, Antoine Lutti, and Siawoosh

Mohammadi. "Estimating the Apparent Transverse Relaxation Time (R2*) from Images with

Different Contrasts (ESTATICS) Reduces Motion Artifacts." *Frontiers in Neuroscience* 8

(September 10, 2014). https://doi.org/10.3389/fnins.2014.00278.

[39] Preibisch, C., and R. Deichmann. "Influence of RF Spoiling on the Stability and Accuracy

of T_1 Mapping Based on Spoiled FLASH with Varying Flip Angles: Influence of RF Spoiling on T

$_1$ Mapping." *Magnetic Resonance in Medicine* 61, no. 1 (January 2009): 125–35.

https://doi.org/10.1002/mrm.21776.

[40] Ashburner, John. "Symmetric Diffeomorphic Modeling of Longitudinal Structural MRI."

Frontiers in Neuroscience 6 (2013). https://doi.org/10.3389/fnins.2012.00197.

[41] Ashburner, John, and Karl J. Friston. "Unified Segmentation." *NeuroImage* 26, no. 3 (July

2005): 839–51. https://doi.org/10.1016/j.neuroimage.2005.02.018.

[42] Lorio, S., S. Fresard, S. Adaszewski, F. Kherif, R. Chowdhury, R.S. Frackowiak, J.

Ashburner, et al. "New Tissue Priors for Improved Automated Classification of Subcortical

Brain Structures on MRI." *NeuroImage* 130 (April 2016): 157–66.

https://doi.org/10.1016/j.neuroimage.2016.01.062.

[43] Andersen, Sarah M., Steven Z. Rapcsak, and Pélagie M. Beeson. "Cost Function Masking

during Normalization of Brains with Focal Lesions: Still a Necessity?" *NeuroImage* 53, no. 1

(October 2010): 78–84. https://doi.org/10.1016/j.neuroimage.2010.06.003.

[44] Moon, Nathan, Elizabeth Bullitt, Koen van Leemput, and Guido Gerig. "Automatic Brain

and Tumor Segmentation." In *Medical Image Computing and Computer-Assisted Intervention*

— *MICCAI 2002*, edited by Takeyoshi Dohi and Ron Kikinis, 2488:372–79. Lecture Notes in

Computer Science. Berlin, Heidelberg: Springer Berlin Heidelberg, 2002.

https://doi.org/10.1007/3-540-45786-0_46.

[45] Pandit, Lekha. "No Evidence of Disease Activity (NEDA) in Multiple Sclerosis - Shifting

the Goal Posts." *Annals of Indian Academy of Neurology* 22, no. 3 (2019): 261.

https://doi.org/10.4103/aian.AIAN_159_19.

[46] Anderson, Marti J. "Permutation Tests for Univariate or Multivariate Analysis of

Variance and Regression." *Canadian Journal of Fisheries and Aquatic Sciences* 58, no. 3

(March 1, 2001): 626–39. https://doi.org/10.1139/f01-004.

[47] Yoav Benjamini and Yosef Hochberg. "Controlling the False Discovery Rate: a Practical

and Powerful Approach to Multiple Testing" *Journal of the Royal Statistical Society. Series B*

(Methodological), Vol. 57, No.1 (1995), 289-300.

[48] Leutritz, Tobias, Maryam Seif, Gunther Helms, Rebecca S Samson, Armin Curt, Patrick

Freund, and Nikolaus Weiskopf. "Multiparameter Mapping of Relaxation (R1 , R2 *), Proton

Density and Magnetization Transfer Saturation at 3 T▯: A Multicenter Dual-vendor

Reproducibility and Repeatability Study." *Human Brain Mapping* 41, no. 15 (October 15,

2020): 4232–47. https://doi.org/10.1002/hbm.25122.

[49] Andica, Christina, Akifumi Hagiwara, Koji Kamagata, Kazumasa Yokoyama, Keigo Shimoji,

Asami Saito, Yuki Takenaka, et al. "Gray Matter Alterations in Early and Late Relapsing-

Remitting Multiple Sclerosis Evaluated with Synthetic Quantitative Magnetic Resonance

Imaging." *Scientific Reports* 9, no. 1 (December 2019): 8147.

https://doi.org/10.1038/s41598-019-44615-3.

[50] Bonnier, Guillaume, Alexis Roche, David Romascano, Samanta Simioni, Djalel Meskaldji,

David Rotzinger, Ying-Chia Lin, et al. "Advanced MRI Unravels the Nature of Tissue

Alterations in Early Multiple Sclerosis." *Annals of Clinical and Translational Neurology* 1, no.

6 (June 2014): 423–32. https://doi.org/10.1002/acn3.68.

[51] Engström, Maria, Jan B. M. Warntjes, Anders Tisell, Anne-Marie Landtblom, and Peter

Lundberg. "Multi-Parametric Representation of Voxel-Based Quantitative Magnetic

Resonance Imaging." Edited by Friedemann Paul. *PLoS ONE* 9, no. 11 (November 13, 2014):

e111688. https://doi.org/10.1371/journal.pone.0111688.

[52] Gracien, René-Maxime, Alina Jurcoane, Marlies Wagner, Sarah C. Reitz, Christoph

Mayer, Steffen Volz, Stephanie-Michelle Hof, et al. "Multimodal Quantitative MRI

Assessment of Cortical Damage in Relapsing-Remitting Multiple Sclerosis: Cortical

Quantitative MRI in RRMS." *Journal of Magnetic Resonance Imaging* 44, no. 6 (December

2016): 1600–1607. https://doi.org/10.1002/jmri.25297.

[53] Neema, Mohit, James Stankiewicz, Ashish Arora, Venkata S.R. Dandamudi, Courtney E.

Batt, Zachary D. Guss, Ali Al-Sabbagh, and Rohit Bakshi. "T1- and T2-Based MRI Measures of

Diffuse Gray Matter and White Matter Damage in Patients with Multiple Sclerosis." *Journal*

of Neuroimaging 17 (April 2007): 16S-21S. https://doi.org/10.1111/j.1552-

6569.2007.00131.x.

[54] Reitz, Sarah C., Stephanie-Michelle Hof, Vinzenz Fleischer, Alla Brodski, Adriane Gröger,

René-Maxime Gracien, Amgad Droby, et al. "Multi-Parametric Quantitative MRI of Normal

Appearing White Matter in Multiple Sclerosis, and the Effect of Disease Activity on T2." *Brain*

Imaging and Behavior 11, no. 3 (June 2017): 744–53. https://doi.org/10.1007/s11682-016-

9550-5.

[55] Stevenson, V.L, G.J.M Parker, G.J Barker, K Birnie, P.S Tofts, D.H Miller, and A.J

Thompson. "Variations in T1 and T2 Relaxation Times of Normal Appearing White Matter

and Lesions in Multiple Sclerosis." *Journal of the Neurological Sciences* 178, no. 2 (September

2000): 81–87. https://doi.org/10.1016/S0022-510X(00)00339-7.

[56] Bonnier, Guillaume, Benedicte Maréchal, Mário João Fartaria, Pavel Falkowskiy, José P.

Marques, Samanta Simioni, Myriam Schluep, et al. "The Combined Quantification and

Interpretation of Multiple Quantitative Magnetic Resonance Imaging Metrics Enlightens

Longitudinal Changes Compatible with Brain Repair in Relapsing-Remitting Multiple Sclerosis

Patients." *Frontiers in Neurology* 8 (September 27, 2017): 506.

https://doi.org/10.3389/fneur.2017.00506.

[57] Elkady, Ahmed M., Dana Cobzas, Hongfu Sun, Gregg Blevins, and Alan H. Wilman.

"Discriminative Analysis of Regional Evolution of Iron and Myelin/Calcium in Deep Gray

Matter of Multiple Sclerosis and Healthy Subjects: Analysis of Iron and Myelin in MS."

Journal of Magnetic Resonance Imaging 48, no. 3 (September 2018): 652–68.

https://doi.org/10.1002/jmri.26004.

[58] Elkady, Ahmed M., Dana Cobzas, Hongfu Sun, Peter Seres, Gregg Blevins, and Alan H.

Wilman. "Five Year Iron Changes in Relapsing-Remitting Multiple Sclerosis Deep Gray Matter

Compared to Healthy Controls." *Multiple Sclerosis and Related Disorders* 33 (August 2019):

107–15. https://doi.org/10.1016/j.msard.2019.05.028.

[59] Khalil, M., C. Langkammer, A. Pichler, D. Pinter, T. Gattringer, G. Bachmaier, S. Ropele, S.

Fuchs, C. Enzinger, and F. Fazekas. "Dynamics of Brain Iron Levels in Multiple Sclerosis: A

Longitudinal 3T MRI Study." *Neurology* 84, no. 24 (June 16, 2015): 2396–2402.

https://doi.org/10.1212/WNL.0000000000001679.

[60] Gracien, René-Maxime, Sarah C. Reitz, Stephanie-Michelle Hof, Vinzenz Fleischer,

Amgad Droby, Mathias Wahl, Helmuth Steinmetz, Sergiu Groppa, Ralf Deichmann, and

Johannes C. Klein. "Longitudinal Quantitative MRI Assessment of Cortical Damage in Multiple

Sclerosis: A Pilot Study: Longitudinal Cortical QMRI in MS." *Journal of Magnetic Resonance*

Imaging 46, no. 5 (November 2017): 1485–90. https://doi.org/10.1002/jmri.25685.

[61] Laule, Cornelia, I. M. Vavasour, K. P. Whittall, J. Oger, D. W. Paty, D. K. B. Li, A. L.

MacKay, and D. L. Arnold. "Evolution of Focal and Diffuse magnetisation Transfer

Abnormalities in Multiple Sclerosis." *Journal of Neurology* 250, no. 8 (August 2003): 924–31.

https://doi.org/10.1007/s00415-003-1115-z.

[62] Hayton, T., J. Furby, K. J. Smith, D. R. Altmann, R. Brenner, J. Chataway, K. Hunter, D. J.

Tozer, D. H. Miller, and R. Kapoor. "Longitudinal Changes in Magnetisation Transfer Ratio in

Secondary Progressive Multiple Sclerosis: Data from a Randomised Placebo Controlled Trial

of Lamotrigine." *Journal of Neurology* 259, no. 3 (March 2012): 505–14.

https://doi.org/10.1007/s00415-011-6212-9.

[63] Rocca, Maria A, Giovanna Mastronardo, Mariemma Rodegher, Giancarlo Comi, and

Massimo Filippi. "Long-Term Changes of Magnetization Transfer-Derived Measures from

Patients with Relapsing-Remitting and Secondary Progressive Multiple Sclerosis," *AJNR Am J*

Neuroradiol 20 (1999):821–827

[64] Weiskopf, Nikolaus, Siawoosh Mohammadi, Antoine Lutti, and Martina F. Callaghan.

"Advances in MRI-Based Computational Neuroanatomy: From Morphometry to in-Vivo

Histology." *Current Opinion in Neurology* 28, no. 4 (August 2015): 313–22.

https://doi.org/10.1097/WCO.0000000000000222.

[65] Hametner, Simon, Verena Endmayr, Andreas Deistung, Pilar Palmrich, Max Prihoda,

Evelin Haimburger, Christian Menard, et al. "The Influence of Brain Iron and Myelin on

Magnetic Susceptibility and Effective Transverse Relaxation - A Biochemical and Histological

Validation Study." *NeuroImage* 179 (October 2018): 117–33.

https://doi.org/10.1016/j.neuroimage.2018.06.007.

[66] Mangeat, G., S.T. Govindarajan, C. Mainero, and J. Cohen-Adad. "Multivariate

Combination of Magnetization Transfer, T 2 * and B0 Orientation to Study the Myelo-

Architecture of the in Vivo Human Cortex." *NeuroImage* 119 (October 2015): 89–102.

https://doi.org/10.1016/j.neuroimage.2015.06.033.

[67] Dutta, R., and B. D. Trapp. "Pathogenesis of Axonal and Neuronal Damage in Multiple

Sclerosis." *Neurology* 68, no. Issue 22, Supplement 3 (May 29, 2007): S22–31.

https://doi.org/10.1212/01.wnl.0000275229.13012.32.

[68] Frischer, Josa M., Stephen D. Weigand, Yong Guo, Nilufer Kale, Joseph E. Parisi, Istvan

Pirko, Jay Mandrekar, et al. "Clinical and Pathological Insights into the Dynamic Nature of the

White Matter Multiple Sclerosis Plaque: Dynamic Nature of MS Plaque." *Annals of Neurology*

78, no. 5 (November 2015): 710–21. https://doi.org/10.1002/ana.24497.

[69] Lassmann, Hans, Wolfgang Brück, and Claudia Lucchinetti. "Heterogeneity of Multiple

Sclerosis Pathogenesis: Implications for Diagnosis and Therapy." *Trends in Molecular*

Medicine 7, no. 3 (March 2001): 115–21. https://doi.org/10.1016/S1471-4914(00)01909-2.

[70] Lucchinetti, Claudia, Wolfgang Brück, Joseph Parisi, Bernd Scheithauer, Moses

Rodriguez, and Hans Lassmann. "Heterogeneity of Multiple Sclerosis Lesions: Implications for

the Pathogenesis of Demyelination." *Annals of Neurology* 47, no. 6 (June 2000): 707–17.

https://doi.org/10.1002/1531-8249(200006)47:6<707::AID-ANA3>3.0.CO;2-Q.

[71] Dousset, Vincent, Annick Gayou, Bruno Brochet, and Jean-Marie Caille. "Early Structural

Changes in Acute MS Lesions Assessed by Serial Magnetization Transfer Studies." *Neurology*

51, no. 4 (October 1998): 1150–55. https://doi.org/10.1212/WNL.51.4.1150.

[72] Levesque, Ives R., Paul S. Giacomini, Sridar Narayanan, Luciana T. Ribeiro, John G. Sled,

Doug L. Arnold, and G. Bruce Pike. "Quantitative Magnetization Transfer and Myelin Water

Imaging of the Evolution of Acute Multiple Sclerosis Lesions: Quantitative MT and T_2 in

Acute MS Lesions." *Magnetic Resonance in Medicine* 63, no. 3 (March 2010): 633–40.

https://doi.org/10.1002/mrm.22244.

[73] Elskamp, Ij van den, Dl Knol, H. Vrenken, G. Karas, A. Meijerman, M. Filippi, L. Kappos, et

al. "Lesional Magnetization Transfer Ratio: A Feasible Outcome for Remyelinating Treatment

Trials in Multiple Sclerosis." *Multiple Sclerosis Journal* 16, no. 6 (June 2010): 660–69.

https://doi.org/10.1177/1352458510364630.

[74] Filippi, Massimo, Maria A. Rocca, Gianvito Martino, Mark A. Horsfield, and Giancarlo

Comi. "Magnetization Transfer Changes in the Normal Appering White Matter Precede the

Appearance of Enhancing Lesions in Patients with Multiple Sclerosis." *Annals of Neurology*

43, no. 6 (June 1998): 809–14. https://doi.org/10.1002/ana.410430616.

[75] Fazekas, F, S Ropele, C Enzinger, T Seifert, and S Strasser-Fuchs. "Quantitative

Magnetization Transfer Imaging of Pre-Lesional White-Matter Changes in Multiple Sclerosis."

Multiple Sclerosis Journal 8, no. 6 (December 2002): 479–84.

https://doi.org/10.1191/1352458502ms860oa.

[76] Zheng, Yufan, Jar-Chi Lee, Richard Rudick, and Elizabeth Fisher. "Long-Term

Magnetization Transfer Ratio Evolution in Multiple Sclerosis White Matter Lesions: Long-

Term MTR Evolution in MS WM Lesions." *Journal of Neuroimaging* 28, no. 2 (March 2018):

191–98. https://doi.org/10.1111/jon.12480.

[77] Chawla, Sanjeev, Ilya Kister, Tim Sinnecker, Jens Wuerfel, Jean-Christophe Brisset,

Friedemann Paul, and Yulin Ge. "Longitudinal Study of Multiple Sclerosis Lesions Using Ultra-

High Field (7T) Multiparametric MR Imaging." Edited by Quan Jiang. *PLOS ONE* 13, no. 9

(September 13, 2018): e0202918. https://doi.org/10.1371/journal.pone.0202918.

[78] Lieury, Alice, Marie Chanal, Géraldine Androdias, Richard Reynolds, Sylvie Cavagna,

Pascale Giraudon, Christian Confavreux, and Serge Nataf. "Tissue Remodeling in Periplaque

Regions of Multiple Sclerosis Spinal Cord Lesions: Glia Remodeling in MS Spinal Cord." *Glia*

62, no. 10 (October 2014): 1645–58. https://doi.org/10.1002/glia.22705.

[79] Kuhlmann, Tanja, Samuel Ludwin, Alexandre Prat, Jack Antel, Wolfgang Brück, and Hans

Lassmann. "An Updated Histological Classification System for Multiple Sclerosis Lesions."

Acta Neuropathologica 133, no. 1 (January 2017): 13–24. https://doi.org/10.1007/s00401-

016-1653-y.

[80] Lassmann, Hans, Wolfgang Brück, and Claudia F. Lucchinetti. "The Immunopathology of Multiple Sclerosis: An Overview." *Brain Pathology* 17, no. 2 (April 2007): 210–18. https://doi.org/10.1111/j.1750-3639.2007.00064.x.

[81] Dal-Bianco, Assunta, Günther Grabner, Claudia Kronnerwetter, Michael Weber, Romana Höftberger, Thomas Berger, Eduard Auff, et al. "Slow Expansion of Multiple Sclerosis Iron Rim Lesions: Pathology and 7 T Magnetic Resonance Imaging." *Acta Neuropathologica* 133, no. 1 (January 2017): 25–42. https://doi.org/10.1007/s00401-016-1636-z.

[82] Elliott, Colm, Jerry S Wolinsky, Stephen L Hauser, Ludwig Kappos, Frederik Barkhof, Corrado Bernasconi, Wei Wei, Shibeshih Belachew, and Douglas L Arnold. "Slowly Expanding/Evolving Lesions as a Magnetic Resonance Imaging Marker of Chronic Active Multiple Sclerosis Lesions." *Multiple Sclerosis Journal* 25, no. 14 (December 2019): 1915–25. https://doi.org/10.1177/1352458518814117.

[83] De Stefano, N., A. Giorgio, M. Battaglini, M. Rovaris, M. P. Sormani, F. Barkhof, T. Korteweg, et al. "Assessing Brain Atrophy Rates in a Large Population of Untreated Multiple Sclerosis Subtypes." *Neurology* 74, no. 23 (June 8, 2010): 1868–76. https://doi.org/10.1212/WNL.0b013e3181e24136.

[84] Eshaghi, Arman, Ferran Prados, Wallace J. Brownlee, Daniel R. Altmann, Carmen Tur, M. Jorge Cardoso, Floriana De Angelis, et al. "Deep Gray Matter Volume Loss Drives Disability

914 Worsening in Multiple Sclerosis: Deep Gray Matter Volume Loss." *Annals of Neurology* 83,

915 no. 2 (February 2018): 210–22. https://doi.org/10.1002/ana.25145.

917 [85] Bermel, Robert A, and Rohit Bakshi. "The Measurement and Clinical Relevance of Brain

918 Atrophy in Multiple Sclerosis." *The Lancet Neurology* 5, no. 2 (February 2006): 158–70.

919 https://doi.org/10.1016/S1474-4422(06)70349-0.

921 [86] Zivadinov, R., A. T. Reder, M. Filippi, A. Minagar, O. Stuve, H. Lassmann, M. K. Racke, M.

922 G. Dwyer, E. M. Frohman, and O. Khan. "Mechanisms of Action of Disease-Modifying Agents

923 and Brain Volume Changes in Multiple Sclerosis." *Neurology* 71, no. 2 (July 8, 2008): 136–44.

924 https://doi.org/10.1212/01.wnl.0000316810.01120.05.

926 [87] De Stefano, Nicola, Maria Laura Stromillo, Antonio Giorgio, Maria Letizia Bartolozzi,

927 Marco Battaglini, Mariella Baldini, Emilio Portaccio, Maria Pia Amato, and Maria Pia Sormani.

928 "Establishing Pathological Cut-Offs of Brain Atrophy Rates in Multiple Sclerosis." *Journal of*

929 *Neurology, Neurosurgery & Psychiatry*, April 22, 2015, jnnp-2014-309903.

930 https://doi.org/10.1136/jnnp-2014-309903.

932 [88] on behalf of the MAGNIMS study group, Christian Enzinger, Frederik Barkhof, Olga

933 Ciccarelli, Massimo Filippi, Ludwig Kappos, Maria A. Rocca, et al. "Nonconventional MRI and

934 Microstructural Cerebral Changes in Multiple Sclerosis." *Nature Reviews Neurology* 11, no.

935 12 (December 2015): 676–86. https://doi.org/10.1038/nrneurol.2015.194.

[89] Bodini, Benedetta, Mattia Veronese, Daniel García-Lorenzo, Marco Battaglini, Emilie

Poirion, Audrey Chardain, Léorah Freeman, et al. "Dynamic Imaging of Individual

Remyelination Profiles in Multiple Sclerosis." *Annals of Neurology* 79, no. 5 (May 2016): 726–

38. https://doi.org/10.1002/ana.24620.

[90] Patrikios, P., C. Stadelmann, A. Kutzelnigg, H. Rauschka, M. Schmidbauer, H. Laursen, P.

S. Sorensen, W. Bruck, C. Lucchinetti, and H. Lassmann. "Remyelination Is Extensive in a

Subset of Multiple Sclerosis Patients." *Brain* 129, no. 12 (June 9, 2006): 3165–72.

https://doi.org/10.1093/brain/awl217.

[91] Gracien, RM., Nürnberger, L., Hok, P. *et al.* Evaluation of brain ageing: a quantitative

longitudinal MRI study over 7 years. *Eur Radiol* **27,** 1568–1576 (2017).

https://doi.org/10.1007/s00330-016-4485-1

Tables

	All Patients (n = 17)	
Age, y, median (range)	36 (25-65)	
Sex, F/M	7/10	
MS phenotype (RRMS/MS)	11/6	
Baseline disease duration, y, median (range)	3.4 (0.3-28)	
Baseline EDSS, median (range)	2.5 (1-6.5)	
Baseline number of relapses, median (range)	RRMS: 2 (1-5)	PMS: N/A
Disease-modifying treatment	RRMS: First line, n: 5 Second line, n: 6	PMS: Ocrelizumab, n: 2 None, n: 4

Table 1: Demographic data of the study sample

	NAWM	NACGM	NADGM
MT	**0.039 (.011)***	**0.017 (.007)***	0.004 (.749)
PD	-0.018 (.670)	0.405 (.225)	0.250 (.552)
R1	0.009 (.139)	0.004 (.471)	0.010 (.111)
R2*	**0.295 (.002)***	0.121 (.092)	0.066 (.770)
BPF	-0.884 (.1562)		
LF	21.23 (.1082)		

Table 2: Regression coefficients and their associated *p*-values (in parentheses) for the effects of X_{status} on the individual ARoC for each qMRI parameter (MTsat, PD, R1 and R2*) and for volumetric measurements (BPF and LF).

* Results significant at $p < .05$, FDR corrected.

Legends of figures

Figure 1: Chartflow of data creation and processing (see text). MPM maps were created with

the hMRI-toolbox, FLAIR images were directly acquired for both sessions (T0 and T1). A

preliminary mask was constructed based on T0 FLAIR. All images (MPM and FLAIR, T0 and

T1) were co-registered to the MPM T0 space. Segmentation using USwL allowed to isolate

968   the different tissue classes.

970   **Figure 2:** Schematic illustration of the NAWM and 3 lesions-related areas: focal FLAIR lesion

971   (dark gray area), initial peripheral lesion detected at T0 (medium gray area), later peripheral

972   lesion detected at T1 (dashed, left, and light gray, right, area).

974   **Figure 3:** Line plots illustrating individual ARoC's for MTsat (left) and R2* (right) in NAWM.

975   Each line corresponds to one subject. Dotted lines represent increasing rates.

977   **Figure 4:** Violin plots of significant change rates in microstructure with respect to X_{status}.

978   From left to right: MT in NAWM, MT in NACGM, R2* in NAWM. * $P < .05$.

980   **Figure 5**: Microstructural parameters in NAWM and the 3 lesion-related areas, for each scanning time

981   T0 and T1. P-values were obtained with *post-hoc* tests on the tissue area effect. * $P < .05$.

984   **Appendix A: Supplementary data legends**

985   **Supplementary data 1:** Multi-echo 3D FLASH acquisition parameters for Siemens Magnetom
986   PRISMA MRI

988   **Supplementary data 2:** Extended demographic data. Age, disease duration, EDSS and
989   relapses values were taken at baseline.

991   **Supplementary data 3:** Longitudinal clinical information and derived disease status score

993   **Supplementary data 4**: Line plots illustrating individual ARoC's for PD (left) and R1 (right) in
994   NAWM. Each line corresponds to one subject. Dotted lines represent increasing rates.

996   **Supplementary data 5:** Differences of lesion class Least Squares Means. First two columns
997   correspond to tissue class labels (0 = FLAIR lesion, 1 = Later peripheral lesion, 2 = Initial
998   peripheral lesion, 3 = NAWM).

1000

1001

Figure 1: Chartflow of data creation and processing (see text). MPM maps were created with the hMRI-toolbox, FLAIR images were directly acquired for both sessions (T0 and T1). A preliminary mask was constructed based on T0 FLAIR. All images (MPM and FLAIR, T0 and T1) were co-registered to the MPM T0 space. Segmentation using USwL allowed to isolate the different tissue classes.

Figure 2: Schematic illustration of the NAWM and 3 lesions-related areas: focal

FLAIR lesion (dark gray area), initial peripheral lesion detected at T0 (medium

gray area), later peripheral lesion detected at T1 (dashed, left, and light gray,

right, area)

Figure 3: Line plots illustrating individual ARoC's for MTsat (left) and R2* (right) in NAWM.

Each line corresponds to one subject. Dotted lines represent increasing rates.

Figure 4: Violin plots of significant change rates in microstructure with respect to X_{status}.

From left to right: MT in NAWM, MT in NACGM, R2* in NAWM. $* P < .05$.

Figure 5: Microstructural parameters in NAWM and the 3 lesion-related areas, for each scanning time T0 and T1. P-values were obtained with *post-hoc* tests on the tissue area effect. * $P < .05$.