1   **Title:** Comparison of total and neutralizing SARS-CoV-2 spike antibodies against omicron and
2   other variants in paired samples after two or three doses of mRNA vaccine
3
4

Amanda K. Debes, PhD, MS[1]

Shaoming Xiao, MSPH[2]

Emily R. Egbert MPH, MAT[2]

Patrizio Caturegli MD, MPH[2]

Ioannis Sitaras, PhD, MRes[1]

Andrew Pekosz, PhD[1,2]

Aaron M. Milstone, MD, MHS[1,2]

[1] Johns Hopkins University Bloomberg School of Public Health , Baltimore, MD, USA

[2] Johns Hopkins University School of Medicine , Baltimore, MD, USA

Correspondence:
15
16   Amanda K. Debes PhD, MS
17   Department of International Health
18   615 N. Wolfe Street, E5036
19   Baltimore, MD 21205
20   adebes1@jhu.edu
21

22   **Manuscript Word Count:** 594

Abstract

Recognizing that anti-SARS-CoV-2 antibody levels wane over time following the 2-dose

SARS-CoV-2 mRNA series, the FDA approved a booster dose for people greater than 12 years

old. Limited data exist on whether a booster dose of the mRNA vaccine results in greater

antibody protection than the primary series. We examined total and neutralizing antibodies to

the spike protein of SARS-CoV-2, and neutralizing antibodies against Washington-1 (WA-1) and

variants of concern (VOC) including Beta, Delta and Omicron in a longitudinal cohort.

Healthcare workers (HWs) were included in the analysis if serum was collected 1) within 14-44

days post-dose2 of an mRNA SARS-CoV-2 vaccine (Timepoint 1, TP1), or 2) at least 8 months

post-dose2 (Timepoint 2, TP2), or 3) within 14-44 days following mRNA booster (Timepoint 3,

TP3). HWs with prior covid-positive PCR were excluded. We found that there is little to no

neutralizing capability following a 2-dose mRNA vaccine series against the omicron variant, and

neutralizing capacity to any variant strain tested has been lost by 8-months post two-dose

vaccination series. However, the mRNA booster series eliminates the immune escape observed

by the omicron variant with the two-dose series. Neutralizing titers were significantly higher for

all variants post-boost compared to the titers post two-dose series. The longitudinal nature of

our cohort facilitated the analysis of paired samples pre and post boost, showing a greater than

15-fold increase in neutralization against omicron post-boost in these paired samples. An mRNA

booster dose provides greater quantity and quality of antibodies compared to a two-dose

regimen and is critical to provide any protection against the omicron variant.

Introduction

SARS-CoV-2 antibody levels wane following two-dose mRNA vaccination and infection.[1] mRNA booster doses are available and protect against hospitalization and death, but booster uptake remains low.[2] Our objective was to compare total and neutralizing SARS-CoV-2 spike antibodies against against Washington-1 (WA-1) and variants of concern (VOC) in a longitudinal cohort.

Methods

Healthcare workers (HWs) were consented into a seroprevalence cohort beginning June 2020 and followed through November 2021.[1] HWs provided serum samples longitudinally and were included in this analysis if serum was collected 1) within 14-44 days post-dose2 of an mRNA SARS-CoV-2 vaccine (Timepoint 1, TP1), or 2) at least 8 months post-dose2 (Timepoint 2, TP2), or 3) within 14-44 days following mRNA booster (Timepoint 3, TP3). HWs with prior covid-positive PCR were excluded. To determine if the increase in magnitude of antibody response to the mRNA vaccine strain, as measured by enzyme-linked immunosorbent assay (ELISA) [Euroimmun], led to an increased recognition of VOC, neutralizing antibody titer (NT) assays were performed against the vaccine strain (WA-1) and the Beta, Delta and Omicron variants as previously described.[1,3] For NT assays, 45 HWs were selected, prioritizing paired samples from TP1 and TP3, and others were selected at random.

Wilcoxon rank sum test was used for unpaired analyses, and Wilcoxon signed rank test and Friedman test for paired analyseis. The Johns Hopkins University Institutional Review Board approved this study. Analyses were performed in R, version 4.1.2.

Results

Of 3032 HWs originally enrolled in the longitudinal cohort, 1353 contributed serum to at least one of the 3 groups: 507 within in TP1, 879 in TP2, and 273 in TP3. Of these 1353 participants, 81% were women, 96% were Non-Hispanic/Latino, and 81% were White. The median (IQR) age of participants was 41.8 (33.8 - 53.3) years.

High levels of antibodies in TP1 waned to lower levels in TP2 and then boosted to much

higher levels in TP3. Of the TP3 samples tested, 94% demonstrated spike IgG assay saturation

compared to 59% in TP1 (Figure 1). Spike IgG measurements correlated with NT against WA-1.

TP1 samples had lower NT activity across VOC compared to WA-1 (Figure 2). By TP2, there

was little NT to Beta and Delta, and none to omicron (titer<20).[4] At TP3, NT activity against all

viruses was boosted and the fold reductions between WA-1 and VOC were less than those

observed in TP1 among paired samples.

100
Discussion

After two-dose vaccination, HWs developed high spike IgG and NT to vaccine strains but

significantly lower NT against VOC was evident. By 8-month post-dose-2, all antibody

measures had waned and NT against VOC had dropped to nearly undetectable levels. Boosting

proved critical, as NT were significantly higher for all VOC compared to post-dose2 titers,

including a >15 fold increase in neutralization against omicron in paired samples.

Limitations include that paired serum was only available post-dose2 and post-boost, and

nABs titers were only performed on a small subset. The corresponding antibody trends between

serum IgG and NT supports that despite only testing a subset of the participants for virus

neutralization, all participants likely had robust broad NT post-boost, findings supported by

recently published pre and post-boost studies.[5]

This study demonstrates that in paired samples an mRNA vaccine booster produces

greater quantity and function of spike antibodies and NT as compared to primary SARS-CoV-2

mRNA immunization and was necessary to restore measurable NT to VOC. The booster dose

eliminated the immune escape observed by Omicron following two-dose mRNA immunization.

The variable NT to mRNA booster and whether breakthrough omicron infection boosts NT must

be investigated to understand durable immunity against future VOC.

Acknowledgements: The authors would like to thank members of the Johns Hopkins Hospital

Clinical Immunology Laboratory, Danielle Koontz and Ani Voskertchian of the Johns Hopkins

Division of Pediatric Infectious Diseases, Elizabeth Colantuoni from the Johns Hopkins

Bloomberg School of Public Health, and Avinash Gadala from the Johns Hopkins Health

System. Research reported in this publication was supported in part by the National Institute of

Allergy and Infectious Diseases of the National Institutes of Health (NIH) under award number

K24AI141580 (A.M.), the Johns Hopkins Center of Excellence in Influenza Research and

Surveillance (NIH/NIAID N2772201400007C AP) and the generosity of the collective community

of donors to the Johns Hopkins University School of Medicine and the Johns Hopkins Health

System for Covid-19 research.

Conflict of Interest Disclosure: A.M. reports grant support from Merck for work unrelated to

this study. Other authors report no conflicts.

References

1.   Zhong D, Xiao S, Debes AK, et al. Durability of Antibody Levels After Vaccination With mRNA
SARS-CoV-2 Vaccine in Individuals With or Without Prior Infection. *Jama.* 2021.

2.   Grannis SJ, Rowley EA, Ong TC, et al. Interim Estimates of COVID-19 Vaccine Effectiveness
Against COVID-19-Associated Emergency Department or Urgent Care Clinic Encounters and
Hospitalizations Among Adults During SARS-CoV-2 B.1.617.2 (Delta) Variant Predominance -
Nine States, June-August 2021. *MMWR Morbidity and mortality weekly report.* 2021;70(37):1291-
1293.
3.   Karaba AH, Zhu X, Liang T, et al. A third dose of SARS-CoV-2 vaccine increases neutralizing
antibodies against variants of concern in solid organ transplant recipients. *Am J Transplant.* 2021.
4.   Klein SL, Pekosz A, Park HS, et al. Sex, age, and hospitalization drive antibody responses in a
COVID-19 convalescent plasma donor population. *The Journal of clinical investigation.*
2020;130(11):6141-6150.
5.   Wu M, Wall EC, Carr EJ, et al. Three-dose vaccination elicits neutralising antibodies against
omicron. *The Lancet.* 2022.

Figure 1: Spike IgG serum antibodies and live-virus neutralizing antibody titers (NT) against the

vaccine strain (WA-1). Data shown at threetime points: within 14-44 days post-dose2 (Timepoint

1), at least 8 months post-dose2 (Timepoint 2), and within 14-44 post-booster (Timepoint 3). IgG

antibody measurements estimated optical density ratios with a lower threshold of 1.23 and

upper threshold of 11.00 based on assay saturation. NTs were reported using NT50, and a

positive threshold was defined as NT≥20.[4] Timepoint 1: 396(78%) females, 484(95%) non-

Hispanic/Latino, 383(76%) got Pfizer for primary dose, 395 (78%) whites, median age(IQR):

39.9(32.4, 51.9) Timepoint 2: 739(84%) females, 845(96%) non-Hispanic/Latino, 657(75%) got

Pfizer for primary dose, 723 (82%) whites, median age(IQR): 43.0(35.1, 53.5) Timepoint 3:

215(79%) females, 263(96%) non-Hispanic/Latino, 225(82%) got Pfizer for primary dose, 238

(87%) whites, median age(IQR): 44.9(34.3, 55.6). NT were performed at Timepoint 1 (n=15),

Timepoint 2 (n=14), Timepoint 3 (n=16).

199
Figure 2: Comparison of neutralizing antibody titers (NT) to SARS-CoV-2 vaccine strain (WA-1),

Beta, Delta, and Omicron VOC from healthcare workers with paired serum samples in a

longitudinal cohort. Data shown at three timepoints: within 14-44 days post-dose-2 (Timepoint

1), at least 8 months post-dose-2 (Timepoint 2), and within 14-44 post-booster (Timepoint 3).

Top panel shows NT titer for each variant across the three timepoints with connecting lines

illustrating 15 paired samples in Timepoints 1 and 3. Bottom panel shows NT at each timepoint

for each VOC. Fold change (increase/difference) represents geometric median fold change. P-

values have been corrected for multiple comparisons using Bonferroni methods.