CADM2 is implicated in impulsive personality and numerous other traits by genome- and phenome-wide association studies in humans and mice

Sandra Sanchez-Roige*[1,2], Mariela V Jennings[1], Hayley H A Thorpe[3], Jazlene E Mallari[1], Lieke C van der Werf[1], Sevim B Bianchi[1], Calvin Lee[1], Travis T Mallard[4], Samuel A Barnes[1], Jin Yi Wu[1], Amanda M Barkley-Levenson[1], Ely C Boussaty[1], Cedric E Snethlage[1], Danielle Schafer[1], Zeljana Babic[1], Boyer D Winters[5], Katherine E Watters[6,7], Thomas Biederer[6], on behalf of the 23andMe Research Team[8], James Mackillop[9], David N Stephens[10], Sarah L Elson[8], Pierre Fontanillas[8], Jibran Y Khokhar[3], Jared W Young[1], Abraham A Palmer*[1,11]

[1]Department of Psychiatry, University of California San Diego, San Diego, CA, USA. [2]Department of Medicine, Vanderbilt University Medical Center, Nashville, Tennessee, USA. [3]Department of Biomedical Sciences, Ontario Veterinary College, University of Guelph, Ontario, Canada. [4]Psychiatric and Neurodevelopmental Genetics Unit, Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, 02114, USA. [5]Department of Psychology, University of Guelph, Guelph, ON, Canada. [6]Department of Neuroscience, Tufts University School of Medicine, Boston, MA, USA. [7]Department of Neurology, Yale School of Medicine, New Haven, CT, USA. [8]23andMe, Inc., Sunnyvale, CA, USA. [9]Peter Boris Centre for Addictions Research, McMaster University and St. Joseph's Healthcare Hamilton, Hamilton, ON, Canada and Homewood Research Institute, Guelph, ON, Canada. [10]Laboratory of Behavioural and Clinical Neuroscience, School of Psychology, University of Sussex, Brighton, UK. [11]Institute for Genomic Medicine, University of California San Diego, San Diego, CA, USA.

*Co-corresponding Authors: Sandra Sanchez-Roige, sanchezroige@ucsd.edu, Abraham A. Palmer, aap@ucsd.edu

Keywords: Cadm2, UPPS-P, BIS, impulsivity, GWAS, PheWAS, animal models, cross-species translation

ABSTRACT

Impulsivity is a multidimensional heritable phenotype that broadly refers to the tendency to act prematurely and is associated with multiple forms of psychopathology, including substance use disorders. We performed genome-wide association studies (**GWAS**) of eight impulsive personality traits from the Barratt Impulsiveness Scale and the short UPPS-P Impulsive Personality Scale (N=123,509-133,517 23andMe research participants of European ancestry), and a measure of Drug Experimentation (N=130,684). Because these GWAS implicated the gene *CADM2*, we next performed single-SNP phenome-wide studies (**PheWAS**) of several of the implicated variants in *CADM2* in a multi-ancestral 23andMe cohort (N=3,229,317, European; N=579,623, Latin American; N=199,663, African American). Finally, we produced *Cadm2* mutant mice and used them to perform a Mouse-PheWAS ("MouseWAS") by testing them with a battery of relevant behavioral tasks. In humans, impulsive personality traits showed modest chip-heritability (~6-11%), and moderate genetic correlations (r_g=.20-.50) with other personality traits, and various psychiatric and medical traits. We identified significant associations proximal to genes such as *TCF4* and *PTPRF*, and also identified nominal associations proximal to *DRD2* and *CRHR1*. PheWAS for *CADM2* variants identified associations with 378 traits in European participants, and 47 traits in Latin American participants, replicating associations with risky behaviors, cognition and BMI, and revealing novel associations including allergies, anxiety, irritable bowel syndrome, and migraine. Our MouseWAS recapitulated some of the associations found in humans, including impulsivity, cognition, and BMI. Our results further delineate the role of *CADM2* in impulsivity and numerous other psychiatric and somatic traits across ancestries and species.

INTRODUCTION

Impulsivity is a multifaceted psychological construct that has been broadly defined as thoughts or actions that are "poorly conceived, prematurely expressed, unduly risky or inappropriate to the situation, and that often result in undesirable consequences" [1]. Impulsivity has been repeatedly associated with numerous psychiatric diseases, including ADHD and substance use disorders [2, 3]. We previously performed genome-wide association studies (**GWAS**) of impulsive personality traits (n=21,806-22,861) using two of the most widely used impulsivity questionnaires, the Barratt Impulsiveness Scale (**BIS-11**; 3 traits) and the Impulsive Personality Scale (**UPPS-P**; 5 traits), as well as a measure of Drug Experimentation [4]. These traits were partially genetically correlated, suggesting that each impulsivity domain is governed by overlapping but distinct biological mechanisms [4, 5]. Our work also identified significant genetic correlations between impulsivity and numerous psychiatric and substance use traits, in line with the NIMH Research Domain Criteria (**RDoC**), proposing impulsivity as a transdiagnostic endophenotype for psychopathology [6].

The cell adhesion molecule 2 (*CADM2*) gene, which was the most robustly implicated gene in our prior GWAS of impulsivity [4], has also been extensively implicated in other risky and substance use behaviors [7]. *CADM2* mediates synaptic plasticity and is enriched in the frontal cortex and striatum, which are regions that regulate reward and inhibitory processes. We and others have implicated this gene in traits that may underlie disinhibition in humans, supporting the observed genetic correlations between impulsivity and personality [8], educational attainment [9], cognition [10], risk-taking [11], substance use [4, 10, 12–15], externalizing psychopathology [16], neurodevelopmental disorders [17, 18], physical activity [19], reproductive health [20, 21], metabolic traits [22], and BMI [23], among others (see GWAS Catalog www.ebi.ac.uk/gwas/). *Cadm2* knockout mice have previously been assessed for body weight and energy homeostasis [24] but have never been behaviorally characterized for measures of impulsivity or related behaviors.

Here, we took three approaches to elucidate genetic factors related to impulsivity. First, we collaborated with 23andMe, Inc., to extend upon our earlier GWAS of impulsivity [4] by increasing our sample size approximately 6-fold (n=123,509-133,517). Second, we performed single-SNP phenome-wide studies (**PheWAS**) of the 5 single nucleotide polymorphisms (**SNPs**) in and around *CADM2* that have been most strongly implicated by the current and prior GWAS. PheWAS were conducted in three ancestral groups (N=3,229,317, European; N=579,623, Latin American; N=199,663, African American) from the 23andMe research cohort, examining close

83   to 1,300 traits, most with no published GWAS. Finally, we performed a mouse-PheWAS

84   ("MouseWAS") by creating and phenotyping mice harboring a *Cadm2* mutant allele in a broad

85   battery of behavioral tasks that included analogous human measures of risk-taking and

86   impulsivity, substance use, cognition and BMI.

MATERIALS AND METHODS

Human Studies

GWAS cohort and phenotypes

We analyzed data from a cohort of up to 133,517 male and female research participants of European ancestry, a subset of which were analyzed in our prior publications [4, 13, 13, 25, 26]. All participants were drawn from the research participant base of 23andMe, Inc., a direct-to-consumer genetics company, and were not compensated for their participation. Participants provided informed consent and participated in the research online, under a protocol approved by the external AAHRPP-accredited IRB, Ethical & Independent Review Services (www.eandireview.com). During 4 months in 2015 and 14 months from 2018-2020, participants responded to a survey that included up to 139 questions pertaining to aspects of impulsivity and substance use and misuse. To measure impulsive personality, we used five subscales from the UPPS-P ([27, 28]; a 20-item that measures (lack of) Premeditation, (lack of) Perseverance, Positive Urgency, Negative Urgency, and Sensation Seeking; **Table S1**). We also administered the BIS-11 ([29]; a 30-item questionnaire that measures Attentional, Motor, and Nonplanning impulsiveness; **Table S1**). Lastly, we measured Drug Experimentation, defined as the number of substances an individual has used (adapted from the PhenX toolkit [30]; **Table S1**). We scored UPPS-P, BIS-11 and Drug Experimentation as previously described [4]. We used quantile normalization, since some scores were not normally distributed (**Figures S1-3**). Only individuals identified as European ancestry based on empirical genotype data [31] were included in this study. Basic demographic information about this sample is presented in **Table S2**. We used Pearson correlation coefficients (*r*) to measure the phenotypic relationships between impulsivity subscales and demographics.

Genome-wide association and secondary analyses

DNA extraction and genotyping were performed on saliva samples by CLIA-certified and CAP-accredited clinical laboratories of Laboratory Corporation of America. Quality control, imputation, and genome-wide analyses were performed by 23andMe [**Table S3**; (32,33)]. 23andMe's analysis pipeline performs linear regression assuming an additive model for allelic effects (**Supplementary Material**). Covariates included age (inverse-normal transformed), sex, the top five principal genotype components, and indicator variables for genotyping platforms. *p*-values were corrected for genomic control. We examined genotype*sex interactions for

suggestive loci. In addition, for the top loci, we examined African American and Latin American

23andMe research participants who had responded to the same survey.

We used the FUMA web-based platform (version 1.3.6a) and MAGMA v1.08 [32, 33] to

explore the functional consequences of the GWAS loci and to conduct gene-based analyses.

We used LDSC [34] to calculate genetic correlations (r_g) between UPPS-P, BIS and

Drug Experimentation, and 96 selected traits informed by prior literature.

Phenome-wide association scan (PheWAS) in 23andMe

We performed single-SNP PheWAS for 5 *CADM2* SNPs (rs993137, rs62263923,

rs11708632, rs818219, rs6803322) using up to 1,291 well-curated self-reported phenotypes

from a separate cohort of 23andMe research participants of European (N≤3,229,317), Latin

American (N≤579,623) and African American (N≤199,663) ancestries. We excluded traits with

<1,000 responses, based on a prior simulation study for PheWAS power analysis [35]. Ancestry

was determined by analyzing local ancestry ([31] **Supplementary Material**). The variants were

selected based on our GWAS results and previous literature (**Table S4**, and **Supplementary**

Material). Genotyped and imputed variant statistics in PheWAS are shown in **Table S5**.

An overview of the data collection process has been previously described [36]. All

regression analyses were performed using R version 3.2.2. We assumed additive allelic effects

and included covariates for age (as determined by participant date of birth), sex, and the top five

ancestry-specific principal components. We used a 5% FDR correction for multiple testing.

MouseWAS

Subjects, behavioral characterization, and statistical analyses

Our *Cadm2* mutant mice were produced at the University of California San Diego,

Moores Cancer Center, Transgenic Mouse Core. We used the JM8.N4 cryosperm line

(CSD70565 KOMP), which carries a floxed null allele in the *Cadm2* gene (**Figure S30**), on a

C57BL/6N background. We crossed the floxed null allele line with a constitutive CRE driver line

(Stock# 014094; The Jackson Laboratory), yielding a global constitutive null allele. We used a

heterozygous x heterozygous (**HET**) breeding scheme, which produced homozygous (**HOM**)

mutant *Cadm2* mice and their HET and wildtype (**WT**) littermates. Mice were genotyped using

allele-specific polymerase chain reaction on ear notch tissue followed by gel electrophoresis

[37]. CADM2 protein expression levels were quantified by western blotting (**Figure S31**).

Five separate cohorts of male and female mice were used for these studies. See

Supplementary Material for a more detailed description of the tasks and analyses of main

variables. Procedures were approved by the University of California San Diego Institutional

Animal Care and Use Committee. The UCSD animal facility meets all federal and state

requirements for animal care and was approved by the American Association for Accreditation

of Laboratory Animal Care. Procedures from cohort 2 were conducted in accordance with the

Canadian Council on Animal Care and were approved by the University of Guelph Institutional

Animal Care and Use Committee.

RESULTS

Genome-wide association analyses and secondary analyses

Self-reported impulsivity and Drug Experimentation scores are shown in **Table S6**. We found that ~6-11% of the phenotypic variation of these traits can be explained by common variants (**Table S7**). We identified 21 genome-wide significant associations ($p<5.0E-08$) for UPPS-P (5 traits), BIS (3 traits), and Drug Experimentation (**Figure 1**; **Figures S4-21**; **Table S8**). Although we tested 9 traits, in keeping with the standards of the field, we did not adjust the significance threshold. We also detected several nominal associations ($p<1.0E-06$, **Table S8**); we discuss some of them in the **Supplementary Material**.

GWAS of UPPS-P

Premeditation. We detected one significant hit (rs2958162, $p=2.50E-10$), located on chromosome 18 in the *TCF4* gene, which encodes a helix-loop-helix transcription factor and is widely expressed throughout the body and during development. Polymorphisms in *TCF4* have been associated with risk-taking and adventurousness [15], alcohol consumption [38], schizophrenia [39], depression [40, 41], and neuroticism [42, 43] (**Table S9**); *TCF4* is also a non-GWAS candidate gene for other psychiatric and neurological conditions [44].

Perseverance. We detected one significant association (rs5943997, $p=1.50E-8$) in the *POLA1* gene on the X chromosome. *POLA1* has been related to blood traits [44] and neurodevelopmental disorders [45], but its association with impulsivity is novel.

Positive Urgency. We identified one significant hit (rs143987963, $p=4.30E-08$) on chromosome 12, near the genes *MDM1* and *RAP1B*; however, inspection of the locus zoom plot (**Figure S9**) does not support a robust association.

Negative Urgency. We detected three significant hits: rs4840542 ($p=1.60E-09$), on chromosome 8, in the *XKR6* gene; rs5008475 ($p=4.90E-09$), on chromosome 5, near *TMEM161B* and *MEF2C*; and rs7829975, on chromosome 8, near *SGK223* and *CLDN23* ($p=5.00E-09$). Variants in strong LD with rs4840542 and rs7829975 are highly pleiotropic, and have been previously associated with several traits (**Table S9**), including body mass index (**BMI**) [46, 47], neuroticism [48, 49], depression [50], blood pressure, and alcohol consumption [51]. *XKR6* was also implicated in a recent GWAS of externalizing [16], and a GWAS of anxiety and depression [50].

Sensation Seeking. We detected 5 significant associations. First, we again observed a
previously reported [4] association with a SNP near *CADM2* (rs11288859, *p*=2.10E-09) on
chromosome 3. We also detected an association with a SNP in *TCF4* (rs2958178, *p*=3.80E-12).
We identified a significant hit in *CACNA2D1* (rs38547, *p*=2.10E-08) on chromosome 18.
CACNA2D1 has been previously associated with feeling nervous [48], and levels of sex
hormone-binding globulin [52]. Furthermore, we found a significant association (rs1605379*,*
p=3.80E-08) on chromosome 16, near *CYLD* and *SALL1*. SNPs in strong LD with rs1605379
have been previously identified for risk-taking, adventurousness, and smoking initiation (**Table
S9**). Lastly, we found a significant association (rs12600879*, p*=4.10E-08) on chromosome 17,
near *TBX21* and *OSBPL7*. Variants in strong LD with rs12600879 have been associated with
BMI [53], but the finding in relation to impulsivity is novel.

GWAS of BIS-11

Attentional. We identified one significant association (rs10196237, *p*=1.10E-08) on
chromosome 2, near the genes *SPHKAP and PID1*. *SPHKAP* has been previously associated
with educational attainment [9], but the association with impulsivity is novel.

Motor. We detected one significant association near *CADM2* (rs35614735, *p*=3.20E-11).
We also identified an association (rs111502401, *p*=2.00E-08), on chromosome 19, near the
genes *ZNF229* and *ZNF180*; however, inspection of the regional association is not supportive of
a strong association (**Figure S17**).

Nonplanning. We detected 2 variants: rs35614735 (*p*=4.70E-12) near *CADM2*, which
was the same SNP identified for Motor impulsivity; and rs6872863 (*p*=1.20E-08) in the gene
ELOVL7 on chromosome 5. Variants in strong LD with rs6872863 have been reported for a
variety of traits including educational attainment, mathematical ability [9], household income
[54], and brain morphology, such as cortical surface area [55] (**Table S9**). However, there is
extensive LD in this region, making the association difficult to interpret.

GWAS of Drug Experimentation

We previously reported [4] a suggestive association (rs2163971, *p*=3.00E-07) near the
CADM2 gene. In the present study, we identified a nearby SNP that was genome-wide
significant (rs35614735, *p*=2.80E-15). We also report 4 novel hits (rs951740, *p*=9.70E-10,
PTPRF on chromosome 1; rs12713405, *p*=9.70E-09, *BLC11A* on chromosome 2*;* rs67660520,

p=7.60E-09, *CADPS2* on chromosome 7*;* rs7128648, *p*=2.50E-09, *NCAM1* on chromosome

11). Intriguingly, *PTPRF* has been recently associated with problematic prescription opioid use

[25] and opioid use disorder [56], as well as smoking initiation/cessation [57], cognition [58], and

educational attainment [9] (**Table S9**). Variants in strong LD with rs67660520 have been

associated with ADHD [59], smoking initiation [57], number of sexual partners [15] and BMI [47]

(**Table S9**). *NCAM1* variants have been previously associated with alcohol, cannabis and

smoking behaviors [66, 85], mathematical ability [9], and anxiety and depression [50], among

other traits.

Gene-based analyses

Similar to the GWAS results, gene-based analyses using MAGMA identified an

association (Bonferroni *p*<2.53E-06; **Table S10**) between *CADM2* and 6 of the 9 traits

examined in this paper: Premeditation, Sensation Seeking (UPPS-P); Attentional, Motor and

Nonplanning (BIS-11); and Drug Experimentation. *TCF4*, which was significantly associated

with Premeditation and Sensation Seeking in the GWAS, was significantly associated with these

traits in the gene-based analysis. *MAPT*, which has been previously associated with many traits

including multiple alcohol-related behaviors [13], was implicated in Negative Urgency. Lastly,

KDM4A, which was recently related to problematic opioid use and interacts with selective

serotonin reuptake inhibitors and dopaminergic agents [25], was significantly associated with

Drug Experimentation.

Phenotypic and genetic correlations

A phenotypic and genetic correlation matrix of all 9 traits is shown in **Figure S22** and

Tables S11-12. Consistent with the literature and our prior work [4, 5, 89, 90], both phenotypic

and genetic inter-correlations among the UPPS-P and BIS subscales were high and positive,

with the exception of Sensation Seeking and Perseverance, suggesting that these traits may

represent relatively different constructs [5, 13, 89]. Drug experimentation was positively and

significantly associated with all impulsive personality traits.

All impulsivity traits were phenotypically associated (*r*=-0.34-0.11) with demographic

variables (**Table S12**), impulsivity scores being greater in male and younger research

participants, compared to female and older participants; and in participants with higher BMI,

lower household income, and fewer years of education, as we previously reported [13].

Figure 2 shows a genetic correlation matrix of BIS, UPPS-P, Drug Experimentation and
several other phenotypes (full results in **Table S13**). As anticipated, we found positive moderate
to high genetic correlations (r_g=0.25-0.79) between virtually all UPPS-P (except Perseverance
and Sensation Seeking) and BIS subscales, and Drug Experimentation, and substance use
disorders **Table S13**).

We also observed moderate to strong associations between all impulsive subscales
(except UPPS-P Perseverance) and other personality traits, such as risk-taking (r_g=0.15-0.65),
neuroticism (r_g=-0.23-0.84), and loneliness (r_g=0.17-0.54), particularly for Positive and Negative
Urgency. Extraversion was positively associated with Sensation Seeking (r_g=0.34). Externalizing
psychopathology, which represents disorders and behaviors characterized by deficits in
inhibition, was strongly associated with all impulsivity facets (r_g=0.28-0.92), except
Perseverance.

We also identified positive associations with an array of psychiatric phenotypes,
including ADHD (r_g=0.20-0.47), depression (r_g=-0.13-0.47) and anxiety (r_g=-0.38-0.61)
disorders, and cross-disorder (r_g=0.12-0.44). The associations were again primarily significant
for all except Perseverance and Sensation Seeking. Other disorders showed weaker
associations (e.g., schizophrenia, r_g=-0.09-0.15) or were only significantly associated with one
impulsivity facet [e.g., anorexia nervosa (Perseverance, r_g=-0.16); bipolar disorder (Motor,
r_g=0.22)].

Most impulsivity subscales were genetically correlated with lower socioeconomic
variables [e.g., educational attainment (r_g=-0.49 to -0.16), income (r_g=-0.38 to -0.16), Townsend
index (r_g=0.18-0.58)].

Metabolic and medical phenotypes, such as BMI (r_g=0.18-0.28), chronic pain (r_g=0.22-
0.46), insomnia (r_g=0.20-0.42), and coronary artery disease (r_g=0.18-0.30), were genetically
correlated with all impulsive subscales (except Perseverance and Sensation Seeking). We also
noted negative genetic associations with parental longevity (r_g=-0.17 to -0.32).

PheWAS

To explore the impact of specific variants in and around *CADM2*, we performed single-
SNP PheWAS using 5 of the most implicated SNPs, independently, against 1,291 traits (**Figure
3**). The list of PheWAS association results using the 23andMe cohort after 5% FDR correction is

available in **Tables S14** (summary)**, S15** (Europeans), **S16** (Latin American) and **S17** (African
Americans).

In European cohorts, *CADM2* variants had been previously identified to be significantly
associated with numerous traits (**Table S18**). Most SNPs were highly correlated (R^2>0.1) and
tagged similar traits (**Figure S23**), but the overlap was incomplete (**Figure S24** and **Table S19**).
rs993137, located at 85,449,885 bp on chromosome 3, showed the highest number of
associations (378), which we describe below.

We replicated all previously known associations in 23andMe participants of European
ancestry, identifying signals across all categories tested (**Table S15**). These included negative
associations with risky behavior (e.g., lower risk for adventurousness [β=-0.05, *p*=1.33E-08],
risk-taking tendencies [β=-0.02, *p*=1.13E-07]) and substance use behaviors (e.g., lower risk for
alcohol consumption [β=-0.03, *p*=2.05E-09] and tobacco initiation (β=-0.02, *p*=3.66E-12; but see
packs per day, β=0.01, *p*=1.05E-03), as well as negative associations with psychiatric disorders
characterized by deficits in impulsivity, such as lower risk for ADHD (β=-0.05, *p*=2.17E-41).
Furthermore, we found positive associations with educational outcomes (e.g., higher
educational attainment (β=0.03, *p*=1.67E-12). Novel findings included positive associations with
allergies (β=0.04, *p*=4.51E-03), anxiety (e.g., panic [β=0.02, *p*=6.82E-08]), and medical
conditions (e.g., IBS [β=0.02, *p*=8.89E-07]), anemia (β=0.01, *p*=8.30E-74), hepatitis C (β=-0.06,
p=8.36E-10). Intriguingly, we also detected positive associations with pain phenotypes (β=0.02,
p=8.37E-12) and a need for a higher dose of pain medication (β=0.01, *p*=1.02E-06).

For the overlapping phenotypes, UK Biobank PheWAS results [91] largely supported the
23andMe PheWAS findings (except for smoking behaviors). For example, we identified
associations with dietary traits (e.g., daily fruit and vegetable intake (β=-0.01, *p*=4.23E-11),
pastry frequency (β=0.01, *p*=7.36E-06), sleep quality (β=-0.01, *p*=2.53E-03), and number of
pregnancies (β=-0.01, *p*=7.69E-04), among others (**Table S15**, [12]).

For the PheWAS of the Latin American cohort, 47 traits were significantly associated
with *CADM2* variants (**Table S16**). The highest number of associations were again observed for
rs993137 (47), which are described below. Similarly, although some of the SNPs were
correlated (R^2>0.1; **Figure S24**), the overlap was incomplete (**Figure S26**, **Table S20**). The
pattern of associations was consistent with those described in the European cohort. The
strongest associations were with risky behaviors, such as adventurousness (β=-0.04, *p*=1.76E-

17), risk-taking (β=-0.02, p=5.90E-07), alcohol consumption (β=-0.03, p=1.41E-12), and

disorders characterized by high levels of impulsivity, such as ADHD (β=-0.04, p=4.74E-10). The

novel findings were, again, with multiple forms of allergies (e.g., seasonal allergies, β=0.03,

p=3.0E-04), migraine (β=0.04, p=1.56E-04), sleep behaviors (e.g., sleep apnea, β=-0.03,

p=6.76E-04), among others.

All findings that were in common between the European and Latin American cohorts

showed the same direction of effect and similar effect sizes. We did not identify FDR-significant

associations in the African American cohort (**Table S17**). The effect sizes were generally

extremely small (**Figures S27-28**), as is expected for a single gene and complex traits.

MouseWAS

Figure 4 summarizes the MouseWAS results across the five cohorts tested. Full

statistics and additional secondary measures are described in the **Supplementary Material** and

Table S20.

Cohort 1 - Motivation, inhibition, and risk-taking behavior. No differences in motivation

were found between WT and HET mice during the Progressive Breakpoint task [$F_{(1,51)}$=0.003,

p=9.57E-01; **Figure 4A**]. However, we noted significant genotype differences in behavioral

flexibility in the Probabilistic Reversal Learning (**PRL**) task, as indexed by the number of trials to

first reversal [$F_{(1,42)}$=4.27, p=4.50E-02; **Figure 4B**], and risky behavior in the mouse Iowa

Gambling Task (**IGT**; $F_{(1,51)}$=4.70, p=3.50E-02; **Figure 4C**], HET mice requiring fewer trials to

reach criterion and choosing risky options less frequently than WT mice (p<0.05), respectively.

The number of premature responses, on the contrary, were higher in HET mice [$F_{(1,51)}$=5.78,

p=2.00E-02] compared to WT mice (p<0.05; **Figure 4D**). In the Behavioral pattern monitor

(**BPM**), HET mice exhibited greater exploratory behavior, as shown by an increase in hole-

pokes [$F_{(1,53)}$= 4.88, p=3.20E-02; **Figure 4E**], compared to WT mice (p<0.05), but general levels

of activity, such as distance traveled ($F_{(1,53)}$= 0.42, p=5.21E-01; **Figure 4F**), were similar across

the genotypes. Lastly, although the startle response was equal across the groups (**Figure 4G**),

prepulse inhibition (**PPI**) was larger in HET mice compared to WT mice (p<0.05; **Figure 4H**),

particularly at ISI 25 and 100 in HET mice [$F_{(1,53)}$=8.23, p=6.00E-03, $F_{(1,53)}$=4.50, p=3.90E-02,

respectively].

Cohort 2 - Motoric impulsivity. The main outcome tested in cohort 2 were premature

responses via the 5-choice serial reaction time task (**5CSRTT**; **Figure 4J-M**). Premature

339 responses were lower in HOM ($p<0.001$) and WT ($p<0.02$) mice compared to HET mice under

340 standard conditions ($F_{(2,36)}=8.74$, $p=8.06\text{E-}04$; **Figure 4J**), and compared to both HET ($p<0.001$)

341 and WT ($p<0.01$) mice during a long ITI session ($H_{(2)}=16.10$, $p=3.19\text{E-}04$; **Figure 4L**). HOM

342 mice were faster at learning the 5CSRTT, requiring fewer days for adequate baseline

343 performance ($F_{(2,36)}=7.42$, $p=2.00\text{E-}03$; **Figure 4I**), compared to WT mice ($p<0.01$).

344   *Cohort 3 - General locomotion, anxiety-like behavior, and ethanol consumption.* We

345 found a significant effect of genotype on the distance traveled in the Open Field [**OF**;

346 $F_{(2,70)}=7.525$, $p=1.00\text{E-}03$; **Figure 4N**], HOM mice showing higher levels of locomotor activity

347 than WT mice ($p=1.40\text{E-}02$). No differences in anxiety-like behavior were detected across WT,

348 HET or HOM mice in the Elevated Plus Maze (**EPM**) or Light-Dark Box (**LDB**) tests (**Figure 4O-**

349 **P**, **Table S20**). The total amount of ethanol consumed during the drinking-in-the-dark (**DID**)

350 paradigm did not differ between the groups ([$F_{(2,78)}=1.084$, $p=3.44\text{E-}01$]; **Figure 4Q**).

351   *Cohort 4 - Body weight.* Relative to WT mice, there was a significant reduction in body

352 weight in HOM mice from week 21 onwards ($\beta=-3.74\pm1.27$, $p=4.00\text{E-}03$; **Figure 4R**).

353   *Cohort 5 - Dendrite morphology.* Quantitative analyses of MSN in the NAc revealed no

354 difference in dendritic spine density across the groups (**Figure 5S**).

DISCUSSION

In this study, we performed the largest GWAS of impulsive personality traits to date, we conducted the first multi-ancestral PheWAS exploring the role of *CADM2* on a diverse array of traits, and we performed a corresponding MouseWAS using *Cadm2* mutant mice to assess its role in impulsivity and other relevant behaviors. We extend on prior findings [4, 5] showing that the genetic architecture of impulsivity facets only partially overlap, providing further support to the idea of impulsivity being a multifaceted construct even at the genetic level. We identified positive genetic correlations across multiple domains, particularly substance use disorders, confirming that NIMH RDoC transdiagnostic domains [6], or endophenotypes, such as impulsive personality traits, can be used to dissect the genetic basis of psychiatric illness and normal functioning. RDoC or transdiagnostic traits are beneficial because they enable translational research and provide a more granular biological understanding of psychiatric disorders. Using mouse and human correlates, we provide further evidence that *CADM2* is a robust candidate gene for impulsivity and an important modulator of numerous other psychiatric and somatic traits.

We increased the sample size of our prior GWAS of impulsivity by almost 6-fold and identified 21 genome-wide significant loci implicated in impulsive personality and Drug Experimentation. For instance, SNPs located in the gene *TCF4* were implicated in 3 subscales; this gene is also highly pleiotropic for other psychiatric conditions. Furthermore, we identified associations with *NCAM1*, which, intriguingly, is a critical member of the NTAD (*NCAM1-TTC12-ANKK1-DRD2*) gene cluster [66] and variants correlated with *NCAM1* in that cluster have been associated with differences in D2 receptor density [67]. We also detected associations near *XKR6* and *AFF3*, which have been recently implicated in externalizing psychopathology [16], and *PTPRF* and *KDM4A*, recently implicated in problematic opioid use [25] and opioid use disorder [56]. Although in this report we focused on *CADM2*, functional studies of those genes are also warranted. Furthermore, we found nominal evidence for candidate gene studies implicating monoamine neurotransmitters in impulsivity and Drug Experimentation (*DRD2*, *HTR3B*). High impulsivity depends on a neural network that includes the ventral striatum (subsuming the NAc) with top-down control from prefrontal cortical regions, and is modulated by monoamine neurotransmitters including dopamine and serotonin [68]; this is the first GWAS to implicate genes modulating these systems as robust candidate genes for impulsivity.

Recent studies have implicated the *CADM2* gene in impulsivity and traits associated with

reward sensitivity and multiple domains of human health. We confirmed numerous previously

reported associations and extended our findings of variants related to *CADM2*. *CADM2* was

significantly associated with 4 out of the 9 traits that we measured in GWAS and 6 out of the 9

traits that we measured in gene-based analyses. In the PheWAS, *CADM2* variants were

associated with decreased risk for externalizing psychopathology, but also increased risk for

internalizing psychopathology (anxiety, depression, OCD). We also observed novel associations

with migraines and various allergies. Others using a similar approach with UK Biobank data

have found that this enrichment of associations is higher than expected [12] compared to other

genes. These results provide evidence that *CADM2* variants are associated with broad health

outcomes, but whether this gene affects human health via disruptions in inhibitory control or

reward systems, or whether it acts via multiple pathways [69], is still not fully understood.

A relatively unique feature of our study is that, to follow up on the *CADM2* loci implicated

in human studies, we generated a *Cadm2* mutant mouse line and used it to perform a PheWAS-

like study in mice, which we have termed a MouseWAS to emphasize its conceptual similarity to

human PheWAS studies. These functional experiments provided information about the causality

and directionality of effects in its reported associations. We found evidence that loss of *Cadm2*

resulted in *less* risky behavior and *improved* information processing, extending on prior work in

humans [4, 10, 16, 60, 67].

Cadm2 expression may uniquely contribute to the different domains of impulsivity. The

IGT assays preference for high risk, high reward (disadvantageous) choices vs low risk, low

reward (advantageous) choices [70]. HET mice exhibited a greater preference for selecting the

safe option vs their WT littermates. This finding can be contrasted with the *elevated* premature

responses in the 5CSRTT seen in HET vs WT mice, reflective of motoric impulsivity. However,

premature responses have also been linked to temporal discrimination, wherein mice and

humans overestimating the passage of time exhibit higher premature responses [71, 72]. The

preference for less risky options of HET mice in the IGT could reflect their misjudgment of time –

resulting in higher premature responses – and thus avoidance of higher temporal punishment in

the IGT.

We also observed genotype differences in performance that could be indicative of

Cadm2 function in information processing. HET mice exhibiting *better* PPI at the shortest

temporal window (25 ms) supports the premise that these mice have faster processing speeds.

HET mice also showed small increases in hole-poking in the BPM test, which is thought to

reflect exploration of an environment and information gathering. Finally, we observed that HOM

mice acquired 5CSRTT faster than WT littermates. Taken together, these results suggest that

Cadm2 reduction may improve some facets of information processing.

Findings from the 5CSRTT provide evidence that *Cadm2* deletion improves some

information processing and impulsivity outcomes, while being detrimental to others. HET mice

were the most likely to commit 5CSRTT premature responses, although HOM mice were

surprisingly the least likely to make premature responses. Interestingly, although not significant,

there was a consistent elevation in the number of premature responses committed by the HOM

mice as the stimulus duration was reduced. This could suggest that HOM mice, like HET mice,

may show motoric impulsivity deficits when performing tasks that require greater attentional

demand. Compared with WT, HOM mice also showed impaired accuracy performance under

RSD conditions, in line with our human findings of *CADM2* association with BIS Attentional, and

cognitive function by others [10]. The heterogeneity of performance outcomes in the HOM mice

further supports a unique but overlapping contribution of genetics across impulsivity domains.

In this paper, we have translated measures from human to mice. These studies begin

the process of understanding the biological basis of associations identified by GWAS. The

methods for measuring impulsivity in humans and mice are fundamentally different. Despite

these differences, our MouseWAS identified several measures of impulsivity that were

influenced by *Cadm2*, consistent with our observations in humans. Furthermore, *CADM2* has

been shown to be implicated in BMI in humans [24, 73] and energy homeostasis in mice [24];

extending on this, we found novel evidence of body weight reductions in adult mutant mice.

Interestingly, *Cadm2* did not have more general effects on mouse behavior; for instance, we did

not observe deficits in anxiety-like behavior or general motivation, as some of the human

PheWAS findings revealed. A few other measures were also inconsistent across species,

particularly measures of alcohol consumption, where *CADM2* showed a role in humans [7, 13,

74, 75] but not mice. Lastly, some measures identified by our human PheWAS (e.g., allergies

and other medical conditions) were not examined in our MouseWAS. This approach highlights

the challenges of using mouse models to further investigate the role of specific genes in

behavioral traits.

CADM2 encodes the immunoglobulin adhesion protein SynCAM 2, which is part of the

family of synaptic adhesion molecules known as SynCAMs. Studies have shown the influence

of SynCAMs on synaptogenesis [76–80], axon guidance [81], and neuron myelination [82–84],

processes that have direct effects on the pathology of neurodevelopmental diseases [54].

CADM2 is strongly expressed in the striatum and frontal cortex, which are core regions that

regulate impulsivity [68]. We did not observe changes in spine density in the Nac, which

suggests that *Cadm2* may not have a role as a postsynaptic organizer of spines in this region,

or may have redundant functions that are compensated in the mutant mice by other molecules.

Based on *in-silico* analyses in humans, *CADM2* expression seems to be greater at earlier

stages of development (**Figure S29**); whether *Cadm2* may affect earlier stages of development

(prenatal and early postnatal) that are compensated in adulthood has not been investigated in

this study.

Several limitations of this study are worth noting. The discovery GWAS only includes

male and female participants of European ancestry. While we provided exploratory analyses of

top variants in other ancestries and by broken down by sex (**Supplementary Table 22**), larger

sample sizes would be needed to perform GWAS separately in males and females. Our results

are also biased by potential ascertainment and characteristics of the sample; the 23andMe

participant population is more educated and has higher socioeconomic status and lower levels

of drug use and impulsivity than the general US population [84]. Replication in additional cohorts

with different characteristics is warranted. Moreover, although the traits we studied are extracted

via well-established questionnaires, they are self-reported measures, which are different from

behavioral phenotypes [85, 86]. Another issue is that, although we tested multiple variants in the

CADM2 loci, further conditional analyses are required to determine if this signal and previously

reported associations implicating *CADM2* loci, including a large non-coding rare deletion in the

first intron of *CADM2* [69], may tag the same underlying genetic effect. We are also unaware of

the sequence of events, and whether there is true pleiotropy or mediation effects has not been

examined. The analyses were well-powered for moderate and large effect sizes. Still, for

unclear reasons, despite similar minor allele frequencies and imputation quality of the SNPs we

tested across all ancestries, we identified no significant associations in the African American

cohort. Finally, although our mouse studies detected some discordant cross-species effects of

Cadm2 on behavior, background strain effects [87] or subtle allelic variations (vs whole KO)

may explain those differences. While some results are suggestive of additive effects, we were

unable to evaluate different genetic models due to lack of sufficient sample sizes for HOM mice.

Future multivariate analyses examining paths of commonality and specificity across impulsivity

facets may provide further insights not herein examined.

In conclusion, we show that impulsivity facets are extremely polygenic, but of very high
transdiagnostic significance. Genetic studies using research participants not ascertained for
neuropsychiatric disorders may represent an efficient and cost-effective strategy for elucidating
the genetic basis and etiology of genetically complex psychiatric diseases. Using homologous
measures of impulsivity in mice and humans across three ancestral backgrounds, we provide
evidence of the overarching role of *CADM2* on impulsivity, and a much broader impact on
human health.

Data availability

We will provide summary statistics for the top 10,000 SNPs upon publication (**Tables**
S22-30). Full GWAS summary statistics will be made available through 23andMe to qualified
researchers under an agreement with 23andMe that protects the privacy of the 23andMe
participants. Please visit https://research.23andme.com/collaborate/#dataset-access/ for more
information and to apply to access the data.

Acknowledgements

We would like to thank the research participants and employees of 23andMe for making
this work possible. The following members of the 23andMe Research Team contributed to this
study: Stella Aslibekyan, Adam Auton, Elizabeth Babalola, Robert K. Bell, Jessica Bielenberg,
Katarzyna Bryc, Emily Bullis, Daniella Coker, Gabriel Cuellar Partida, Devika Dhamija,
Sayantan Das, Teresa Filshtein, Kipper Fletez-Brant, Will Freyman, Karl Heilbron, Pooja M.
Gandhi, Karl Heilbron, Barry Hicks, David A. Hinds, Ethan M. Jewett, Yunxuan Jiang, Katelyn
Kukar, Keng-Han Lin, Maya Lowe, Jey C. McCreight, Matthew H. McIntyre, Steven J. Micheletti,
Meghan E. Moreno, Joanna L. Mountain, Priyanka Nandakumar, Elizabeth S. Noblin, Jared
O'Connell, Aaron A. Petrakovitz, G. David Poznik, Morgan Schumacher, Anjali J. Shastri, Janie
F. Shelton, Jingchunzi Shi, Suyash Shringarpure, Vinh Tran, Joyce Y. Tung, Xin Wang, Wei
Wang, Catherine H. Weldon, Peter Wilton, Alejandro Hernandez, Corinna Wong, Christophe
Toukam Tchakouté.

We would also like to thank The Externalizing Consortium for sharing the GWAS
summary statistics of externalizing. The Externalizing Consortium: Principal Investigators:
Danielle M. Dick, Philipp Koellinger, K. Paige Harden, Abraham A. Palmer. Lead Analysts:
Richard Karlsson Linnér, Travis T. Mallard, Peter B. Barr, Sandra Sanchez-Roige. Significant
Contributors: Irwin D. Waldman. The Externalizing Consortium has been supported by the

National Institute on Alcohol Abuse and Alcoholism (R01AA015416 -administrative supplement),
and the National Institute on Drug Abuse (R01DA050721). Additional funding for investigator
effort has been provided by K02AA018755, U10AA008401, P50AA022537, as well as a
European Research Council Consolidator Grant (647648 EdGe to Koellinger). The content is
solely the responsibility of the authors and does not necessarily represent the official views of
the above funding bodies. The Externalizing Consortium would like to thank the following groups
for making the research possible: 23andMe, Add Health, Vanderbilt University Medical Center's
BioVU, Collaborative Study on the Genetics of Alcoholism (COGA), the Psychiatric Genomics
Consortium's Substance Use Disorders working group, UK10K Consortium, UK Biobank, and
Philadelphia Neurodevelopmental Cohort.

Disclosures

MVJ, SBB, YH, SSR and AAP were supported by funds from the California Tobacco-
Related Disease Research Program (TRDRP; Grant Number 28IR-0070, T29KT0526 and
T32IR5226). SBB was also supported by P50DA037844, SSR was also supported by NIH/NIDA
DP1DA054394. JM and SSR were supported by the Brain and Behavior Foundation (grant
27676) and the Families for Borderline Personality Disorder Research (Beth and Rob Elliott)
2018 NARSAD Young Investigator Grant. The content is solely the responsibility of the authors
and does not necessarily represent the official views of the National Institutes of Health. SAB
was supported by NIH/NIMH grants R01MH108653 and R21MH117518. AMBL was supported
by NIH/NIAAA grant K99AA027835. TB acknowledges support by NIH/NIDA R01 DA018928.
JM is supported by the Peter Boris Chair in Addictions Research. HHAT is funded through a
Natural Science and Engineering Research Council PGS-D scholarship, and studies in Cohort 2
were supported by the Canadian Institutes of Health Research Project Grant (PJT-173442 to
JYK).

PF and SLE are employees of 23andMe, Inc., and hold stock or stock options in
23andMe. JY reports having received grant support funding from Sunovion, Heptares, and
Gilgamesh, as well as honoraria from Marvel Biotech, none of which were involved in the
current project. The other authors report no conflict of interest.

REFERENCES

1.   Daruna JH, Barnes PA. A neurodevelopmental view of impulsivity. Impuls. Client Theory Res. Treat., Washington, D.C.: American Psychological Association; 1993. p. 23-27.
2.   Coskunpinar A, Dir AL, Cyders MA. Multidimensionality in Impulsivity and Alcohol Use: A Meta-Analysis Using the UPPS Model of Impulsivity. Alcohol Clin Exp Res. 2013;37:1441–1450.
3.   Jackson JNS, MacKillop J. Attention-Deficit/Hyperactivity Disorder and Monetary Delay Discounting: A Meta-Analysis of Case-Control Studies. Biol Psychiatry Cogn Neurosci Neuroimaging. 2016;1:316–325.
4.   Sanchez-Roige S, Fontanillas P, Elson SL, Gray JC, de Wit H, MacKillop J, et al. Genome-wide association studies of impulsive personality traits (BIS-11 and UPPSP) and drug experimentation in up to 22,861 adult research participants identify loci in the *CACNA1I* and *CADM2* genes. J Neurosci. 2019:2662–18.
5.   Gustavson DE, Friedman NP, Fontanillas P, Elson SL. The Latent Genetic Structure of Impulsivity and Its Relation to Internalizing Psychopathology. Psychol Sci. 2020;31:1025-1035.
6.   Insel TR. The NIMH Research Domain Criteria (RDoC) Project: Precision Medicine for Psychiatry. Am J Psychiatry. 2014;171:395–397.
7.   Arends RM, Pasman JA, Verweij KJH, Derks EM, Gordon SD, Hickie I, et al. Associations between the CADM2 gene, substance use, risky sexual behavior, and self-control: A phenome‐wide association study. Addict Biol. 2021;26:e13015.
8.   Boutwell B, Hinds D, Tielbeek J, Ong KK, Day FR, Perry JRB, et al. Replication and characterization of CADM2 and MSRA genes on human behavior. Heliyon. 2017;3:e00349.
9.   Lee JJ, Wedow R, Okbay A, Kong E, Maghzian O, Zacher M, et al. Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. Nat Genet. 2018;50:1112–1121.
10.   Ibrahim-Verbaas CA, Bressler J, Debette S, Schuur M, Smith AV, Bis JC, et al. GWAS for executive function and processing speed suggests involvement of the CADM2 gene. Mol Psychiatry. 2016;21:189–197.
11.   Strawbridge RJ, Ward J, Cullen B, Tunbridge EM, Hartz S, Bierut L, et al. Genome-wide analysis of self-reported risk-taking behaviour and cross-disorder genetic correlations in the UK Biobank cohort. Transl Psychiatry. 2018;8:39.
12.   A. Pasman J, Chen Z, Vink JM, Van Den Oever MC, Pattij T, Vries TJD, et al. The *CADM2* gene and behavior: A phenome-wide scan in UK-Biobank. medRxiv. 2021.
13.   Sanchez-Roige S, Palmer AA, Fontanillas P, Elson SL, the 23andMe Research Team, the Substance Use Disorder Working Group of the Psychiatric Genomics Consortium, Adams MJ, et al. Genome-Wide Association Study Meta-Analysis of the Alcohol Use Disorders Identification Test (AUDIT) in Two Population-Based Cohorts. Am J Psychiatry. 2019;176:107–118.
14.   Pasman JA, Verweij KJH, Gerring Z, Stringer S, Sanchez-Roige S, Treur JL, et al. GWAS of lifetime cannabis use reveals new risk loci, genetic overlap with psychiatric traits, and a causal effect of schizophrenia liability. Nat Neurosci. 2018;21:1161–1170.
15.   Karlsson Linnér R, Biroli P, Kong E, Meddens SFW, Wedow R, Fontana MA, et al. Genome-wide association analyses of risk tolerance and risky behaviors in over 1 million individuals identify hundreds of loci and shared genetic influences. Nat Genet. 2019;51:245–257.
16.   Linnér RK. Multivariate analysis of 1.5 million people identifies genetic associations with traits related to self-regulation and addiction. Nat Neurosci. 2021;24:27.

17. Casey JP, Magalhaes T, Conroy JM, Regan R, Shah N, Anney R, et al. A novel approach
of homozygous haplotype sharing identifies candidate genes in autism spectrum disorder.
Hum Genet. 2012;131:565–579.
18. Demontis D, Walters RK, Martin J, Mattheisen M, Als TD, Agerbo E, et al. Discovery of
the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. Nat
Genet. 2019;51:63–75.
19. Klimentidis YC, Raichlen DA, Bea J, Garcia DO, Wineinger NE, Mandarino LJ, et al.
Genome-wide association study of habitual physical activity in over 377,000 UK Biobank
participants identifies multiple variants including CADM2 and APOE. Int J Obes.
2018;42:1161–1176.
20. Day FR, Helgason H, Chasman DI, Rose LM, Loh P-R, Scott RA, et al. Physical and
neurobehavioral determinants of reproductive onset and success. Nat Genet.
2016;48:617–623.
21. Mills MC, Tropf FC, Brazel DM, van Zuydam N, Vaez A, eQTLGen Consortium, et al.
Identification of 371 genetic variants for age at first sex and birth linked to externalising
behaviour. Nat Hum Behav. 2021;5:1717–1730.
22. Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, et al. Genetic studies of
body mass index yield new insights for obesity biology. Nature. 2015;518:197–206.
23. Justice AE, Winkler TW, Feitosa MF, Graff M, Fisher VA, Young K, et al. Genome-wide
meta-analysis of 241,258 adults accounting for smoking behaviour identifies novel loci for
obesity traits. Nat Commun. 2017;8:14977.
24. Yan X, Wang Z, Schmidt V, Gauert A, Willnow TE, Heinig M, et al. Cadm2 regulates body
weight and energy homeostasis in mice. Mol Metab. 2018;8:180–188.
25. Sanchez-Roige S, Fontanillas P, Jennings MV, Bianchi S, Huang Y, Hatoum A, et al.
Genome-wide association study of problematic opioid prescription use in 132,113
23andMe research participants of European ancestry. Mol Psychiatry; 2021;26:6209-17.
26. Sanchez-Roige S, Fontanillas P, Elson SL, 23andMe Research Team, Pandit A, Schmidt
EM, et al. Genome-wide association study of delay discounting in 23,217 adult research
participants of European ancestry. Nat Neurosci. 2018;21:16–18.
27. Cyders MA, Littlefield AK, Coffey S, Karyadi KA. Examination of a short English version of
the UPPS-P Impulsive Behavior Scale. Addict Behav. 2014;39:1372–1376.
28. Whiteside SP, Lynam DR. The Five Factor Model and impulsivity: using a structural
model of personality to understand impulsivity. Personal Individ Differ. 2001;30:669–689.
29. Patton JH, Stanford MS, Barratt ES. Factor structure of the barratt impulsiveness scale.
Wiley Period Inc. 1995;51:768–774.
30. VanderBroek L, Acker J, Palmer AA, de Wit H, MacKillop J. Interrelationships among
parental family history of substance misuse, delay discounting, and personal substance
use. Psychopharmacology (Berl). 2016;233:39–48.
31. Durand EY, Do CB, Mountain JL, Macpherson JM. Ancestry Composition: A Novel,
Efficient Pipeline for Ancestry Deconvolution. bioRxiv. 2014.
32. Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and
annotation of genetic associations with FUMA. Nat Commun. 2017;8:1826.
33. Leeuw CA de, Mooij JM, Heskes T, Posthuma D. MAGMA: Generalized Gene-Set
Analysis of GWAS Data. PLOS Comput Biol. 2015;11:e1004219.
34. Bulik-Sullivan BK, Loh P-R, Finucane HK, Ripke S, Yang J, Patterson N, et al. LD Score
regression distinguishes confounding from polygenicity in genome-wide association
studies. Nat Genet. 2015;47:291–295.
35. Verma A, Bradford Y, Dudek S, Lucas AM, Verma SS, Pendergrass SA, et al. A
simulation study investigating power estimates in phenome-wide association studies.
BMC Bioinformatics. 2018;19:120.

36. Ehm MG, Aponte JL, Chiano MN, Yerges-Armstrong LM, Johnson T, Barker JN, et al.
Phenome-wide association study using research participants' self-reported data provides
insight into the Th17 and IL-17 pathway. PLOS ONE. 2017;12:e0186405.

37. Lloyd KCK. A knockout mouse resource for the biomedical research community: The
KOMP Repository. Ann N Y Acad Sci. 2011;1245:24–26.

38. Evangelou E, Gao H, Chu C, Ntritsos G, Blakeley P, Butts AR, et al. New alcohol-related
genes suggest shared genetic mechanisms with neuropsychiatric disorders. Nat Hum
Behav. 2019;3:950–961.

39. Ikeda M, Takahashi A, Kamatani Y, Momozawa Y, Saito T, Kondo K, et al. Genome-Wide
Association Study Detected Novel Susceptibility Genes for Schizophrenia and Shared
Trans-Populations/Diseases Genetic Effect. Schizophr Bull. 2019;45:824–834.

40. Howard DM, Adams MJ, Clarke T-K, Hafferty JD, Gibson J, Masoud S, et al. Genome-
wide meta-analysis of depression identifies 102 independent variants and highlights the
importance of the prefrontal brain regions. Nat Neurosci. 2019;22:343–352.

41. Yang H, Liu D, Zhao C, Feng B, Lu W, Yang X, et al. Mendelian randomization integrating
GWAS and eQTL data revealed genes pleiotropically associated with major depressive
disorder. Transl Psychiatry. 2021;11:225.

42. Baselmans BML, Jansen R, Ip HF, van Dongen J, Abdellaoui A, van de Weijer MP, et al.
Multivariate genome-wide analyses of the well-being spectrum. Nat Genet. 2019;51:445–
451.

43. Hill WD, Weiss A, Liewald DC, Davies G, Porteous DJ, Hayward C, et al. Genetic
contributions to two special factors of neuroticism are associated with affluence, higher
intelligence, better health, and longer life. Mol Psychiatry. 2020;25:3034–3052.

44. Teixeira JR, Szeto RA, Carvalho VMA, Muotri AR, Papes F. Transcription factor 4 and its
association with psychiatric disorders. Transl Psychiatry. 2021;11:19.

45. Van Esch H, Colnaghi R, Freson K, Starokadomskyy P, Zankl A, Backx L, et al. Defective
DNA Polymerase α-Primase Leads to X-Linked Intellectual Disability Associated with
Severe Growth Retardation, Microcephaly, and Hypogonadism. Am J Hum Genet.
2019;104:957–967.

46. Christakoudi S, Evangelou E, Riboli E, Tsilidis KK. GWAS of allometric body-shape
indices in UK Biobank identifies loci suggesting associations with morphogenesis,
organogenesis, adrenal cell renewal and cancer. Sci Rep. 2021;11:10688.

47. Pulit SL, Stoneman C, Morris AP, Wood AR, Glastonbury CA, Tyrrell J, et al. Meta-
analysis of genome-wide association studies for body fat distribution in 694 649
individuals of European ancestry. Hum Mol Genet. 2019;28:166–174.

48. Nagel M, Watanabe K, Stringer S, Posthuma D, van der Sluis S. Item-level analyses
reveal genetic heterogeneity in neuroticism. Nat Commun. 2018;9:905.

49. Lo M-T, Hinds DA, Tung JY, Franz C, Fan C-C, Wang Y, et al. Genome-wide analyses for
personality traits identify six genomic loci and show correlations with psychiatric
disorders. Nat Genet. 2017;49:152–156.

50. Thorp JG, Campos AI, Grotzinger AD, Gerring ZF, An J, Ong JS, et al. Symptom-level
modelling unravels the shared genetic architecture of anxiety and depression. Nat Hum
Behav. 2021;5:1432-44.

51. Feitosa MF, Kraja AT, Chasman DI, Sung YJ, Winkler TW, Ntalla I, et al. Novel genetic
associations for blood pressure identified via gene-alcohol interaction in up to 570K
individuals across multiple ancestries. PLOS ONE. 2018;13:e0198166.

52. Ruth KS, Day FR, Tyrrell J, Thompson DJ, Wood AR, Mahajan A, et al. Using human
genetics to understand the disease impacts of testosterone in men and women. Nat Med.
2020;26:252–258.

53.   Rask-Andersen M, Karlsson T, Ek WE, Johansson Å. Genome-wide association study of
694   body fat distribution identifies adiposity loci and sex-specific genetic effects. Nat
695   Commun. 2019;10:339.
696   54.   Hill WD, Davies NM, Ritchie SJ, Skene NG, Bryois J, Bell S, et al. Genome-wide analysis
697   identifies molecular systems and 149 genetic loci associated with income. Nat Commun.
698   2019;10:5741.
699   55.   Zhao B, Luo T, Li T, Li Y, Zhang J, Shan Y, et al. Genome-wide association analysis of
700   19,629 individuals identifies variants influencing regional brain volumes and refines their
701   genetic co-architecture with cognitive and mental health traits. Nat Genet. 2019;51:1637–
702   1644.
703   56.   Gaddis N, Mathur R, Marks J, Zhou L, Quach B, Waldrop A, et al. Multi-trait genome-wide
704   association study of opioid addiction: *OPRM1* and Beyond. medRxiv. 2021.
705   57.   Liu M, Jiang Y, Wedow R, Li Y, Brazel DM, Chen F, et al. Association studies of up to 1.2
706   million individuals yield new insights into the genetic etiology of tobacco and alcohol use.
707   Nat Genet. 2019;51:237–244.
708   58.   Davies G, Lam M, Harris SE, Trampush JW, Luciano M, Hill WD, et al. Study of 300,486
709   individuals identifies 148 independent genetic loci influencing general cognitive function.
710   Nat Commun. 2018;9:2098.
711   59.   ADHD Working Group of the Psychiatric Genomics Consortium, 23andMe Research
712   team, Rovira P, Demontis D, Sánchez-Mora C, Zayats T, et al. Shared genetic
713   background between children and adults with attention deficit/hyperactivity disorder.
714   Neuropsychopharmacology. 2020;45:1617–1626.
715   60.   Karlsson Linnér R, Biroli P, Kong E, Meddens SFW, Wedow R, Fontana MA, et al.
716   Genome-wide association analyses of risk tolerance and risky behaviors in over 1 million
717   individuals identify hundreds of loci and shared genetic influences. Nat Genet.
718   2019;51:245–257.
719   61.   Pritikin JN, Neale MC, Prom-Wormley EC, Clark SL, Verhulst B. GW-SEM 2.0: Efficient,
720   Flexible, and Accessible Multivariate GWAS. Behav Genet. 2021;51:343–357.
721   62.   MacKillop J, Weafer J, C. Gray J, Oshri A, Palmer A, de Wit H. The latent structure of
722   impulsivity: impulsive choice, impulsive action, and impulsive personality traits.
723   Psychopharmacology (Berl). 2016;233:3361–3370.
724   63.   Gray JC, MacKillop J, Weafer J, Hernandez KM, Gao J, Palmer AA, et al. Genetic
725   analysis of impulsive personality traits: Examination of a priori candidates and genome-
726   wide variation. Psychiatry Res. 2018;259:398–404.
727   64.   Leppert B, Millard LAC, Riglin L, Davey Smith G, Thapar A, Tilling K, et al. A cross-
728   disorder PRS-pheWAS of 5 major psychiatric disorders in UK Biobank. PLOS Genet.
729   2020;16:e1008185.
730   65.   Metha JA, Brian ML, Oberrauch S, Barnes SA, Featherby TJ, Bossaerts P, et al.
731   Separating Probability and Reversal Learning in a Novel Probabilistic Reversal Learning
732   Task for Mice. Front Behav Neurosci. 2020;13:270.
733   66.   Mota NR, Araujo-Jnr EV, Paixão-Côrtes VR, Bortolini MC, Bau CHD. Linking dopamine
734   neurotransmission and neurogenesis: the evolutionary history of the NTAD (NCAM1-
735   TTC12-ANKK1-DRD2) gene cluster. Genet Mol Biol. 2012;35:912–918.
736   67.   Greenbaum L, Ravona-Springer R, Livny A, Shelly S, Ganmore I, Alkelai A, et al. The
737   CADM2 gene is associated with processing speed performance – evidence among
738   elderly with type 2. Worl J Biol Psychiatry. 2019;20:577-83.
739   68.   Dalley JW, Everitt BJ, Robbins TW. Impulsivity, Compulsivity, and Top-Down Cognitive
740   Control. Neuron. 2011;69:680–694.
741   69.   Collins RL, Glessner JT, Porcu E, Niestroj L-M, Ulirsch J, Kellaris G, et al. A cross-
742   disorder dosage sensitivity map of the human genome. medRxiv. 2021.

70. van Enkhuizen J, Henry BL, Minassian A, Perry W, Milienne-Petiot M, Higa KK, et al.
Reduced Dopamine Transporter Functioning Induces High-Reward Risk-Preference
Consistent with Bipolar Disorder. Neuropsychopharmacology. 2014;39:3112–3122.

71. Cope ZA, Halberstadt AL, van Enkhuizen J, Flynn AD, Breier M, Swerdlow NR, et al.
Premature responses in the five-choice serial reaction time task reflect rodents' temporal
strategies: evidence from no-light and pharmacological challenges. Psychopharmacology
(Berl). 2016;233:3513–3525.

72. Hsu T-Y, Lee H-C, Lane TJ, Missal M. Temporal Preparation, Impulsivity and Short-Term
Memory in Depression. Front Behav Neurosci. 2019;13:258.

73. Morris J, Bailey MES, Baldassarre D, Cullen B, de Faire U, Ferguson A, et al. Genetic
variation in CADM2 as a link between psychological traits and obesity. Transl Psychiatry.
2019;9:7339.

74. Clarke T-K, Adams M, Davies G, Howard D, Hall L, Padmanabhan S, et al. Genome-wide
association study of alcohol consumption and genetic overlap with other health-related
traits in UK Biobank. Mol Psychiatry. 2017;22:1376-84.

75. Kranzler HR, Zhou H, Kember RL, Smith RV, Justice AC, Damrauer S, et al. Genome-
wide association study of alcohol consumption and use disorder in 274,424 individuals
from multiple populations. Nat Commun. 2019;10:1499.

76. Giza JI, Jung Y, Jeffrey RA, Neugebauer NM, Picciotto MR, Biederer T. The Synaptic
Adhesion Molecule SynCAM 1 Contributes to Cocaine Effects on Synapse Structure and
Psychostimulant Behavior. Neuropsychopharmacology. 2013;38:628–638.

77. Biederer T, Sara Y, Mozhayeva M, Atasoy D, Liu X, Kavalali ET, et al. SynCAM, a
Synaptic Adhesion Molecule That Drives Synapse Assembly. Science. 2002;297:1525–
1531.

78. Fogel AI, Akins MR, Krupp AJ, Stagi M, Stein V, Biederer T. SynCAMs Organize
Synapses through Heterophilic Adhesion. J Neurosci. 2007;27:12516–12530.

79. Robbins EM, Krupp AJ, Perez de Arce K, Ghosh AK, Fogel AI, Boucard A, et al. SynCAM
1 Adhesion Dynamically Regulates Synapse Number and Impacts Plasticity and
Learning. Neuron. 2010;68:894–906.

80. Yamada A, Inoue E, Deguchi-Tawarada M, Matsui C, Togawa A, Nakatani T, et al. Necl-
2/CADM1 interacts with ErbB4 and regulates its activity in GABAergic neurons. Mol Cell
Neurosci. 2013;56:234–243.

81. Niederkofler V, Baeriswyl T, Ott R, Stoeckli ET. Nectin-like molecules/SynCAMs are
required for post-crossing commissural axon guidance. Development. 2010;137:427–435.

82. Maurel P, Einheber S, Galinska J, Thaker P, Lam I, Rubin MB, et al. Nectin-like proteins
mediate axon–Schwann cell interactions along the internode and are essential for
myelination. J Cell Biol. 2007;178:861–874.

83. Spiegel I, Adamsky K, Eshed Y, Milo R, Sabanay H, Sarig-Nadir O, et al. A central role
for Necl4 (SynCAM4) in Schwann cell–axon interaction and myelination. Nat Neurosci.
2007;10:861–869.

84. Park J, Liu B, Chen T, Li H, Hu X, Gao J, et al. Disruption of Nectin-Like 1 Cell Adhesion
Molecule Leads to Delayed Axonal Myelination in the CNS. J Neurosci. 2008;28:12815–
12819.

85. Enticott PG, Ogloff JRP, Bradshaw JL. Associations between laboratory measures of
executive inhibitory control and self-reported impulsivity. Personal Individ Differences.
2006;41:285-94.

86. Lane SD, Cherek DR, Rhoades HM, Pietras CJ, Tcheremissine OV. Relationships
Among Laboratory and Psychometric Measures of Impulsivity: Implications in Substance
Abuse and Dependence. Addict Dis & Their Treatment. 2003;2:33-40.

87.   Sittig LJ, Carbonetto P, Engel KA, Krauss KS, Barrios-Camacho CM, Palmer AA. Genetic
793   Background Limits Generalizability of Genotype-Phenotype Relationships. Neuron.
794   2016;91:1253–1259.
795
796

Figure Legends

Figure 1. Porcupine plot displaying 21 genome-wide significant hits for all impulsivity facets and Drug Experimentation. *CADM2* was consistent across 3/8 impulsivity facets [Sensation Seeking (UPPS-P), Motor and Nonplanning impulsivity (BIS-11)] at a genome-wide association level, and with 3 more impulsivity facets [Attentional (BIS-11), Negative Urgency and Premeditation (UPPS-P]), at a gene-based level (**Table S8**). *CADM2* was also associated with risky behavior, such as Drug Experimentation.

Figure 2. Genetic correlations (r_g) between UPPS-P, BIS, and Drug Experimentation, and other substance use, psychiatric, personality, cognitive, metabolic, health, pain, longevity and sleep traits (see **Table S9** for full results). All values survive 5% FDR correction.

Figure 3. FDR-significant associations from *CADM2* single-SNP PheWAS in individuals of European ancestry (**A**) and Latin American ancestry (**B**). Results for the SNP with the highest number of significant PheWAS associations (rs993137) are presented; results for additional SNPs are included in Supplementary Tables 15-17. No FDR-significant findings were detected in individuals of African American ancestry. The size of the dots represents the magnitude of the effect size for each trait. The effect sizes ranged from -0.14 to 0.13 in the European cohort and from -0.08 to 0.16 in the Latin American cohort. Dotted line denotes Bonferroni significance ($p<3.79E-05$).

Figure 4. MouseWAS examined the complete and partial loss of *Cadm2* function and behavioral consequences in the Probability Breakpoint Ratio Task (PBRT, **A**), Probabilistic Reversal Learning Task (PRLT, **B**), Iowa gambling task (IGT, **C-D**), Behavioral Pattern Monitor task (BPM, **E-F**), Prepulse Inhibition (PPI, **G-H**), 5-Choice Serial Reaction Time Task (5CSRTT) performance (**I-M**), Locomotor (LCM) activity (**N**), Elevated Plus Maze (EPM, **O**), Light-Dark Box (LDB, **P**), and Drinking in the Dark (DID, **Q**); longitudinal body weight changes (**R**), and dendrite morphology (DSM) in the nucleus accumbens (**S**). Western blot (**WB**) analysis (**T**) of Cadm2 protein in whole brain, frontal cortex and striatum. WT, wildtype, HET, heterozygote, HOM, homozygote. Sample size by cohort: cohort 1 (WT=25, HET=30, HOM=3), cohort 2 (WT=13; HET=14, HOM=12), cohort 3 (WT=22; HET=44, HOM=12), cohort 4 (WT=29, HT=54, HOM=17), cohort 5 (WT=3, HET=3, HOM=3). In cohort 1, the sample size of the HOM mice deviated from the expected Mendelian frequency for unknown reasons; these animals were excluded from the analyses. Males are represented in circles, females in triangles. * $p<0.05$, ** $p<0.01$, *** $p<0.001$.