Logistic regression with machine learning sheds light on the problematic sexual behavior

phenotype

Shui Jiang (MSc)[1], Keanna Wallace (BSc)[1,2], Esther Yang (MSc)[1,2,3], Leslie Roper (MSc)[1,2],

Garima Aryal (MSc)[1,2], Dawon Lee (MD)[1,2,4], Rohit J Lodhi (MD, PhD, MRCPsych,

FRANZCP)[2,5], Rick Isenberg (MD)[6], Bradley Green (PhD)[7], David Wishart (PhD)[8], Katherine J

Aitchison (BM BCh, PhD, FRCPsych)[1,2,9,10*]

[1]Department of Medical Genetics, Faculty of Medicine and Dentistry, College of Health Sciences,

University of Alberta, Canada

[2]Department of Psychiatry, Faculty of Medicine and Dentistry, College of Health Sciences, University of

Alberta, Canada

[3]Department of Emergency Medicine, Faculty of Medicine and Dentistry, College of Health Sciences,

University of Alberta, Canada

[4]Department of Medicine, Faculty of Medicine and Dentistry, College of Health Sciences, University of

Alberta, Canada

[5]Department of Psychiatry, University of Western Ontario, Canada

[6]Psychological Counselling Services, Ltd., Scottsdale, USA

[7]Department of Psychology and Counseling, University of Texas at Tyler, USA

[8]Department of Biological Sciences, Faculty of Science, College of Natural and Applied Sciences,

University of Alberta, Canada

[9]Neuroscience and Mental Health Institute, University of Alberta, Canada

[10]Division of Clinical Sciences, Psychiatry Section, Northern Ontario School of Medicine

*Corresponding author:

Email: kaitchis@ualberta.ca (KJA) Phone number: 780-492-4018

Sources of support: The study was funded by an Alberta Centennial Addiction and Mental

Health Research Chair and transitional funding (to KJA), and by the American Foundation for

Addiction Research (to KJA). Several scholarships, including a University of Alberta Dean's

Doctoral Student Award 2021-2022, a University of Alberta Medical Science Graduate Program

Scholarship 2020, a University of Alberta Doctoral Recruitment Scholarship 2017 and an

Alberta Graduate Excellence Scholarship 2020 provided additional funding for the first author

(SJ). A Fulbright-Canada-Palix Foundation grant to Dr. Patrick Carnes facilitated his

contributions to study design.

Conflict of interest: Dr. Carnes has a conflict of interest with the American Foundation for

Addiction Research, and for this reason, he is not an author on the paper. Dr. Aitchison has

received two research grants from Janssen Inc., Canada (fellowship grants for trainees). The

analysis and data interpretation were conducted independently.

Abstract word count: 250

Manuscript word count: 3124

Number of references: 40

Abstract

Objectives: There has been a longstanding debate about whether the mechanisms involved in

problematic sexual behavior (PSB) are similar to those observed in addictive disorders, or related

to impulse control or to compulsivity. The aim of this report was to contribute to this debate by

investigating the association between PSB, addictive disorders (internet addiction, compulsive

buying), measures associated with the construct known as reward deficiency (RDS), and

obsessive-compulsive disorder (OCD) in young adults in post-secondary education.

Methods: A Canadian university Office of the Registrar invited 68,846 eligible students and

postdoctoral fellows. Out of 4710 expressing interest in participating, 3359 completed on-line

questionnaires, and 1801 completed the Mini-International Neuropsychiatric Interview (MINI).

PSB was measured by combining those screening positive (≥ 6) on the Sexual Addiction

Screening Test-Revised (SAST-R) Core with those self-reporting PSB. Current mental health

condition(s) and childhood trauma were measured by self-report. OCD was assessed by a

combination of self-report and MINI data.

Results: 12.18% (407/3341) of participants screened positive on the SAST-R Core. On logistic

regression, OCD, attention deficit, internet addiction, a family history of PSB, childhood trauma,

compulsive buying and male gender were associated with PSB. On multiple correspondence

analysis (MCA), OCD appeared to cluster separately from the other measures, and the pattern of

data differed by gender.

Conclusions: Factors that have previously been associated with RDS and OCD are both

associated with increased odds of PSB. The factors associated with RDS appear to contribute to a

separate data cluster from OCD, and to lie closer to PSB.

INTRODUCTION

A theory of hypersexuality with dependence was first proposed by Orford in 1978.[1] The term sex addiction was used by Carnes in 1983, 1990, and 1991[2] and by Goodman in 1998.[3] Continuation of a sexual behavior despite adverse consequences and/or distress caused or worsened by the sexual behavior has been a consistent theme in the literature,[1,3,4] despite differences of opinion as to whether the condition should be classified as an addiction or as an impulse control disorder. Diagnostic criteria proposed by Carnes in 2005[4] led to sex addiction being included in the Diagnostic and Statistical Manual of Mental Disorders (DSM)-III-R (302.87). However, despite draft proposed criteria for the inclusion of sex addiction and internet addiction in DSM-5,[5] the only behavioral addiction currently included in the main body of DSM-5 (American Psychiatric Association 2013) is gambling disorder (312.31), which was first included in DSM as an impulse control disorder.[6] In the International Statistical Classification of Diseases (ICD), compulsive sexual behavior disorder was proposed for inclusion as an impulse control disorder in 2014,[7] which occurred in 2018.[8] However, the scientific discussion about whether or not compulsive sexual behavior disorder could constitute a manifestation of a behavioral addiction was acknowledged,[9] and it was predicted (as in the case of problematic gambling),[6] that understanding would evolve as research elucidated the phenomenology and neurobiological underpinnings of the condition.

In this article, the term problematic sexual behavior (PSB) is used because it focuses on the behavior rather than on the potentially heterogeneous etiology and pathology.[10] PSB is characterized by repetitive sexual behaviors and it is associated with uncontrolled sexual urges/impulses and distress,[11] and social and functional impairments.[8,11] It has been estimated that the prevalence of PSB is 3-6% in adults,[4,12-14] with higher frequencies in specific populations,

89   including those in post-secondary education. Reid (2011) reported that 19% of college men met

90   criteria for hypersexuality,[15] while Giordano and Cecil (2014) found 11.1% of men and women

91   in an undergraduate sample met these criteria.[16] Cashwell et al. (2015) reported that out of 379

92   undergraduates, 21.2% of men and 6.7% of women screened positive, i.e., scored in a range

93   indicating that they should be offered further assessment for sex addiction.[17]

94   Various vulnerability/risk factors have been associated with PSB, including those

previously associated with the construct known as reward deficiency syndrome (RDS)

(substance[18] and non-substance use disorders,[19] ADHD,[20] and personality disorder),[21] as well as

other psychiatric disorders,[22] and childhood trauma.[23] In terms of psychiatric disorders, owing to

the classification of PSB within ICD-10 as an impulse control disorder, we specifically included

obsessive-compulsive disorder. Mick and Hollander (2006) hypothesized that impulsivity

initiated the early stage of PSB, with compulsivity being involved in repetitive behaviors and

hence in the maintenance of PSB.[24] The magnitude of the contribution of the various different

postulated mechanisms, and whether PSB encompasses various syndromes with different

etiologies is not at present known. Moreover, systematic data on the prevalence of PSB and

associated sociodemographic factors across diverse populations, including non-treatment seekers

remain to be provided.[8]

106   In light of the above, this paper aims to further elucidate the prevalence,

107   sociodemographic features, phenomenology, and neurobiological underpinnings of the construct

108   known as PSB in a diverse sample of adults in post-secondary education. We hypothesized that

109   in this sample, firstly, screening positive for factors previously associated with RDS (such as

110   internet addiction, compulsive buying, nicotine dependence, and pathological gambling) and

111   secondly, having OCD and childhood trauma would be associated with increased odds of PSB.

METHODS

The study inclusion criteria were as follows: at least 18 years of age, undergraduates, graduates,

postdoctoral fellows, and recently convocated students registered in at least one course in the

preceding year, except for those who had completely withdrawn after registration without any

reasons to not return in the next academic term or not be on campus (such as suspension), and

being able to answer in English. A Canadian university Office of the Registrar invited 68,846

eligible students and postdoctoral fellows by email. Students interested in participating emailed

the study team. These students were then sent an email invitation to review the participant

information and complete consent on-line, followed by the screening measures (hosted by the

Qualtrics platform), with a subsequent email inviting them to complete the Mini-International

Neuropsychiatric Interview (MINI) version 7.0.2 on-line.

Measures

The Sexual Addiction Screening Test-Revised (SAST-R)[25] is a 45-item screening tool

comprising several subscales designed to detect potentially problematic sexual behavior. Each

question is answered in a binary manner (yes/no=0/1). Sexual activities deemed to be

problematic by participants in the past 30 days were also separately assessed by asking the

following question: "Within the last 30 days, how often have you participated in activities of a

sexual nature that you would regard as problematic?"

Measures used to screen for internet addiction, compulsive buying, personality disorders,

ADHD, nicotine dependence, and pathological gambling were: the Internet Addiction Test

(IAT),[26] the Richmond Compulsive Buying Scale (RCBS),[27] the 8-item Standardized Assessment

of Personality–Abbreviated Scale as a Self-Administered Screening Test (SA-SAPAS),[28] the full

Adult ADHD Self-Report Scale (ASRS-v1.1),[29] the Fagerström Test for Nicotine Dependence

(FTND),[30] and the DSM-V Pathological Gambling Diagnostic Form (PGDF),[31] respectively.

Gender, sexual orientation, a family history of domestic violence or of sex addiction, and having

received a diagnosis of OCD, or any other mental health condition(s), childhood physical,

emotional/verbal, or sexual trauma were additionally collected by self-report. The MINI was

used to output psychiatric disorders by DSM-5 criteria.

Data analysis

Demographic data and logistic regression

Data were analyzed by STATA (Stata/SE 16) and R (Version 3.6.3) after dropping the missing

data. The PSB variable was created by combining those screening positive on the SAST-R Core

with those endorsing monthly self-reported PSB, after subtracting the two SAST-R Core

questions referred to below. Previous or current physical/verbal, emotional/sexual childhood

trauma and a family history of domestic violence (11.77% 387/3288) were regrouped as total

childhood trauma due to the relatively low rates of endorsement for the former group (physical

trauma: 2.98%, 99/3322; emotional trauma: 6.87%, 232/3322; and sexual trauma: 3.31%,

110/3212). The SAST-R Core question 1 (which enquires about a history of sexual abuse in

childhood or adolescence, SASTC1) was combined with total childhood trauma. Self-reported a

family history of sex addiction ("yes"/ "no") data were combined with SAST-R Core question 2

(which enquires about parental trouble with sexual behavior, SASTC2) to create a new family

history of PSB variable. IAT was regrouped as "yes" (N=1174, combining mild, moderate, and

severe groups) and "no" (N=2160, the "normal" group).

Differences in data distributions between those with and without PSB were assessed by

Pearson's χ^2 test for categorical variables and by non-parametric testing (Mann-Whitney U test)

for linear non-normally distributed variables. For correlational analyses, tetrachoric (for binary

variables) and Kendall's τ (for categorical variables) correlations were utilized. SA-SAPAS,

OCD, ASRS, IAT, childhood trauma, a family history of PSB and gender were used as

covariates in the regression analyses. All reported *P* values are unadjusted for multiple testing.

Multiple correspondence analysis (MCA) was used for data clustering.

Machine learning approaches

The "Haven" and "dplyr" packages were used for data input. The "forcats" and "creditmodel"

packages were used for recoding and splitting the dataset (into 80% training and 20% testing),

respectively. The Synthetic Minority Oversampling Technique (SMOTE) from the "Caret"

package was used to address data imbalance in the training set.[32] Leave-one-out cross-validation

(LOOCV) in the "Caret" package was used for resampling.[33] Logistic regression (two-class

summary) in the "caret" package was used for regression. Accuracy, area under the receiver

operating characteristic (ROC) curve, sensitivity, specificity, and F score were used to evaluate

the models. The "pROC" package was used for the area under the ROC curve.

Multiple correspondence analysis (MCA)

Multiple correspondence analysis was conducted on OCD, RCBS, SA-SAPAS, ASRS, IAT, and

childhood trauma, stratifying by gender (women N=2084 and men N=1012). As a family history

of PSB was correlated with the combined childhood trauma variable (tetrachoric ϱ=0.69,

P<0.001, N=3019, Figure 1, Supplemental Digital Content, Table 3), these variables were

combined as a supplementary childhood trauma-related variable. The "FactoMineR,"

"factoextra," and "GDAtools" packages in R were used for analysis and data visualization.

RESULTS

Out of 4710 interested in participating, 237 withdrew from the study, 3359 completed online

questionnaires using the Qualtrics platform, and 1801 completed the MINI (Supplemental Digital

Content, Figure 1).

Demographic data

The mean score on the SAST-R Core Scale among all participants was 2.23 (SD=2.73), with

12.18% (407/3341) of participants scoring at least six (the threshold for screening positive in the

general population.[25] Of the 20 items composing the SAST-R Core questions, item 11 (Do you

hide some of your sexual behavior from others?) was the most endorsed (1234/3341,

endorsement rate: 36.78%), while item 9 ("Are any of your sexual activities against law?") was

the least endorsed (35/3342; endorsement rate: 1.05%) (Supplemental Digital Content, Table 1).

The frequency of participants included in the PSB variable was 16.53% (532/3219). There was a

significant difference in the distribution of the following variables by those with and without

PSB: gender, sexual orientation, OCD, RCBS, a family history of PSB, SA-SAPAS, current

mental health condition(s), childhood trauma, ASRS, and IAT (Table 1, and Supplemental

Digital Content, Table 2).

Correlational analyses in the whole dataset after dropping the missing values showed that

PSB was correlated with non-heterosexual sexual orientation (baseline: heterosexual), current

mental health condition(s), childhood trauma, RCBS, OCD, SA-SAPAS, a family history of

PSB, ASRS, IAT and gender (baseline: women) (Figure 1, Supplemental Digital Content, Table

3). Interestingly, a family history of PSB was shown to be strongly correlated with childhood

trauma in our dataset (tetrachoric $\varrho=0.69$, $P<0.001$, N=3019), and both were correlated with

PSB. For example, of those responding "yes" to a family history of sex addiction, 32.65%

(16/49) screened positive on the SAST-R Core. Our data also showed that: (A) PSB was

correlated with unadjusted SAST-R Core (tetrachoric $\varrho=0.97$, $P<0.001$, N=3219) and self-

reported monthly PSB (tetrachoric $\varrho=0.70$, $P<0.001$, N=3271), (B) self-reported OCD was

correlated with OCD MINI diagnosis (tetrachoric $\varrho=0.47$, $P<0.001$, N=1785) (Supplemental

Digital Content, Table 4).

Regression models

Logistic regression

Outliers were removed from the logistic regression (N=47, with 8 categorized as "yes" to having

PSB, Supplemental Digital Content, Figure 2) if their Pearson standardized residual had an

absolute value of more than 2,[34] or Hosmer-Lemeshow $\Delta\hat{\beta}$ more than four.[35] After outlier

removal as above, no further outliers needed to be removed on the basis of leverage values (a

measure of the distance between independent variable observations). The following variables

were associated with PSB (pseudo $R^2=0.16$): OCD ($P=0.015$), ASRS ($P<0.001$), IAT ($P<0.001$),

a family history of PSB ($P<0.001$), SA-SAPAS ($P<0.001$), childhood trauma ($P<0.001$), RCBS

($P<0.001$) and gender (men) ($P<0.001$) (Figure 2).

Machine Learning (ML) logistic regression

Our ML logistic regression showed that RCBS ($P=0.0030$), SA-SAPAS ($P<0.001$), OCD

($P<0.001$), a family history of PSB ($P<0.001$), ASRS ($P<0.001$), childhood trauma ($P<0.001$),

IAT ($P<0.001$), and gender (men) ($P<0.001$) were all associated with increased odds of PSB

(Figure 2B). A validation of the existing model was also undertaken. The validation model was

trained with SAST-R Core (subtracting SASTC1 and SASTC2, N=3141) and tested with self-

reported PSB (N=3145, Figure 2C). Sensitivity analyses were performed using unadjusted

SAST-R Core and family history of sexual addiction by logistic regression and ML logistic

regression (Supplemental Digital Content, Figure 3-4).

Multiple correspondence analysis (MCA)

The MCA results showed that in both women and men, the first two dimensions accounted for the most explained variance (Supplemental Digital Content, Figure 5). PSB was strongly represented in both men and women in dimensions 1, 2, 4 and 5, although showing slightly different data clustering patterns by gender in these dimensions (Figure 3-4). In dimension 1, PSB clustered with all variables in women and men. In dimension 2 (Figure 3 and Supplemental Digital Content, Figure 5), PSB clustered with IAT and ASRS in men, and with RCBS, IAT and ASRS in women. In women and men, dimensions 4 and 5 illustrate that PSB was clustered relatively closely to the composite childhood trauma variable. (Figure 4, Supplemental Digital Content, Figure 5). The quality of representation and contributions of each variable are shown in Supplemental Digital Content, Table 6.

DISCUSSION

The frequency of problematic sexual behavior, as defined by screening positive on the SAST-R Core, was 8.49% in the women and 19.7% in the men in this post-secondary education sample, consistent with prior reports in this population.[15-17] Factors consistently associated with increased odds of PSB were gender, mental health conditions (ADHD, personality disorder, and OCD), other problematic behaviors such as internet addiction (IAT) and compulsive buying (RCBS), childhood trauma (including a family history of domestic violence), and a family history of PSB. Our results also showed that the self-reported data were correlated with the screening tool and/or MINI data (Supplemental Digital Content, Table 3).

Model validation (training with SAST-R Core subtracting the first two questions, and testing with self-reported PSB) resulted in very similar model fitting statistics (Figures 2B and 2C). A sensitivity analysis (Supplemental Digital Content, Figure 3-4) was undertaken using the

original variables, namely SAST-R Core (without subtracting the first two question), childhood

trauma (without adding the first question of SAST-R Core or a family history of domestic

violence) and a family history of sex addiction (without incorporating SASTC2). The logistic

regression model using these variables (pseudo R^2=0.13) revealed that RCBS (P=0.033), ASRS

(P<0.001), IAT (P<0.001), OCD (P<0.001), childhood trauma (P<0.001), and gender (baseline:

women, P<0.001) remained significant. A family history of sex addiction was the only variable

that became non-significant (P=0.17), probably due to the lower frequency of endorsement of

this variable (N=49) compared to that of SASTC2. The combined variable (a family history of

sex addiction with SASTC2, 6.55% endorsement: 214/3268), was, however, significant in all

models. Using the new PSB variable, including childhood trauma and a family history of PSB

but excluding sexual orientation as well as current mental health condition(s) improved the

model fit slightly in the logistic regression models (e.g., sensitivity: unadjusted 77.20% vs

adjusted 79.70%, F score: unadjusted 0.42 vs. adjusted 0.45). In general, the ML logistic

regression models showed a higher sensitivity and F score compared to conventional logistic

regression.

The variables previously associated with the RDS construct, such as SA-SAPAS, ASRS,

and IAT, were associated with increased odds of PSB. In the final logistic (Figure 2) and ML

logistic regressions (Figure 2), RCBS was associated with increased odds of PSB. Of note, there

were gender differences regarding RCBS: RCBS was negatively correlated with men (Figure 1,

baseline: women), and PSB displayed a weaker association with RCBS in men on MCA (Figure

3).

In addition, interestingly, the distribution of PSB differed significantly between OCD-

positive and -negative groups (P<0.001, Table 1), and PSB was correlated with OCD (ϱ=0.27,

P<0.001, Figure 1, Supplemental Digital Content, Table 3). In our final regression models and

sensitivity analyses (by both conventional and ML logistic regression), OCD was associated with

increased odds of PSB (OR 1.79, *P*=0.015, and OR 1.85, *P*=0.013 for conventional and ML

logistic regression respectively; sensitivity models: OR 2.28, *P*<0.001, and OR 1.73, *P*=0.033,

respectively).

Limitations

Limitations of this study include the following. Firstly, as respondents to the email from

the registrar knew the subject area of the study, there is a potential selection bias. However, of

note, the frequency of PSB in our dataset was comparable to that seen in other studies of adults

in post-secondary education as mentioned above.[12,17] Subsample sizes were another limiting

factor. Specifically, as the frequency of endorsement of OCD was relatively low (120/3206,

3.74%), although it appears on the MCA that PSB clusters more closely to the measures of

reward deficiency than to OCD, this should not be overinterpreted. For sexual orientation and

gender, the smaller groups were combined in the former case and dropped in the latter. For the

former, bisexual (366/3371, 10.86%), homosexual (129/3371, 3.83%), and other (91/3371,

2.70%) were regrouped as "non-heterosexual sexual orientation." Due to the small numbers in

the gender minority groups (trans and other: 47/3375, 1.39%), only men and women were

included in the logistic regression analysis. Another potential limitation is that we dropped

missing values (rather than using imputation). However, if the proportion of missing data is

relatively small (5% or less), this should not be problematic.[36] We also dropped two measures,

the FTND (0.45%, 15/3332) and PGDF (2.28%, 76/3332) due to very low endorsement rates

(which were lower than in another university student sample).[37] Although various methods may

be used to impute data, some such methods (e.g., monotone imputation, or chained equations)

may introduce bias, and perfect (inaccurate) predictions, as the methods themselves involve the

fitting of a regression model for a categorical outcome.[38] In our sample, the feature with the most

missing data in the SAST-R Core scale was question 14, which is about feeling depressed after

having sex, with only 16 out of 3341 (0.48%) endorsed (and out of those 16 participants, only 1

scored as SAST-R Core positive). Another limitation is that logistic regression is sensitive to

high correlations between predictor variables. Although several variables were correlated with

each other in our dataset, the correlation coefficients were less than 0.80,[39] and the mean variance

inflation factor (VIF) and the condition number were at acceptable levels (VIF<10, condition

number<10) in our final models,[40] indicating that multi-collinearity was not a significant

concern. A power calculation using STATA using the parameters for OCD (the weakest

predictor, P1=0.16, P2=0.24, pseudo R^2=0.16) that for an α of 0.05, a sample size of only 264

would give a power of 0.8. Finally, the proportion of the variance (pseudo R^2=0.16, Figure 2)

explained by our analysis was relatively low. Of note, the variables that we included in our

models were driven by the prior literature. Future research could include feature selection

techniques, alternative machine learning models, and consideration of genetic and other

biological variables. Validation through additional independent datasets with overlapping

variables would also advantageous.

CONCLUSIONS

In summary, we have identified associations between problematic sexual behavior in adults in

post-secondary education and the following: compulsive buying, personality disorder, OCD,

ADHD, childhood trauma, a family history of PSB, internet addiction, and male gender. Our

results not only suggest that measures associated with reward deficiency syndrome are associated

with PSB, but also that OCD is a vulnerability factor. Childhood trauma and a family history of

domestic violence also showed associations with PSB, implying that emotional regulation and

attachment style may be involved in the development of PSB. Our data are consistent with

different subgroups within PSB with slightly somewhat contrasting etiological mechanisms by

gender.

Acknowledgments

322   We would like to acknowledge the contribution of Dr. Patrick Carnes to the study design, and

323   volunteer input of Penny Carnes, Grace Li, Hana Graham, and Chanelle Martens to data

324   collection. SJ would like to thank Afia Anjum from DW's lab for her assistance with R. There

325   was no pharmaceutical or industry support for the work reported herein.

REFERENCES

1. Orford J. Hypersexuality: implications for a theory of dependence. *Br J Addict Alcohol Other Drugs*. Sep 1978;73(3):299-10. doi:http://dx.doi.org/10.1111/j.1360-0443.1978.tb00157.x

2. Carnes P. *Don't call it love : recovery from sexual addiction*. Bantam Books; 1991:439.

3. Goodman A. Sexual addiction: terminology and theory. *Psychiatric Times* 1998;15(7):22-26.

4. Carnes P. *Sexual addiction*. 8 ed. vol I. Lippincott, Williams & Wilkins: A Wolters Kluwer Company; 2005.

5. Carnes PJ, Hopkins TA, Green BA. Clinical relevance of the proposed sexual addiction diagnostic criteria: relation to the Sexual Addiction Screening Test-Revised. *Journal of addiction medicine*. Nov-Dec 2014;8(6):450-61. doi:http://dx.doi.org/10.1097/ADM.0000000000000080

6. Rosenthal RJ. Inclusion of pathological gambling in DSM-III, its classification as a disorder of impulse control, and the role of Robert Custer. *International Gambling Studies*. 2020;20(1):151-170. doi:https://doi.org/10.1080/14459795.2019.1638432

7. Rehm J, Probst C, Kraus L, Lev-Ran S. *The addiction concept revisited*. Reframing addiction: Policies, processes and pressures. The ALICE RAP project; 2014:103-117.

8. Kraus SW, Krueger RB, Briken P, et al. Compulsive sexual behaviour disorder in the ICD-11. *World Psychiatry*. Feb 2018;17(1):109-110. doi:https://doi.org/10.1002/wps.20499

9. Griffiths MD. Compulsive sexual behaviour as a behavioural addiction: The impact of the internet and other issues. *Addiction*. 2016;111(12):2107-2108. doi:http://dx.doi.org/10.1111/add.13315

10. Joannides P. The challenging landscape of problematic sexual behaviors, including "sexual addiction" and "hypersexuality.*" New directions in sex therapy: Innovations and alternatives*. 2012:69-83. doi:https://doi.org/10.4324/9780203122556

11. Dickenson JA, Gleason N, Coleman E, Miner MH. Prevalence of Distress Associated With Difficulty Controlling Sexual Urges, Feelings, and Behaviors in the United States. *JAMA Netw Open*. Nov 2 *2018*;*1*(*7*):e184468. doi:https://doi.org/10.1001/jamanetworkopen.2018.4468

12. Carnes PJ, Green BA, Merlo LJ, Polles A, Carnes S, Gold MS. PATHOS: a brief screening application for assessing sexual addiction. *J Addict Med*. Mar *2012*;*6*(*1*):*29-34*. doi:https://doi.org/10.1097/ADM.0b013e3182251a28

13. Carnes PJ. *Facing the Shadow: Starting Sexual and Relationship Recovery: a Gentle Path to Beginning Recovery from Sex Addiction*. Gentle Path Press; 2015.

14. Klein V, Rettenberger M, Briken P. Self-reported indicators of hypersexuality and its correlates in a female online sample. *J Sex Med*. Aug 2014;11(8):1974-81. doi:https://doi.org/10.1111/jsm.12602

15. Reid RC, Li DS, Gilliland R, Stein JA, Fong T. Reliability, validity, and psychometric development of the pornography consumption inventory in a sample of hypersexual men. *J Sex Marital Ther*. 2011;37(5):359-85. doi:https://doi.org/10.1080/0092623x.2011.607047

16. Giordano AL, Cecil AL. Religious coping, spirituality, and hypersexual behavior among college students. *Sexual Addiction & Compulsivity*. 2014;21(3):225-239. doi:https://doi.org/10.1080/10720162.2014.936542

17. Cashwell C, L. Giordano A, Lewis T, A. Wachtel K, L. Bartley J. Using the PATHOS Questionnaire for Screening Sexual Addiction Among College Students: A Preliminary

Exploration. *Sexual Addiction & Compulsivity*. 2015; 22:2, 154-166.
doi:https://doi.org/10.1080/10720162.2015.1037481
18.   Deneke E, Knepper C, Green BA, Carnes PJ. Comparative study of three levels of care in
a substance use disorder inpatient facility on risk for sexual addiction. *Sexual Addiction &*
Compulsivity. 2015;22(2):109-125. doi:https://doi.org/10.1080/10720162.2014.979382
19.   Carnes PJ. The Sexual Addiction Screening Process. *Clinical Management of Sex*
Addiction. 2019:21-39.
20.   Blankenship R, Laaser M. Sexual Addiction and ADHD: Is There A Connection? *Sexual*
Addiction & Compulsivity. 2004;11(1-2):7-20. doi:https://doi.org/10.1080/10720160490458184
21.   Ballester-Arnal R, Castro-Calvo J, Giménez-García C, Gil-Juliá B, Gil-Llario M.
Psychiatric comorbidity in compulsive sexual behavior disorder (CSBD). *Addictive behaviors*.
2020;107:106384. doi:https://doi.org/10.1016/j.addbeh.2020.106384
22.   Grant JE, Lust K, Chamberlain SR. Body dysmorphic disorder and its relationship to
sexuality, impulsivity, and addiction. *Psychiatry Research*. 2019;273:260-265.
doi:https://doi.org/10.1016/j.psychres.2019.01.036
23.   London S, Quinn K, Scheidell JD, Frueh BC, Khan MR. Adverse Experiences in
Childhood and Sexually Transmitted Infection Risk From Adolescence Into Adulthood. *Sexually*
transmitted diseases. 2017;44(9):524-532. doi:https://doi.org/10.1097/OLQ.0000000000000640
24.   Mick TM, Hollander E. Impulsive-compulsive sexual behavior. *CNS Spectr*.
2006;11(12):944-55. doi:https://doi.org/10.1017/s1092852900015133
25.   Carnes P, Green B, Carnes S. The same yet different: Refocusing the Sexual Addiction
Screening Test (SAST) to reflect orientation and gender. *Sexual Addiction & Compulsivity*.
2010;17(1):7-30. doi:https://doi.org/10.1080/10720161003604087
26.   Young KS. Internet addiction: The emergence of a new clinical disorder.
Cyberpsychology & behavior. 1998;1(3):237-244.
27.   Ridgway NM, Kukar-Kinney M, Monroe KB. An expanded conceptualization and a new
measure of compulsive buying. *Journal of consumer Research*. 2008;35(4):622-639. doi:
https://doi.org/10.1086/591108
28.   Merlhiot G, Mondillon L, Vermeulen N, Basu A, Mermillod M. Adaptation and
Validation of the Standardized Assessment of Personality–Abbreviated Scale as a Self-
Administered Screening Test (SA-SAPAS). *Journal of Psychology and Psychotherapy*.
2014;4:6. doi:http://dx.doi.org/10.4172/2161-0487.1000164
29.   Kessler RC, Adler L, Ames M, *et al*. The World Health Organization Adult ADHD Self-
Report Scale (ASRS): a short screening scale for use in the general population. *Psychol Med*.
Feb 2005;35(2):245-56. doi:https://doi.org/10.1017/s0033291704002892
30.   Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO. The Fagerström test for
nicotine dependence: a revision of the Fagerstrom Tolerance Questionnaire. *British journal of*
addiction. 1991;86(9):1119-1127.
31.   Rennert L, Denis C, Peer K, Lynch KG, Gelernter J, Kranzler HR. DSM-5 gambling
disorder: prevalence and characteristics in a substance use disorder sample. *Experimental and*
clinical psychopharmacology. 2014;22(1):50-56. doi:https://doi.org/10.1037/a0034518
32.   Benedan L, Monti GS. Predicting the Risk of Gambling Activities in Adolescence: A
Case Study. Springer; 2019:47-57. doi:https://dx.doi.org/10.1007/978-3-030-51222-4_5
33.   Nwanganga F, Chapple M. *Practical Machine Learning in R*. John Wiley & Sons; 2020.

34.   Chen CY, Yang HCP, Chen CW, Chen TH. Diagnosing and revising logistic regression
models: effect on internal solitary wave propagation. *Engineering Computations*.
2008;25(2):121-139. doi:https://dx.doi.org/10.1108/02644400810855940
35.   Hosmer Jr DW, Lemeshow S, Sturdivant RX. *Applied logistic regression*. vol 398. John
Wiley & Sons; 2013.
36.   Momeni A, Pincus M, Libien J. Imputation and Missing Data. *Introduction to Statistical*
Methods in Pathology. Springer International Publishing; 2018:185-200.
37.   Öncel SY, Gebizlioğlu ÖL, Alioğlu FA. Risk factors for smoking behavior among
university students. *Turkish Journal of Medical Sciences*. 2011;41(6):1071-1080.
doi:https://dx.doi.org/10.3906/sag-1009-1122
38.   White IR, Daniel R, Royston P. Avoiding bias due to perfect prediction in multiple
imputation of incomplete categorical variables. *Comput Stat Data Anal*. 2010;54(10):2267-2275.
doi:https://doi.org/10.1016/j.csda.2010.04.005
39.   Shrestha N. Detecting Multicollinearity in Regression Analysis. *American Journal of*
Applied Mathematics and Statistics. 2020/06/16 2020;8(2):39-42.
doi:http://dx.doi.org/10.12691/ajams-8-2-1
40.   Kim JH. Multicollinearity and misleading statistical results. *Korean J Anesthesiol*.
2019;72(6):558-569. doi:https://doi.org/10.4097/kja.19087

Figure 1. Correlation matrix for problematic sexual behavior (PSB) with demographic and clinical variables

P values are provided in Supplemental Digital Content, Table 3.

MH_condition: mental health condition; CT_FHDV: childhood trauma and a family history of domestic violence (including physical, verbal/emotional and sexual childhood trauma and SAST-R Core question 1); RCBS: Richmond Compulsive Buying Scale; OCD: obsessive-compulsive disorder (including self-reported and MINI-diagnosis); SA-SAPAS: Standardized Assessment of Personality–Abbreviated Scale as a Self-Administered Screening Test; FHPSB: a family history of problematic sexual behavior (including a family history of sexual addiction and SAST-R Core question 2); ASRS: the Adult ADHD Self-Report Scale (v1.1); IAT: the Internet Addiction Test

Figure 2. Forest plots of problematic sexual behavior (PSB) logistic regression analysis

A. By conventional logistic regression

B. By machine learning (ML) logistic regression

C. Validation ML logistic regression model

A. The log likelihood of the model was -1149.98 (N=3049, after removing outliers, with a

pseudo R^2=0.16, P<0.00001). The model passed the linktest with a significant P(hat)

(P<0.001) and a non-significant P(hatsq) (P=0.12). The probability of the Hosmer-

Lemeshow χ^2 test was insignificant (P=0.64), suggesting a good model fit. The mean

variance inflation factor (VIF) and the condition number were 1.08 and 6.95, respectively.

The model accuracy, area under the receiver operating (ROC) curve, sensitivity, specificity,

and F score were 84.32%, 77.37%, 14.46%, 98.19% and 0.23, respectively.

B. The machine learning logistic regression was conducted on the dataset without removing

the outliers (training, N=2477; testing, N=619; train to test ratio: 0.80). The model

accuracy, area under the ROC curve, sensitivity, specificity and F score was

74.64%.73.90%, 53.55%, 79.76% and 0.45, respectively.

C. The validation model was trained by adjusted PSB, N=3141, and was tested by monthly

PSB, N=3145. The model accuracy, area under the ROC curve, sensitivity, specificity and F

score was 75.04%.74.26%, 59.11%, 76.97% and 0.34, respectively.

OCD: obsessive-compulsive disorder (including self-reported and MINI-diagnosis); ASRS: the

Adult ADHD Self-Report Scale (v1.1); IAT: the Internet Addiction Test; FHPSB: a family

history of problematic sexual behavior (including a family history of sex addiction and SAST-R

Core question 2); SA-SAPAS: Standardized Assessment of Personality–Abbreviated Scale as a

Self-Administered Screening Test; CT_FHDV: childhood trauma and a family history of

domestic violence (including physical, verbal/emotional and sexual childhood trauma and SAST-

R Core question 1); RCBS: Richmond Compulsive Buying Scale

476
477

Figure 3. Results of multiple correspondence analysis (dimensions 1 and 2) in women (A)

and men (B)

A. Women

B. Men

The maps of data binarized into two categories (yes/no, for each variable) in dimensions 1 and 2

in women and men are shown. OCD: obsessive-compulsive disorder (including self-reported and

MINI-diagnosis); SA-SAPAS: Standardized Assessment of Personality–Abbreviated Scale as a

Self-Administered Screening Test; ASRS: the full Adult ADHD Self-Report Scale (v1.1); IAT:

the Internet Addiction Test; RCBS: Richmond Compulsive Buying Scale; CT_FHPSB:

childhood trauma and a family history of problematic sexual behavior

Figure 4. Results of multiple correspondence analysis (dimensions 4 and 5) in women (A) and men (B)

A. Women

B. Men

The maps of data binarized into two categories (yes/no, for each variable) in dimensions 4 and 5 in women and men are shown. OCD: obsessive-compulsive disorder (including self-reported and MINI-diagnosis); SA-SAPAS: Standardized Assessment of Personality–Abbreviated Scale as a Self-Administered Screening Test; ASRS: the full Adult ADHD Self-Report Scale (v1.1); IAT: the Internet Addiction Test; RCBS: Richmond Compulsive Buying Scale; CT_FHPSB: childhood trauma and a family history of problematic sexual behavior

Table 1. Distribution of demographic and clinical variables by problematic sexual behavior (PSB) group

Categories			PSB "yes" 532/3219 (16.53%)	PSB "no" 2687/3219 (83.47%)	Total number	*P* value
SAST-R Core score			6.58±3.22 (N=532)	1.38±1.56 (N=2687)	3219	*P*<0.001[a]
Age			22.32±5.13 (N=532)	22.34±5.05 (N=2686)	3218	*P*=0.81[a]
Gender	**Women**		232 (10.82%)	1913 (89.18%)	2145	*P*<0.001[b]
	Men		290 (28.27%)	736 (71.73%)	1026	
	Total		522 (16.46%)	2649 (83.54%)	3171	
Sexual Orientation	**Heterosexual**		420 (15.85%)	2230 (84.15%)	2650	*P*=0.034[b]
	Non-heterosexual		110 (19.50%)	454 (80.50%)	564	
	Total		530 (16.49%)	2684 (83.51%)	3214	
OCD	**No**		490 (15.88%)	2596 (84.12%)	3086	*P*<0.001[b]
	Yes		40 (33.33%)	80 (66.67%)	120	
	Total		530 (16.53%)	2676 (83.47%)	3206	
RCBS	**No**		471 (15.54%)	2559 (84.46%)	3030	*P*<0.001[b]
	Yes		61 (32.28%)	128 (67.72%)	189	
	Total		532 (16.53%)	2687 (83.47%)	3219	

FHPSB	**No**	453 (15.38%)	2492 (84.62%)	2945	*P*<0.001[b]
	Yes	71 (34.63%)	134 (65.37%)	205	
	Total	524 (16.63%)	2626 (83.37%)	3150	
SA-SAPAS	**No**	425 (14.76%)	2454 (85.24%)	2879	*P*<0.001[b]
	Yes	107 (31.47%)	233 (68.53%)	340	
	Total	532 (16.53%)	2687 (83.47%)	3219	
Mental health condition(s)	**No**	399 (15.35%)	2200 (84.65%)	2599	*P*<0.001[b]
	Yes	131 (21.87%)	468 (78.13%)	599	
	Total	530 (16.57%)	2668 (83.43%)	3198	
Childhood trauma	**No**	339 (13.94%)	2092 (86.06%)	2431	*P*<0.001[b]
	Yes	185 (25.45%)	542 (74.55%)	727	
	Total	524 (16.59%)	2634 (83.41%)	3158	
ASRS	**No**	321 (12.97%)	2153 (87.03%)	2474	*P*<0.001[b]
	Yes	211 (28.32%)	534 (71.68%)	745	
	Total	532 (16.53%%)	2687 (83.47%)	3219	
IAT	**Normal (No)**	241 (11.50%)	1855 (88.50%)	2096	*P*<0.001[b]
	Mild, moderate or severe (Yes)	291 (25.91%)	832 (74.09%)	1123	
	Total	532 (16.53%)	2687 (83.47%)	3219	

[a]Mann-Whitney U test

[b]Pearson's χ^2 test

OCD: obsessive-compulsive disorder (including self-reported and MINI-diagnosis); RCBS:

Richmond Compulsive Buying Scale; FHPSB: a family history of problematic sexual behavior

(including a family history of sex addiction and SAST Core question 2); SA-SAPAS:

Standardized Assessment of Personality–Abbreviated Scale as a Self-Administered Screening

515   Test; Childhood trauma: childhood trauma (including physical, verbal/emotional and sexual

516   childhood trauma and SAST Core question 1) and a family history of domestic violence; ASRS:

517   the Adult ADHD Self-Report Scale (v1.1); IAT: the Internet Addiction Test