1

Case-control investigation of invasive Salmonella disease in Africa – comparison of

human, animal and household environmental isolates find no evidence of environmental

or animal reservoirs of invasive strains

Leonard Koolman[1,2 #], Reenesh Prakash[1,2 #], Yohane Diness[1], Chisomo Msefula[4], Tonney S.

Nyirenda[4], Franziska Olgemoeller[1,3], Blanca Perez-Sepulveda[2], Jay C. D. Hinton[2], Siân V.

Owen[2], Nicholas A. Feasey[1,3], Philip M. Ashton[1,2*], Melita A. Gordon[1,2,4]

[1] Malawi-Liverpool Wellcome Programme, Blantyre, Malawi

[2] Institute of Infection, Veterinary & Ecological Sciences, University of Liverpool, United

Kingdom

[3] Department of Clinical Sciences, Liverpool School of Tropical Medicine, United Kingdom

[4] Kamuzu University of Health Sciences, Blantyre, Malawi

# contributed equally

*Corresponding author

Email: philip.ashton@liverpool.ac.uk

Abstract

Background

Invasive *Salmonella* infections are a major cause of morbidity and mortality in Sub-Saharan Africa (SSA), but the sources and transmission routes are uncertain. We investigated potential sources for cases of invasive disease by sampling healthy people, animals, and the environment in index-case and geographically-matched control households.

Methods

Sixty index cases of human invasive *Salmonella* infection were recruited (28 invasive Non-Typhoidal *Salmonella* (iNTS) disease, and 32 typhoid). Each index-case household was geographically matched to a control household. Extensive microbiological sampling included stool sampling from healthy household members, stool or rectal swabs from household-associated animals and boot-sock sampling of the household environment.

Results

1203 samples were taken from 120 households, yielding 43 *Salmonella* isolates from 25 households (overall sample positivity 3.6%). Isolates from households were all NTS and spanned 15 Sequence Types (STs). iNTS disease was caused by 3 STs of Salmonella Typhimurium, mainly ST313. Two *S.* Typhimurium isolates from index cases closely matched isolates from their respective asymptomatic household members (2 and 3 SNPs different respectively). There was no overlap of STs causing iNTS disease with environmental or animal isolates, despite recovery of diverse NTS.

Conclusions

The finding of NTS strains from index cases matching household members, coupled with lack of overlap with either animal or environmental isolates, supports a hypothesis that healthy humans are the source of iNTS infections in the household. The breadth of NTS strains found in the household environment across all sites demonstrated the robustness of

47   sampling and methodology to detect NTS, and suggests a diverse ecology of *Salmonella* in

48   this setting. The lack of *S.* Typhi isolated from the household environment may suggest a

49   need for further methodological development to culture sources of typhoid.

Author summary

Invasive *Salmonella* infections lead to millions of disability adjusted life years (DALYs) lost every year globally. There are two main types of invasive *Salmonella* infections in Africa; i) typhoid fever, caused by *Salmonella* Typhi, and ii) invasive Non-Typhoidal Salmonella (iNTS) disease, primarily caused in our setting by *Salmonella* Typhimurium. Despite their high disease burden, and despite observed differences in the epidemiology of typhoid compared to iNTS disease, we do not have a good understanding of their reservoirs and transmission. Therefore, we carried out extensive microbiological sampling of the household members and living environments of patients with invasive *Salmonella* infections, and of geographically-matched control households, and investigated the genetic relationships between household *Salmonella* and index-case blood-stream isolates using whole genome sequencing (WGS). The only sample type where we identified *Salmonella* that matched strains causing iNTS disease was the stool of healthy people from the case household, suggesting either person-to-person spread or infection from a common source. Boot-sock sampling of the household environment resulted in the highest yield of *Salmonella* of any of our sampling strategies. None of the other 41 environmental *Salmonella* isolates from non-human sources, including 4 domestic animal-associated isolates, matched the disease-causing types. Our findings are consistent with the reservoir of Typhimurium iNTS infections being the human gastrointestinal tract and transmission occurring within households.

Introduction

Salmonella infections are a substantial cause of morbidity and mortality in many low- and middle-income countries and are an important cause of enteric infections leading to invasive bloodstream infections (BSI) [1,2]. Both typhoidal and Non-Typhoidal *Salmonella* (NTS) contribute extensively to the burden of BSI. Globally in 2017, invasive NTS (iNTS) disease was estimated to cause 535,000 illnesses and 77,500 deaths [2]. Sub-Saharan Africa accounts for 85.8% of global iNTS deaths [2]. There were an estimated 14.3 million cases of typhoid and paratyphoid fever in 2017, resulting in 135,900 deaths, 15.8% of which were in Sub-Saharan Africa [1].

Salmonella Typhi has emerged as a major cause of BSI in southern and eastern Africa since 2012 [3–7], and Malawi has a very high incidence of 444/100000 person-years of observation [8]. The pathways by which people are exposed to *S*. Typhi are uncertain [9]. Improved understanding of reservoirs and transmission routes would help with the design of effective interventions, but delineating the relative importance of long (i.e. waterborne or environmentally mediated) and short cycle (i.e. within household) transmission routes has remained difficult for a variety of faecal oral pathogens [9,10].

In High Income Countries (HICs), NTS are predominantly associated with self-limiting enterocolitis. In Sub-Saharan Africa (SSA), however, the epidemiological picture is dominated by iNTS disease, diagnosed by positive blood-culture, which carries a high case-fatality [11–15]. iNTS in Africa is associated with host susceptibilities, including HIV in adults and malaria infections, malnutrition or HIV in young children [13,15–17]. Two *Salmonella* serotypes, Typhimurium and Enteritidis, have been reported to be the commonest causes of iNTS in Malawi, South Africa, Kenya, Mozambique and Mali, with *S*. Typhimurium Sequence Type

(ST) 313 (ST313) and S. Enteritidis ST11 being the most frequently identified [17–24]. In the

city of Blantyre, Malawi, there have been two epidemics of iNTS disease, the first caused by

S. Enteritidis between 1999-2002, and the second caused by *S.* Typhimurium between 2002-

2008, with *S.* Typhimurium continuing to cause a significant burden of disease [17]. Both

epidemics were associated with multi-drug resistance [17].

While NTS infection in HICs is generally associated with zoonotic transmission, the reservoirs

and transmission route of the pathogens responsible for iNTS in SSA remains unknown.

Seasonal patterns of iNTS, and the close domestic association of humans and animals in parts

of SSA suggest a possible role for environmental factors [25,26]. Serotypes associated with

iNTS have been isolated from animals in South Africa, although no genetic characterisation

was carried out on these samples [27,28]. In contrast, microbiological investigations in Kenya,

based either on serotype and Pulsed Field Gel Electrophoresis or more recently WGS, found

isolates in the stool of household members which were closely related to index cases, and only

rarely found linked serotypes in environmental samples such as water from nearby rivers,

household and market vendor food, or household animals [29,30]. The Global Enteric

Multicentre Study found iNTS associated sub-types of *Salmonella* in the stool of 42 children

with diarrhoea and 17 healthy children, highlighting the potential role of the human gut as a

source of iNTS infections [31]. In The Gambia, there was no overlap observed between STs

observed in iNTS cases and those observed in rectal swabs from domestic animals from the

households of cases [32], while in Tanzania and Kenya, extensive sampling of "meat pathway"

isolates and human disease isolates only isolated ST313 from human samples [33]. ST313 was

isolated from pigs in Nairobi, but was shown to group with ST313 from the UK, which are not

associated with invasive disease [34,35]. Genomic evidence suggests that *S.* Typhimurium

ST313, one of the commonest causes of iNTS, is under-going genome degradation [22,23], a

phenomenon associated with narrower host range (i.e. becoming human restricted) and

increased invasiveness in a range of bacteria, including *S.* Typhi and *S.* Paratyphi A [36,37].

Transcriptomic and phenotypic evidence also points towards specific niche adaptations to

human hosts among African isolates of NTS causing invasive disease [38,39].

To address the knowledge gap around sources for strains of Salmonella causing invasive

Salmonella infections, we conducted extensive human, animal, and household environmental

sampling of iNTS and typhoid index-case households and geographically matched control

households in Blantyre, Malawi. This study is the first to combine extensive bootsock

environmental sampling alongside sampling of humans and animals, and to use WGS of the

resulting isolates, aiming primarily to investigate overlap of invasive index strains with

household sources, and also to comprehensively determine broader household patterns of

bacterial relatedness in relation to invasive Salmonella diseases.

Methods

Setting

Queen Elizabeth Central Hospital (QECH) provides free healthcare to the approximately 1.3 million inhabitants of Blantyre District, Malawi, and is the tertiary referral hospital for southern Malawi. Since 1998, the Malawi-Liverpool-Wellcome Trust Clinical Research Programme has provided a quality controlled diagnostic blood culture service for febrile adult and paediatric medical patients admitted to QECH since 1998. This service is provided for admitted adults (>16 years old) with axillary temperature over 37.5°C or clinical suspicion of sepsis, and for children (<16 years old) who were malaria slide negative, or positive and critically ill, or with clinical suspicion of sepsis.

Case and control recruitment

Sequential blood culture confirmed adult or paediatric cases with invasive *Salmonella* infections (either iNTS or typhoid fever) presenting at QECH between March 2015 and October 2016 were approached as index cases. We sought written informed consent from adults, and guardian consent for minors. Patients living outside the Blantyre district and those with recurring iNTS disease were excluded. Index case households were sampled within a maximum of 14 days following initial presentation to QECH. Following recruitment, the field team visited the index cases in their households, where GPS co-ordinates were taken and a household socio-demographic questionnaire was completed. Control households were then selected by random bottle-spin and pacing 100 m from the index case household, recruited with informed consent, and GPS and questionnaire data were also collected. Exclusion criteria for control households were current treatment for invasive *Salmonella* disease for any family member, or declining consent. In this event, the next-nearest house in the same direction was selected.

Sampling Methodology

We carried out a microbiological survey of the index-case household and the control

household, comprising stool from household members, stool or rectal swabs from domestic

and household-associated animals and systematic boot-sock sampling of the living

environment (latrine, rubbish area, bedroom, cooking area and the house perimeter).

Stool (humans): Samples (5 to 10 g) were collected from all available family members in the

index case and control households. Sample containers were provided 24-48 hours before the

main sample-collection visit, and household members were requested to collect up to three

consecutive stools from each individual. The three samples belonging to each person were

pooled, to maximise recovery of *Salmonella*.

Stool (animals): Fresh stool or rectal swab samples were collected from domestic or household-

associated animals (cows, chickens, goats, pigs, dogs, cats, rats, doves, guinea pigs or gecko

lizards).

Environmental: The investigator's shoe was first covered with a waterproof sterile bag to

prevent cross-contamination, then a sterile bootsock was put over this, moistened with sterile

water, and single bootsocks were used to collect samples from each of the follow sites: pit-

latrine, outdoor rubbish area, cooking and bedroom areas, and from the outside perimeter of

the household, using a standardised protocol for location and number of paces (10 paces), and

a step and twist motion. Bootsocks were then removed and placed into a sterile plastic bag for

transport.

All samples were transferred from the field to the laboratory in coolboxes, and processed on

the same day.

 Sample Enrichment and culture

Supplementary Figure 1 shows a schematic of sample processing. A portion (~1g) of stool was

emulsified in 9ml of Selenite F selective enrichment broth (Oxoid, UK) and incubated at 37°C

for 18-24 hours. For samples other than stool (i.e. bootsocks, swabs), we used a pre-enrichment

step; these samples were incubated in Buffered Peptone Water (BPW) at 37°C for 18-24 hours.

One ml of the BPW sample was then transferred into 9 ml Selenite broth and incubated for a

further 18-24 hours at 37°C.

After incubation, 100µl of bacterial culture taken from the surface of the Selenite broth was

sub-cultured onto Xylose Lysine Deoxycholate (XLD) agar, incubated at 37°C for 18-24 hours

and examined for suspected colonies of *Salmonella* (red/pink colonies with black centres). Pure

candidate colonies were biochemically confirmed as *Salmonella* species using API 20E

(BioMerieux). Serotyping was performed by the slide agglutination technique. Putative

Salmonella isolates were sub-cultured from beads onto Nutrient agar (Oxoid, UK) and

incubated for 24 hours at 37°C. Following growth, a drop of 0.85% saline was placed onto a

microscope slide and a single colony from the Nutrient agar was emulsified into the saline.

Positive serotyping was confirmed if agglutination occurred in antigens O4, O9, Vi, Hd, Hg,

Hi, Hm. If an organism was positive for Vi antigen but negative for O9 antigen, a dense

suspension of the organism was prepared in 0.85% saline, autoclaved at 121°C for 15 minutes

and the agglutination repeated. *S.* Typhi was differentiated from NTS isolates if agglutination

occurred for O9, Vi and Hd antigens. All API 20E confirmed *Salmonella* isolates which were

not *S.* Typhi were sent to the University of Liverpool, UK, for WGS.

Whole Genome Sequencing (WGS)

Bacterial genomic DNA was extracted from overnight cultures in LB (1% tryptone, 0.5% yeast

extract, 0.5% NaCl; pH 7.0) using the DNeasy Blood and Tissue kit (Qiagen), as per the Gram-

negative bacteria protocol for the Quick-DNA^{TM} Universal Kit (Zymo Research), Biological

Fluids & Cells protocol. Sequencing libraries were constructed using the TruSeq DNA PCR-

free library preparation kit (Illumina) or the TruSeq Nano DNA HT library preparation kit

(Illumina) using a target insert size of 550bp. Libraries were sequenced on an Illumina MiSeq

instrument using a 2x250bp paired end protocol at the Centre for Genomic Research,

University of Liverpool, UK. Raw sequencing reads were trimmed to remove Illumina adapter

sequences using Cutadapt [40] and further trimmed to remove poor quality sequence using

Sickle [41] version 1.2 with a minimum window quality score of 20, discarding reads which

were less than 10 bp after trimming. Trimmed reads were submitted to Enterobase

(https://enterobase.warwick.ac.uk/) for draft genome assembly, MLST assignment, *in silico*

serotype prediction and phylogenetic analysis [42]. SNP identification within Enterobase was

carried out via a mapping and SNP calling pipeline, and phylogenetic tree construbtion done

with RAxML [43–46] Determinants of antimicrobial resistance genes were identified using the

amr-finder-plus software version 3.9.8, using database version 2020-12-17.1 [47].

Data availability

All sequencing data is available from ENA in the BioProject X. Per sample BioSample

accessions are listed in Supplementary Table 1.

    Results

Sixty-seven index-case households (35 typhoid and 32 iNTS) were eligible for enrolment

between March 2015 and October 2016. Seven index-case households declined to participate,

leaving 32 typhoid index case households, with 32 controls, and 28 iNTS index case

households, with 28 controls recruited for the study, giving a total of 120 households. All

households were in high-density low-income housing locations within Blantyre.

Isolation of Salmonella by culture: Table 1 shows sampling and positivity rates for Salmonella

by sample-type and by household. We collected a total of 1203 samples from the 120 index-

case and control households (mean 10 samples per household), yielding 43 isolates (all NTS)

which were confirmed as *Salmonella* by WGS, giving an overall sample positivity of 3.6%

(Table 1). Overall, sample positivity was very similar across human, animal and bootsock

environment samples (human stool 16/491, 3.3%; animals 4/110, 3.6%; bootsocks 23/620,

3.8%), and across different household categories (Table 1). Animal sampling was from 110

animals in 71/120 (59%) households, and included both domesticated and non-domesticated

species (cows, chickens, pigs, dogs, cats, rats, doves, guinea pigs and geckos).

Salmonella was found in 25/120 households (overall household positivity of 21%). There was

a significantly higher household positivity in typhoid households (28%) and their

geographically-matched neighbours, compared to NTS cases and control households (13%)

(Chi-squared P-value = 0.036), particularly in the environmental bootsock category of samples.

No *S.* Typhi were isolated from any category of household samples, including from people

living in the same house as index cases.

Table 1: Isolation of Salmonella from different sample types in iNTS and typhoid case and paired control households

	iNTS Index Case Households (HH) (n=28)						iNTS Control Households (HH) (n=28)					
	Total samples	No. samples with Salmonella	% positive samples	No. HH tested	No. HH with Salmonella	% Positive HH	Total samples	No. samples with Salmonella	% positive samples	No. HH tested	No. HH with Salmonella	% Positive HH
Human Stool	116	9	7.8	28	4	14.3	97	0	0.0	28	0	0.0
Animal stool/swab	27	0	0.0	20	0	0.0	38	0	0.0	17	0	0.0
HH environment	137	0	0.0	28	0	0.0	131	6	4.6	28	3	10.7
Total	**280**	**9**	**3.2**	**28.0**	**4**	14.3	**266**	**6**	**2.3**	**28**	**3**	**10.7**
	Typhoid Index Case households (HH) (n=32)						Typhoid Control households (HH) (n=32)					
Human Stool	157	4	2.5	32	2	6.3	121	3	2.5	32	3	9.4
Animal stool/swab	22	2	9.1	18	2	11.1	23	2	8.7	16	2	12.5
HH environment	168	5	3.0	32	4	12.5	166	12	7.2	32	9	28.1
Total	**347**	**11**	**3.2**	**32**	**7**	**21.9**	**310**	**17**	**5.5**	**32**	**11**	**34.4**

Relatedness of Sequence Types from different sample categories: Figure 1 demonstrates the

wide diversity and relationships of Sequence Types (STs) isolated from different sample

categories (invasive disease, healthy human stool, animals, and the household environment).

Table 2 shows the individual household origins of each isolate, and the relationship between

ST and imputed serovar for each isolate. Among invasive NTS disease cases, 8 of 27 index

case NTS isolates could not be resuscitated from the archive for sequencing. The remaining

19 typed index NTS isolates were all *S*. Typhimurium from just 3 STs; 13 were ST313, 5

were ST3257 and 1 was ST19. These latter STs were each only one locus variant away from

ST313. One representative typhoid index strain (ST2) is also included for reference. Among

healthy household members there were 16 isolates from 8 STs; among animals there were 4

isolates from 3 STs; and from environmental samples there were 23 isolates from 9 different

STs.

Importantly, in two instances, there was ST overlap between invasive index-case isolates and

healthy human household samples; these were 2 cases of invasive disease caused by *S*.

Typhimurium ST313 and ST3257, that were linked to isolates from 2 healthy household

members in their respective households, who also had also ST313 and ST3257 isolated from

their stools (Figure 1). Interestingly, although stool isolates from household members were

found equally among adults (n=7) and children (n=8), in both cases the household member

carrying a matching isolate was an adult. The genetic relatedness between index and human

household strains was explored at higher resolution by WGS phylogenetic analysis (Figure 2

and below).

Also importantly, there was no overlap in STs found between isolates causing invasive

human disease and any animal or environmental isolates, despite a good rate of isolation from

all 3 sample-types. There was, however, non-co-localised overlap in some STs between

healthy humans and household-associated animals. A monophasic ST3262 was found in a

cat, a gecko, and a human, but these were all in different households. Similarly, ST3261 was

found in 2 healthy humans and a chicken, but these 3 events were also not from the same

households (Table 1). Significantly, ST3261 nor ST3262 are not known to have caused

human invasive disease in Blantyre. ST325 was found only in an animal (gecko).

As might be expected in this low-income setting, where household sanitation is limited and

faecal contamination of the household may occur, there were STs isolated from the

environment that overlapped with those from healthy humans, and household animals. Of 9

STs found in the environment, 3 (ST3262, ST3263, ST2152) showed some overlap with

healthy human or animal isolates. However, a further 6 STs were found only in the

household environment, revealing the diversity of Salmonella strains in this setting.

Figure 1: Minimum spanning tree of all isolates. Each circle represents a sequence type, the size of the circle is proportional to the number of isolates of that ST. Red text indicates the number of locus variants between two STs. Two invasive disease index isolates had a matched isolate from a healthy human sample of the same ST from the same household. These isolates are indicated by black and red outlines.

297   Table 2: Index case and household isolates described in this study

Household (HH) ID S=case HH C=control HH	Index case NTS or typhoid?	Household case or control?	Household sample type	Sequence Type (ST)	Imputed serovar	Match to Index case blood culture?	Household (HH) ID S=case HH C=control HH
1C	Typhoid	Control	Animal stool (chicken)	3261	Agoueve|Cubana		1C
2S	iNTS	Case household	Index case blood	3257	Typhimurium		2S
3C	Typhoid	Control	Family member stool	3261	Agoueve|Cubana		3C
4S	iNTS	Case household	Index case blood	313	Typhimurium		4S
6S	Typhoid	Case household	Family member stool	3261	Agoueve|Cubana		6S
6C	Typhoid	Control	Boot sock perimeter	3263	Havana|II		6C
6C	Typhoid	Control	Family member stool	3263	Havana|II		6C
8C	Typhoid	Control	Boot sock bedroom	3263	Havana|II		8C
9S	iNTS	Case household	Index case blood	313	Typhimurium		9S
9S	iNTS	Case household	Family member stool	3263	Havana|II		9S
9S	iNTS	Case household	Family member stool	3263	Havana|II		9S
9S	iNTS	Case household	Family member stool	3263	Havana|II		9S
9S	iNTS	Case household	Family member stool	3263	Havana|II		9S
9S	iNTS	Case household	Family member stool	3263	Havana|II		9S
9S	iNTS	Case household	Family member stool	3263	Havana|II		9S
12S	iNTS	Case household	Index case blood	3257	Typhimurium		12S
14S	iNTS	Case household	Index case blood	313	Typhimurium		14S
15S	Typhoid	Case household	Animal stool (gecko)	3262	II 42:r:-|IIIb		15S
15S	Typhoid	Case household	Boot sock perimeter	3262	II 42:r:-|IIIb		15S
16S	iNTS	Case household	Index case blood	19	Typhimurium		16S
16S	iNTS	Case household	Family member stool	3262	II 42:r:-|IIIb		16S
16C	iNTS	Control	Boot sock latrine	2347	Mgulani		16C
16C	iNTS	Control	Boot sock cooking area	2347	Mgulani		16C
17S	Typhoid	Case household	Boot sock latrine	3262	II 42:r:-|IIIb		17S
18S	iNTS	Case household	Index case blood	3257	Typhimurium		18S
22C	Typhoid	Control	Boot sock latrine	3609	Djama|II 42:z29:-		22C
23S	Typhoid	Case household	Animal stool (gecko)	316	Montevideo		23S
25S	iNTS	Case household	Family member stool	313	Typhimurium	match1	25S
25S	iNTS	Case household	Index case blood	313	Typhimurium	match1	25S

25C	iNTS	Control	Boot sock latrine	473	Hadar		25C
25C	iNTS	Control	Boot sock bedroom	473	Hadar		25C
25C	iNTS	Control	Boot sock cooking area	473	Hadar		25C
26S	iNTS	Case household	Index case blood	313	Typhimurium		26S
27S	iNTS	Case household	Index case blood	313	Typhimurium		27S
27C	iNTS	Control	Boot sock bedroom	3262	II 42:r:-|IIIb		27C
28C	Typhoid	Control	Animal rectal swab	3262	II 42:r:-|IIIb		28C
28C	Typhoid	Control	Boot sock latrine 1	3262	II 42:r:-|IIIb		28C
28C	Typhoid	Control	Boot sock latrine 1	3262	II 42:r:-|IIIb		28C
29C	Typhoid	Control	Boot sock latrine	2152	Gaminara		29C
29C	Typhoid	Control	Family member stool	3608	Ogbete|II		29C
31S	Typhoid	Case household	Index case blood	2	Typhi		31S
32S	iNTS	Case household	Index case blood	313	Typhimurium		32S
33S	iNTS	Case household	Index case blood	313	Typhimurium		33S
36S	iNTS	Case household	Index case blood	3257	Typhimurium		36S
39S	iNTS	Case household	Index case blood	313	Typhimurium		39S
41C	Typhoid	Control	Boot sock latrine	3262	II 42:r:-|IIIb		41C
42S	iNTS	Case household	Index case blood	313	Typhimurium		42S
44S	iNTS	Case household	Family member stool	3257	Typhimurium	match2	44S
44S	iNTS	Case household	Index case blood	3257	Typhimurium	match2	44S
46S	iNTS	Case household	Index case blood	313	Typhimurium		46S
47S	iNTS	Case household	Index case blood	313	Typhimurium		47S
48S	Typhoid	Case household	Boot sock bedroom	3262	II 42:r:-|IIIb		48S
50S	Typhoid	Case household	Boot sock bedroom	3262	II 42:r:-|IIIb		50S
50S	Typhoid	Case household	Boot sock cooking area	3262	II 42:r:-|IIIb		50S
50C	Typhoid	Control	Boot sock perimeter	3611	II 40:b:-		50C
53C	Typhoid	Control	Boot sock bedroom	3262	II 42:r:-|IIIb		53C
56C	Typhoid	Control	Boot sock bedroom	293	Amager		56C
56C	Typhoid	Control	Boot sock perimeter	293	Amager		56C
56C	Typhoid	Control	Boot sock cooking area	3610	Aberdeen		56C
57S	iNTS	Case household	Index case blood	313	Typhimurium		57S
59S	Typhoid	Case household	Family member	14	Senftenberg		59S
59S	Typhoid	Case household	Family member stool	2152	Gaminara		59S
59S	Typhoid	Case household	Family member stool	2152	Gaminara		59S

Sequence types and WGS-inferred serovars from different sample categories:

The relationship between STs and WGS-inferred serovars for all isolates in the study is

documented in Table 2 (genome database accession numbers are available in Supplementary

Table 1). Of 19 index iNTS case isolates from blood culture, 19/19 were *S.* Typhimurium, of

3 different STs; ST313 (n=13), ST3257 (a Single Locus Variant of ST313, n=5) and ST19 (n

= 1) (Supplementary Table 1, Figure 1).

Serovars that were associated with healthy humans were Havana (7), Typhimurium (2),

Gaminara (2), Agoueve/Cubana (2), Seftenberg (1) and Ogbete (1). Serovars associated with

animals were II42:r:-|IIIb 42:r:-:[z50] (1 gecko, 1 cat), Montevideo (1 gecko) and

Agoueve/Cubana (1 chicken). The wide range of serovars found in the household environment

included II42:r:-|IIIb 42:r:-:[z50] (10), Hadar (3), Havana (3), Amager (2), Mgulani (2) Djama

(1) and Aberdeen (1).

Reflecting the broad diversity of isolates, many STs/serovars were represented by just a single

isolate. In addition, where there were multiple isolations of the same serovar, a large proportion

were localised from within the same household. Striking examples of this are: of 9 isolates of

S. Havana, 6 arose from one household (9S), from 6 out of 7 children sampled; of 3 isolates of

S Hadar, all arose from the same household (25C, latrine, bedroom and cooking area); of 12

isolates of II42:r:-|IIIb 42:r:-:[z50] there were multiple positive samples in three households,

including from animal and environmental samples from the same households. Similarly, 2

isolates each of S. Mgulani and S. Amager isolates were linked to the same respective

households (16C and 56C). This indicates a very marked geographical diversity between

households, and a degree of homogeneity within households among NTS that are not linked to

invasive disease.

Whole genome phylogeny of index/household pairs: We employed whole genome-derived

phylogeny to infer at high resolution the relationship between the human invasive disease

cases of *S.* Typhimurium ST313 & ST3257 and the stool carriage isolates of their respective

index household members. In both cases, the isolate that caused the invasive disease formed a

monophyletic group with the carriage isolate from the same index household (Figure 2). The

SNP distance was 2 SNPs in one case, and 3 SNPs in the other (Figure 2).

Figure 2: Whole genome maximum likelihood phylogenetic tree of S. Typhimurium which
shows the close relationship between the index case and family member isolates from
households 25 and 44. Sequencing read sets were compared against the reference genome
D23580.

Antimicrobial resistance genes: We had WGS data for 63 isolates from index cases of human

invasive disease and household sampling. Antimicrobial resistance gene results are shown in

Supplementary Table 1. There were no antimicrobial resistance genes associated with 52 of

them, including 68% of index blood-stream *S.* Typhimurium isolates (9 ST313s, 3 ST3257

and 1 ST19), and 89% of the animal and bootsock isolates. Two bootsock (both *S.* Mgulani)

and one animal stool (*S.* Montevideo) isolate(s) carried *fosA7* genes associated with

resistance to fosfomycin. Six index-case blood-stream Typhimurium isolates and two

Typhimurium from the stool of household members encoded all or some of the *blaTEM-1,*

catA1, sul1, sul2 and dfrA1 resistance genes which are typically associated with invasive *S.*

Typhimurium in Malawi.

 Discussion

Our findings successfully address our primary question, and support the hypothesis that

humans may be the primary source of the organisms responsible for iNTS in sub-Saharan

Africa. All index cases were caused by 3 STs of *S.* Typhimurium, and matching *S.*

Typhimurium STs were isolated from the stool of their respective household members. In

both cases, the isolates from household members were of the same ST and within 2-3 SNPs

distance of the isolate from the index case. On the other hand, despite successfully culturing

27 *Salmonella* isolates from the environment and animals, representing 13 different

STs/serovars, none were iNTS-associated *Salmonella* serovars.

Our results and interpretation are consistent with and extend those of several previous reports.

Firstly, in Kenya, iNTS infections caused by *Salmonella* Typhimurium or Enteritidis in an

informal urban settlement were not found to be associated with the rearing of any domestic

animals [48]. Secondly, in Burkina Faso the only isolates that matched the index cases were

obtained from the stool of household members, despite extensive animal and water sampling

in the households of iNTS patients [49]. Finally, analysis of cases of moderate to severe

diarrhoea and healthy controls in Africa and Asia found ST313 in both sick and healthy

children, supporting the finding that *Salmonella* are found in the stool of healthy children

[31].

Our study also extends the understanding of the sources of iNTS infection in several

additional ways. Most importantly, we sampled from a broad range of locations. Previous

case-control studies have sampled adults and children in the same household, and household-

associated animals, but this is the first study to also sample the household environment. This

yielded a similar culture positivity-rate to other sample-types, but also yielded a much wider

range of STs/serovars, indicating that *Salmonella* has a more diverse ecology in the

household environment than has previously been appreciated. The previously unappreciated

broad diversity of Salmonella in the household environment in this setting is in contrast to the

narrow range of STs causing human invasive disease, and the lack of overlap between these

categories is very striking.

Ours is also the most comprehensive study of its type, finding the greatest number of non-

invasive disease isolates of any publication addressing this topic to date. We isolated a total

of 27 animal and environmental *Salmonella,* almost double the number isolated from a

comparable study [49]. Animal-ownership is not high in overcrowded urban areas, but we

nevertheless sampled 110 household-associated animals from 59% of households, yielding 3

STs from 4 animals of 3 species (1 chicken, 1 cat and 2 geckos). Bootsock sampling provided

53% of the non-invasive disease isolates in our study. This excellent yield from bootsocks for

detecting *Salmonella* in other contexts has been reported in other contexts [50], and we

recommend that similar studies carried out in the future also use this method.

Our methodology was thus able to resolve microbiological connections between people and

their household environments at both the individual household level and the community

level. The presence of some STs in multiple sample types was reassuring, indicating that we

were able to resolve a degree of connectivity between the different sample source types in

this setting, which might be expected as a result of faecal contamination of the household in a

setting with poor sanitation infrastructure. However, the dominant picture was a diversity of

non-invasive disease *Salmonella*, and unique distribution of different STs by sample-type. 6/9

STs found in the environment were found in neither animals nor healthy humans. Three STs

were only found in one sample type, and even when there were multiple isolates of a single

ST, these were often concentrated in the same household or a limited number of households.

This suggests that some serovars may occupy a specific niche.

S. Typhi is human-restricted in its ability to cause disease, but is thought to inhabit the

environment during long-cycle transmission in the community. While we failed to isolate *S.*

Typhi from the environment of either case or control households, this is known to be

technically challenging, and our results highlight the need for further methodological

development in this area [51].

There was no definite broader discernible pattern of STs/serovars observed between different

categories of households. There was, however, higher household-positivity rates in typhoid-

linked case and control household pairs, compared to iNTS-linked case and control

household pairs. Case and matched control houses were 100m apart, and this was chosen as a

distance that meant households were geographically close, but would not be likely to share

latrines or experience direct surface-water cross-contamination, since at that distance there

would usually be several intervening plots in this setting. Poor sanitation is known to be a

risk factor for typhoid because of the long-cycle transmission cycle, while the main

epidemiological risks described for iNTS disease are found in the host; malaria, malnutrition,

anaemia and HIV. It is possible that a higher rate of household contamination with NTS

serovars in case/control household pairs in proximity to typhoid cases reflects worse general

water and sanitation, and possibly higher poverty in that neighbourhood, compared to iNTS

case/control household pairs, but this would require further investigation.

While antimicrobial resistance is a major concern in our setting, only 12% of our household

isolates had one or more genes known to be associated with antimicrobial resistance, in

contrast to 39% of our human invasive disease isolates. The fact that 62% of the ST313

isolated here carried no antimicrobial resistance genes suggests that they could belong to the

recently described ST313 Lineage 3 [52]. We did not carry out AMR phenotyping for the

isolates obtained in this study, but AMR profile inference from genome sequence is highly

accurate for *Salmonella* [53].

A major limitation to our study is that both the index case and the ST313 colonised household

member could have been infected from an unsampled environmental reservoir, either outside

the household or within the household. We do not report on food or water sampling. While

we had hoped to sample food, we found that in practice most households had no refrigeration,

and there were very rarely "leftovers" as meals were eaten at one sitting, meaning this had

limited feasibility. An easier approach in this setting might be to sample from local markets

and food-supply networks. Even in investigations of food-borne outbreaks in HICs, food

testing results are difficult to interpret and have inherent limitations [54]. Epidemiological

analysis identified no vehicle of infection in 38% of putative foodborne outbreaks in the UK,

and microbiological confirmation would only occur in a minority of those 38% [55]. In

addition to a contaminated foodstuff, water sources, both drinking and washing/cleaning,

could be involved in the transmission of organisms causing iNTS, as is seen for *S.* Typhi

[9,56], and further methodological development would be required for these studies. If the

index-case were infected outside the home e.g. at work, school, or nursery, which is a

possible risk factor in our setting [9], then our study would not identify that microbiological

source.

Conclusions

In summary, we conducted comprehensive environmental and household contact sampling of the homes of patients with invasive Salmonella disease, and geographically matched control households. The only household isolates of serotypes associated with iNTS came from the stool of household members of iNTS index cases. This study contributes to accumulating evidence that the source of iNTS infections is the human gastrointestinal tract. Significantly, we found no strains causing human iNTS disease among any household-associated animals, despite extensive animal sampling and a comparable NTS positivity rate. We also uncovered a much wider range of NTS isolates that are not associated with invasive disease, across all sampling sites, than has previously been appreciated, indicating a diverse ecology for Salmonella in the household environment.

Acknowledgements

We are grateful to Alistair Darby, John Kenny and staff at the Centre for Genomic Research at the University of Liverpool for assistance with genome sequencing. Jay Hinton is grateful for funding by a Wellcome Trust Senior Investigator Award (106914/Z/15/Z). The Malawi-Liverpool-Wellcome Programme is core-funded by a grant from Wellcome (206545/Z/17/Z). Melita Gordon is supported by a Research Professorship from the UK Department of Health National Institute of Health Research (NIHR300039)

468 References
469

1. Stanaway JD, Reiner RC, Blacker BF, Goldberg EM, Khalil IA, Troeger CE, et al. The
global burden of typhoid and paratyphoid fevers: a systematic analysis for the Global
Burden of Disease Study 2017. The Lancet Infectious Diseases. 2019;19: 369–381.
doi:10.1016/S1473-3099(18)30685-6

2. Stanaway JD, Parisi A, Sarkar K, Blacker BF, Reiner RC, Hay SI, et al. The global
burden of non-typhoidal salmonella invasive disease: a systematic analysis for the
Global Burden of Disease Study 2017. The Lancet Infectious Diseases. 2019;19: 1312–
1324. doi:10.1016/S1473-3099(19)30418-9

3. Feasey NA, Gaskell K, Wong V, Msefula C, Selemani G, Kumwenda S, et al. Rapid
Emergence of Multidrug Resistant, H58-Lineage Salmonella Typhi in Blantyre, Malawi.
PLOS Neglected Tropical Diseases. 2015;9: e0003748.
doi:10.1371/journal.pntd.0003748

4. Hendriksen RS, Leekitcharoenphon P, Lukjancenko O, Lukwesa-Musyani C,
Tambatamba B, Mwaba J, et al. Genomic Signature of Multidrug-Resistant Salmonella
enterica Serovar Typhi Isolates Related to a Massive Outbreak in Zambia between 2010
and 2012. Munson E, editor. J Clin Microbiol. 2015;53: 262–272.
doi:10.1128/JCM.02026-14

5. Marks F, von Kalckreuth V, Aaby P, Adu-Sarkodie Y, El Tayeb MA, Ali M, et al.
Incidence of invasive salmonella disease in sub-Saharan Africa: a multicentre
population-based surveillance study. Lancet Glob Health. 2017;5: e310–e323.
doi:10.1016/S2214-109X(17)30022-0

6. Neil KP, Sodha SV, Lukwago L, O-tipo S, Mikoleit M, Simington SD, et al. A Large
Outbreak of Typhoid Fever Associated With a High Rate of Intestinal Perforation in
Kasese District, Uganda, 2008–2009. Clinical Infectious Diseases. 2012;54: 1091–1099.
doi:10.1093/cid/cis025

7. N'cho HS, Masunda KPE, Mukeredzi I, Manangazira P, Govore E, Duri C, et al. *Notes*
from the Field: Typhoid Fever Outbreak — Harare, Zimbabwe, October 2017–February
2018. MMWR Morb Mortal Wkly Rep. 2019;68: 44–45.
doi:10.15585/mmwr.mm6802a5

8. Meiring JE, Shakya M, Khanam F, Voysey M, Phillips MT, Tonks S, et al. Burden of
enteric fever at three urban sites in Africa and Asia: a multicentre population-based
study. The Lancet Global Health. 2021;9: e1688–e1696. doi:10.1016/S2214-
109X(21)00370-3

9. Gauld JS, Olgemoeller F, Nkhata R, Li C, Chirambo A, Morse T, et al. Domestic River
Water Use and Risk of Typhoid Fever: Results From a Case-control Study in Blantyre,
Malawi. Clin Infect Dis. 2020;70: 1278–1284. doi:10.1093/cid/ciz405

10. Phelps MD, Azman AS, Lewnard JA, Antillón M, Simonsen L, Andreasen V, et al. The
importance of thinking beyond the water-supply in cholera epidemics: A historical
urban case-study. PLOS Neglected Tropical Diseases. 2017;11: e0006103.
doi:10.1371/journal.pntd.0006103

11. Gilks CF, Brindle RJ, Newnham RS, Watkins WM, Waiyaki PG, Were JBO, et al. Life-
threatening bacteraemia in HIV-1 seropositive adults admitted to hospital in Nairobi,
Kenya. The Lancet. 1990;336: 545–549. doi:10.1016/0140-6736(90)92096-Z

12. Gordon MA, Walsh AL, Chaponda M, Soko D, Mbvwinji M, Molyneux ME, et al.
Bacteraemia and Mortality Among Adult Medical Admissions in Malawi –
Predominance of Non-typhi Salmonellae andStreptococcus pneumoniae. Journal of
Infection. 2001;42: 44–49. doi:10.1053/jinf.2000.0779

13. Berkley JA, Lowe BS, Mwangi I, Williams T, Bauni E, Mwarumba S, et al. Bacteremia
among Children Admitted to a Rural Hospital in Kenya. New England Journal of
Medicine. 2005;352: 39–47. doi:10.1056/NEJMoa040275

14. Reddy EA, Shaw AV, Crump JA. Community-acquired bloodstream infections in
Africa: a systematic review and meta-analysis. The Lancet Infectious Diseases. 2010;10:
417–432. doi:10.1016/S1473-3099(10)70072-4

15. Feasey NA, Dougan G, Kingsley RA, Heyderman RS, Gordon MA. Invasive non-
typhoidal salmonella disease: an emerging and neglected tropical disease in Africa. The
Lancet. 2012;379: 2489–2499. doi:10.1016/S0140-6736(11)61752-2

16. Gordon MA, Graham SM, Walsh AL, Wilson L, Phiri A, Molyneux E, et al. Epidemics
of Invasive Salmonella enterica Serovar Enteritidis and S. enterica Serovar
Typhimurium Infection Associated with Multidrug Resistance among Adults and
Children in Malawi. Clin Infect Dis. 2008;46: 963–969. doi:10.1086/529146

17. Feasey NA, Masesa C, Jassi C, Faragher EB, Mallewa J, Mallewa M, et al. Three
Epidemics of Invasive Multidrug-Resistant Salmonella Bloodstream Infection in
Blantyre, Malawi, 1998–2014. Clin Infect Dis. 2015;61: S363–S371.
doi:10.1093/cid/civ691

18. Keddy KH, Musekiwa A, Sooka A, Karstaedt A, Nana T, Seetharam S, et al. Clinical
and microbiological features of invasive nontyphoidal Salmonella associated with HIV-
infected patients, Gauteng Province, South Africa. Medicine (Baltimore). 2017;96.
doi:10.1097/MD.0000000000006448

19. Akullian A, Montgomery JM, John-Stewart G, Miller SI, Hayden HS, Radey MC, et al.
Multi-drug resistant non-typhoidal Salmonella associated with invasive disease in
western Kenya. PLOS Neglected Tropical Diseases. 2018;12: e0006156.
doi:10.1371/journal.pntd.0006156

20. Mandomando I, Macete E, Sigaúque B, Morais L, Quintó L, Sacarlal J, et al. Invasive
non-typhoidal Salmonella in Mozambican children. Trop Med Int Health. 2009;14:
1467–1474. doi:10.1111/j.1365-3156.2009.02399.x

21. Tapia MD, Tennant SM, Bornstein K, Onwuchekwa U, Tamboura B, Maiga A, et al.
Invasive Nontyphoidal Salmonella Infections Among Children in Mali, 2002–2014:
Microbiological and Epidemiologic Features Guide Vaccine Development. Clin Infect
Dis. 2015;61: S332–S338. doi:10.1093/cid/civ729

22. Kingsley RA, Msefula CL, Thomson NR, Kariuki S, Holt KE, Gordon MA, et al.
Epidemic multiple drug resistant Salmonella Typhimurium causing invasive disease in

sub-Saharan Africa have a distinct genotype. Genome Res. 2009;19: 2279–2287.
doi:10.1101/gr.091017.109

23.   Okoro CK, Kingsley RA, Connor TR, Harris SR, Parry CM, Al-Mashhadani MN, et al.
554   Intra-continental spread of human invasive Salmonella Typhimurium pathovariants in
555   sub-Saharan Africa. Nat Genet. 2012;44: 1215–1221. doi:10.1038/ng.2423

556   24.   Feasey NA, Hadfield J, Keddy KH, Dallman TJ, Jacobs J, Deng X, et al. Distinct
557   Salmonella Enteritidis lineages associated with enterocolitis in high-income settings and
558   invasive disease in low-income settings. Nat Genet. 2016;48: 1211–1217.
559   doi:10.1038/ng.3644

560   25.   Morpeth SC, Ramadhani HO, Crump JA. Invasive non-Typhi Salmonella disease in
561   Africa. Clin Infect Dis. 2009;49: 606–611. doi:10.1086/603553

562   26.   Tack B, Vita D, Phoba M-F, Mbuyi-Kalonji L, Hardy L, Barbé B, et al. Direct
563   association between rainfall and non-typhoidal Salmonella bloodstream infections in
564   hospital-admitted children in the Democratic Republic of Congo. Sci Rep. 2021;11:
565   21617. doi:10.1038/s41598-021-01030-x

566   27.   Gelaw AK, Nthaba P, Matle I. Detection of Salmonella from animal sources in South
567   Africa between 2007 and 2014. Journal of the South African Veterinary Association.
568   2018;89. doi:10.4102/jsava.v89i0.1643

569   28.   Thomas KM, de Glanville WA, Barker GC, Benschop J, Buza JJ, Cleaveland S, et al.
570   Prevalence of Campylobacter and Salmonella in African food animals and meat: A
571   systematic review and meta-analysis. International Journal of Food Microbiology.
572   2020;315: 108382. doi:10.1016/j.ijfoodmicro.2019.108382

573   29.   Kariuki S, Revathi G, Kariuki N, Kiiru J, Mwituria J, Muyodi J, et al. Invasive
574   multidrug-resistant non-typhoidal Salmonella infections in Africa: zoonotic or
575   anthroponotic transmission? J Med Microbiol. 2006;55: 585–591.
576   doi:10.1099/jmm.0.46375-0

577   30.   Kariuki S, Mbae C, Puyvelde SV, Onsare R, Kavai S, Wairimu C, et al. High
578   relatedness of invasive multi-drug resistant non-typhoidal Salmonella genotypes among
579   patients and asymptomatic carriers in endemic informal settlements in Kenya. PLOS
580   Neglected Tropical Diseases. 2020;14: e0008440. doi:10.1371/journal.pntd.0008440

581   31.   Kasumba IN, Pulford CV, Perez-Sepulveda BM, Sen S, Sayed N, Permala-Booth J, et
582   al. Characteristics of Salmonella recovered from stools of children enrolled in the
583   Global Enteric Multicenter Study. Clinical Infectious Diseases. 2021 [cited 10 Feb
584   2021]. doi:10.1093/cid/ciab051

585   32.   Dione MM, Ikumapayi UN, Saha D, Mohammed NI, Geerts S, Ieven M, et al. Clonal
586   Differences between Non-Typhoidal Salmonella (NTS) Recovered from Children and
587   Animals Living in Close Contact in The Gambia. PLOS Neglected Tropical Diseases.
588   2011;5: e1148. doi:10.1371/journal.pntd.0001148

589   33.   Crump JA, Thomas KM, Benschop J, Knox MA, Wilkinson DA, Midwinter AC, et al.
590   Investigating the Meat Pathway as a Source of Human Nontyphoidal Salmonella

Bloodstream Infections and Diarrhea in East Africa. Clinical Infectious Diseases. 2020 [cited 10 Feb 2021]. doi:10.1093/cid/ciaa1153

34. Wilson CN, Pulford CV, Akoko J, Sepulveda BP, Predeus AV, Bevington J, et al. Salmonella identified in pigs in Kenya and Malawi reveals the potential for zoonotic transmission in emerging pork markets. PLOS Neglected Tropical Diseases. 2020;14: e0008796. doi:10.1371/journal.pntd.0008796

35. Ashton PM, Owen SV, Kaindama L, Rowe WPM, Lane CR, Larkin L, et al. Public health surveillance in the UK revolutionises our understanding of the invasive Salmonella Typhimurium epidemic in Africa. Genome Medicine. 2017;9: 92. doi:10.1186/s13073-017-0480-7

36. Parkhill J, Dougan G, James KD, Thomson NR, Pickard D, Wain J, et al. Complete genome sequence of a multiple drug resistant Salmonella enterica serovar Typhi CT18. Nature. 2001;413: 848–852. doi:10.1038/35101607

37. McClelland M, Sanderson KE, Clifton SW, Latreille P, Porwollik S, Sabo A, et al. Comparison of genome degradation in Paratyphi A and Typhi, human-restricted serovars of Salmonella enterica that cause typhoid. Nature Genetics. 2004;36: 1268–1274. doi:10.1038/ng1470

38. Aulicino A, Rue-Albrecht KC, Preciado-Llanes L, Napolitani G, Ashley N, Cribbs A, et al. Invasive Salmonella exploits divergent immune evasion strategies in infected and bystander dendritic cell subsets. Nat Commun. 2018;9: 4883. doi:10.1038/s41467-018-07329-0

39. Hammarlöf DL, Kröger C, Owen SV, Canals R, Lacharme-Lora L, Wenner N, et al. Role of a single noncoding nucleotide in the evolution of an epidemic African clade of Salmonella. PNAS. 2018;115: E2614–E2623. doi:10.1073/pnas.1714718115

40. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal; Vol 17, No 1: Next Generation Sequencing Data Analysis. 2011. Available: http://journal.embnet.org/index.php/embnetjournal/article/view/200/479

41. Joshi N, Fass J. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files. 2011. Available: https://github.com/najoshi/sickle

42. Alikhan N-F, Zhou Z, Sergeant MJ, Achtman M. A genomic overview of the population structure of Salmonella. PLOS Genetics. 2018;14: e1007261. doi:10.1371/journal.pgen.1007261

43. Zhou Z, Alikhan N-F, Mohamed K, Fan Y, Group the AS, Achtman M, et al. The EnteroBase user's guide, with case studies on Salmonella transmissions, Yersinia pestis phylogeny, and Escherichia core genomic diversity. Genome Res. 2020;30: 138–152. doi:10.1101/gr.251678.119

44. Creating a SNP Tree — EnteroBase documentation. [cited 26 Jan 2022]. Available: https://enterobase.readthedocs.io/en/latest/enterobase-tutorials/snp-tree-tutorial.html

45. Zhou Z. EToKi (Enterobase Tool Kit). 2022. Available: https://github.com/zheminzhou/EToKi

46.   Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast, scalable
and user-friendly tool for maximum likelihood phylogenetic inference. Bioinformatics.
2019;35: 4453–4455. doi:10.1093/bioinformatics/btz305

47.   Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I, et al. Validating the
AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance
Genotype-Phenotype Correlations in a Collection of Isolates. Antimicrob Agents
Chemother. 2019;63. doi:10.1128/AAC.00483-19

48.   Mbae C, Mwangi M, Gitau N, Irungu T, Muendo F, Wakio Z, et al. Factors associated
with occurrence of salmonellosis among children living in Mukuru slum, an urban
informal settlement in Kenya. BMC Infectious Diseases. 2020;20: 422.
doi:10.1186/s12879-020-05134-z

49.   Post AS, Diallo SN, Guiraud I, Lompo P, Tahita MC, Maltha J, et al. Supporting
evidence for a human reservoir of invasive non-Typhoidal Salmonella from household
samples in Burkina Faso. PLOS Neglected Tropical Diseases. 2019;13: e0007782.
doi:10.1371/journal.pntd.0007782

50.   Berghaus RD, Thayer SG, Law BF, Mild RM, Hofacre CL, Singer RS. Enumeration of
Salmonella and Campylobacter spp. in Environmental Farm Samples and Processing
Plant Carcass Rinses from Commercial Broiler Chicken Flocks. Appl Environ
Microbiol. 2013;79: 4106–4114. doi:10.1128/AEM.00836-13

51.   Rigby J, Elmerhebi E, Diness Y, Mkwanda C, Tonthola K, Galloway H, et al.
Optimized methods for detecting Salmonella Typhi in the environment using validated
field sampling, culture and confirmatory molecular approaches. Journal of Applied
Microbiology. 2021;132: 1503–1517. doi:10.1111/jam.15237

52.   Pulford CV, Perez-Sepulveda BM, Canals R, Bevington JA, Bengtsson RJ, Wenner N,
et al. Stepwise evolution of Salmonella Typhimurium ST313 causing bloodstream
infection in Africa. Nature Microbiology. 2020; 1–12. doi:10.1038/s41564-020-00836-1

53.   Neuert S, Nair S, Day MR, Doumith M, Ashton PM, Mellor KC, et al. Prediction of
Phenotypic Antimicrobial Resistance Profiles From Whole Genome Sequences of Non-
typhoidal Salmonella enterica. Front Microbiol. 2018;9. doi:10.3389/fmicb.2018.00592

54.   Guidelines for Foodborne Disease Outbreak Response. Council to Improve Foodborne
Outbreak Response; 2009.

55.   Adak GK, Meakins SM, Yip H, Lopman BA, O'Brien SJ. Disease Risks from Foods,
England and Wales, 1996–2000. Emerg Infect Dis. 2005;11: 365–372.
doi:10.3201/eid1103.040191

56.   Crump JA, Sjölund-Karlsson M, Gordon MA, Parry CM. Epidemiology, Clinical
Presentation, Laboratory Diagnosis, Antimicrobial Resistance, and Antimicrobial
Management of Invasive Salmonella Infections. Clin Microbiol Rev. 2015;28: 901–937.
doi:10.1128/CMR.00002-15