Case-control investigation of invasive Salmonella disease in Africa reveals no

evidence of environmental or animal reservoirs of invasive strains

Leonard Koolman[1,2 #], Reenesh Prakash[1,2 #], Yohane Diness[1], Chisomo Msefula[4],

Tonney S. Nyirenda[4], Franziska Olgemoeller[1,3], Paul Wigley[2], Blanca Perez-

Sepulveda[2], Jay C. D. Hinton[2], Siân V. Owen[2], Nicholas A. Feasey[1,3], Philip M.

Ashton[1,2*], Melita A. Gordon[1,2,4]

[1] Malawi-Liverpool Wellcome Programme, Blantyre, Malawi

[2] Institute of Infection, Veterinary & Ecological Sciences, University of Liverpool,

United Kingdom

[3] Department of Clinical Sciences, Liverpool School of Tropical Medicine, United

Kingdom

[4] Kamuzu University of Health Sciences, Blantyre, Malawi

# contributed equally

*Corresponding author

Email: philip.ashton@liverpool.ac.uk

Abstract

Background

Invasive *Salmonella* infections are a major cause of morbidity and mortality in Sub-Saharan Africa (SSA). However, the sources and modes of transmission are uncertain. To shed light on potential transmission routes of invasive Salmonellosis, we obtained *Salmonella* isolates from healthy people, animals, and the environment of index-cases and geographically-matched control households.

Methodology

Sixty index cases of human invasive *Salmonella* infection were recruited. Twenty eight invasive Non-Typhoidal *Salmonella* (iNTS) disease and 32 typhoid patients consented to household sampling. Each index-case household was geographically matched to a control household. Extensive microbiological sampling included stool sampling from healthy household members, stool or rectal swabs from household-associated animals and boot-sock sampling of the household environment.

Findings

1203 samples were taken from 120 households, yielding 43 non-Typhoidal *Salmonella* (NTS) isolates from 25 households (overall sample positivity 3.6%). In the 28 iNTS patients, disease was caused by 3 STs of Salmonella Typhimurium, mainly ST313. In contrast, the isolates from households spanned 15 sequence types (STs). Two *S.* Typhimurium isolates from index cases closely matched isolates from their respective asymptomatic household members (2 and 3 SNP differences respectively). Despite the recovery of a diverse range of NTS, there was no overlap between the STs causing iNTS disease and any environmental or animal isolates,

Conclusions

The finding of NTS strains from index cases that matched household members, coupled with lack of related animal or environmental isolates, supports a hypothesis that healthy humans were the source of iNTS infections in the household. The breadth of NTS strains found in the household environment across all sites demonstrated the robustness of NTS sampling methodology, and suggests a diverse ecology of *Salmonella* in this setting. The lack of *S.* Typhi isolates from the household environment suggests that further methodological development may be needed to culture *S.* Typhi from the environment.

Author summary

Invasive *Salmonella* infections cause the loss of millions of disability adjusted life years (DALYs) every year globally. The two main types of invasive *Salmonella* infections in Africa are i) typhoid fever, caused by *Salmonella* Typhi, and ii) invasive Non-Typhoidal Salmonella (iNTS) disease, primarily caused in our setting by *Salmonella* Typhimurium. Despite the high disease burden, and the observed differences between the epidemiology of typhoid and iNTS disease, we lack an understanding of the reservoirs and transmission routes of iNTS. Therefore, we carried out extensive microbiological sampling of the household members and living environments of patients with invasive *Salmonella* infections, and of geographically-matched control households, and investigated the genetic relationships between household *Salmonella* and index-case blood-stream isolates by whole genome sequencing (WGS). We only identified *Salmonella* that matched iNTS-associated strains in the stool of healthy people from the same households. Our findings suggest that either iNTS-associated strains are spread person-to-person, or the infections arose from a common source. Boot-sock sampling of the household environment gave the highest yield of *Salmonella* of any of our sampling strategies. None of the 41 environmental *Salmonella* isolates from non-human sources, including 4 domestic animal-associated isolates, matched the disease-causing sequence types. Our findings are consistent with the reservoir of Typhimurium iNTS infections being the human gastrointestinal tract, and transmission occurring within households.

Introduction

Salmonella infections are a substantial cause of morbidity and mortality in many low- and middle-income countries and are an important cause of both enteric infections and invasive bloodstream infections [1,2]. Both typhoidal and Non-Typhoidal *Salmonella* (NTS) contribute extensively to the burden of BSI. Globally in 2017, invasive NTS (iNTS) disease was estimated to cause 535,000 illnesses and 77,500 deaths [2]. Here, the term invasive refers to the ability to cross the gut barrier to infect areas of the human body that are normally sterile. Sub-Saharan Africa accounts for 85.8% of global iNTS deaths [2]. There were an estimated 14.3 million cases of typhoid and paratyphoid fever in 2017, resulting in 135,900 deaths, 15.8% of which were in Sub-Saharan Africa [1].

Salmonella Typhi is a major cause of bloodstream infections in southern and eastern Africa, with multiple outbreak reports since 2012 [3–7], and Malawi has a very high incidence of 444/100,000 person-years of observation [8]. The most important pathways by which people are exposed to *S*. Typhi are uncertain [9]. Improved understanding of the relative importance of different transmission paths would help with the design of effective interventions, but delineating this for long (i.e. waterborne or environmentally mediated) and short cycle (i.e. within household) transmission routes has remained difficult for a variety of faecal oral pathogens [9,10].

In High Income Countries (HICs), NTS are predominantly associated with self-limiting enterocolitis. In Sub-Saharan Africa (SSA), however, the epidemiological picture is dominated by iNTS disease [11–16], diagnosed by positive blood-culture, which carries an estimated case-fatality of 14.7% [17]. In Africa, iNTS is particularly

associated with HIV in adults and malaria infections, malnutrition or HIV in young

children [13,15,18,19]. Two *Salmonella* serotypes, Typhimurium and Enteritidis, have

been reported to be the most common causes of iNTS in Malawi, South Africa, Kenya,

Mozambique and Mali, with *S.* Typhimurium Sequence Type (ST) 313 (ST313) and S.

Enteritidis ST11 being the most frequently identified [19–27].

While NTS infection in HICs is generally associated with zoonotic transmission, the

reservoirs and transmission route of the pathogens responsible for iNTS in SSA

remains unknown. Seasonal patterns of iNTS, and the close domestic association of

humans and animals in parts of SSA suggest that environmental factors could play a

role [28,29]. Serotypes associated with iNTS have been isolated from animals in

South Africa, although no genetic characterisation was carried out on these samples

[30,31]. In contrast, microbiological investigations in Kenya, based on WGS, found

isolates in the stool of household members that were closely related to index cases

[32]. An early study from Kenya using PFGE found that iNTS linked serotypes were

only rarely found in environmental samples such as water from nearby rivers,

household and market vendor food, or household animals [33].

The Global Enteric Multicentre Study found iNTS associated sub-types of *Salmonella*

in the stool of 42 children with diarrhoea and 17 healthy children, highlighting the

potential role of the human gut as a source of iNTS infections [34]. In The Gambia, no

overlap was observed between STs from iNTS cases and those from rectal swabs

from domestic animals from the households of cases [35], while in Tanzania and

Kenya, extensive sampling of "meat pathway" isolates and human disease isolates

only isolated ST313 from human samples [36]. Although ST313 was isolated from pigs

in Nairobi, this was related to an ST313 variant from the UK that is not associated with

invasive disease [37,38].

To address the knowledge gap concerning the reservoir of Salmonella strains

responsible for invasive *Salmonella* infections, we conducted extensive human,

animal, and household environmental sampling of iNTS and typhoid index-case

households and geographically matched control households in Blantyre, Malawi. This

study is the first to combine extensive bootsock environmental sampling, sampling of

humans and animals, and WGS of the resulting isolates to investigated the relationship

between isolates causing invasive disease with isolates from household sources.

Methods

Setting

Queen Elizabeth Central Hospital (QECH) provides free healthcare to the approximately 1.3 million inhabitants of Blantyre District, Malawi, and is the tertiary referral hospital for the southern region of Malawi. Since 1998, the Malawi-Liverpool-Wellcome Trust Clinical Research Programme has provided a quality assured diagnostic blood culture service for febrile adult and paediatric medical patients admitted to QECH. This service is provided for admitted adults (>16 years old) with axillary temperature over 37.5°C or clinical suspicion of sepsis, and for children (<16 years old) who were malaria slide negative, or positive and critically ill, or with clinical suspicion of sepsis.

Case and control recruitment

Sequential blood culture confirmed adult or paediatric cases with invasive *Salmonella* infections (either iNTS or typhoid fever) presenting at QECH between March 2015 and October 2016 were recruited. Patients living outside the Blantyre district and those with recurring iNTS disease were excluded. Index case households were sampled within a maximum of 14 days following initial presentation to QECH. Following recruitment, the field team visited the index cases in their households, where GPS co-ordinates were taken and a household socio-demographic questionnaire was completed. Control households were then selected by random bottle-spin and pacing 100 m from the index case household, recruited with informed consent, and GPS and questionnaire data were also collected. Exclusion criteria for control households were current treatment for invasive *Salmonella* disease for any family member, or declining consent. In this event, the next-nearest house in the same direction was selected.

Sampling Methodology

We carried out a microbiological survey of the index-case household and the control household, comprising stool from household members, stool or rectal swabs from domestic and household-associated animals and systematic boot-sock sampling of the living environment (latrine, rubbish area, bedroom, cooking area and the house perimeter).

Stool (humans): Samples were collected from all available family members in the index case and control households. Sample containers were provided 24-48 hours before the main sample-collection visit, and household members were requested to collect up to three consecutive stools from each individual. The three samples belonging to each person were pooled, to maximise recovery of *Salmonella*.

Stool (animals): Fresh stool or rectal swab samples were collected from domestic or household-associated animals (cows, chickens, goats, pigs, dogs, cats, rats, doves, guinea pigs or gecko lizards).

Environmental: The investigator's shoe was first covered with a waterproof sterile bag to prevent cross-contamination, then a sterile bootsock was put over this, moistened with sterile water, and single bootsocks were used to collect samples from each of the follow sites: pit-latrine, outdoor rubbish area, cooking and bedroom areas, and from the outside perimeter of the household, using a standardised protocol for location and number of paces (10 paces), and a step and twist motion. Bootsocks were then removed and placed into a sterile plastic bag for transport.

All samples were transferred from the field to the laboratory in cool boxes and processed on the same day.

Sample Enrichment and culture

Supplementary Figure 1 shows a schematic of sample processing. A portion (~1g) of stool was emulsified in 9ml of Selenite F selective enrichment broth (Oxoid, UK) and incubated at 37°C for 18-24 hours. For samples other than stool (i.e. bootsocks, swabs), we used a pre-enrichment step; these samples were incubated in Buffered Peptone Water (BPW) at 37°C for 18-24 hours. One ml of the BPW sample was then transferred into 9 ml Selenite broth and incubated for a further 18-24 hours at 37°C.

After incubation, 100µl of bacterial culture taken from the surface of the Selenite broth was sub-cultured onto Xylose Lysine Deoxycholate (XLD) agar, incubated at 37°C for 18-24 hours and examined for suspected colonies of *Salmonella* (red/pink colonies with black centres). Pure candidate colonies were biochemically confirmed as *Salmonella* species using API 20E (BioMerieux). Serotyping was performed by the slide agglutination technique. Putative *Salmonella* isolates were sub-cultured onto Nutrient agar (Oxoid, UK) and incubated for 24 hours at 37°C. Following growth, a drop of 0.85% saline was placed onto a microscope slide and a single colony from the Nutrient agar was emulsified into the saline. Positive serotyping was confirmed if agglutination occurred in antigens O4, O9, Vi, Hd, Hg, Hi, Hm. If an organism was positive for Vi antigen but negative for O9 antigen, a dense suspension of the organism was prepared in 0.85% saline, autoclaved at 121°C for 15 minutes and the agglutination repeated. *S.* Typhi was differentiated from NTS isolates if agglutination occurred for O9, Vi and Hd antigens. All API 20E confirmed *Salmonella* isolates which were not *S.* Typhi were sent to the University of Liverpool, UK, for WGS.

Whole Genome Sequencing (WGS)

Bacterial genomic DNA was extracted from overnight cultures in LB (1% tryptone, 0.5% yeast extract, 0.5% NaCl; pH 7.0) using the DNeasy Blood and Tissue kit (Qiagen), as per the Gram-negative bacteria protocol for the Quick-DNA™ Universal Kit (Zymo Research), Biological Fluids & Cells protocol. Sequencing libraries were constructed using the TruSeq DNA PCR-free library preparation kit (Illumina) or the TruSeq Nano DNA HT library preparation kit (Illumina) using a target insert size of 550bp. Libraries were sequenced on an Illumina MiSeq instrument using a 2x250bp paired end protocol at the Centre for Genomic Research, University of Liverpool, UK. Raw sequencing reads were trimmed to remove Illumina adapter sequences using Cutadapt [44] and further trimmed to remove poor quality sequence using Sickle [45] version 1.2 with a minimum window quality score of 20, discarding reads which were less than 10 bp after trimming. Trimmed reads were submitted to Enterobase (https://enterobase.warwick.ac.uk/) for draft genome assembly, MLST assignment, *in silico* serotype prediction and phylogenetic analysis [46]. SNP identification within Enterobase was carried out via a mapping and SNP calling pipeline, and phylogenetic tree construction done with RAxML [47–50] Determinants of antimicrobial resistance genes were identified using the amr-finder-plus software version 3.9.8, using database version 2020-12-17.1 [51].

Data availability

All sequencing data is available from ENA in the BioProject PRJNA818086. Per sample BioSample accessions are listed in Supplementary Table 1.

Ethics

235 We sought written informed consent from adults, and guardian consent for minors.

236 Ethical permission for this study was granted by the Univeristy of Malawi College of

237 Medicine Research Ethics Committee, application number COMREC P.08/14/1617.

Results

Sixty-seven index-case households (i.e. the households of people diagnosed with invasive Salmonelloses) associated with 35 typhoid and 32 iNTS patients were eligible for enrolment between March 2015 and October 2016. Seven index-case households declined to participate, leaving 32 typhoid index-case households with 32 paired control households, and 28 iNTS index case households, with 28 paired control households recruited for the study, giving a total of 120 households. All households were in high-density low-income housing locations within Blantyre.

Isolation of Salmonella by culture: Table 1 shows sampling and positivity rates for *Salmonella* by sample-type and by household. We collected a total of 1203 samples from the 120 households (a mean of 10 samples per household), yielding 43 isolates (all NTS) which were confirmed as *Salmonella* by WGS, giving an overall sample positivity of 3.6% (Table 1). Overall, sample positivity was similar for human, animal and bootsock environment samples (human stool 16/491, 3.3%; animals 4/110, 3.6%; bootsocks 23/620, 3.8%), and across different household categories (Table 1). Animal sampling was from 110 animals in 71/120 (59%) households, and included both domesticated and non-domesticated species (cows, chickens, pigs, dogs, cats, rats, doves, guinea pigs and geckos). *Salmonella* bacteria were isolated from 25/120 (21%) households. However, no *S.* Typhi were isolated from any category of household samples.

Table 1: Sources of *Salmonella* isolates from case and paired control households linked to iNTS and typhoid patients. HH stands for household, and an index case refers a diagnosed case of invasive *Salmonellosis*.

	iNTS Index Case Households (HH) (n=28)						iNTS Control Households (HH) (n=28)					
	Total samples	No. samples with Salmonella	% positive samples	No. HH tested	No. HH with Salmonella	% Positive HH	Total samples	No. samples with Salmonella	% positive samples	No. HH tested	No. HH with Salmonella	% Positive HH
Human Stool	116	9	7.8	28	4	14.3	97	0	0.0	28	0	0.0
Animal stool/swab	27	0	0.0	20	0	0.0	38	0	0.0	17	0	0.0
HH environment	137	0	0.0	28	0	0.0	131	6	4.6	28	3	10.7
Total	**280**	**9**	**3.2**	**28.0**	**4**	14.3	**266**	**6**	**2.3**	**28**	**3**	**10.7**
	Typhoid Index Case households (HH) (n=32)						Typhoid Control households (HH) (n=32)					
Human Stool	157	4	2.5	32	2	6.3	121	3	2.5	32	3	9.4
Animal stool/swab	22	2	9.1	18	2	11.1	23	2	8.7	16	2	12.5
HH environment	168	5	3.0	32	4	12.5	166	12	7.2	32	9	28.1
Total	**347**	**11**	**3.2**	**32**	**7**	**21.9**	**310**	**17**	**5.5**	**32**	**11**	**34.4**

 Relatedness of Sequence Types from different sample categories.

Figure 1 demonstrates the wide diversity and relationships of Sequence Types (STs)

isolated from different sample categories (invasive disease, healthy human stool,

animals, and the household environment). Table 2 shows the individual household

origins of each isolate, and the relationship between ST and serovar for each isolate.

Among invasive NTS disease cases, 8 of 27 index case NTS isolates could not be

resuscitated from the freezer archive and so could not be genome-sequenced. The

remaining 19 typed index NTS isolates were all *S.* Typhimurium and included just 3

STs; 13 were ST313, 5 were ST3257 and 1 was ST19. ST3257 varies from ST313

by only a single MLST locus. One representative typhoid index strain (ST2) was also

included for reference. Among healthy household members, 16 isolates represented

8 STs; among animals there were 4 isolates from 3 STs; and from environmental

samples there were 23 isolates from 9 different STs.

Table 2: Characteristics of isolates from index cases of invasive disease and and household isolates

Household (HH) ID S=case C=control	Index case NTS or typhoid?	Household case or control?	Household sample type	Sequence Type (ST)	Serovar	Serogroup	Match to Index case blood culture?	Household (HH) ID S=case HH C=control HH
1C	Typhoid	Control	Animal stool	3261	Agoueve|Cubana	13,22|1,13,23		1C
2S	iNTS	Case	Index case blood	3257	Typhimurium	1,4,[5],12		2S
3C	Typhoid	Control	Family member stool	3261	Agoueve|Cubana	13,22|1,13,23		3C
4S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		4S
6S	Typhoid	Case	Family member stool	3261	Agoueve|Cubana	1,13,23|1,13,22		6S
6C	Typhoid	Control	Boot sock perimeter	3263	Havana|II 1,13,22:g,t:[1,5]	13,22|1,13,23		6C
6C	Typhoid	Control	Family member stool	3263	Havana|II 1,13,22:g,t:[1,5]	1,13,23|1,13,22		6C
8C	Typhoid	Control	Boot sock bedroom	3263	Havana|II 1,13,22:g,t:[1,5]	1,13,23|1,13,22		8C
9S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		9S
9S	iNTS	Case	Family member stool	3263	Havana|II 1,13,22:g,t:[1,5]	1,13,23|1,13,22		9S
9S	iNTS	Case	Family member stool	3263	Havana|II 1,13,22:g,t:[1,5]	1,13,23|1,13,22		9S
9S	iNTS	Case	Family member stool	3263	Havana|II 1,13,22:g,t:[1,5]	1,13,23|1,13,22		9S
9S	iNTS	Case	Family member stool	3263	Havana|II 1,13,22:g,t:[1,5]	1,13,23|1,13,22		9S
9S	iNTS	Case	Family member stool	3263	Havana|II 1,13,22:g,t:[1,5]	1,13,23|1,13,22		9S
9S	iNTS	Case	Family member stool	3263	Havana|II 1,13,22:g,t:[1,5]	1,13,23|1,13,22		9S
12S	iNTS	Case	Index case blood	3257	Typhimurium	1,4,[5],12		12S
14S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		14S
15S	Typhoid	Case	Animal stool (gecko)	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		15S
15S	Typhoid	Case	Boot sock perimeter	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		15S
16S	iNTS	Case	Index case blood	19	Typhimurium	1,4,[5],12		16S
16S	iNTS	Case	Family member stool	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		16S
16C	iNTS	Control	Boot sock latrine	2347	Mgulani	38		16C
16C	iNTS	Control	Boot sock cooking	2347	Mgulani	38		16C
17S	Typhoid	Case	Boot sock latrine	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		17S
18S	iNTS	Case	Index case blood	3257	Typhimurium	1,4,[5],12		18S
22C	Typhoid	Control	Boot sock latrine	3609	Djama|II 42:z29:-	1,42|42		22C
23S	Typhoid	Case	Animal stool (gecko)	316	Montevideo	6,7,14,[54]		23S
25S	iNTS	Case	Family member stool	313	Typhimurium	1,4,[5],12	match1	25S
25S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12	match1	25S

25C	iNTS	Control	Boot sock latrine	473	Hadar	6,8		25C
25C	iNTS	Control	Boot sock bedroom	473	Hadar	6,8		25C
25C	iNTS	Control	Boot sock cooking	473	Hadar	6,8		25C
26S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		26S
27S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		27S
27C	iNTS	Control	Boot sock bedroom	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		27C
28C	Typhoid	Control	Animal rectal swab	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		28C
28C	Typhoid	Control	Boot sock latrine 1	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		28C
28C	Typhoid	Control	Boot sock latrine 1	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		28C
29C	Typhoid	Control	Boot sock latrine	2152	Gaminara	16		29C
29C	Typhoid	Control	Family member stool	3608	Ogbete|II 43:z:1,5	43		29C
31S	Typhoid	Case	Index case blood	2	Typhi	9,12[Vi]		31S
32S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		32S
33S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		33S
36S	iNTS	Case	Index case blood	3257	Typhimurium	1,4,[5],12		36S
39S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		39S
41C	Typhoid	Control	Boot sock latrine	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		41C
42S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		42S
44S	iNTS	Case	Family member stool	3257	Typhimurium	1,4,[5],12	match2	44S
44S	iNTS	Case	Index case blood	3257	Typhimurium	1,4,[5],12	match2	44S
46S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		46S
47S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		47S
48S	Typhoid	Case	Boot sock bedroom	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		48S
50S	Typhoid	Case	Boot sock bedroom	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		50S
50S	Typhoid	Case	Boot sock cooking	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		50S
50C	Typhoid	Control	Boot sock perimeter	3611	II 40:b:-	40		50C
53C	Typhoid	Control	Boot sock bedroom	3262	II 42:r:-|IIIb 42:r:-:[z50]	42		53C
56C	Typhoid	Control	Boot sock bedroom	293	Amager	3,{10}{15}		56C
56C	Typhoid	Control	Boot sock perimeter	293	Amager	3,{10}{15}		56C
56C	Typhoid	Control	Boot sock cooking	3610	Aberdeen	11		56C
57S	iNTS	Case	Index case blood	313	Typhimurium	1,4,[5],12		57S
59S	Typhoid	Case	Family member	14	Senftenberg	1,3,19		59S
59S	Typhoid	Case	Family member stool	2152	Gaminara	16		59S
59S	Typhoid	Case	Family member stool	2152	Gaminara	16		59S

There were two instances of the same sequence type of S. Typhimurium being

isolated from both invasive index-case isolates and healthy human household

samples. Specifically, 2 cases of invasive disease caused by *S.* Typhimurium ST313

and ST3257 were linked to isolates from 2 healthy household members who shared

the same address and also had ST313 and ST3257 isolated from their stools (Figure

1). In both cases the household member carrying a matching isolate was an adult,

whereas stool isolates of *Salmonella* from household members were found equally

among adults (n=7) and children (n=8).

Figure 1: Minimum spanning tree based on seven MLST genes showing the genetic similarity between different niches – the only non-clinical niche where STs associated with human invasive disease were present was the gut of household contacts. Each circle represents a sequence type, the size of the circle is proportional to the number of isolates of that ST. STs that vary by one locus are joined by a black bar. Two invasive disease index isolates had a matched isolate from a healthy human sample of the same ST from the same household. These isolates are indicated by black and red outlines.

We found no overlap in STs between isolates causing invasive human disease and

any animal or environmental isolates, despite a good rate of isolation from all 3

sample-types. We did identify non-co-localised overlap between some STs from

healthy humans and household-associated animals. A monophasic ST3262 was

found in a cat, a gecko, and a human, but in geographically distinct households.

Similarly, ST3261 was isolated from 2 healthy humans and a chicken, but were

associated with different households (Table 1). ST325 was found only in an animal

(gecko).

Some STs isolated from the environment overlapped with those from healthy humans,

and household animals. Of 9 STs found in the environment, 3 (ST3262, ST3263,

ST2152) showed some overlap with healthy human or animal isolates. However, a

further 6 STs were found only in the household environment, revealing the diversity of

Salmonella strains in this setting.

Sequence types and WGS-inferred serovars from different sample categories:

The relationship between STs and WGS-inferred serovars for all isolates in the study

is documented in Table 2 (with accession numbers in Supplementary Table 1). Of the

19 index iNTS case isolates from blood culture, 19/19 were *S.* Typhimurium, of 3

different STs; ST313 (n=13), ST3257 (a Single Locus Variant of ST313, n=5) and

ST19 (n = 1) (Supplementary Table 1, Figure 1).

Serovars that were carried by apparently healthy humans were Havana (7),

Typhimurium (2), Gaminara (2), Agoueve/Cubana (2), Seftenberg (1) and Ogbete (1).

Serovars associated with animals were II42:r:-|IIIb 42:r:-:[z50] (1 gecko, 1 cat),

Montevideo (1 gecko) and Agoueve/Cubana (1 chicken). The wide range of serovars

found in the household environment included II42:r:-|IIIb 42:r:-:[z50] (10), Hadar (3),

Havana (3), Amager (2), Mgulani (2) Djama (1) and Aberdeen (1).

Reflecting the broad diversity of isolates, many STs/serovars were represented by just

a single isolate. In addition, where there were multiple isolations of the same serovar,

a large proportion were localised from within the same household. Striking examples

of this are: of 9 isolates of *S*. Havana, 6 arose from a single household (9S), from 6

out of 7 children sampled; of 3 isolates of *S*. Hadar, all arose from the same household

(25C, latrine, bedroom and cooking area); of 12 isolates of II42:r:-|IIIb 42:r:-:[z50] there

were multiple positive samples in three households, originating from both animal and

environmental samples in the same households. Similarly, 2 isolates each of *S*.

Mgulani and *S*. Amager isolates were linked to the same respective households (16C

and 56C). This indicates a very marked geographical diversity between households,

and a degree of homogeneity within households among NTS that are not linked to

invasive disease.

Whole genome phylogeny of index/household pairs

We used whole genome-derived phylogenetics to infer the high-resolution

relationship between the human invasive disease cases of *S*. Typhimurium ST313 &

ST3257 and the stool carriage isolates of their respective index household members.

In both the ST313 and ST3257 pairs, the isolate that caused the invasive disease

formed a monophyletic group with the carriage isolate from the same index

household (Figure 2). The SNP distance was 2 SNPs in one case, and 3 SNPs in the

other (Figure 2).

Figure 2: The close relationship between the index case and family member isolates
from households 25 and 44 shown with a phylogenetic tree. Whole genome
maximum likelihood tree of 19 S. Typhimurium isolates described in Table 1. The
numbers at the end of each branch indicate household, as described in the first
column of Table 1 (all these households are case households). Sequencing read
sets were compared against the reference genome D23580 as described in
Methods.

Antimicrobial resistance genes

We searched the WGS data for 63 isolates from index cases of human invasive

disease and household sampling for antimicrobial resistance genes (Supplementary

Table 1). No antimicrobial resistance genes were carried by 52 genomes,

representing 68% blood-stream infection *S.* Typhimurium isolates (9 ST313s, 3

ST3257 and 1 ST19), and 89% of the animal and bootsock isolates. Two bootsock

isolate (both *S.* Mgulani) and one animal stool isolate (*S.* Montevideo) carried the

fosA7 gene associated with resistance to fosfomycin. Six Typhimurium isolates from

BSI and two Typhimurium from household member stool encoded all or some of the

blaTEM-1, catA1, sul1, sul2 and *dfrA1* resistance genes which are typically

associated with invasive *S.* Typhimurium in Malawi [27].

Discussion

Our findings support the hypothesis that humans are the primary reservoir for iNTS in sub-Saharan Africa. The iNTS patients carried 3 STs of *S.* Typhimurium in their bloodstream. Matching *S.* Typhimurium STs were isolated from stool of their respective household members of two cases, the isolates from household members were within a 2-3 SNP distance of the isolate from the index case. None of the 27 *Salmonella* isolates from the environment and animals could be linked to iNTS-cases.

Our findings are consistent with previous reports. Firstly, in Kenya, iNTS infections caused by *Salmonella* Typhimurium or Enteritidis in an informal urban settlement were not associated with the rearing of any domestic animals [52]. Secondly, in Burkina Faso the only isolates that matched the index cases were obtained from the stool of household members, despite extensive animal and water sampling in the households of iNTS patients [53]. Finally, analysis of cases of moderate to severe diarrhoea and healthy controls in Africa and Asia found ST313 in both sick and healthy children, supporting the finding that *Salmonella* are found in the stool of healthy children without diarrhoeal symptoms [34].

Whilst previous case-control studies have focused on adults and children in the same household, and household-associated animals, here we have additionally studied the household environment. This yielded a similar culture positivity-rate to other sample-types, and yielded a much wider range of STs/serovars. The fact that Salmonellae have a more diverse ecology in the household environment than previously appreciated in this setting contrasts with the narrow range of STs that cause human invasive disease [43]. The lack of overlap between the ecological niches of serovars

causing colonising the household and causing human bloodstream categories is very

striking.

Our methodology successfully identified microbiological connections between people

and their household environments at both the individual household level and the

community level. The presence of the same STs in multiple sample types from the

same household indicates that our methodology can identify this connectivity when it

is present in this setting, this strengthens our negative findings of no link between

Salmonella from the environment and from iNTS cases. This excellent yield from

bootsocks for detecting *Salmonella* in other contexts has been reported elsewhere

[54], and we recommend that similar studies carried out in the future also use this

method.

S. Typhi is human-restricted in its ability to cause disease, but has a poorly defined

environment during long-cycle transmission in the community [55]. While we failed to

isolate *S.* Typhi from the environment of either case or control households, this is

known to be technically challenging, and our results highlight the need for further

methodological development in this area [56].

While antimicrobial resistance is a major concern in our setting, only 12% of our

household isolates carried one or more genes known to be associated with

antimicrobial resistance, in contrast to 39% of our human invasive disease isolates.

The fact that 62% of the ST313 isolated here carried no antimicrobial resistance genes

suggests that they could belong to the recently described ST313 Lineage 3 [27]. We

did not carry out AMR phenotyping for the isolates obtained in this study, but AMR

profile inference from genome sequence is highly accurate for *Salmonella* [57].

There are limitations to our study; both the index case and the ST313 colonised household member could have been simultaneously infected from an unsampled environmental reservoir. While we had hoped to sample food, we found that in practice most households lacked refrigeration, and there was rarely food in the house to sample, as meals were eaten at one sitting. In addition to a contaminated foodstuff, water sources, both drinking and washing/cleaning, could be involved in the transmission of organisms causing iNTS, as is seen for *S.* Typhi [9,58], but further methodological development would be required for these studies. If the index-case were infected outside the home e.g. at work, school, or nursery, which is a possible risk factor in our setting [9], then our study would not identify that microbiological source.

Conclusions

In summary, we conducted comprehensive environmental and contact sampling within the households of patients with invasive *Salmonella* disease, and geographically matched control households. The only household matched isolates NTS came from the stool of household members, whilst no *S.* Typhi were detected outwith the index typhoid case. This study contributes to accumulating evidence that the reservoir of iNTS infections in Africa is the human gastrointestinal tract. Significantly, no strains associated with human iNTS disease were identified among any household-associated animals, despite extensive recovery from animal samples. We uncovered a much wider range of NTS isolates that were not associated with invasive disease, across all sampling sites, than has previously been appreciated, consistent with a diverse ecology for *Salmonella* in the household environment.

Acknowledgements

We are grateful to Alistair Darby, John Kenny and staff at the Centre for Genomic Research at the University of Liverpool for assistance with genome sequencing. The Malawi-Liverpool-Wellcome Programme is core-funded by a grant from Wellcome (206545/Z/17/Z). This work was supported in part by a Wellcome Trust Senior Investigator award [grant number 106914/Z/15/Z] to Jay Hinton. For the purpose of open access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. Melita Gordon is supported by a Research Professorship from the UK Department of Health National Institute of Health Research (NIHR300039)

References

1. Stanaway JD, Reiner RC, Blacker BF, Goldberg EM, Khalil IA, Troeger CE, et al. The global burden of typhoid and paratyphoid fevers: a systematic analysis for the Global Burden of Disease Study 2017. The Lancet Infectious Diseases. 2019;19: 369–381. doi:10.1016/S1473-3099(18)30685-6

2. Stanaway JD, Parisi A, Sarkar K, Blacker BF, Reiner RC, Hay SI, et al. The global burden of non-typhoidal salmonella invasive disease: a systematic analysis for the Global Burden of Disease Study 2017. The Lancet Infectious Diseases. 2019;19: 1312–1324. doi:10.1016/S1473-3099(19)30418-9

3. Feasey NA, Gaskell K, Wong V, Msefula C, Selemani G, Kumwenda S, et al. Rapid Emergence of Multidrug Resistant, H58-Lineage Salmonella Typhi in Blantyre, Malawi. PLOS Neglected Tropical Diseases. 2015;9: e0003748. doi:10.1371/journal.pntd.0003748

4. Hendriksen RS, Leekitcharoenphon P, Lukjancenko O, Lukwesa-Musyani C, Tambatamba B, Mwaba J, et al. Genomic Signature of Multidrug-Resistant Salmonella enterica Serovar Typhi Isolates Related to a Massive Outbreak in Zambia between 2010 and 2012. Munson E, editor. J Clin Microbiol. 2015;53: 262–272. doi:10.1128/JCM.02026-14

5. Marks F, von Kalckreuth V, Aaby P, Adu-Sarkodie Y, El Tayeb MA, Ali M, et al. Incidence of invasive salmonella disease in sub-Saharan Africa: a multicentre population-based surveillance study. Lancet Glob Health. 2017;5: e310–e323. doi:10.1016/S2214-109X(17)30022-0

6. Neil KP, Sodha SV, Lukwago L, O-tipo S, Mikoleit M, Simington SD, et al. A Large Outbreak of Typhoid Fever Associated With a High Rate of Intestinal Perforation in Kasese District, Uganda, 2008–2009. Clinical Infectious Diseases. 2012;54: 1091–1099. doi:10.1093/cid/cis025

7. N'cho HS, Masunda KPE, Mukeredzi I, Manangazira P, Govore E, Duri C, et al. *Notes from the Field:* Typhoid Fever Outbreak — Harare, Zimbabwe, October 2017–February 2018. MMWR Morb Mortal Wkly Rep. 2019;68: 44–45. doi:10.15585/mmwr.mm6802a5

8. Meiring JE, Shakya M, Khanam F, Voysey M, Phillips MT, Tonks S, et al. Burden of enteric fever at three urban sites in Africa and Asia: a multicentre population-based study. The Lancet Global Health. 2021;9: e1688–e1696. doi:10.1016/S2214-109X(21)00370-3

9. Gauld JS, Olgemoeller F, Nkhata R, Li C, Chirambo A, Morse T, et al. Domestic River Water Use and Risk of Typhoid Fever: Results From a Case-control Study in Blantyre, Malawi. Clin Infect Dis. 2020;70: 1278–1284. doi:10.1093/cid/ciz405

10. Phelps MD, Azman AS, Lewnard JA, Antillón M, Simonsen L, Andreasen V, et al. The importance of thinking beyond the water-supply in cholera epidemics: A historical urban case-study. PLOS Neglected Tropical Diseases. 2017;11: e0006103. doi:10.1371/journal.pntd.0006103

11.   Gilks CF, Brindle RJ, Newnham RS, Watkins WM, Waiyaki PG, Were JBO, et al. Life-
threatening bacteraemia in HIV-1 seropositive adults admitted to hospital in Nairobi,
Kenya. The Lancet. 1990;336: 545–549. doi:10.1016/0140-6736(90)92096-Z

12.   Gordon MA, Walsh AL, Chaponda M, Soko D, Mbvwinji M, Molyneux ME, et al.
Bacteraemia and Mortality Among Adult Medical Admissions in Malawi –
Predominance of Non-typhi Salmonellae andStreptococcus pneumoniae. Journal of
Infection. 2001;42: 44–49. doi:10.1053/jinf.2000.0779

13.   Berkley JA, Lowe BS, Mwangi I, Williams T, Bauni E, Mwarumba S, et al. Bacteremia
among Children Admitted to a Rural Hospital in Kenya. New England Journal of
Medicine. 2005;352: 39–47. doi:10.1056/NEJMoa040275

14.   Reddy EA, Shaw AV, Crump JA. Community-acquired bloodstream infections in
Africa: a systematic review and meta-analysis. The Lancet Infectious Diseases. 2010;10:
417–432. doi:10.1016/S1473-3099(10)70072-4

15.   Feasey NA, Dougan G, Kingsley RA, Heyderman RS, Gordon MA. Invasive non-
typhoidal salmonella disease: an emerging and neglected tropical disease in Africa. The
Lancet. 2012;379: 2489–2499. doi:10.1016/S0140-6736(11)61752-2

16.   Uche IV, MacLennan CA, Saul A. A Systematic Review of the Incidence, Risk Factors
and Case Fatality Rates of Invasive Nontyphoidal Salmonella (iNTS) Disease in Africa
(1966 to 2014). PLOS Neglected Tropical Diseases. 2017;11: e0005118.
doi:10.1371/journal.pntd.0005118

17.   Marchello CS, Birkhold M, Crump JA, Martin LB, Ansah MO, Breghi G, et al.
Complications and mortality of non-typhoidal salmonella invasive disease: a global
systematic review and meta-analysis. The Lancet Infectious Diseases. 2022;0.
doi:10.1016/S1473-3099(21)00615-0

18.   Gordon MA, Graham SM, Walsh AL, Wilson L, Phiri A, Molyneux E, et al. Epidemics
of Invasive Salmonella enterica Serovar Enteritidis and S. enterica Serovar
Typhimurium Infection Associated with Multidrug Resistance among Adults and
Children in Malawi. Clin Infect Dis. 2008;46: 963–969. doi:10.1086/529146

19.   Feasey NA, Masesa C, Jassi C, Faragher EB, Mallewa J, Mallewa M, et al. Three
Epidemics of Invasive Multidrug-Resistant Salmonella Bloodstream Infection in
Blantyre, Malawi, 1998–2014. Clin Infect Dis. 2015;61: S363–S371.
doi:10.1093/cid/civ691

20.   Keddy KH, Musekiwa A, Sooka A, Karstaedt A, Nana T, Seetharam S, et al. Clinical
and microbiological features of invasive nontyphoidal Salmonella associated with HIV-
infected patients, Gauteng Province, South Africa. Medicine (Baltimore). 2017;96.
doi:10.1097/MD.0000000000006448

21.   Akullian A, Montgomery JM, John-Stewart G, Miller SI, Hayden HS, Radey MC, et al.
Multi-drug resistant non-typhoidal Salmonella associated with invasive disease in
western Kenya. PLOS Neglected Tropical Diseases. 2018;12: e0006156.
doi:10.1371/journal.pntd.0006156

22. Mandomando I, Macete E, Sigaúque B, Morais L, Quintó L, Sacarlal J, et al. Invasive
non-typhoidal Salmonella in Mozambican children. Trop Med Int Health. 2009;14:
1467–1474. doi:10.1111/j.1365-3156.2009.02399.x

23. Tapia MD, Tennant SM, Bornstein K, Onwuchekwa U, Tamboura B, Maiga A, et al.
Invasive Nontyphoidal Salmonella Infections Among Children in Mali, 2002–2014:
Microbiological and Epidemiologic Features Guide Vaccine Development. Clin Infect
Dis. 2015;61: S332–S338. doi:10.1093/cid/civ729

24. Kingsley RA, Msefula CL, Thomson NR, Kariuki S, Holt KE, Gordon MA, et al.
Epidemic multiple drug resistant Salmonella Typhimurium causing invasive disease in
sub-Saharan Africa have a distinct genotype. Genome Res. 2009;19: 2279–2287.
doi:10.1101/gr.091017.109

25. Okoro CK, Kingsley RA, Connor TR, Harris SR, Parry CM, Al-Mashhadani MN, et al.
Intra-continental spread of human invasive Salmonella Typhimurium pathovariants in
sub-Saharan Africa. Nat Genet. 2012;44: 1215–1221. doi:10.1038/ng.2423

26. Feasey NA, Hadfield J, Keddy KH, Dallman TJ, Jacobs J, Deng X, et al. Distinct
Salmonella Enteritidis lineages associated with enterocolitis in high-income settings and
invasive disease in low-income settings. Nat Genet. 2016;48: 1211–1217.
doi:10.1038/ng.3644

27. Pulford CV, Perez-Sepulveda BM, Canals R, Bevington JA, Bengtsson RJ, Wenner N,
et al. Stepwise evolution of Salmonella Typhimurium ST313 causing bloodstream
infection in Africa. Nature Microbiology. 2020; 1–12. doi:10.1038/s41564-020-00836-1

28. Morpeth SC, Ramadhani HO, Crump JA. Invasive non-Typhi Salmonella disease in
Africa. Clin Infect Dis. 2009;49: 606–611. doi:10.1086/603553

29. Tack B, Vita D, Phoba M-F, Mbuyi-Kalonji L, Hardy L, Barbé B, et al. Direct
association between rainfall and non-typhoidal Salmonella bloodstream infections in
hospital-admitted children in the Democratic Republic of Congo. Sci Rep. 2021;11:
21617. doi:10.1038/s41598-021-01030-x

30. Gelaw AK, Nthaba P, Matle I. Detection of Salmonella from animal sources in South
Africa between 2007 and 2014. Journal of the South African Veterinary Association.
2018;89. doi:10.4102/jsava.v89i0.1643

31. Thomas KM, de Glanville WA, Barker GC, Benschop J, Buza JJ, Cleaveland S, et al.
Prevalence of Campylobacter and Salmonella in African food animals and meat: A
systematic review and meta-analysis. International Journal of Food Microbiology.
2020;315: 108382. doi:10.1016/j.ijfoodmicro.2019.108382

32. Kariuki S, Mbae C, Puyvelde SV, Onsare R, Kavai S, Wairimu C, et al. High
relatedness of invasive multi-drug resistant non-typhoidal Salmonella genotypes among
patients and asymptomatic carriers in endemic informal settlements in Kenya. PLOS
Neglected Tropical Diseases. 2020;14: e0008440. doi:10.1371/journal.pntd.0008440

33. Kariuki S, Revathi G, Kariuki N, Kiiru J, Mwituria J, Muyodi J, et al. Invasive
multidrug-resistant non-typhoidal Salmonella infections in Africa: zoonotic or

anthroponotic transmission? J Med Microbiol. 2006;55: 585–591.
doi:10.1099/jmm.0.46375-0

34.   Kasumba IN, Pulford CV, Perez-Sepulveda BM, Sen S, Sayed N, Permala-Booth J, et
al. Characteristics of Salmonella recovered from stools of children enrolled in the
Global Enteric Multicenter Study. Clinical Infectious Diseases. 2021 [cited 10 Feb
2021]. doi:10.1093/cid/ciab051

35.   Dione MM, Ikumapayi UN, Saha D, Mohammed NI, Geerts S, Ieven M, et al. Clonal
Differences between Non-Typhoidal Salmonella (NTS) Recovered from Children and
Animals Living in Close Contact in The Gambia. PLOS Neglected Tropical Diseases.
2011;5: e1148. doi:10.1371/journal.pntd.0001148

36.   Crump JA, Thomas KM, Benschop J, Knox MA, Wilkinson DA, Midwinter AC, et al.
Investigating the Meat Pathway as a Source of Human Nontyphoidal Salmonella
Bloodstream Infections and Diarrhea in East Africa. Clinical Infectious Diseases. 2020
[cited 10 Feb 2021]. doi:10.1093/cid/ciaa1153

37.   Wilson CN, Pulford CV, Akoko J, Sepulveda BP, Predeus AV, Bevington J, et al.
Salmonella identified in pigs in Kenya and Malawi reveals the potential for zoonotic
transmission in emerging pork markets. PLOS Neglected Tropical Diseases. 2020;14:
e0008796. doi:10.1371/journal.pntd.0008796

38.   Ashton PM, Owen SV, Kaindama L, Rowe WPM, Lane CR, Larkin L, et al. Public
health surveillance in the UK revolutionises our understanding of the invasive
Salmonella Typhimurium epidemic in Africa. Genome Medicine. 2017;9: 92.
doi:10.1186/s13073-017-0480-7

39.   Parkhill J, Dougan G, James KD, Thomson NR, Pickard D, Wain J, et al. Complete
genome sequence of a multiple drug resistant Salmonella enterica serovar Typhi CT18.
Nature. 2001;413: 848–852. doi:10.1038/35101607

40.   McClelland M, Sanderson KE, Clifton SW, Latreille P, Porwollik S, Sabo A, et al.
Comparison of genome degradation in Paratyphi A and Typhi, human-restricted
serovars of Salmonella enterica that cause typhoid. Nature Genetics. 2004;36: 1268–
1274. doi:10.1038/ng1470

41.   Aulicino A, Rue-Albrecht KC, Preciado-Llanes L, Napolitani G, Ashley N, Cribbs A, et
al. Invasive Salmonella exploits divergent immune evasion strategies in infected and
bystander dendritic cell subsets. Nat Commun. 2018;9: 4883. doi:10.1038/s41467-018-
07329-0

42.   Hammarlöf DL, Kröger C, Owen SV, Canals R, Lacharme-Lora L, Wenner N, et al.
Role of a single noncoding nucleotide in the evolution of an epidemic African clade of
Salmonella. PNAS. 2018;115: E2614–E2623. doi:10.1073/pnas.1714718115

43.   Piccini G, Montomoli E. Pathogenic signature of invasive non-typhoidal Salmonella in
Africa: implications for vaccine development. Hum Vaccin Immunother. 16: 2056–
2071. doi:10.1080/21645515.2020.1785791

44.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
621   EMBnet.journal; Vol 17, No 1: Next Generation Sequencing Data Analysis. 2011.
622   Available: http://journal.embnet.org/index.php/embnetjournal/article/view/200/479

623   45.   Joshi N, Fass J. Sickle: A sliding-window, adaptive, quality-based trimming tool for
624   FastQ files. 2011. Available: https://github.com/najoshi/sickle

625   46.   Alikhan N-F, Zhou Z, Sergeant MJ, Achtman M. A genomic overview of the population
626   structure of Salmonella. PLOS Genetics. 2018;14: e1007261.
627   doi:10.1371/journal.pgen.1007261

628   47.   Zhou Z, Alikhan N-F, Mohamed K, Fan Y, Group the AS, Achtman M, et al. The
629   EnteroBase user's guide, with case studies on Salmonella transmissions, Yersinia pestis
630   phylogeny, and Escherichia core genomic diversity. Genome Res. 2020;30: 138–152.
631   doi:10.1101/gr.251678.119

632   48.   Creating a SNP Tree — EnteroBase documentation. [cited 26 Jan 2022]. Available:
633   https://enterobase.readthedocs.io/en/latest/enterobase-tutorials/snp-tree-tutorial.html

634   49.   Zhou Z. EToKi (Enterobase Tool Kit). 2022. Available:
635   https://github.com/zheminzhou/EToKi

636   50.   Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast, scalable
637   and user-friendly tool for maximum likelihood phylogenetic inference. Bioinformatics.
638   2019;35: 4453–4455. doi:10.1093/bioinformatics/btz305

639   51.   Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I, et al. Validating the
640   AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance
641   Genotype-Phenotype Correlations in a Collection of Isolates. Antimicrob Agents
642   Chemother. 2019;63. doi:10.1128/AAC.00483-19

643   52.   Mbae C, Mwangi M, Gitau N, Irungu T, Muendo F, Wakio Z, et al. Factors associated
644   with occurrence of salmonellosis among children living in Mukuru slum, an urban
645   informal settlement in Kenya. BMC Infectious Diseases. 2020;20: 422.
646   doi:10.1186/s12879-020-05134-z

647   53.   Post AS, Diallo SN, Guiraud I, Lompo P, Tahita MC, Maltha J, et al. Supporting
648   evidence for a human reservoir of invasive non-Typhoidal Salmonella from household
649   samples in Burkina Faso. PLOS Neglected Tropical Diseases. 2019;13: e0007782.
650   doi:10.1371/journal.pntd.0007782

651   54.   Berghaus RD, Thayer SG, Law BF, Mild RM, Hofacre CL, Singer RS. Enumeration of
652   Salmonella and Campylobacter spp. in Environmental Farm Samples and Processing
653   Plant Carcass Rinses from Commercial Broiler Chicken Flocks. Appl Environ
654   Microbiol. 2013;79: 4106–4114. doi:10.1128/AEM.00836-13

655   55.   Crump JA. Progress in Typhoid Fever Epidemiology. Clinical Infectious Diseases.
656   2019;68: S4–S9. doi:10.1093/cid/ciy846

657   56.   Rigby J, Elmerhebi E, Diness Y, Mkwanda C, Tonthola K, Galloway H, et al.
658   Optimized methods for detecting Salmonella Typhi in the environment using validated

field sampling, culture and confirmatory molecular approaches. Journal of Applied
Microbiology. 2021;132: 1503–1517. doi:10.1111/jam.15237

57. Neuert S, Nair S, Day MR, Doumith M, Ashton PM, Mellor KC, et al. Prediction of
Phenotypic Antimicrobial Resistance Profiles From Whole Genome Sequences of Non-
typhoidal Salmonella enterica. Front Microbiol. 2018;9. doi:10.3389/fmicb.2018.00592

58. Crump JA, Sjölund-Karlsson M, Gordon MA, Parry CM. Epidemiology, Clinical
Presentation, Laboratory Diagnosis, Antimicrobial Resistance, and Antimicrobial
Management of Invasive Salmonella Infections. Clin Microbiol Rev. 2015;28: 901–937.
doi:10.1128/CMR.00002-15