Identities of women who have an autoimmune rheumatic disease (ARD) during pregnancy

planning, pregnancy and early parenting: A qualitative study

Key words: Identity, motherhood, person-centred care, auto-immune rheumatic disease,

parenting, shared decision-making

Short title: Identities of women with ARD during preconception decision making.

Authors: Denitza Williams[1,6]*, Bethan Pell[2], Aimee Grant[3], Julia Sanders[4], Ann Taylor[5],

Adrian Edwards[6], Ernest Choy[7] & Rhiannon Phillips[1]

[1]Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University

[2]Centre for the Development and Evaluation of Complex Intervention for Public Health

Improvement (DECIPHer), Cardiff University, Cardiff, United Kingdom

[3]Centre for Lactation, Infant Feeding and Translational Research, Swansea University

[4]Division of Population Medicine, School of Medicine, College of Biomedical and Life

Sciences, Cardiff University.

[5]School of Healthcare Sciences, Cardiff University

[6]Centre for Medical Education, School of Medicine, Cardiff University

[7]CREATE Centre, Section of Rheumatology, Division of Infection and Immunity, Cardiff

University

[1]Rhiannon Phillips, Cardiff School of Sport and Health Sciences, Cardiff Metropolitan

University

*Corresponding Author: WilliamsD74@cardiff.ac.uk (DW)

Abstract

Objective: Women of reproductive age who have autoimmune rheumatic diseases (ARDs)

have expressed a need to be better supported with making decisions about pregnancy.

Women with ARDs want their motherhood identities and associated preferences to be

taken into account in decisions about their healthcare. The aim of this study was to explore

the interplay between illness and motherhood identities of women with ARDs during

preconception decision making.

Methods: Timeline-facilitated qualitative interviews with women diagnosed with an ARD

(18-49 years old). Participants were purposively sampled based on the following three

criteria: thinking about getting pregnant, currently pregnant, or had young children.

Interviews were thematically analysed.

Results: Twenty-two women were interviewed face-to-face (N=6) or over the telephone

(N=16). Interview length ranged from 20 minutes to 70 minutes, with a mean length of 48

minutes. Three main themes were identified: prioritisation, discrepancy, and trade-off.

Difficulties in balancing multiple identities in healthcare encounters were reported. Women

used 'self-guides' as a reference for priority setting in a dynamic process that shifted as their

level of disease activity altered and as their motherhood identity became more or less of a

focus at a given point in time. Women's illness and motherhood identities did not present in

isolation but were intertwined.

Conclusions: Findings highlight the need for holistic person-centred care that supports

women with the complex and emotive decisions relating to preconception decision-making.

In practice, health professionals need to consider women's multiple and sometimes

conflicting identities, and include both their condition and family associated goals and

values within healthcare communication.

Introduction

Women of reproductive age who have autoimmune rheumatic diseases (ARDs) such as rheumatoid arthritis (RA) and systemic lupus erythematosus (SLE) have expressed a need to be better supported with making decisions about pregnancy (1-3). Many women with ARDs will have positive pregnancy and parenting outcomes, but there are risks involved that need to be considered, including the potential effects of the disease itself and some disease-modifying treatments on fertility, pregnancy and breastfeeding (1, 4-6). Women with ARDs want their motherhood identities and associated preferences to be taken into account in decisions about their healthcare (2). In this manuscript, we investigate the development and interplay between illness and motherhood identities of women with ARDs as they navigate the healthcare system.

The diagnosis of chronic illness, such as autoimmune rheumatic disease, can lead to changes in an individual's sense of 'self' as the previous identity is replaced by an 'illness identity' once a diagnosis is made and treatment instigated. (7). An illness identity includes perceptions in relation to physical changes, emotional reactions to the physical changes, cognitive constructions of the illness and the impact this has on the self-perceived future (8). When women who have an ARD become mothers, they can be challenged by the symptoms associated with their illness such as fatigue, pain and impaired psychological well-being in addition to the usual challenges presented to women without a long-term condition (1, 9).

The social category of being a mother is not neutral and holds a particular meaning and significance across time and place. There is an ideology of the expectant mother and the identity she must hold (10).   The identity of a 'moral and good' mother is one that is taken

on before birth or even pregnancy and embodies the guiding ethic of 'proper' care of

children (11). In contrast to the 'moral and good mother' is that of the 'immoral and bad

mother'(10). The identity of an 'immoral and bad mother' mother would be that of

someone who fails to live up to the social ideal of a mother that has been imposed by

societal and cultural expectations (10). Developing the identity of a mother involves

internalising perceived ideals of how women ought to function as a mother, whilst

understanding the sometimes contradictory lived reality of mothering (12, 13). Women

who do not reach, or struggle to maintain, this ideal identity have been found to feel

increasing tension over their shortcomings and have reported negative psychological effects

such as feelings of guilt (14, 15).

The Self-Discrepancy Theory (16) suggests that the degree of discrepancy between the

actual self (the attributes a person possesses), ought self (who a person feels they should or

must be), and ideal self (who a person hopes or wishes to be) can lead to negative affect,

guilt and shame (17). The values that people attach to their actual, ought, and ideal

identities act as 'self-guides' in influencing behaviour and motivation. Analysis of

conversations of a UK on-line parenting forum suggested that women can feel pressure to

be an unrealistic 'selfless' and 'ideal' mother (18). Women's perceived shortcomings in

fulfilling their own and others' expectations of their role as a mother can leave them feeling

dissatisfied and questioning their identity (19, 20). In the context of women with ARDs, Self-

Discrepency Theory would suggest that if mothers experience a discrepancy between their

actual self (a woman managing a painful and potentially physically disabling condition), their

ideal self (the mother they wish to be) and their ought self (the mother they feel they

should be), this would lead to negative self-evaluations (17, 21).

Understanding how women's illness and motherhood identities influence, and are

influenced by, interactions with healthcare professionals and services is important in

designing patient-centred approaches to healthcare. In this study, we use participatory

qualitative methods to explore the self-identity of women with ARDs as they consider

pregnancy, are pregnant, and/or have young children.

Methods

Design

Timeline-facilitated qualitative interviews with women with ARDs who took part in the STAR

Family Survey (1).

Ethical approval

Ethical approval for the study, including the consent process, was granted by the Cardiff

University School of Medicine Research Ethics Committee on 20/10/16 REF16/56.

Participants and recruitment

Women who had expressed an interest in taking part in an interview were sampled from

the STAR Family Survey (1). The STAR Family Survey was a United Kingdom-based cross-

sectional on-line survey (n=128) focusing on women with ARDs' experiences of pre-

conception, pregnancy and early parenting.

Interview participants were purposively sampled to broadly achieve equal representation of

those who were: thinking about getting pregnant, currently pregnant, or had young

children. The aim was to explore the identity in women who were at different stages of

starting a family, whilst also managing an ARD. Participants were contacted through e-mail

or telephone, based on the contact details provided in the survey. Those who expressed

interest in being interviewed were sent a study information pack containing participant

information sheet, consent form and stamped return envelope. Participants were provided

with a £20 shopping voucher as a thank you for taking part.

Inclusion criteria

Women aged 18-49 years, who had been diagnosed with an ARD (i.e. inflammatory arthritis

or auto-immune connective tissue disease for which people would normally be under the

care of a rheumatologist). Women also needed to be planning on becoming pregnant in the

next 5 years, currently pregnant or have a child/children who were under the age of 5 years.

Participants had completed the STAR Family Study survey (1) and consented to interview.

Exclusion criteria

Women who were under 18 or over 50 years of age and those who had a disease not

classified as an ARD (e.g. joint hypermobility, fibromyalgia).

Interview procedure

A flexible narrative approach using visual methods was used to encourage participants to

talk about their 'lived experiences' regarding their experiences of identity, motherhood and

healthcare in their own words, focusing on things that were important to them, rather than

being guided by researcher-generated topics (22). Before the interviews, women were sent

a resource pack containing an example timeline outline, stationary, paper, stickers, and an

information sheet outlining examples of the topics we were interested in covering during

the interview. The timelines provided a visual tool which enabled women to map out their

journey so far. The timelines were used in interviews to provide cues, prompt discussion,

provide an opportunity for self-reflection, and help with building a rapport between

participants and the interviewer (23, 24). Preparing a timeline was voluntary and women

were encouraged to use their own formats (e.g. notes, diagrams they had drawn) as an

alternative if they wished to do so. Interviews were conducted face-to-face or over the

telephone. Face-to-face interviews were conducted in the participant's home or at Cardiff

University. Interviews were conducted by Denitza Williams, PhD (DW) and Bethan Pell, BSc

(BP). At the time of the interviews, DW was a post-doctoral researcher and BP was a

research assistant, both identified as female. Neither interviewer identified as having a long-

term condition; DW was a mother to two children, BP did not have any children. The

interviewers had no prior relationships with the participants.

Analysis

All interviews were audio-recorded and transcribed verbatim by a professional transcription

company. DW conducted the analysis. Data were analysed thematically and analysis was

based primarily on social phenomenology (25), focussing on exploring how women made

sense of their identity and motherhood choices. The data were analysed using Braun and

Clarke's Phases of Thematic Analysis (26) which involved familiarisation, generating initial

codes and identifying themes. NViVo10 was used to support the analysis. The protocol did

not include double coding of data. Instead, the qualitative research team (DW, BP, AG, RP)

held regular data analysis meetings to discuss the themes arising from the data, and the

development of the coding framework. Each member of the qualitative research group (RP,

AG, DW, BP) contributed their own perspective, relating to disciplinary backgrounds

including psychology, sociology, social policy and public health. This approach has been

identified as appropriate for enhancing the validity and quality of qualitative research (27).

Analysis was guided by the concept of information power (28) instead of data saturation.

<u>Results</u>

Twenty-two women were interviewed face-to-face (N=6) or over the telephone (N=16).

Interview length ranged from 20 minutes to 70 minutes, with a mean length of 48 minutes.

Through thematic analysis**,** three main themes were identified: prioritisation, discrepancy,

and trade-off (see Figure 1). Each of the main themes and sub-themes will be discussed

below.

<u>Figure 1: Thematic map</u>

Figure 1: Thematic map

Key: ———▶ Sub-theme

———▶ Related theme

184 **<u>Prioritisation of motherhood versus illness identities</u>**

185 Women spoke about prioritisation of the disease versus their desire for motherhood. Two

186 sub-themes of diagnosis and transition were identified. Women were often 'juggling

187 meaningful identities' (29); their identity as a woman, a mother, and a person with a long

188 term condition.

Diagnosis: the disease comes first

Women often reflected on the diagnosis of their ARD and their respective treatment options, contextualising their experiences in terms of their identity as an individual who was at crisis point and in desperate need of effective disease management. As disease management was the priority at this point, when the decision to use disease modifying drugs that might have an impact on fertility was made, their future motherhood plans were often not discussed. A woman's own self-identity and the prioritised identify by the clinician at that point was of a sick patient in need of treatment, and not a woman who might in the future like to have a child:

> *Yeah, I think we did talk about kind of because of the chemotherapy treatments it was likely that possibly I wouldn't be able to conceive. So it was kind of not brushed under the carpet but just one of those ok this has happened and we're talking about it but she's not in the place where she's thinking about having children and actually it's better to be healthy (P1)*

Transition: more than the disease

For women who were diagnosed before becoming a mother, shifts in priority were evident following diagnosis. Women reported that their disease needed to be well-managed to enable them to become a mother. Some women reported concern about the potential effects of their ARD when parenting. Concerns about being able to incorporate children into the 'boundaries of themselves' (30) in terms of the expansion of the self to incorporate children and the associated demands were raised.

if we're kind of successful in that [becoming a mother], the impact of having a child

to bring up and everything that you need to be able to do with that, am I going to be

well enough to do that in terms of sitting and you know being able to provide that

child with everything it needs in terms of energy and everything else. (P2)

Women discussed how the decision to have children needed to be planned well ahead of

time, because many of the drugs used to treat their ARD have teratogenic effects:

Yeah I guess with like children it's always been ingrained that it has to be planned

and stuff like (P2)

Women needed to transition by modifying the drugs that they were taking to manage their

disease before trying for a family. It was felt that this need to plan ahead was forced upon

them due to their ARD and felt that most people do not need to actively transition so far in

advance, or even to consult a health professional at all when planning to become a mother.

The transition signified a motivational and behavioural change between effective

management of disease and actively trying to conceive. The women in our study described

this transition as a difficult time, because they had made the decision that they would like to

have a child but often felt uncertainty about the effect of a pregnancy on their ARD.

I've got my own kind of own concerns about you know being at a point where I'm

well enough to become pregnant, being well during the pregnancy, can I stay well

enough for 9 months that I don't need extra medication that would impact on the

pregnancy (P9)

Self-guides are specific representations of the self or others and can include mental

representations of valued attributes (31). Self-guides can originate from diverse sources

such as general social norms, particularly if the guides have relevance in the terms of the life

experiences of the women (31). Here, the values attached to women's illness identity and

motherhood identity are guiding them in different directions, creating tension as they try to

decide which values are most important at a given time.

Self-discrepancy

Some women identified discrepancies between their perceptions of motherhood ideals and

their actual ability to meet these. Sub-themes included discrepancies in fertility, physical

ability and opportunity.

Fertility

Women reflected on the impact of their ARD, their perceived fertility and the chances of a

pregnancy ending in miscarriage. One woman experienced a miscarriage during a disease

flare. She talked about a discrepancy of feelings because she was informed that her poorly

managed disease contributed to the miscarriage, she was also aware that women with ARDs

are more likely to suffer a miscarriage, but also that miscarriage is common among healthy

women. She reported frustration that she had been advised to restart teratogenic

medication and to postpone planning for a baby until her disease was fully under control.

So yeah following the miscarriage I was all for planning to just get pregnant again,

but on advice from the rheumatologist and the renal specialist, they you know

because I have been so ill that is probably why I miscarried and they have managed

to convince me that I'm better off waiting, I'm back on the Micophenolate effectively

and I'm in a treatment plan that's not pregnancy friendly to try and get my lupus

under control (P15)

This signals a loss of a valued mothering identity, an important part of the self (32). This

signifies a discrepancy between the self-guides held by the health professional and the

woman in terms of pursuing a course of action that is consistent with her wish to become a

mother (31).

Physical ability

The potential impact of physical disability was identified as a factor in motherhood identity.

One woman reflected on her social identity in terms of the practical social help she received

from a volunteer. The availability of this support helped the mother feel that her children

were not missing out on 'normal' childhood due to her long-term condition. This seemed to

ease some of the guilt she felt in relation to limited mobility:

the only thing I have to say that's helped my kids through it all is in [city] there is the

young carer's project they are a charitable organisation (…) they take kids out every

other week to do things like to the cinema or the park, or they go bowling. Once a

year they go on a residential trip as well so it's kind of to give them a break from

caring (P14)

Women reported that their physical disabilities affected their ability to participate in social

group activities relating to motherhood. For example, most women had attended a local

mother and baby group, but found it difficult to participate, mainly because activities were

often floor based. Women reported feeling embarrassed that they were unable to

participate like other mothers, questioning their motherhood identity and highlighting the

discrepancy between their perceived ought and actual self. For example, one woman took

her child to a baby massage group. The activity was floor based and due to the effect of the

ARD on her physical abilities she struggled to get back up from the floor and to pick her baby

up:

If I was on the floor I couldn't get up so the first couple of weeks I went I was sat on a

chair leaning over [daughter] but then I just it started to make my back ache and

everything else and I think for like I wasn't quite bonding with her because it felt like

she was so far away so I thought sod it I'll get on the floor with her but that would

mean that I would have to crawl to the chair – hand on to something to get myself up

and I would literally have to haul myself up off the floor, but yeah I needed somebody

to pick <<daughter1>> up off the floor for me and either carry her to the car, or put

her in the sling that I had (…) you feel like the odd [one] (P4)

This event led to a sense of shame and isolation. Motherhood can have an effect on women;

their positive and negative qualities can be heightened when reflecting on the mother-child

relationship (29).

the biggest thing for me is the kids they miss out on so much because of this stupid

thing with me. I feel rotten because they have to be no holidays forever and all their

pals are going to Spain, or Italy or Portugal or States or holidays and my kids are

saying maybe we should go to Norway or Finland because it's so cold and it's not that

sunny [it wouldn't cause disease flare] ((crying)) (P14)

Women felt that they received discrepancies in recommendation from health professionals,

especially for breastfeeding. They were often encouraged to initiate and maintain

breastfeeding by midwives and health visitors, even though felt it was not the best thing for

themselves or their child at that time due to their condition:

there is such a drive to push you to breastfeed as it is best for the baby and I accepted

it, but I also accept that sometimes the best thing in a research report isn't the best

thing for the individual and in our case me and my two little ones it certainly wasn't

and I think that was why in terms of the health visitor and the midwife and the

breastfeeding advocates they're so hell bent on getting a twin mum to breastfeed,

because it's that extra pressure again that they focus so much on that they forget the

rest of the situation (P8)

Although participants reported that there was a general pressure to continue breastfeeding,

some mothers felt that they were pressured to stop breastfeeding by family members or

health professionals, so that they could restart required certain disease modifying

medications. Many women reported feeling guilty about their inability to breastfeed, or if

they had to stop breastfeeding to re-start medication. Reports of feeling guilty seemed to

stem in part from healthcare professional and peer attitudes towards formula feeding.

Women reported feeling guilty and ashamed of their desire not to breastfeed, inability to

breastfeed or their decision to stop breastfeeding:

"The breastfeeding advocates had been trying to get me to consider alternative

medication and stuff which in an ideal world when we have lots of time maybe that

would be a nice thing to do, but when things go downhill drastically you know and I

was in a lot of pain and I was struggling to hold 2 babies to breastfeed as well

because my joints were sore… I marched on and then at 6 weeks I dropped a child… it

took me actually dropping the child I think that made me think she was worth

listening to" (P8)

Opportunity

One woman reflected on the impact of her physical disability on her actual self as a mother. Her baby had an accident at home which needed hospital treatment. Due to her lack of mobility, the woman could not drive to take her child to the hospital independently and had to rely on a neighbour for help. The woman felt that she did not have the opportunity to behave as a 'normal' mother who could react fast and take immediate action to help her child. This intensified her feelings of self-discrepancy and guilt.

> *"[The] gap underneath the bathroom door was quite big, he [son] must've put his toes underneath it and either pulled the door, or pushed it so he chopped off the end of his big toe it was kind of hanging you can see the bone it was so bad. And I was in my jammies, I was so tired. I asked my neighbour to get my other 2 boys ready for school and get them to school in time and I just took a taxi (…) because I can't drive (…) I was again feeling totally useless because I should have been able to drive him, I should've been able to just get dressed get him in the car and to the hospital and I couldn't put one foot in front of the other I was so tired (…) it was horrible you know that feeling that, that you should be able to do more for your kids" (P14)*

A few women talked about the possibility of adopting a child but felt that other people's perceptions of them due to their ARD limited their opportunities. One woman and her husband felt that managing her ARD, as well as having a successful pregnancy, would be too difficult as altering her medication was likely to have a detrimental impact on her health. She felt that her ability to be a mother was pre-determined by adoption agencies who made judgements in relation to her long-term condition.

I would have to prove my ability to be a parent so they wouldn't take [stepchildren]

at face value, my step children and I would have to go to parenting classes, they'd

have to see evidence of me walking, carrying, being able to, they wanted me to have

a – you know a pretend baby at home that they can download and things to prove

that I was able to meet its needs all throughout the day and night endlessly and it

was really full on [husband] and I got furious and I, I was really pissed off at them I

remember saying to them you know in this day and age where you can't discriminate

against colour, or sexuality you feel you can discriminate against disability you know

(P8)

Here, there appears to be a discrepancy between women's actual and ideal self-identity and

agencies' perceptions of what a mother ought to be.

Trade-offs between competing values

The complex identity changes women go through when planning to become a mother can

require multiple types of trade-offs, including disease management, mothering aspirations

and mothering ideals. If women's identities have developed in ways in which they are

committed to becoming a parent and being part of a family, the threat their illness poses to

their ability to do this means they have to revisit that commitment, which can lead to

uncertainty and psychological distress. At this stage, women needed to reflect on their goals

and values and what the future holds for them as well as their ability to fulfil their perceived

identity of a 'good' mother (10) .

Women reflected on their pre-conception journey and the transitions that needed to be

made in terms of their identity as a person with a disease that needed to be managed and

as a woman who wanted to become a mother. It was often felt that women needed to

balance a trade-off between the self as a patient in need of medication to control their ARD

(the actual self) and the self as an individual who is well enough to be a mother to meet

their own expectations and that of society about mothering (the ought self). However, the

'ideal' self is also present here in the sense that women's own goals and aspirations, which

might be at odds with the 'ought self', that is what they think others expect of them.

Planning a family involved a balance of clinician advice about the trade-offs associated with

the management of an ARD and maternal desire. They had to balance multiple identities

and expected roles, that of a woman who has a long-term condition and that of a woman

who wants to have a child or to expand her family. Some women felt that their motherhood

identity was being threatened by the management of their ARD and healthcare

communication:

> *So they then decided they wanted to start me on floudamide I think it's called which*
>
> *was a disease modifying thing but they told me I couldn't pregnant on it and that the*
>
> *likelihood of me taking it would affect my fertility. So at that point we said well hang*
>
> *on a minute you know I'm only sort of 25, 26, don't really want to start taking stuff*
>
> *that going to affect my fertility, maybe we should just put up with stuff the way it is,*
>
> *try and get pregnant, have another baby because we always knew we wanted a*
>
> *second child then go on the drugs at that point if its needs it. (P3)*

Others spoke of specific trade-offs in relation to the management of their disease and their

desire to have children. One woman reflected on how she had traded optimum disease

management with the chance of conceiving a baby despite an active disease flare. For her,

the priority at that time was to try to ensure that she had a chance of having a child, and her

clinical team seemed to be supportive by putting in place necessary referrals:

things were flaring back up and they said well look, what we can do is we can do a

referral to the fertility clinic, they can take a look and see if there's anything

specifically wrong that's stopping it you know you may be eligible for IVF or you know

with a discussion with them it might be that they're able to take your eggs and freeze

them so that we can start the medication and do it that way. (P3)

Women with ARDs who had successful pregnancies felt that they were "lucky" because they

identified themselves at a higher risk of complications on their journey to motherhood

compared to women who do not have a long-term condition. They spoke of trade-offs in

relation to miscarriage risk and the chance of a successful pregnancy:

I am very very grateful for my two children because there's a lot of people who really

suffer and suffer many miscarriages and stuff whereas I have only suffered one

miscarriage in my life, a lot of people have suffered a lot worse and haven't been able

to have children or have had their fertility robbed because of the medications they've

been on and stuff. (P3)

Discussion

In this study, we set out to investigate the development and interplay of women's self-

identity as a person with an ARD and as a mother as they navigated the healthcare system

during preconception planning, pregnancy, and early parenting. Prioritisation of illness

versus motherhood identities, self-discrepancy, and trade-offs between competing values

were identified as major themes in the discussion surrounding women's experiences. The

difficulties in balancing multiple identities in healthcare encounters was evident, both for

the women and for the healthcare professionals. Women seemed to use 'self-guides' (31) as

a reference for priority setting in a dynamic process that shifted as their level of disease

activity altered and as their motherhood identity became more or less of a focus at a given

point in time. Women often felt that they needed to balance their 'actual' self (individual

with a long-term condition) with their perceptions of the 'ought' self (individual well enough

to meet their own as well as society's expectations of being a mother) as well as perceptions

of the 'ideal self' (women's own goals and aspirations). Self-guides became at odds with

each other when the action that was needed to manage disease well conflicted with the

action required to become a mother or to be a 'good mother' as perceived by the individual.

Motherhood identities were also sometimes in conflict with medical recommendations in

relation to achieving optimum disease management. When these discrepancies in identity

and associated self-guides came into conflict (16), women reported feelings of anxiety,

shame, guilt, and frustration. Our findings highlighted the need to consider women's

multiple and sometimes conflicting identities, with their associated goals and values, in

providing patient-centred healthcare.

 It has been argued that a mother's pregnancy and long-term condition must be viewed as

separate but co-existing entities (33). This suggests that women view their identities

separately, with the identity as a mother working separately to the identity as someone who

has a long-term illness. Women's illness and motherhood identities did not present in

isolation in our study, but rather were intertwined, highlighting the need for holistic patient-

centred care that supports women with the complex and emotive decisions that they need

to make.

Previous research has found that the accommodation and recognition of identity shifts in

healthcare communication are imperative for supportive patient decision-making (34). A

patient-centred communication style can facilitate such discussions in consultations, placing

'what matters most to the patient' first. Person-centred care encompasses 'Shared decision-

making' a process where health professionals provide understandable information, discuss

the pros and cons associated with different treatment options, and fully involve patients in

treatment decisions, considering their personal circumstances and preferences (35, 36). This

enables patients to make more informed decisions that align with their preferences and

thus fit better with their identity, become more active and empowered in their own

healthcare, to have better relationships with their health care professionals, and to feel

more satisfied with the choices that they make (37). There is strong policy support for the

routine use of shared decision-making (SDM) in clinical practice (38, 39). The National

Institute for Health and Care Excellence recently published guidance for the incorporation of

SDM within healthcare (40). Adopting SDM in healthcare teams would enable both illness

and motherhood identities and associated goals and values to be considered by women and

their healthcare teams so that women's preferences and priorities can be better

incorporated in healthcare decisions.

Strengths and Limitations

This study explored women's perceived illness and motherhood identities using in-depth

participant centred interviews. This enabled us to understand the way in which women

perceived their motherhood and illness identities and the way this influenced motivation

and behaviour in relation to the management of disease and becoming (or being) a mother.

The women who took part in our interviews were more highly educated and more likely to

be employed than the overall STAR Family survey participants, which is indicative of a

sampling bias that should be taken into consideration when generalising from the findings.

Furthermore, this study was conducted before the global COVID-19 pandemic, which saw

significant shifts in the delivery of healthcare, with modifications to services including

remote consultations for non-urgent healthcare appointments, and significant changes in

the provision of maternity services (41). Further research is needed to explore how these

changes might have impacted on healthcare communication.

Conclusion

This study explored self-identity during pre-conception, pregnancy and early parenting in

women with an autoimmune rheumatic disease related to healthcare communication and

decision-making. Women's illness and motherhood aspirations were key parts of the

formation of their self-identity. Women were balancing and shifting through intertwined

illness and mothering identities, which sometimes came into direct conflict with each other.

It is imperative that both illness and motherhood identities are acknowledged and explored

during healthcare encounters to facilitate preference-sensitive healthcare decisions. A

person-centred approach to healthcare communication would facilitate this by encouraging

dialogue between healthcare professionals and women about preferences associated with

treatment decision-making that incorporate both their illness and mothering identities.

Additional research will be required to develop and test the feasibility and acceptability of a

person-centred intervention to be used in healthcare encounters.

Declaration of Competing Interest

No conflicts of interest have been declared.

Acknowledgments

The authors wish to thank all of the women that took part in this research. We would like to

thank our patient involvement representatives for their continued support with this work.

We would also like to thank the Wellcome Trust for funding this research.

References

1.   Phillips R, Pell B, Grant A, Bowen D, Sanders J, Taylor A, et al. Identifying the unmet

information and support needs of women with autoimmune rheumatic diseases during pregnancy

planning, pregnancy and early parenting: mixed-methods study (Under review). 2018.

2.   Phillips R, Williams D, Bowen D, Morris D, Grant A, Pell B, et al. Reaching a consensus on

research priorities for supporting women with autoimmune rheumatic diseases during pre-

conception, pregnancy and early parenting: A Nominal Group Technique exercise with lay and

professional stakeholders. Wellcome Open Res. 2018;3:75.

3.   Ackerman IN, Jordan JE, Van Doornum S, Ricardo M, Briggs AM. Understanding the

information needs of women with rheumatoid arthritis concerning pregnancy, post-natal care and

early parenting: A mixed-methods study. BMC Musculoskelet Disord. 2015;16:194.

4.   Williams D, Webber J, Pell B, Grant A, Sanders J, Choy E, et al. "Nobody knows, or seems to

know how rheumatology and breastfeeding works": Women's experiences of breastfeeding whilst

managing a long-term limiting condition – A qualitative visual methods study. Midwifery.

2019;78:91-6.

5.   Ostensen M, Brucato A, Carp H, Chambers C, Dolhain RJEM, Doria A, et al. Pregnancy and

reproduction in autoimmune rheumatic diseases. Rheumatology. 2011;50(4):657-64.

6.   Ostensen M, Cetin I. Autoimmune connective tissue diseases. Best Practice & Research

Clinical Obstetrics & Gynaecology. 2015;29(5):658-70.

7.   Goffman E. Stigma: Notes on the Management of Spoiled Identity. . New York: Simon &

Schusler; 1963.

8.   Charmaz K, Rosenfeld D. Chronic Illness. In: Cockerham W, editor. New Blackwell Companion

to Medical Sociology. London: Blackwe;;; 2010. p. 312-33.

9.   Kristiansen TM, Primdahl J, Antoft R, *al. e*. Everyday life with rheumatoid arthritis and

implications for patient education and clinical practice: a focus group study. Musculoskeletal Care.

2012;10:29-38.

10.   Murphey S. Reclaiming a moral identity: stillbirth, stigma and 'moral mothers'. Midwifery.

2011;28:476-80.

11.   May V. On being a 'good' mother: the moral presentation of self in written live stories.

Sociology. 2008;42(470-486).

12.   Choi P, Henshaw C, Baker S, Tree J. Supermum, superwife, supereverything: Performing

femininity in the

transition to motherhood. . Journal of Reproductive and Infant Psychology. 2005;23:167–80.

13.   Laney E, Hall E, Anderson T, Willingham M. Becoming a Mother: The Influence of

Motherhood on Women's Identity Development Identity: An International Journal of Theory and

Research. 2015(15):126–45,.

14.   Arendell T. Conceiving and investigating motherhood: The decade's scholarship. Journal of

Marriage and the Family. 2000:1192-207.

15.   Phanco ML. Internal tension and faith: A qualitative exploration of mothers working in

Christian academia

(Unpublished doctoral dissertation). Rosemead School of Psychology, Biola University, La Mirada,

CA. 2003.

16.   Higgins ET. Self-discrepancy: a theory relating self and affect. Psychological Review.

1987;94(319-340).

17.   Morley S. The Self in Pain. Review in Pain. 2010;4(1):24-7.

18.   Pederson S. The good, the bad and the 'good enough' mother on the UK parenting forum

Mumsnet. Women's Studies International Forum. 2016;59:32-8.

19.   Feldhusen C, Bjork M, Forsblad-d'Elia H, al. e. Perception, consequences, communication

and strategies for handling fatigue in persons with rheumatoid arthritis of workling age- a focus

group study. Clin Rheumatol. 2013;32(557-566).

20.   Hatcher SL. The psychological experience of nursing mothers upon learning of a toxic

substance in their breast milk. . Psychiatry. 1982;45(172-181).

21.   Goffman E. Stigma: Notes on the Management of Spoiled Identity. London: Penguin; 1963 :

12.

22.   Gabb J, Fink J. Telling Moments and Everyday Experience: Multiple Methods Research on

Couple Relationships and Personal Lives. Sociology. 2015;49(5):970-87.

23.   Goldenberg T, Finneran C, Andes KL, Stephenson R. Using participant-empowered visual

relationship timelines in a qualitative study of sexual behaviour. Global Public Health. 2016;11(5-

6):699-718.

24.   Pell B, Williams D, Phillips R, Sanders J, Edwards A, Choy E, et al. Using Visual Timelines in

Telephone Interviews: Reflections and Lessons Learned From the Star Family Study. International

Journal of Qualitative Methods. 2020;19:1609406920913675.

25.   Fereday J, Muir-Cochrane E. Demonstrating rigor using thematic analysis: A hybrid approach

of inductive and deductive coding and theme development. International journal of qualitative

methods. 2006;5(1):80-92.

26.   Braun V, Clarke V. Using thematic analysis in psychology. Qualitative Research in

Psycholog2006.

27.   Barbour R. Checklists for improving rigour in qualitative research: a case of the tail wagging

dog? BMJ. 2001;322(7294):1115-7.

28.   Malterud K, Siersma V, Guassora A. Sample size in qualitative interview studies:guided by

information power. Qual Health Res. 2016;26:1753-60.

29.   Feddersen H, Kristiansen T, Andersen P, Horslev-Petersen K, Primdahl J. Juggling identities of

rheumatoid arthritis, motherhood and paid work- a grounded theory study. Disability and

Rehabilitation. 2018;https://doi.org/10.1080/09638288.2018.1433723.

30. Laney E, Hall E, Anderson T, Willingham M. Becoming a mother: the influence of

motherhood on women's identity development. International journal of theory and research

2015;15(2):126-45.

31. Moretti M, Higgins ET. Internal representations of others in self-regulation: a new look at a

classic issue. Social Cognition. 1999;17(2):186-208.

32. Wonch Hill P, Cacciatore J, Shreffler KM, Pritchard KM. The loss of self: The effect of

miscarriage, stillbirth, and child death on maternal self-esteem. Death Studies. 2017;41(4):226-35.

33. Thomas H. Pregnancy, illness and the concept of career. Sociol Health Illn. 2003;25(5):383-

407.

34. Palmer-Wackerly AL, Dailey PM, Krok-Schoen JL, Rhodes ND, Krieger JL. Patient Perceptions

of Illness Identity in Cancer Clinical Trial Decision-Making. Health Communication. 2018;33(8):1045-

54.

35. Elwyn G, et al. Shared decision making: a model for clinical practice. . Journal of general

internal medicine. 2012;27`10`:1361-7.

36. Joseph-Williams N, Williams D, Wood F, Lloyd A, Brain K, Thomas N, et al. A descriptive

model of shared decision making derived from routine implementation in clinical practice

('Implement-SDM'): Qualitative study. Patient Education and Counseling. 2019;102(10):1774-85.

37. De Silva D. Evidence: Helping people share decision making. . London; 2012.

38. Barr PJ, Durand M. Implementing shared decision making: The role of incentives. Shared

Decision Making in Health Care: Achieving evidence-based patient choice. 2016.

39. Joseph-Williams N, Lloyd A, Edwards A, Stobbart L, Tomson D, Macphail S, et al.

Implementing shared decision making in the NHS: lessons from the MAGIC programme. Bmj.

2017;357.

40. NICE. Shared decision making 2021 [Available from:

https://www.nice.org.uk/guidance/ng197.

41. Foundation. PI. Babies in Lockdown. 2020 Augusy 2020.

Figure 1: Thematic map

Key: ──────▶ Sub-theme

──────▶ Related theme

Figure 1 Williams et al