Graphical abstract

Prognostic peripheral blood biomarkers at ICU admission predict COVID-19 clinical outcomes

Melina Messing[1], Mypinder S. Sekhon[2], Michael R. Hughes[1], Sophie Stukas[3], Ryan L. Hoiland[4], Jennifer Cooper[3], Nyra Ahmed[3], Mark Hamer[1], Yicong Li[1], Samuel B. Shin[1], Lin Wei Tung[1], Cheryl Wellington[3], Don D. Sin[2,5], *Kevin B. Leslie[1] and *Kelly M. McNagny[1]

[1] The Biomedical Research Centre, Dept of Biomedical Engineering, University of British Columbia, Vancouver, BC, Canada.

[2] UBC Department of Medicine (Division of Respirology), Vancouver, BC, Canada.

[3] UBC Department of Pathology and Laboratory Medicine, Vancouver, BC, Canada.

[4] UBC Department of Anesthesiology, Pharmacology, and Therapeutics, Vancouver, BC, Canada

[5] The University of British Columbia (UBC) Centre for Heart Lung Innovation (HLI), St Paul's Hospital, Vancouver, BC, Canada.

*Corresponding Authors:

Dr. Kelly M. McNagny
Biomedical Research Centre
University of British Columbia
2222 Health Sciences Mall
Vancouver, B.C. CANADA
Ph: 604 822 7824
kelly@brc.ubc.ca

Dr. Kevin B. Leslie
Biomedical Research Centre
University of British Columbia
2222 Health Sciences Mall
Vancouver, B.C. CANADA
cytokine86@gmail.com

Abstract

The COVID-19 pandemic continues to challenge the capacities of hospital ICUs which currently

lack the ability to identify prospectively those patients who may require extended management. In

this study of 90 ICU COVID-19 patients, we used multiplexed cytokine evaluation and, on 42 of

these patients (binned into Initial and Replication Cohorts), CyTOF-based deep

immunophenotyping. This revealed blood prognostic biomarkers that, at time of ICU admission,

prospectively distinguish, with 91% sensitivity and 91% specificity, patients who will

subsequently die or have long ICU stays ($\geq$ 6 days) from those who will have short-stays (< 6

days). This is achieved through a tiered evaluation of serum IL-10 and targeted

immunophenotyping of monocyte subsets (specifically, CD11c^{low} classical monocytes) through

statistical approaches. We have distilled this down to a prognostic test that could prove useful in

guiding clinical resource allocation, treatment regimens and assessment of new therapeutic

interventions.

MAIN

COVID-19 continues to overwhelm effective health care delivery in most parts of the world due

to challenges in achieving sufficiently high vaccination rates, vaccine hesitancy and the emergence

of more virulent viral variants for which current vaccines offer more modest protection. Thus,

waves of rapid outbreaks continue to threaten ICU capacities[1,2].

Individual patient responses to infection by the SARS-CoV-2 virus can vary dramatically ranging

from asymptomatic or mild flu-like symptoms to much more severe symptoms including acute

respiratory distress syndrome (ARDS) and death[3,4]. Individual patient outcomes are remarkably

challenging to predict but severe disease has been broadly linked to advanced age, obesity,

underlying comorbidities and secondary infections[5,6,7,8,9,10]. Neither symptoms nor conventional

clinical measurements (serum C-reactive protein (CRP), blood D-dimers etc.) have sufficient

prognostic power and thus approved interventions for severe COVID-19 (including systemic

corticosteroids and tocilizumab) are administered broadly as clinicians lack the ability to identify

accurately patients at risk of long-term complications and death[11,12,13,14,15]. Immunologically,

"severe" patients have been reported to exhibit lymphopenia, neutrophilia, accumulation of lung

monocytes, emergency myelopoiesis, and substantial changes in serum cytokine and chemokine

profiles likely reflecting a cytokine storm as the result of a delayed, but exuberant, immune

response to infection[16,17,18,19,20,21,22,23]. The latter has been of particular interest for the development

of prognostic tools but while some markers have proven useful in measuring the severity of active

COVID-19, to date, they have lacked the necessary statistical power to prospectively predict the

likelihood of incipient severe disease[24,25,26,27,28,29,30,31]. For example, serum IL-6 (alone, or together

with other inflammatory markers) has most consistently been linked to severe active disease and,

by some groups, was shown to predict the need for subsequent mechanical ventilation as well as

survival[32,33,34,35,36,]. In contrast, other studies have struggled to link tightly IL-6 (or TNFα, IFNγ or

GM-CSF) to an elevated risk of severe disease and instead have proposed various combinations of

serum levels of CCL5, IL-1RA and IL-10, EN-RAGE, TNFSF14 and oncostatin M as indicators

of incipient severe disease and, in some cases predictors, of disease severity[37,38,39]. While,

individually, these studies show several biomarkers capable of triaging patients, these inconsistent

and sometimes contradictory results highlight the need for biomarkers with robust statistical power

to be clinically useful.

Studies focused on cellular changes have linked a decreased frequency of monocytes (and, more

variably, alterations in the frequency of natural killer cells (NK), plasmacytoid dendritic cells

(pDC), type-2 conventional dendritic cells (DC2), mucosal associated invariant T (MAIT) cells

and other lineages) with active severe disease and poor outcomes[17,19,34,40,41,42,43,44]. While these

global cellular profiling efforts provide important insights into the immune response to SARS-

CoV-2 infection, they have yet to be translated into prognostic tools to assist with individualized

care.

Here we focused on the development of an immunological biomarker screen that, at ICU admission

for COVID-19, predicts length of ICU stay or death. Strikingly, we find that, at ICU admission,

measurements of serum IL-10 and simple monocyte subset surface signatures, specifically,

CD11c[low] classical monocytes, can predict with 91% sensitivity and 91% specificity patients who

will either die or have a longer stay in ICU. We offer these biomarkers as a model clinical

laboratory test with future potential in gaining insights into variable responses to SARS-CoV-2

infection.

Results

COVID-19 patient group selection and optimization of high dimensional immune profiling

To identify potential prognostic markers of COVID-19, we collected PBMCs from 8 healthy controls and, within the first 48 hours of ICU admission, serum samples from 90 ICU COVID-19 patients (the "Cytokine Cohort") admitted during the "second wave" of COVID-19 (November, 2020–February, 2021) together with serial serum samples across different timepoints during the course of their ICU admission. PBMCs collected from 14 of these 90 ICU COVID-19 patients (the "Initial Cohort") were analyzed by mass cytometry (CyTOF) with a training set of 35 monoclonal antibodies (Supplementary Table 1) designed to detect broad shifts in levels of normal PBMC lineages as well as their activation status, and the possible mobilization of tissue resident innate immune cells and bone marrow progenitors into peripheral blood. Based on these data we developed a refined, second-generation, 38-antibody CyTOF panel (Supplementary Table 1), which was used on PBMCs collected from a further 28 of the 90 ICU COVID-19 patients (the "Replication Cohort"). Data from the Replication Cohort were used to validate observations from the Initial Cohort on early alterations in immune responses that could effectively differentiate patients likely to recover after a short ICU stay from those who would either die or have prolonged stays in ICU. The ICU admission sera from all patients in the Cytokine Cohort (which includes all patients in the Initial Cohort and the Replication Cohort) were analyzed for levels of four cytokines: IL-1β, IL-6, IL-10 and TNFα (Fig. 1a, b).

Clinical and demographic details of all patients and healthy controls are presented in Table 1 and include age, sex, body mass index (BMI), requirement for ventilation during admission and ICU admission levels of serum CRP, blood D-dimer levels, and white blood cell counts along with their

differentials. The average age of the ICU patients was 63.5 years with a 2-to-1 bias towards male

patients, consistent with previous patient demographic reports linking more severe COVID-19

with older male patients[45]. Table 1 also bins patients into two clinical outcome groups of "Short-

Stay" and "Long-Stay/Died" based on the length of time in the ICU and survival: "Short-Stay"

patients are classified as those spending < 6 days in the ICU and were survivors, while "Long-

Stay/Died" patients are defined as patients who spent 6 or more days in the ICU or died during

their stay in ICU. The choice of 6 days as the cut-off was based upon iterative empirical analyses

of immune data that divided, with greatest statistical significance (by p-value), patients into two

sub-groups with distinct clinical outcomes (Fig. 1a, c).

Importantly, we found no significant differences between the two clinical outcome groups with

respect to mean age, BMI, blood clotting parameters (D-dimer levels) or serum CRP levels (Fig.

1d, e). At admission, the mean total PMN counts were significantly increased in the Long-

Stay/Died group compared to the healthy controls ($p<0.0001$) and compared to the Short-Stay

group ($p=0.025$) (Fig. 1f). In our separate analyses of just the Initial Cohort and Replication

Cohorts, however, differences in PMN counts were not statistically significant and thus we did not

consider this measurement as a useful prognostic biomarker of clinical outcomes in the context of

smaller cohort numbers. PBMC counts were also not significantly changed between healthy

controls and patients or between our two clinical outcome groups (Fig. 1f). Thus, while these

routine clinical tests follow a broad spectrum of parameters including inflammation, coagulopathy,

hypo-immunity and autoimmunity, none consistently proves prognostic in identifying patients

who would require an extended stay in ICU or die. Accordingly, we conducted more detailed

immunological examinations focused on a single, specific COVID-19-associated process, namely

inflammation.

Serum cytokine analyses as prognostic screens for predicting clinical outcome

We began by examining serum levels of IL-1β, IL-6, IL-10 and TNFα at ICU admission in all

serum samples from our Cytokine Cohort of 90 COVID-19 patients. Strikingly, we found that the

mean ICU admission levels of serum IL-10 (p = 0.004) and TNFα (p = 0.0005) were significantly

elevated in the Long-Stay/Died group relative to the Short-Stay group (Fig. 1g). In the Cytokine

Cohort, 43% (39/90) of patients had serum ICU admission levels of IL-10 levels $\geq$ 15pg/ml and

79% (31/39) of these patients fell into the Long-Stay/Died group. Similarly, 42% (38/90) of

patients in the Cytokine Cohort had serum ICU admission levels of TNFα $\geq$ 10pg/ml and 79%

(30/38) of these patients were members of the Long-Stay/Died group. Interestingly, serum IL-10

and TNFα only showed a weak correlation with each other (Pearson correlation coefficient $R^2 =$

0.12) suggesting that each may represent a different aspect or chronology of the inflammatory

process.

Given the clear prognostic value of ICU ***admission levels*** of TNFα and IL-10, we also examined

the subsequent mean maximum serum cytokine levels in ***post admission*** samples from patients in

the Short-Stay and the Long-Stay/Died groups and found an even more significant difference

between the two groups for both serum TNFα (p < 0.0001) and serum IL-10 (p = 0.0009) (Fig.

1h). Intriguingly, many patients in the Long-Stay/Died group who demonstrated modest admission

levels of serum IL-10 and TNFα subsequently developed high levels during their stay in ICU,

further reinforcing the importance of these two cytokines as predictive measures of patient

outcomes and monitoring the trajectory of disease.

While admission levels of serum IL-6 were also significantly different between the two clinical

groups (p = 0.007) (Fig. 1i), we excluded this cytokine from further analyses due to the potential

confounding effects of anti-IL-6 receptor antibody (tocilizumab) treatments, which has routinely

been administered to COVID-19 patients in British Columbia during ICU admission since

February 2021 and such treatments could complicate the interpretation of our results. Finally, there

were no significant differences between the two clinical outcome groups with respect to mean

serum IL-1β levels at ICU admission (p = 0.205) and thus this cytokine was also not analyzed

further (Fig. 1i). In summary, we found that ICU admission levels of serum IL-10 and TNFα were

useful and statistically powerful prognostic markers for clinical outcomes in severe COVID-19.

Major PBMC subsets fail to distinguish Short-Stay from Long-Stay/Died patients

We examined whether parallel CyTOF analyses of peripheral immune cells sampled at the time of

ICU admission could reveal additional prognostic biomarkers that identify patients in the Long-

Stay/Died group, particularly among those that had serum IL-10 levels <15pg/ml and/or serum

TNFα levels <10pg/ml. PBMC samples were available for 42/90 of the Cytokine Cohort patients,

and these 42 samples were divided into an Initial Cohort (14 samples) and a Replication Cohort

(28 samples).

Using a 35-marker CyTOF panel on the Initial Cohort (Supplemental Figure 1) and a 38-marker

CyTOF panel on the Replication Cohort (Fig 1.), we saw no differences between the Short-Stay

patients and Long-Stay/Died patients with respect to major peripheral blood immune populations.

The more-focussed and larger 38-marker CyTOF panel, used to analyze immune cell subsets in

the Replication Cohort, permitted clear identification of broad blood cell lineages (B, T, NK, and

myelomonocytic) as well as major subsets within each cell lineage leading to 41 distinct cell subset

clusters based on the variable expression of these cell-surface markers (Fig. 2a-c). While these

analyses of the Replication Cohort samples and the Initial Cohort samples confirmed previous

reports[5,46] of general lymphopenia in COVID-19 patients relative to healthy controls with respect

to both total CD4 T cells and total CD8 T cells, these markers failed to discriminate between the

Short-Stay and Long-Stay/Died patient groups. Also consistent with previous reports, we observed

no significant differences in total B cells in COVID-19 patients compared to healthy controls or

between the two clinical outcome groups (Fig. 2d). Although mean total NK cells, MAIT cells, γδ

T cells, DC2/3 and pDC were depleted in COVID-19 patients relative to healthy controls these,

too, failed to distinguish the Short-Stay group from the Long-Stay/Died group (Fig. 2e). Finally,

while mean total monocytes and stem cell levels were significantly increased in COVID-19

patients relative to healthy controls they failed, individually, to distinguish the Short-Stay from the

Long-Stay/Died patient groups (Fig. 2f). In summary, broad immune subset analyses were

insufficient to predict COVID-19 patient clinical outcomes with respect to the length of stay in

ICU and/or death in either the Initial Cohort or the Replication Cohort. We, therefore, performed

more detailed analyses of immune cell subsets within these broad cell categories to identify more

subtle potential differences between the two clinical outcome groups that could assist in the

prospective identification of Long-Stay/Died patients.

Levels of a distinct monocyte subset at the time of ICU admission predicts subsequent clinical

outcome

To reveal a larger diversity of specific immune cell subsets we performed more focused cluster

analyses on patient PBMC samples from the Initial Cohort and the Replication Cohort after pre-

gating for selected major cellular subsets. To restrict the clustering to the myelomonocytic

compartment we performed gated analyses on GM-CSFR$^+$(CD116$^+$) CD19$^-$ CD3$^-$ cells (Fig. 3a)

and restricted clustering to shared marker channels to enable direct comparison between the Initial

and the Replication cohorts. These analyses did not reveal subsets that separated Short-Stay from

Long-Stay/Died patients with respect to absolute cell counts. To focus more specifically on the

monocytic subsets as well as to simplify the cluster analyses, after pre-gating on CD116+ CD19-

CD3- cells, we restricted the marker channels selected for clustering to a set of 7 markers useful

in defining monocytic subsets (CD45, CD14, CD16, CD11c, HLA-DR, CD123, CD56, see Figs.

3b,c). Interestingly, this strategy revealed a CD11c^{low} classical monocytic subset (CD45$^+$ CD116$^+$

CD3ϵ^- CD11c^{low} HLA-DR$^+$ CD14$^+$ CD16$^{-/low}$ CD123$^{-/low}$) that, in both the Initial and the

Replication Cohorts, was consistently enriched in COVID-19 patients relative to healthy controls

(p = 0.001, Replication Cohort) and was preferentially enriched in the Long-Stay/Died group

relative to the Short-Stay group (p = 0.019, Replication Cohort) (Fig. 3d). The prognostic value of

the CD11c^{low} classical monocytic marker was restricted to this subset of classical monocytes in

both the Initial and Replication Cohorts and did not reflect underlying changes of total classical

monocytes which were unchanged in the two clinical outcome groups (Fig. 3e). Moreover, for

both Initial and Replication Cohorts, total intermediate monocytes (CD14$^+$ CD16^{int}) and total non-

classical monocytes (CD14^{low} CD16$^+$), as well as observed subpopulations of these types of

monocytes, did not prove useful prognostically (Fig. 3f). Focussing the analyses further on

classical monocytes, we found that a three-marker gating strategy was sufficient to identify the

CD11c^{low} classical monocyte population identified by our multi-dimensional analyses (shown here

for one representative sample from each group in the Replication Cohort, where intensity is

proportional to relative frequency of cells) (Fig. 3g). Thus, the prognostically useful biomarker of

CD11c^{low} classical monocytes was detectable in two dimensions in both the Initial and the

Replication Cohorts using antibodies to a small set of cell-surface markers.

Because lymphopenia has been a consistent feature of severe COVID-19 and T cell subset

alterations have been described, we performed similar in-depth analyses of the T cell

compartments by gating on CD3^{+} cells prior to clustering. Consistent with our analyses of major

subsets in the previous section, we were able to confirm and extend our and other groups' findings

that more subtle T cell subsets are significantly depleted in patients relative to healthy controls

including subsets in the CD4^{+}, CD8^{+}, MAIT and γδ T cell compartments (Supplementary Figure

2). However, while we gained valuable insight into the altered T cell response in COVID-19

patients, none of these T cell subsets were prognostically useful in separating the Long-Stay/Died

patients from the Short-Stay patients.

Combined evaluation of immune parameters as a tool to predict clinical outcome

Since deeper analyses of multiple cytokines and cell subsets at ICU admission revealed significant

differences between the Long-Stay/Died and Short-Stay groups, we sought to combine these

findings to generate a streamlined prognostic tool that could accurately predict whether a patient,

newly admitted to ICU, was likely to have a subsequent longer stay in ICU or die. Although both

serum TNFα and serum IL-10 were significantly elevated in Long-Stay/Died patients relative to

Short-Stay patients in the Cytokine Cohort, using Pearson analyses, the length of stay in ICU

correlated more significantly with serum IL-10 levels ($R^2 = 0.48$) and maximum IL-10 levels (R^2

$= 0.54$) than with serum levels of TNFα ($R^2 = 0.14$). Thus, we proceeded with serum IL-10, only,

as our cytokine-based pre-screen portion of a stepwise integrated prognostic tool. As the first step,

using a cut-off value of 15pg/ml for serum IL-10, data from the 90-sample Cytokine Cohort

demonstrated a 79% specificity and 55% sensitivity in predicting that a patient newly admitted to

ICU would have a longer stay in ICU or die (Fig. 4g). This prognostic specificity of 79% is

somewhat comparable with that seen in the smaller subsets of the Cytokine Cohort, namely the

14-sample Initial Cohort (86%) and the 28-sample Replication Cohort (100%). Similarly, the

prognostic sensitivity of 55% in the Cytokine Cohort is somewhat comparable with that observed

in the Initial Cohort (86%) and the Replication Cohort (56%) (Figs. 4a-c,g). The variations in

estimates of prognostic sensitivity and specificity between cohorts (the Cytokine Cohort and its

two subsets of the Initial Cohort and Replication Cohort), however, likely demonstrate variations

that reflect the influences of random patient sampling and, very importantly, cohort size. These

results validate serum IL-10 levels as a pre-screen for patients likely to die or to experience a long

ICU stay.

We then explored the utility of combining serum IL-10 levels (with a cut-off value of 15pg/ml)

with the levels of CD11c^{low} classical monocytes (with a cut-off value of 2.7×10^7/ml) as a stepwise

integrated diagnostic tool. With this approach, 100% of the Long-Stay/Died patients were correctly

identified in the Initial Cohort and 88% of Long-Stay/Died patients were correctly identified in the

Replication Cohort, the latter with a specificity of 100% (Figs. 4d,e,g). These analyses of all 42

patients in the combined Initial and Replication Cohorts (n = 42) allowed us to predict with 91%

sensitivity and 91% specificity the clinical outcome of COVID-19 patients newly admitted to ICU

with respect to the likelihood of extended stay or death in ICU (Figs. 4f,g). Thus, our results

suggest that a simple screen of two biomarkers at the time of ICU admission allows for rapid

identification of those patients who are likely to die or require extended ICU care (Fig. 4h) and has

clear implications for patient care and health care delivery.

Discussion

The goal of the present study was to identify prognostic biomarkers that, at time of ICU admission,

could predict subsequent outcome of COVID-19. Such markers are in urgent need and, with further

testing and refinement, could serve to triage patients into specific groups for timely and appropriate

care while, at the same time, offer insights into the immune-mediated determinants of disease

response. Like many previous studies, we found that although severe COVID-19 is linked to broad

shifts in peripheral blood immune subsets (increased PMNs and T cell lymphopenia) and increased

blood inflammatory markers (CRP, D-dimer, etc.), none of these proved prognostic with respect

to subsequent length of ICU stay and/or death. Therefore, we used CyTOF-based PBMC

immunophenotyping and serum cytokine analyses on samples drawn at ICU admission to focus

our attention on more subtle shifts in inflammatory parameters with a view to identifying

prognostic biomarkers. Through iterative empirical analyses of these data, we identified two

groups of ICU patients who would subsequently have clinically distinguishable disease outcomes:

those who would be discharged from ICU within 6 days and those who would require a longer

ICU stay or die. We then used retrospective analyses to generate a simple set of biomarkers that

could easily be applied in the clinic to identify, at the time of ICU admission, those patients at

greater risk of death or lengthy stay in ICU.

A high admission level of serum IL-10 ($\geq$ 15pg/ml), alone, was (in a cohort of 90 patients) a

singular powerful biomarker that identified patients in the Long Stay/Died group with 79%

specificity, though with a lower sensitivity of 55%. Additional high dimensional cell surface

protein analyses of 42 patients revealed a simple set of monocyte markers, specifically those

identifying CD11c^{low} classical monocytes, that when combined with admission serum IL-10 levels

accurately predicted with 91% specificity and 91% sensitivity at the time of ICU admission

patients who would subsequently either have a longer stay in ICU or who would die (validated in

initial and replication cohorts). Thus, based on the information from our evaluation of 4 serum

cytokines and 38 surface markers and validated on two separate clinical cohorts, we have distilled

our prognostic screen down to a composite test of one cytokine and one monocyte subset as

predictive biomarkers that could be evaluated in most clinical laboratories. Indeed, our

demonstration that the cellular biomarker can be detected and visualized in two dimensional

analyses (Fig 3f) using limited markers reinforces the likelihood that this biomarker will be

detectable using conventional clinical flow cytometry.

Although individually several of the biomarkers examined here have been investigated previously

and described as markers of disease severity, there has been a lack of clear consensus on their

prognostic utility in the published literature. For example, both IL-6 and IL-10 emerged early as

candidate clinical markers of disease severity, but to our knowledge are not widely used in standard

prognostic testing at hospital or ICU admission[32,33,36,37]. This likely reflects the fact that, used in

isolation and without a detailed quantitative evaluation of threshold levels predicting outcome,

their presence or absence provides a more superficial indication of current inflammatory status and

fails to predict the temporal trajectory of clinical disease (increasing or decreasing severity). This

also may explain why anti-IL-6 receptor therapy has shown only limited efficacy as a broad-

spectrum therapeutic for severe COVID-19 and fails to reduce overall mortality[13,47]. Similarly,

although corticosteroids have emerged as a standard-of-care for COVID-19 ICU patients and

undoubtedly provide improved recovery after infection, they are widely recognized as "double-

edged swords": while they are effective at suppressing excessive inflammation, they also potently

suppress adaptive immune responses, potentially reducing viral clearance and increasing

susceptibility to secondary infections[12,48]. With that backdrop, a benefit of the streamlined ICU

biomarker panel described here is that it provides a direct prognostic link to patient outcome and

may also serve as a biomarker panel for monitoring patient response to therapies.

In our study, deep immunophenotyping of the myeloid compartment in COVID-19 patients proved

pivotal in defining markers to predict patient outcomes. While we saw no significant early changes

in total monocyte numbers or total classical monocyte numbers or frequencies, a prognostically

useful monocytic subset was contained within these broader subsets which highlights the need for

a high-dimensional evaluation to identify subtle, but informative, changes in immune

subpopulations that might otherwise have been overlooked. After identification of such subtle

biomarkers using high-dimensional analytic technologies, simpler two-dimensional technologies

(using limited markers) can then be used to measure the biomarker clinically. It is noteworthy that

previous studies have linked both increased peripheral blood monocytes and increased numbers of

inflammatory macrophages in the lung to severity of COVID-19[49,50,51,52]. Other studies reported

subtle, monocyte subset-specific changes in severe COVID-19, including dysfunctional pro-

inflammatory cytokine production, reduction in HLA-DR transcripts, accumulation of HLA-DR^{low}

monocytes and reduction of non-classical monocytes[16,17,34,38,42]. Further, CD11c^{low} monocytic

enrichment has been described in severe COVID-19[17]. The data presented here confirm and extend

these observations in a manner that facilitates accurate prognostication. They also reveal CD11c[low]

classical monocytes as new target populations for more focused mechanistic studies in future

research. While the combination of these two biomarkers certainly provides prognostic

information on disease outcome in COVID-19, there is a possible parallel interpretation of the

results: since all patients received corticosteroids at the time of admission, the biomarkers

described here may actually be identifying those patients who are, in fact, more responsive to

corticosteroid therapy. We leave this intriguing possibility for future investigation.

Although not specifically addressed here, we believe that these prognostic biomarkers provide a

roadmap for future studies aimed at guiding and monitoring response to therapy. Such monitoring

is particularly important in the context of therapies that have the acknowledged potential of

exacerbating clinical disease if given in a temporally inappropriate manner in the COVID-19 cycle

of stimulation and progression to clearance and resolution. While we have focused here on the

utility of these markers at the time of ICU admission it is possible that these may prove even more

valuable as temporal monitoring tools for revealing disease trajectory on this continuum and

responses to therapeutic intervention.

Methods

Patients, controls, and clinical information

This study was approved by the University of British Columbia Clinical Research Ethics board (H20-00685) and patient blood was collected at St. Paul's Hospital and Vancouver General Hospital in Vancouver, BC following informed consent. All COVID-19 patients had a positive nasal or tracheal real time reverse transcription polymerase chain reaction (RT-PCR) SARS-CoV-2 test. To avoid unnecessary virus exposure, patient blood was collected in concert with routine care. Patient samples (n = 90) were collected within 48 hours of ICU admission between November 2020 and February 2021. Patient demographics and clinical information are listed in table 1. Patient samples were transported to the main campus of the University of British Columbia for further processing. Healthy control blood samples (n = 8) were collected from age-matched volunteers who showed no COVD-19 symptoms or other illnesses and that had no history of COVID-19.

Specimen collection and isolation

Blood designated for peripheral blood mononuclear cell (PBMC) analysis was collected into citrate coated BD Vacutainer™ Glass Mononuclear Cell Preparation (CPT) tubes and PBMCs were isolated within four hours following collection according to manufacturer guidelines. Red blood cell lysis was performed with ACK lysing buffer (Gibco) for 10 minutes. Isolated PBMCs were frozen in fetal bovine serum (Gibco) with 10% dimethyl sulfoxide and stored in liquid nitrogen. Blood designated for serum analysis was collected into BD Vacutainer™ Serum Separation Tubes (SST) and allowed to clot for at least 30 minutes prior to centrifugation at 1200 rcf for 10 minutes and serum collection and storage at -80 °C.

Antibody staining and CyTOF data collection

Frozen PBMCs were thawed at 37 °C and washed with RPMI 1640 containing 10% FBS and 25U

nuclease (Thermo Fisher Scientific). Between 1- 4 x 10^6 cells per sample were used for antibody

staining. Prior to fixation, all centrifugation steps were performed at 500 rcf and 4 °C. All reagent

dilutions were prepared according to manufacturer instructions unless stated otherwise. For

live/dead cell analysis, PBMCs were stained with Cell-ID™ Intercalator-Rh (Fluidigm) for 15

minutes at 37 °C and washed with MaxPar® MCSB. Prior to surface staining, cells were incubated

with human TruStain FcX™ (Biolegend) for 15 minutes at 4 °C and stained with a surface antibody

cocktail for 30 minutes at RT (see Supplemental Table 2 for complete list of antibodies). The MR1-

5-OP-RU tetramer was incubated together with the antibody cocktail. After incubation, the cells

were washed and incubated for 30 minutes at RT with the secondary anti-APC antibody. Prior to

fixation and nuclear staining, PBMCs were washed with MaxPar® MCSB and incubated in

MaxPar® Fix and Perm Buffer (Fluidigm) and Cell-ID™ Intercalator-IR (Fluidigm) for 1 hour.

Post fix, all centrifugation steps were performed at 900 rcf and 4 °C. To prepare for CyTOF

acquisition, PBMCs were washed with MilliQ water and resuspended in EQ™ Four Element

Calibration Beads (Fluidigm). Samples were acquired with a CyTOF®2 mass cytometer. An

average of 400,000 events were collected for each sample at a flow rate of 45µl/min.

Cytokine data collection

374 Serum cytokines IL-6, IL-10, TNFα and IL-1β were quantified using the Simoa HD-1 platform

375 from Quanterix (Billerica, MA) according to manufacturing guidelines and as specified by Stukas

376 *et. al.*[53].

Data processing

All data files were normalized (https://github.com/nolanlab/bead-normalization) and events of

interest were manually gated with the FlowJo gating software (BD Biosciences). Dimensionality

reduction and clustering were performed with Uniform Manifold Approximation and Projection

(UMAP) and Rphenograph respectively, as provided in the bioconductor package Cytofkit

(https://github.com/JinmiaoChenLab/cytofkit2). The input files were equally down sampled and

cytofAsinh was used as the transformation method. The Rphenograph k was set to the default of

30. The dimensionality reduction and clustering were both performed on the entire data set as well

as separately on manually pre-gated populations. Populations were identified manually based on

marker expression (e.g., T cells were identified based on expression of CD3, CD4 or CD8).

Statistical analysis and figures

Sample size and statistical tests are indicated in figure legends and all graphs and statistical tests

were generated using GraphPad Prism (GraphPad Software, La Jolla California, USA). A test was

considered statistically significant at a probability of $< 5\%$ ($p < 0.05$) and we did not assume a

Gaussian distribution. Error bars represent mean $\pm$ SD. UMAP plots and heatmaps were exported

from Cytofkit and experimental outline figures, including the graphical abstract were created using

BioRender.com. Figures were assembled in Microsoft PowerPoint.

Acknowledgements

This study was supported by the St. Paul's Foundation. We thank the healthcare teams from St. Paul's Hospital and Vancouver General Hospital as well as the technical support from the Biomedical Research Centre core facility members Michael Williams (AbLab) and Andy Johnson (ubcFLOW cytometry). We acknowledge that the following reagent was obtained through the NIH Tetramer Core Facility: APC coupled MR1-5-OP-RU tetramer. Finally, we thank Jeff Biernaskie and Bryan Yipp at the University of Calgary for helpful discussions and generous seed funding from the Thistledown Foundation.

Contributions

M.M. was responsible for patient sample processing and mass cytometry experiments. K.M.M., K.B.L. and M.M. were responsible for study design, data analyses, interpretation, and manuscript preparation. M.S.S., R.L.H. and D.D.S. provided patient samples and clinical information. M.R.H. managed ethics approvals and assisted with data analyses and interpretation. S.S., C.W., J.C. and N.A. processed serum samples and acquired serum data. S.B.S., Y.L., L.W.T. and M.H. provided technical assistance for patient sample processing and mass cytometry data collection and processing. All authors contributed to editing the manuscript.

Data availability

All data produced in the present study are available upon reasonable request to the authors.

References

1. Cevik, M., Kuppalli, K., Kindrachuk, J. & Peiris, M. Virology, transmission, and pathogenesis of SARS-CoV-2. *BMJ* **371**, (2020).
2. Wiersinga, W. J., Rhodes, A., Cheng, A. C., Peacock, S. J. & Prescott, H. C. Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019 (COVID-19): A Review. *JAMA* **324**, 782–793 (2020).
3. Wang, D. *et al.* Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* **323**, 1061–1069 (2020).
4. Matthay, M. A. *et al.* Acute respiratory distress syndrome. *Nat. Rev. Dis. Prim.* **5**, (2019).
5. Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497–506 (2020).
6. Chen, G. *et al.* Clinical and immunological features of severe and moderate coronavirus disease 2019. *J. Clin. Invest.* **130**, 2620–2629 (2020).
7. Lavezzo, E. *et al.* Suppression of a SARS-CoV-2 outbreak in the Italian municipality of Vo'. *Nat. 2020 5847821* **584**, 425–429 (2020).
8. Carotti, M. *et al.* Chest CT features of coronavirus disease 2019 (COVID-19) pneumonia: key points for radiologists. *Radiol. Med.* **125**, 1 (2020).
9. Parekh, M., Donuru, A., Balasubramanya, R. & Kapur, S. Review of the Chest CT Differential Diagnosis of Ground-Glass Opacities in the COVID Era. *https://doi.org/10.1148/radiol.2020202504* **297**, E289–E302 (2020).
10. Feng, Z. *et al.* Early prediction of disease progression in COVID-19 pneumonia patients with chest CT and clinical characteristics. *Nat. Commun.* **11**, (2020).
11. Chen, J. S. *et al.* Nonsteroidal Anti-inflammatory Drugs Dampen the Cytokine and Antibody Response to SARS-CoV-2 Infection. *J. Virol.* **95**, (2021).
12. Jung, C. *et al.* Steroid use in elderly critically ill COVID-19 patients. *Eur. Respir. J.* 2100979 (2021) doi:10.1183/13993003.00979-2021.
13. Salama, C. *et al.* Tocilizumab in Patients Hospitalized with Covid-19 Pneumonia. *N. Engl. J. Med.* **384**, 20–30 (2021).
14. Mehta, P. *et al.* COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* **395**, 1033–1034 (2020).
15. Combes, A. J. *et al.* Global absence and targeting of protective immune states in severe COVID-19. *Nat. 2021 5917848* **591**, 124–130 (2021).
16. Wilk, A. J. *et al.* A single-cell atlas of the peripheral immune response in patients with severe COVID-19. *Nat. Med. 2020 267* **26**, 1070–1076 (2020).
17. J, S.-S. *et al.* Severe COVID-19 Is Marked by a Dysregulated Myeloid Cell Compartment. *Cell* **182**, 1419-1440.e23 (2020).
18. ML, M. *et al.* A neutrophil activation signature predicts critical illness and mortality in COVID-19. *Blood Adv.* **5**, 1164–1177 (2021).
19. Wilk, A. J. *et al.* Multi-omic profiling reveals widespread dysregulation of innate immunity and hematopoiesis in COVID-19. *J. Exp. Med.* **218**, (2021).
20. Del Valle, D. M. *et al.* An inflammatory cytokine signature predicts COVID-19 severity and survival. *Nat. Med.* **26**, 1636–1643 (2020).
21. Overmyer, K. A. *et al.* Large-Scale Multi-omic Analysis of COVID-19 Severity. *Cell Syst.* **12**, 23-40.e7 (2021).
22. Park, J. *et al.* In-depth blood proteome profiling analysis revealed distinct functional

characteristics of plasma proteins between severe and non-severe COVID-19 patients. *Sci. Rep.* **10**, (2020).

23.	Lee, J. S. *et al.* Immunophenotyping of COVID-19 and influenza highlights the role of type I interferons in development of severe COVID-19. *Sci. Immunol.* **5**, (2020).

24.	Luo, X. *et al.* Prognostic Value of C-Reactive Protein in Patients With Coronavirus 2019. *Clin. Infect. Dis.* **71**, 2174–2179 (2020).

25.	Shu, T. *et al.* Plasma Proteomics Identify Biomarkers and Pathogenesis of COVID-19. *Immunity* **53**, 1108-1122.e5 (2020).

26.	Caricchio, R. *et al.* Preliminary predictive criteria for COVID-19 cytokine storm. *Ann. Rheum. Dis.* **80**, 88–95 (2021).

27.	Heller, R. A. *et al.* Prediction of survival odds in COVID-19 by zinc, age and selenoprotein P as composite biomarker. *Redox Biol.* **38**, (2021).

28.	Henry, B. M., De Oliveira, M. H. S., Benoit, S., Plebani, M. & Lippi, G. Hematologic, biochemical and immune biomarker abnormalities associated with severe illness and mortality in coronavirus disease 2019 (COVID-19): a meta-analysis. *Clin. Chem. Lab. Med.* **58**, 1021–1028 (2020).

29.	Fouladseresht, H. *et al.* Predictive monitoring and therapeutic immune biomarkers in the management of clinical complications of COVID-19. *Cytokine Growth Factor Rev.* **58**, 32–48 (2021).

30.	Majumder, J. & Minko, T. Recent Developments on Therapeutic and Diagnostic Approaches for COVID-19. *AAPS J.* **23**, (2021).

31.	Demichev, V. *et al.* A time-resolved proteomic and prognostic map of COVID-19. *Cell Syst.* **12**, 780-794.e7 (2021).

32.	T, H. *et al.* Elevated levels of IL-6 and CRP predict the need for mechanical ventilation in COVID-19. *J. Allergy Clin. Immunol.* **146**, 128-136.e4 (2020).

33.	Han, H. *et al.* Profiling serum cytokines in COVID-19 patients reveals IL-6 and IL-10 are disease severity predictors. (2020) doi:10.1080/22221751.2020.1770129.

34.	Silvin, A. *et al.* Elevated Calprotectin and Abnormal Myeloid Cell Subsets Discriminate Severe from Mild COVID-19. *Cell* **182**, 1401-1418.e18 (2020).

35.	Laing, A. G. *et al.* A dynamic COVID-19 immune signature includes associations with poor prognosis. *Nat. Med. 2020 2610* **26**, 1623–1635 (2020).

36.	Mazzoni, A. *et al.* Impaired immune cell cytotoxicity in severe COVID-19 is IL-6 dependent. *J. Clin. Invest.* **130**, 4694–4703 (2020).

37.	Y, Z. *et al.* Longitudinal COVID-19 profiling associates IL-1RA and IL-10 with disease severity and RANTES with mild disease. *JCI insight* **5**, (2020).

38.	Arunachalam, P. S. *et al.* Systems biological assessment of immunity to mild versus severe COVID-19 infection in humans. *Science (80-.).* **369**, 1210–1220 (2020).

39.	Mahler, M., Meroni, P. L., Infantino, M., Buhler, K. A. & Fritzler, M. J. Circulating Calprotectin as a Biomarker of COVID-19 Severity. *Expert Rev. Clin. Immunol.* **17**, 431–443 (2021).

40.	Maucourant, C. *et al.* Natural killer cell immunotypes related to COVID-19 disease severity. *Sci. Immunol.* **5**, 6832 (2020).

41.	S, D. *et al.* Mucosal-Associated Invariant T (MAIT) Cells Are Highly Activated and Functionally Impaired in COVID-19 Patients. *Viruses* **13**, (2021).

42.	Bonnet, B. *et al.* Severe COVID-19 is characterized by the co-occurrence of moderate cytokine inflammation and severe monocyte dysregulation. *EBioMedicine* **73**, (2021).

43. Chevrier, S. *et al.* A distinct innate immune signature marks progression from mild to severe COVID-19. *Cell Reports Med.* **2**, 100166 (2021).
44. Wang, W. *et al.* High-dimensional immune profiling by mass cytometry revealed immunosuppression and dysfunction of immunity in COVID-19 patients. *Cell. Mol. Immunol. 2020 176* **17**, 650–652 (2020).
45. Peckham, H. *et al.* Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ITU admission. *Nat. Commun. 2020 111* **11**, 1–10 (2020).
46. Guan, W. *et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N. Engl. J. Med.* **382**, 1708–1720 (2020).
47. B, F. COVACTA trial raises questions about tocilizumab's benefit in COVID-19. *Lancet. Rheumatol.* **2**, e592 (2020).
48. Ghanei, M. *et al.* The efficacy of corticosteroids therapy in patients with moderate to severe SARS-CoV-2 infection: a multicenter, randomized, open-label trial. *Respir. Res.* **22**, 1–14 (2021).
49. Liao, M. *et al.* Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19. *Nat. Med. 2020 266* **26**, 842–844 (2020).
50. Melms, J. C. *et al.* A molecular single-cell lung atlas of lethal COVID-19. *Nat. 2021 5957865* **595**, 114–119 (2021).
51. Wauters, E. *et al.* Discriminating mild from critical COVID-19 by innate and adaptive immune single-cell profiling of bronchoalveolar lavages. *Cell Res.* **31**, 272–290 (2021).
52. Xu, G. *et al.* The differential immune responses to COVID-19 in peripheral and lung revealed by single-cell RNA sequencing. *Cell Discov.* **6**, (2020).
53. Stukas, S. *et al.* The Association of Inflammatory Cytokines in the Pulmonary Pathophysiology of Respiratory Failure in Critically Ill Patients With Coronavirus Disease 2019. *Crit. Care Explor.* **2**, e0203 (2020).

Table 1. Cohort clinical information

Characteristics	Healthy Controls (n = 8)	COVID-19 Patients (n=90)[1]
Age (mean years)	50.1	63.5
Sex (M:F)	5:3	56:34
Diagnosis	Asymptomatic/ healthy	SARS-COV-2 +
Collection timepoint (days)	NA	0-2 post admission
Severity	NA	ICU
BMI (mean)	NA	30.9
CRP (mean mg/L)	NA	75.6
D-dimer (mean ng/ml)	NA	3064.3
Length of stay (Short-Stay:Long-Stay/Died)[2]	NA	34:56
Days in ICU (mean)	NA	9.3
Outcome (n; recovered:died)	NA	74:16
Mechanically ventilated (n, %)	NA	45, 60
PBMCs (mean 10^9/ml)	2.5	2.1
PMNs (mean 10^9/ml)	3.9	8.6

[1]BMI, D-dimer, CRP and ventilation information was not available for 16/90 patients; calculations were adjusted accordingly.
[2]Short-Stay: <6 days; Long-Stay/Died: ≥ 6 days in ICU or Died.
BMI: Body mass index; CRP: C-reactive protein; PBMCs: Peripheral blood mononuclear cells; PMNs: Polymorphonuclear leukocytes

Fig. 1: Patient cohort selection, characteristics and cytokine analyses.

a, Experimental design overview: peripheral blood was collected from COVID-19 patients within 48h of ICU admission; immune cells and serum were isolated and stored followed by immune cell subset and cytokine analyses and clinical data integration. **b,** patient cohorts overview. **c,** patient outcome groups based on length of stay in ICU. (**d-e),** Patient age, body mass index (BMI), C-reactive protein (CRP) levels and D-dimer levels. **f,** Complete blood counts of patients and healthy controls. (**g-i),** Serum cytokine levels of IL-10, TNFα, maximum IL-10, maximum TNFα, IL-6 and IL-1β. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ns, $p \geq 0.05$ by two-tailed, two-sample unequal variance Student's t-Test.

Fig. 2: Major PBMC subsets fail to identify Long-Stay/Died patients in the Replication Cohort.

a, UMAP projection of ungated CyTOF-derived data from the replication cohort (n=28). **b,** proportion of immune cell subsets in Healthy Controls (HC), Short-Stay (SS) and Long-Stay/Died (LS/D) patient outcome groups **c,** Mean marker expression heatmap of clusters shown in a. **d,** Absolute counts of adaptive PBMC subsets (CD4 T, CD8 T, B), **e,** innate and unconventional subsets (NK, MAIT, γδ T, pDC, DC2/3), **f,** monocytes and stem cells. *, p < 0.05; **, p < 0.01; ***, p < 0.001; ns, p ≥ 0.05 by two-tailed, two-sample unequal variance Student's t-Test.

Fig. 3: CD11c^low Classical Monocytes are predictive of clinical outcome.

a, Representative gating of CD116+CD3-CD19- cells. **b,** Initial Cohort UMAP projections of CD116+ CD3-CD19- gated cells (all samples combined; limited clustering channels) and mean marker expression heatmap. **c,** same as in b but for the Replication Cohort. **d,** Initial (top) and Replication Cohorts (bottom) absolute counts of monocyte subset predictive of clinical outcome. **e-f,** Absolute counts of non-significant monocyte subset identified based on gated clustering (top row: Initial Cohort, bottom row: Replication Cohort). **g,** Manual gating strategy to view predictive monocyte subset. *, p < 0.05; **, p < 0.01; ***, p < 0.001; ns, p ≥ 0.05 by two-tailed, two-sample unequal variance Student's t-Test.

Fig. 4: Prognostic cytokine and cellular biomarkers predict clinical outcome.

g	Sensitivity	Specificity	False Positives	False Negatives
IL-10 screen (Cytokine Cohort; n = 90)	55% (31/56)	79% (31/39)	21% (8/39)	45% (25/56)
IL-10 screen (Initial Cohort; n = 14)	86% (6/7)	86% (6/7)	14% (1/7)	14% (1/7)
IL-10 screen (Replication Cohort; n = 28)	56% (9/16)	100% (9/9)	0% (0/9)	44% (7/16)
IL-10 screen (Combined; n = 42)	61% (14/23)	93% (14/15)	7% (1/15)	39% (9/23)
IL-10 and monocyte screen (Initial Cohort; n = 14)	100% (7/7)	78% (7/9)	22% (2/9)	0% (0/7)
IL-10 and monocyte screen (Replication Cohort; n = 28)	88% (14/16)	100% (14/14)	0% (0/14)	12% (2/16)
IL-10 and monocyte screen (Combined Cohorts; n = 42)	91% (21/23)	91% (21/23)	9% (2/23)	9% (2/23)

a-c, Cytokine levels scatter plots of Initial (left), Replication (middle) and combined Cohorts (right) with dashed lines at cut-off value of 15pg/ml for serum IL-10. **d-f,** Cytokine and cellular levels scatter plots for Initial (left) Replication (middle) and combined cohorts (right) with dashed lines at cut-off values of 15pg/ml and 2.7 (x10^7/ml) of serum IL-10 and CD11c^{low} classical monocytes respectively. **g,** Sensitivity and specificity calculations for each screen and cohort. **h,** Prognostic patient screening chart based on serum IL-10 and CD11c^{low} monocyte subset measurement. *, p < 0.05; **, p < 0.01; ***, p < 0.001; ns, p ≥ 0.05 by two-tailed, two-sample unequal variance Student's t-Test and R^2 by two-tailed Pearson correlation with 95% confidence interval.

Fig. S1: Major PBMC subsets fail to identify Long-Stay/Died patients in the Initial Cohort.

a, UMAP projection of ungated CyTOF-derived data from the Initial cohort (n=14). **b,** proportion of immune cell subsets in Healthy Controls (HC), Short-Stay (SS) and Long-Stay/Died (LS/D) patient outcome groups **c,** Mean marker expression heatmap of clusters shown in a. **d,** Absolute counts of adaptive PBMC subsets (CD4 T, CD8 T, B), **e,** innate and unconventional subsets (NK, MAIT, γδ T, pDC, DC2/3), **f,** monocytes and stem cells. *, p < 0.05; **, p < 0.01; ***, p < 0.001; ns, p ≥ 0.05 by two-tailed, two-sample unequal variance Student's t-Test.

Fig. S2: T cell subsets fail to identify Long-Stay/Died patients.

a, Representative gating of CD3+ cells. **b,** Initial Cohort UMAP projections of CD3+ gated cells (all samples combined; limited clustering channels) and mean marker expression heatmap. **c,** same as in b but for the Replication Cohort. **d,** Initial Cohort absolute counts of T cell subsets identified based on gated clustering. **e,** same as in d but for the Replication Cohort. *, p < 0.05; **, p < 0.01; ***, p < 0.001; ns, p ≥ 0.05 by two-tailed, two-sample unequal variance Student's t-Test.

Table S1. List of CyTOF antibodies

Antibody/Tetramer	Metal/Fluorophore*	Clone	Company	Used in Cohort
CD1c	151Eu	L161	**Biolegend**	Replication
CD3ε	143Nd	OKT3	**Biolegend**	both
CD4	174Yb	SK3	**Biolegend**	both
CD8α	168Er	SK1	**Biolegend**	both
CD8β	141Pr	SIDI8BEE	**eBioscience**	both
CD11c	147Sm	Bu15	**Biolegend**	both
CD14	153Eu	M5E2	**Biolegend**	both
CD16	158Gd	3G8	**Biolegend**	both
CD19	142Nd	HIB19	**Biolegend**	both
CD25	169Tm	BC96	**Biolegend**	both
CD27	175Lu	O323	**Biolegend**	Replication
CD31	145Nd	WM59	**Biolegend**	both
CD32	160Gd	IV.3	**Stemcell Technologies**	both
CD34	156Gd	581	**Biolegend**	both
CD38	106Cd	HIT2	**Biolegend**	both
CD45	89Y	HI30	**Biolegend**	both
CD45RA	110Cd	HI100	**Biolegend**	both
CD45RO	112Cd	UCHL1	**Biolegend**	both
CD56	148Nd	NCAM16.2	**BD Bioscience**	both
CD94 (NKG2C)	161Dy	DX22	**Biolegend**	both
CD116	150Nd	4H1	**Biolegend**	both
CD117	171Yb	104D2	**Biolegend**	Initial
CD123	164Dy	6H6	**Biolegend**	both
CD127	165Ho	A019D5	**Biolegend**	both
CD161	159Tb	HP-3G10	**Biolegend**	both
CD197/CCR7	171Dy	G043H7	**Biolegend**	Replication
CD200R	173Yb	0X-108	**BD Bioscience**	both
CD294 (CRTH2)	163Dy	BM16	**Biolegend**	both
CD301 (CLEC10A)	154Sm	H037G3	**Biolegend**	Replication
CD304 (NRP1)	172Yb	12C2	**Biolegend**	Replication
FcεRα1	176Yb	AER-37	**Biolegend**	both
HLA-DR	170Er	L243	**Biolegend**	both
IgD	116Cd	IA6-2	**Biolegend**	both
IL-18Rα	162Dy	H44	**Biolegend**	both
KLRG1	144Nd	SA231A2	**Biolegend**	both
LAG3	175Lu	11C3C65	**Biolegend**	Initial
TCRαβ	155Gd	T10B9.1A-31	**BD Bioscience**	both
TCRγδ	152Sm	B1	**Biolegend**	both
TRAV1-2	115Ln	3C10	**Biolegend**	both
Anti-APC (Secondary)	149Sm	APC003	**Biolegend**	both
MR1-5-OP-RU (Primary)	APC	NA	**NIH Tetramer Facility**	both