

1 **Recommendations for an Optimal Model of integrated**  
2 **case detection, referral, and confirmation of Neglected**  
3 **Tropical Diseases: A case study in Bong County, Liberia**  
4

5 **Tiawanlyn G. Godwin-Akpan**<sup>1\*</sup>, Shahreen Chowdhury<sup>2</sup>, Emerson J. Rogers<sup>3</sup>, Karsor K.  
6 Kollie<sup>3</sup>, Fasseneh Z. Zaizay<sup>4</sup>, Anna Wickenden<sup>5</sup>, Georgina V.K. Zawolo<sup>6</sup>, Colleen B.M.C.  
7 Parker<sup>7</sup>, Laura Dean<sup>2</sup>

- 8  
9 1. AIM Initiative/American Leprosy Mission, Greenville, South Carolina, USA  
10 2. Department of International Public Health, Liverpool School of Tropical Medicine, Liverpool,  
11 United Kingdom  
12 3. Neglected Tropical Disease Program, Ministry of Health, Oldest Congo Town, Monrovia,  
13 Liberia  
14 4. Neglected Tropical Disease Program, Ministry of Health, Oldest Congo Town, Monrovia,  
15 Liberia. **Current Address: REDRESS, Oldest Congo Town, Monrovia, Liberia**  
16 5. AIM Initiative/American Leprosy Mission, Greenville, South Carolina, USA. **Current**  
17 **Address:** Effect Hope, Markham, ON, Canada  
18 6. Countdown Liberia, Pacific Institute for Research and Evaluation, Capitol Hill, University of  
19 Liberia, Monrovia, Liberia  
20 7. Research Unit, Ministry of Health, Liberia, Oldest Congo Town, Monrovia, Liberia

## 21 Abstract

### 22 Background

23 People affected by Neglected Tropical Diseases (NTDs), specifically leprosy, Buruli ulcer (BU),  
24 yaws, and lymphatic filariasis, experience significant delays in accessing health services, often  
25 leading to catastrophic physical, psychosocial, and economic consequences. Global health actors  
26 have recognized that Sustainable Development Goal 3:3 is only achievable through an integrated  
27 inter and intra-sectoral response. This study evaluated existing case detection and referral  
28 approaches in Liberia, utilizing the findings to develop and test an Optimal Model for integrated  
29 community-based case detection, referral, and confirmation. Finally, this study evaluates the  
30 efficacy of implementing the Optimal Model in improving the early diagnosis of NTDs.

### 31 Methodology/Principal Findings

32 The study used mixed methods, including key informant interviews, focus group discussions,  
33 participant observation, quantitative analysis, and reflexive sessions to evaluate the  
34 implementation of an Optimal Model developed through this study. The quantitative results from  
35 the testing of the optimal model are of limited utility. The annual number of cases detected  
36 increased in the twelve months of implementation in 2020 compared to 2019 (pre-intervention)  
37 but will require observation over a more extended period to be of significance. Qualitative data  
38 revealed essential factors that impact the effectiveness of integrated case detection. Data  
39 emphasized the gendered dynamics in communities that shape the case identification process,  
40 such as men and women preferring to see health workers of the same gender. Furthermore, the  
41 qualitative data showed an increase in knowledge of the transmission, signs, symptoms, and  
42 management options amongst CHW, which enabled them to dispel misconceptions and stigma  
43 associated with NTDs.

## 44 Conclusion/Significance

45 This study demonstrates the opportunity for greater integration in training, case detection,  
46 rereferral, and confirmations. However, the effectiveness of this approach depends on a high  
47 level of collaboration, joint planning, and implementation embedded within existing health  
48 systems infrastructure. Together, these approaches improve access to health services for NTDs.

## 49 Author Summary

50 Global health professionals and stakeholders have advocated for integration across diseases and  
51 sectors to improve the success of public health interventions. This advocacy has also impacted  
52 NTDs programs globally. NTDs interventions are becoming more integrated than disease-  
53 specific activities to maximize limited resources, improve coverage and access to healthcare  
54 services. However, documentation on the effectiveness of integrated approaches to improve  
55 access to healthcare services is minimal. This study evaluated existing case detection and referral  
56 approaches in Liberia, utilizing the findings to develop and test an Optimal Model for integrated  
57 community-based case detection, referral, and confirmation. Finally, this study evaluates the  
58 efficacy of implementing the Optimal Model in improving the early diagnosis of NTDs. The  
59 results provide evidence of the benefits of an integrated approach and the programmatic  
60 challenges to achieve the goal of improving access to health services for persons affected by  
61 NTDs.

## 62 Introduction

63 Globally, people affected by Neglected Tropical Diseases (NTDs), specifically leprosy, Buruli  
64 ulcer (BU), yaws, and lymphatic filariasis, experience significant delays in receiving diagnosis  
65 and treatment; this is due to multiple factors, including stigma, a lack of knowledge,

66 accessibility, and gender (1). However, compounding these challenges is the lack of the  
67 availability of appropriate disease management and disability inclusion (DMDI) services of  
68 reasonable quality in the community available through the health system (1,2). Social  
69 stigmatization and discrimination are very high for people with NTDs due to the visible physical  
70 impairments caused by many NTDs. Therefore, remaining hidden in their communities is a  
71 typical response to stigma (3,4). This stigma and lack of awareness of NTDs further contribute to  
72 the poor health-seeking behavior of people with NTDs (2,5–7). The time between symptoms,  
73 diagnosis, and treatment determines the treatment regimen and illness outcome (8). Early  
74 diagnosis and management reduce the risk of significant physical, economic, and psychosocial  
75 impacts of NTDs (9). Therefore, addressing the barriers to early case detection is essential to  
76 attaining universal health coverage and reducing the disease burden caused by NTDs; this  
77 includes a shift from a passive approach to case detection to an active model. The World Health  
78 Organization (WHO) has emphasized active case finding as part of the five essential intervention  
79 packages to prevent, control, and eliminate NTDs (10,11). Integrating active case finding into  
80 national NTD programs will promote active surveillance instead of passive data collection to  
81 ensure no one is left behind.

82  
83 NTDs affect more than a billion people worldwide, impacting low-income populations in Africa,  
84 Asia, and the Americas (8). NTDs are a diverse group of 20 tropical diseases broken into two  
85 major groups: preventive chemotherapy and transmission control (PCT) and innovative and  
86 intensified disease management (IDM). PCT NTDs are managed at the community level by mass  
87 drug administration (MDA). At the same time, IDM NTDs require early diagnosis and  
88 management on an individual basis (12). NTDs are an equity tracer for the Sustainable

89 Development Goals (SDGs), with specific targets. SDG sub-goal 3:3 states, "to end the  
90 epidemics of AIDS, tuberculosis, malaria, and neglected tropical diseases and combat hepatitis,  
91 water-borne diseases, and other communicable diseases by 2030" (13). Global health actors have  
92 recognized that this is only achievable through an integrated inter and intra sectoral response  
93 (14).

94  
95 This article's integration of NTDs refers to the joint planning, implementation, and evaluation of  
96 activities across programs to achieve common goals (15). Following the launch of the first WHO  
97 Roadmap for NTDs (8), emphasis and prioritization have been placed on providing preventative  
98 Mass Drug Administration (MDA) for PCT NTDs. There is far less emphasis and attention on  
99 diagnostic, curative, and morbidity management services for IDM NTDs (excluding leprosy).  
100 The (PCT) NTDs have been integrated since the early 2010s through MDA with evidence of  
101 success. However, the (IDM) NTDs lag as they require considerable resources, including  
102 personnel with skills and financial support (9,16,17).

103 Consequently, with restricted funding to specific IDM disease conditions (e.g., Buruli ulcer or  
104 leprosy), IDM NTD interventions have been vertical and cannot reach affected persons (10). As  
105 a result, the global commitment to "leave no one behind" cannot be achieved without support to  
106 NTD programs to identify, diagnose, and manage IDM NTDs(18). Furthermore, delay in  
107 diagnosis, treatment, and referral can lead to the ongoing transmission of those NTDs of  
108 epidemic concern, disability and morbidity challenges, and increased economic hardship for  
109 underprivileged families.

110

111 Liberia was one of the first countries to adopt an integrated approach to the case management of  
112 (IDM) NTDs to improve access and universal health coverage for people with NTDs. This  
113 approach responded to lessons learned from the Ebola Virus Disease (EVD) outbreak and two  
114 national policies: “Essential Package of Health Services” and “Investment Plan for building a  
115 resilient health system,” which were developed to build resilient health system (19,20). In line  
116 with these policies, the NTDs and the National Tuberculosis and leprosy control programs, along  
117 with several national and international partners, developed a "National Strategic Plan for the  
118 Integration of Case Management of NTDs (CM-NTDs) in Liberia (2016-2020)"(21). For Liberia  
119 context, CM-NTDs are synonymous with (IDM) NTDs. The plan extends the National NTDs  
120 Masterplan, which follows the WHO NTDs Masterplan template for planning PCT and IDM  
121 NTDs interventions. The strategic plan for integrated case management aligns with the four  
122 strategic priorities of the WHO NTDs Masterplan. These priorities are: 1) government  
123 ownership, advocacy, coordination, and partnerships, 2) resource mobilization and planning for  
124 results, 3) access to interventions and treatment improved and system capacity building, and 4)  
125 monitoring, surveillance, and operational research enhanced (21). The plan includes strategies to  
126 integrate BU, leprosy, clinical manifestations of lymphatic filariasis (hydrocele and  
127 lymphedema), and yaws for case detection, referral, confirmation, and management control to  
128 eliminate NTDs by 2020 (21,22). The fifteen counties in Liberia are all co-endemic for the  
129 targeted diseases requiring case management (21). Due to funding restraints in 2017, the NTD  
130 program launched the integrated plan for case management in five counties: Bong, Bomi, Lofa,  
131 Maryland, and Nimba, alongside the integration of national-level activities including budgeting,  
132 performance monitoring, data management, and drug supply chain. The NTD Program  
133 implemented multiple and diverse community-based case detection models in the pilot counties

134 to increase early case detection, as shown in Fig 1. However, the effectiveness of these differing  
135 models in increasing patient access to care and early diagnosis had not been documented.

<b>Three models aimed at improving early case detection in 2017</b>		
<b>(1)</b> The Community Health Assistant Model (CHA Model) recruited semi-literate community volunteers to identify and refer people with clinical manifestations in the skin typical to NTDs as part of their routine work. CHAs operate in communities outside a 5 kilometers radius from the nearest health facility and receive monthly incentives of \$70.00USD.	<b>(2)</b> The Community Health Volunteer Motivation Model (CHV Model) encouraged community volunteers to identify and report suspected DMDI cases within their communities. The CHVs received USD 5 per confirmed positive case identified.	<b>(3)</b> The Integrated Mass Drug Administration and DMDI Active Case Search Model (MDA-CM model) trained Community Directed Distributors (who usually include CHVs and CHAs) to conduct active case searches for DMDI cases while distributing NTD medications during MDA campaigns.

136  
137 **Fig 1.** NTDs program community-based case detection models implemented through the  
138 integrated case management approach in the five pilot counties in 2017.  
139

140 This study aimed to evaluate existing approaches to case detection and referral in Liberia using  
141 an action research cycle, use the findings to develop an optimal model for integrated community-  
142 based case detection, referral, and confirmation. Finally, this study evaluates the efficacy of the  
143 optimal model in improving early diagnosis NTDs. This paper presents findings from the  
144 formative research phase, describes the optimal model produced, and the key learnings from its  
145 implementation over twelve months.

## 146 147 **The Optimal Model**

148 The Optimal Model was developed in partnership with the National Ministry of Health NTD  
149 team, the Bong County Health Team, people affected by NTDs, and community health workers  
150 with support from national NTD program partners (American Leprosy Missions, Liverpool

151 School of Tropical Medicine, University of Liberia Pacific Institute for Research and Evaluation)  
152 through COR-NTD funding. There was a collaboration between the national program, lower  
153 levels of the health system, and persons affected by NTDs during the entire development  
154 process. The model was developed in two phases over nine months: phase one, the formative  
155 phase, and phase two, the planning phase. During phase one, we conducted a participatory  
156 qualitative evaluation of the program's three models of case detection to identify strengths and  
157 weaknesses. We used the results to design an Optimal Model for case detection. This phase drew  
158 on qualitative participatory methods, including key informant interviews and focus group  
159 discussions with health workers and affected persons on understanding their experiences of  
160 existing case detection approaches. During phase two, the planning phase, a two-day workshop  
161 was held in Monrovia, Liberia, with stakeholders including persons affected by NTDs,  
162 community health workers (CHWs), Bong County Health team, NTD program, and other health  
163 stakeholders from the central, county, and district levels to collaboratively develop the optimal  
164 model based on the findings from phase one. The findings included vital themes such as  
165 inconsistencies and delays in remuneration, geographical barriers, opportunity costs throughout  
166 pocket expenditure of CHWs on transport and calls, and insufficient training and supervision of  
167 DMDI services. Challenges regarding gender, disability, and other axes of social disadvantage  
168 patients face were also explored to understand how they affect case detection; poverty, social  
169 isolation, stigma, and gendered access were highlighted as critical issues. The optimal model  
170 included comprehensive training on integrated approaches to identify, refer, diagnose and  
171 manage NTDs at the community level. The training included a referral process, supervision  
172 structure, and incentive packages. The Optimal Model developed is summarized in Fig 2 and  
173 detailed in S1 File.

### **TRAINING**

A cascaded model of training was developed for CHWs, peer advocates (people affected by NTDs), and health workers as supervisors. An integrated training manual and visual aids on disease signs and symptoms were collaboratively developed to include context specific language. CHWs participated in one-day refresher training to identify, refer and report suspected NTD cases using the visual aids. The CHVs who participated in MDA training used the integrated manual. A one-day peer advocate training focused on peer support, case detection, and referral. Certificates were provided to all trainees on completion of training.

### **SUPERVISION**

A comprehensive supervision plan was developed that included tools, personnel, and responsibilities at all levels (national, county, district, Health Facility, community). A cascaded supervision structure was created to monitor and support the skills transferred during the training. This supervision focused on ensuring effective implementation of the training at all levels. The national NTD program provided a one-day supervision training for Community Health Services Supervisors (CHSS), District Health Officers (DHO), and NTDs Focal Persons.

### **REFERRAL**

Tools were developed to document persons making the referral and patients referred. CHWs received a USD 5 performance-based incentive per confirmed case identified. Furthermore, a communication structure (feedback/referral form) was implemented for the CHWs to inform the HW receiving the referred patient at the facility. In return, the CHW received feedback about the outcome (diagnosis) of the patient referred. This structure promoted the follow-up of referred patients by CHWs to ensure they visit the health facility and receive proper diagnosis and care.

### **LOGISTICS**

All trained case detectors received a logistics toolkit. The toolkit included a flashlight, batteries, T. shirt, and backpacks.

### **GENDER BALANCE**

A balance in gender was considered when selecting CHWs and peer advocates to promote equity and increase patient accessibility and acceptability as many women and men preferred to see CHWs of the same gender. the burden of shame incurred by both women and men during case identification and referral. (Was there gender balance?)

174

175 **Fig 2.** The optimal model for case detection, referral, and confirmation of NTDs

## 176 **Materials and Methods**

### 177 **Ethics Statement**

178 This assessment was approved by the University of Liberia-Pacific Institute for Research &  
179 Evaluation (UL-PIRE) Institutional Review Board (Protocol number: NTDS 167D) (S2 File).

### 180 **Consent and Data Collection Procedure**

181 Informed consent was obtained from each participant in the assessment before FGDs, KIIs, and  
182 reflexive sessions. The participants had an opportunity to ask questions about the study and the  
183 implications of their participation. Participants' confidentiality was assured, and the right to  
184 withdraw from the research at any time of the interview. Upon completing the consent process,  
185 each participant received a copy of their signed consent form (S3 File). Anonymizing names and  
186 other identifiers maintained confidentiality.

187  
188  
189  
190  
191  
192  
193  
194  
195  
196

## Study Setting

Bong County was purposively selected as the study setting for this research based on the following criteria: Bong County is one of the five counties of the integrated CM-NTDs pilot; co-endemic for all priority diseases; has mixed success in terms of case reporting across its nine health districts (i.e., some have identified more cases than others); and all three models of case detection (Fig 1) have been implemented in this setting. Furthermore, Bong shares borders with two integrated CM-NTDs implementation counties (Nimba and Lofa). Patients may travel between these counties for treatment (23). The study was conducted in Bong County, Liberia (Fig 3).




197  
198  
199

**Fig 3.** Map of Liberia showing the study site, Bong County, and other integrated CM-NTDs pilot counties

## 200 Study Design

201 The study uses a Community-Based Participatory Research (CBPR) approach comprising phases  
202 depicted in Fig 4. The CBPR approach promotes the equitable participation of community  
203 members, researchers, and all stakeholders throughout the research process (24). During phases  
204 1-2 (nine months period), the optimal model was developed (as described above) and  
205 implemented during phase 3 (one-year period). The findings presented in this study were  
206 identified in phase four (four months period), the observation and reflection phase, based on the  
207 activities in phases 1,2, and 3. Phase four used a mixed-methods approach to evaluate the  
208 implementation of the optimal model, which is both a process and outcome evaluation.  
209



210

211 **Fig 4.** Conceptual Framing, Theory of Change and Community Based Participatory Research  
 212 Phases

213

## 214 Qualitative Methods

215 During phase four, we conducted a mixed-method evaluation which included key informant  
 216 interviews, focus group discussions, participant observation, reflexive sessions. Participants were  
 217 purposively selected based on their role within delivering the optimal model to affected  
 218 populations and for maximum variation in gender and age. Furthermore, participants were  
 219 selected from all health system levels and had experience in integrated case detection, referral,  
 220 and confirmation of NTDs or benefited from NTD services.

221

222 **Data Collection**

223 Table 1 presents the different data collection methods for the evaluation and participants.

224 **Table 1. Overview of Data Collection**

<b>Data Collection</b>	<b>Number of Sessions</b>	<b>Number of Participants</b>	<b>Total Participants</b>
<i><b>Process Evaluation</b></i>			
A reflexive session with the Research team	1	5	5
A reflexive session with the County Health team	1	2	2
Participant observation at CHW training	1	760	1
Participant observation at supervision training	1	48	1
Participant observation at patient advocate training	1	36	1
<i><b>Outcome Evaluation</b></i>			
Key Informant Interviews with CHSS	4	1	4
FGD with CHWs at health facilities	4	5	20
FGD with patient advocates after training	1	5	5
Quantitative Analysis of Health Systems Data	N/A	N/A	N/A
<b>Total</b>	<b>14</b>	<b>21</b>	<b>39</b>

225

226 **Key Informant Interviews**

227 Four health facilities in Bong were purposively selected based on the geographical variation. A

228 CHSS trained to supervise CHW in the community in each facility was interviewed as a key

229 informant, using an interview guide. We were particularly interested in their experiences and

230 perspectives of training and supervision, specifically, how the optimal model has impacted case

231 identification, referral, and confirmation at different health system levels.

232

## 233 Focus Group Discussions

234 Five FGDs were conducted with community health workers and peer advocates based on their  
235 roles and training. We were interested in their training experience and how cases were identified,  
236 referred to, and confirmed using the optimal model at different health system levels. Separate  
237 group discussions were held with CHWs and patient advocates. Each focus group discussion had  
238 between 5-8 participants.

## 239 Participant Observation

240 Participant observation was conducted at the training of CHWs, supervisors, and peer advocates.  
241 A structured observation grid was used to document reflections and observations of interactions,  
242 participation, and training content.

## 243 Reflexive Sessions

244 Reflexive sessions involving a participatory group discussion to reflect on the optimal model's  
245 design, delivery, and evaluation were conducted with the core research team and the County  
246 Health Team. The core research team remotely participated in a reflexive session over Skype. In  
247 contrast, a reflexive session with the county health team was conducted at the health facility,  
248 including the NTD focal person and the Community Health focal person in Bong County. The  
249 reflexive session included key themes on collaboration, participation and power, the use of  
250 evidence, and reflections on the challenges, strengths, and sustainability of the optimal model.

## 251 Data Analysis

252 All interviews were digitally recorded and transcribed in English. All data were anonymized and  
253 stored securely in a password-protected database. Debriefing sessions with data collectors  
254 focused on key findings, identifying saturation themes, and refining inquiry lines. Data were  
255 analyzed thematically using a framework approach which involved the research team reading

256 and re-reading the transcribed data. A coding framework was developed, transcripts were coded  
257 using NVivo software, with emerging themes discussed and coding refined. The coded  
258 transcripts were charted and summarized in narratives for each theme.

## 259 Quantitative Methods

260 During the optimal model pilot test period (January-December 2020), pre-existing and routine  
261 NTDs program data reporting tools were used to collect quantitative data on the utilization of  
262 NTDs services. Existing health systems data was disaggregated by gender and other socio-  
263 demographic factors and utilized to collate information about the impact of the optimal model.  
264 Descriptive comparisons are made between the optimal model intervention period (2020) and the  
265 preceding year (2019) against key SMART performance-based evaluation indicators. These  
266 indicators include; the number of confirmed cases for NTD morbidity (number of confirmed  
267 hydrocele cases, number of confirmed lymphedema cases, percentage of BU cases identified  
268 with categories 1 and 2 lesions, percentage of newly identified Leprosy patients presenting with  
269 grades two disabilities), and number of confirmed CM-NTD cases per disease.

## 270 Data Analysis

271 The research team compiled a list of NTD patients diagnosed over the twelve months recorded in  
272 the health facility registers in an excel spreadsheet. The individual line list of patients was  
273 analyzed using SPSS version 24.

## 274 Results

### 275 A. QUALITATIVE DATA: Optimal model design and delivery

#### 276 Inclusion and collaboration

277 Partnership and collaboration promoted ownership of the process of developing, implementing,  
278 and evaluating the optimal model as stakeholders were involved in all stages. The research and  
279 implementation team all agreed that collaboration with the County teams, community health  
280 workers, and persons affected by NTDs strengthened the design of the optimal model in order to  
281 ensure the language was context-specific and easily understandable at the community level.

282 *'We could bring everyone on board, and everyone gave their collective views*  
283 *on how the intervention phase should be implemented, which gave birth to the*  
284 *development of training materials, development of IEC materials, how to*  
285 *motivate the community and health workers...search for cases.'* (National  
286 *Program Staff, Reflexive Session)*

287

#### 288 Training

289 Key strengths of the training included pictorial guides and accessible language, which aided case  
290 detection at the community level. The inclusion of stigma education helped CHWs improve  
291 disease understandings and dispel misconceptions and stigma associated with NTDs. The  
292 training also increased knowledge of NTD transmission, signs and symptoms, and management  
293 options. The additional training provided has expanded the program's reach as health facilities  
294 that were previously dormant diagnosed NTDs cases in 2020.

295 *"Yes, from the training, I began to realize that if you have a patient with NTDs, people will begin*  
296 *to stigmatize them; oh, this person got elephantiasis I don't want to get some. But from the*  
297 *training, my thoughts began to tell me that that person is important no matter of their conditions,*  
298 *so I should learn to live with the person just as how I have to live as human beings" (CHW,*  
299 *FGD, Bong)*

300

301 *'During the development of the revised training modules, it was really nice to*  
302 *have the community of workers make inputs...the language really changed...*  
303 *the community health workers gave a lot of, consideration of all of the diseases*  
304 *in that 'you have to say it in this way, and that really made the revised modules*  
305 *really rich' (Researcher, Reflexive Session)*

306

307 The length of training was a key challenge. One day for training was too short, and many CHWs  
308 requested longer training sessions or refresher training. Within this pilot, the duration of the  
309 COR-NTD training deviated from the original optimal model, which was initially planned to last  
310 for two days due to budget constraints. As a result, participants reported feeling rushed and  
311 needed more time on the referral forms to practice filling them out correctly.

312 *"The training is for...it's not supposed to be for twenty-four hours. It is*  
313 *supposed to be for forty-eight hours or seventy-two hours because the notes'*  
314 *training is so bulky. You can't use one day to put everything in your brain. You*  
315 *have to take some time to study. Even though they had pretest and posttest. It*

316 *was okay but it has to take some days to get your brain reminding you of it.”*

317 *(CSS, KII)*

### 318 **Trust and Rapport with Communities: The essential role of peer advocates**

319 Peer advocates are affected by NTDs who have received DMDI services and played a unique  
320 role in case-finding activities. Many peer advocates expressed how sharing their experiences has  
321 increased trust and encouraged more people to attend the health facility. Peer advocates reported  
322 relating well to other persons affected to address stigma and encourage them to seek treatment.

323 *'My experience is I talk with the person, and I tell him I too have suffered from these*  
324 *kinds of diseases and the Phebe hospital took care of it easily, and he should not be*  
325 *scared...and I too have gone under such an operation, and nothing happened....The*  
326 *good thing I experienced about the training was to talk with those that got the sickness*  
327 *like me, and they got treated and got well like me because I have been in the hospital*  
328 *myself before.'* (Peer Advocates, FGD, Bong)

### 329 **Gender Balance**

330 Gendered dynamics shaped the case identification process in communities. Men and women felt  
331 more comfortable discussing their conditions and being screened by CHWs of the same gender.  
332 However, the training sensitized case detectors to consider gender preferences by involving  
333 relatives or friends during case detection; many CHWs described that this resulted in affected  
334 persons being more likely to cooperate.

335 *'Though the person will find it very difficult to express their condition to you as*  
336 *an opposite gender because traditionally they do not feel secure.*

337 *'In identifying cases with women, it's kind of a little quite different. If she's married and perhaps*  
338 *she has a problem, if you must be successful in handling that problem, you must firstly engage*  
339 *the husband making sure to let him know what the problem the woman has so he can know what*  
340 *is your role too in the community. So, in my experience on the field, before engaging somebody*  
341 *wife, if she has a problem, I must also let the husband know that I came on the field and this is*  
342 *my job. This is what your wife has and this is what I want to do. He must approve before I can go*  
343 *on with my function.'* (CHWs, FGD, Bong)

## 344 Supervision

345 The supervision cascade was effective, as CHWs and CHAs reported feeling supported and  
346 encouraged due to the feedback received from supervisors. Some forms were not filled out  
347 correctly in the first supervision round, but this improved with the second round. According to  
348 many CHSS, supervisions were conducted daily in different communities where community  
349 awareness meetings were held. Coordination amongst the different levels of health cadres in the  
350 supervision cascade was a key implementation strength. This coordinated approach through the  
351 use of supervision forms at all levels and a detailed supervision plan was not included in the  
352 previous models.

353 *'Those who can come visit us we can really enjoy them because some of the*  
354 *thing them there I can't understand it. So when he comes he can teach me and*  
355 *it can go well with me. For example, I just use to put the date I will not put*  
356 *nothing on it. But my supervisor when she comes she can say no that's not how*  
357 *you can do it. Do it so. When even you do not find the case, you are supposed*  
358 *to do it so. So it can really encourage me. It building me up to do the work*  
359 *now.'* (CHW, FGD, Bong)

360 However, logistical challenges such as limitations in transportation, gasoline, and  
361 communication cards were mentioned by supervisors as limitations to optimal supervision.  
362 Financial constraints in paying for drugs and lack of drugs in facilities were crucial challenges.  
363 However, this was highlighted by NTD Program staff as being a national issue and not reflective  
364 of the optimal model.

365  
366 *‘Our CHSS does not have support like gasoline for his bike and he most often used his own*  
367 *money to purchase gasoline to go on the field to check on us. I think the government needs to*  
368 *find ways to assist him because that’s the only way he too will be able to supervise us.’ (CHW,*  
369 *FGD, Bong)*

## 370 Referral

371 Improvements in referral and reporting as a result of the optimal model were demonstrated  
372 through the referral structure. Tools such as referral forms, counter referral forms, and the NTDs  
373 ledger improved reporting. The referral process entailed CHWs and peer advocates filling in  
374 referral forms in the community and referring the patient to the health facility for confirmation  
375 and treatment. When confirmed positive, the patients were put on treatment, with a counter  
376 referral form filled by the health facility and sent back to the community as feedback to the  
377 CHW who referred the case.

378 *‘But for now, things being better for the past year. Because cases that I have*  
379 *being getting in recent time, as soon I call the county focal person if that case*  
380 *specimen supposed to be collected I get in touch. He will try to connect and try*  
381 *to collect the specimen. As soon that case is confirmed, the drugs will*

382 *immediately come. As we even speak I get patient in the facility the facility that*  
383 *are on their treatment and they are improving.' (CHSS, KII, Bong)*

384 However, some CHSS and CHWs described challenges in the referral of patients, which  
385 included limitations of forms, financial constraints, and lack of drugs at the health facilities.

## 386 **Remuneration**

387 The performance-based incentive package was motivating, as mentioned by many CHWs  
388 although some CHWs experienced delays in receiving the USD 5 and incentive packages.  
389 Certificates based on the number of cases referred were also considered a source of  
390 encouragement. Other sources of motivation include pride in their roles, success stories of  
391 patients receiving treatment, and recognition in the community.

392 *'Anytime we find case for me I brought cases here I got bag. I got five dollars. So those are*  
393 *motivating us in order for us to move further. I was motivated really. Because if that after the*  
394 *training, my first case that I found I received bag, five dollars with flashlight.' (CHW, FGD,*  
395 *Bong)*


396  
397  
398 *'What we think could work better is that because most of the time, this five-dollar issue can't*  
399 *really become a problem in the facility. When the CHV report the cases, the focal person will*  
400 *call you and say the case is confirmed; you will give your feedback to the CHV in the community.*  
401 *They expect their five-dollar at that time but at time it will take long time before the five dollar.*

402 *All those can make them feel reluctant with the work' (CHSS, KII, Bong)*


## 403 B. Quantitative Data: Optimal model delivery

404 Across all diseases, the annual number of cases diagnosed based on clinical manifestation has  
405 increased compared to the previous year (Fig 5). This was particularly noticeable for hydrocele  
406 and suspected Buruli ulcer cases, with more than twice the number of cases reported in 2020  
407 compared to 2019 (Fig 6). Cases reported in 2020 are from across all health facilities within  
408 Bong County. This is the first year that geographic coverage appears to have reached all health  
409 facilities since the original rollout of integrated case detection in 2016/17. Cases were detected at  
410 an earlier stage of disease progression. A significant proportion of Buruli ulcer cases have been  
411 identified with grade I and grade II lesions in 2020 compared to 2019 (Fig 7). Grade 1 lesions  
412 would indicate greater capacity and earlier diagnosis in Buruli ulcers. The proportion previously  
413 was almost entirely grade II lesions. However, results related to Buruli ulcer should be  
414 interpreted with caution due to the limitations in PCR confirmation due to the COVID-19  
415 pandemic and reduced supplies to Liberia.

416




417  
418 **Fig 5.** Comparison in cases reported in 2019 (year proceeding the optimal model) and 2020 (year  
419 of optimal model implementation)  
420




421  
422 **Fig 6.** Annual comparison of the number of CM NTDs Cases per disease

423 The Optimal Model also shows an increase in the number of health facilities throughout the  
424 county that reported cases (Fig 8). However, many patients reported some level of movement  
425 limitations at diagnosis due to ulceration.



426  
427 **Fig 7.** Percentage of newly diagnosed BU patients per year with category I, II, or III lesions at  
428 point of detection  
429

430 By 2020, more health facilities reported CM NTDs cases referred by CHVs who have been  
431 trained and supported through the Optimal Model package than in 2019. Although the number of  
432 cases reported in 2019 and 2020 per health facility was deficient, several health facilities  
433 reported CM-NTDs cases for the first time in 2020 due to the Optimal Model (Fig 8).



434

435 **Fig 8.** Comparison of the Annual number of CM-NTDs Cases detected per Health Facility in  
 436 Bong County.

437


438 Impact of gender on equitable access – a higher proportion of male cases than females was  
 439 identified across diseases (Fig 9). Five hundred ninety-five males and 109 females participated in  
 440 the case detection and referral training from the nine health districts in Bong County (Table 2).  
 441 The CHW training included both males and females between the ages of 25-60 from catchment  
 442 communities of all the health facilities in Bong County. However, significantly more men than  
 443 women were in this training, as reflected in the cases detected (Fig 9). A more significant  
 444 proportion of cases identified as males is shared across all disease conditions across 2019 and  
 445 2020, excluding leprosy, where case detection appears equal (Fig 9).

446

447 **Table 2. Composition of participants trained, disaggregated by role and gender**

Role of Trainee	Male	Female	Total
CHWs	595	109	704
CSS	29	28	57
Peer Advocates	35	15	50
<b>Total</b>	<b>659</b>	<b>152</b>	<b>811</b>

448  
449  
450



451  
452 **Fig 9.** Case Management of NTDs cases disaggregated by sex per disease condition in 2020,  
453 Bong County, Liberia  
454

455

## 456 Discussion and Conclusion

457 One of the main objectives of the Optimal Model was to support the earlier identification of  
458 NTDs requiring case management to reduce morbidity and disability. For Buruli ulcer, this was  
459 successful, with a significant proportion of cases being identified with grade I and grade II  
460 lesions in 2020 compared to 2019. This result emphasizes that cases are detected at an earlier  
461 stage of disease progression. The training improved case finding across all diseases as the annual  
462 number of cases diagnosed had increased in 2020 compared to 2019 in Bong County.

463 This study, to our knowledge, is the first to focus on identifying an optimal model for integrated  
464 case detection, referral, and confirmation of NTDs in Liberia. Several studies have been  
465 conducted on the integrated case management of NTDs strategy in Liberia, focusing on patient  
466 perception, the outcome of care, and case detection (22,25,26). There have been findings on the  
467 effectiveness of active case search in increasing the number of cases compared to reliance on  
468 passive surveillance at the health facility for specific diseases (27,28). However, the findings  
469 from this study support improved access to care for affected persons through early case  
470 detection.

471 Integrating different disease conditions within a single training reduced barriers to care for  
472 persons affected by NTDs. The collaboration between different cadres of health workers to  
473 develop a comprehensive training manual supported enhanced and early case detection for NTDs  
474 affected persons with poor health-seeking behavior. Our study found that integrating these  
475 disease conditions into one manual increased CHWs' knowledge to conduct community  
476 awareness to reduce stigmatization of persons affected by NTDs. The observations in this study  
477 are similar to Mitja *et al.*, who found that "the success of an integrated approach will rely on  
478 well-trained health workers and village volunteers being able to correctly identify multiple

479 skin conditions" within one interaction (11). The CHWs benefited from the collaboration across  
480 diseases and amongst the different levels of the health system. The knowledge gained and  
481 confidence in their ability to provide valuable information on NTDs to community members and  
482 identify suspect cases was due to the participation of various stakeholders in the development of  
483 the training tools and the design of the optimal model.

484 Alongside integrated training, ensuring the right people receive this training and are well  
485 motivated is essential. For example, our study found that Buruli ulcer and hydrocele cases  
486 reported in 2020 were twice reported in 2019. CHWs reported that this was significantly due  
487 to the integrated training and the frequent supervision and incentives, which motivated them  
488 to continue active case search and community awareness. The on-spot mentoring and follow-  
489 up during supervision by CHSS provided an opportunity for CHWs to gain more knowledge  
490 and confidence to identify cases. This result supports the findings of studies conducted in the  
491 Republic of Benin, Cameroon, and Nigeria who also found an increase in the number of cases  
492 reported with early signs of the disease through referral by trained community health workers  
493 (29–32). In addition, our study prioritized the inclusion of peer advocates as case detectors to  
494 build trust with community members and motivate affected persons to visit the health facility  
495 for proper diagnosis. Youtsu also found that integrated surveillance requires the training of local  
496 health workers (16). Peer advocates' personal experience with the disease conditions and their  
497 understanding of the local community helps reduce stigma and encourage positive healthcare-  
498 seeking behaviors. Program decision-makers need to carefully select who delivers the activities  
499 to impact program outcomes.

500 Gendered access to patients to identify and refer was a key challenge emphasized by  
501 qualitative and quantitative data. The gender disparity in the number of cases diagnosed was

502 linked to the sex of those who received the training. CHWs reported seeking permission from  
503 spouses or family members to screen suspected cases of the opposite sex due to cultural  
504 norms and fear of stigmatization. This finding is consistent with findings from India that  
505 women delayed hospital visits awaiting permission from their husbands or guardians (33). In  
506 order to address gender inequities in CM-NTDs interventions, programs must invest in  
507 frontline health workers to improve communication techniques to reduce shame and stigma as  
508 barriers to healthcare seeking (34). Despite the commitment to gender equity within the  
509 research, it was impossible to achieve gender parity in training as the community health  
510 workforce in Liberia is predominantly male. National programs must proactively advocate for  
511 gender balance in the recruitment of CHWs to ensure that those conducting activities truly  
512 represent the gender makeup of such a community. Additionally, integrating gender in all stages  
513 of planning, tool development, testing, implementation, and evaluation will improve the equity  
514 and impact of case finding.

## 515 Limitations

516 The delivery of this study was limited by the impact of COVID -19 due to partial lockdowns and  
517 social distancing restrictions. Liberia confirmed its first COVID-19 case on March 16, 2020. On  
518 April 8, 2020, the President declared a 21-day renewable national state of emergency for  
519 residents of all fifteen counties until further notice. This meant no inter-county travels between  
520 the capital and other counties and a 3 p.m. to 6 a.m. curfew rather than a total lockdown. During  
521 this time, residents were permitted to leave their homes only to procure food or health items, an  
522 activity limited to one person per household for a maximum of one hour within his or her local  
523 area. This state of emergency was extended two weeks until July 21, 2020. Therefore, case  
524 detection activities were suspended, resulting in the low number of cases reported in the

525 quantitative results. The fluctuation in the number of cases detected between February to  
526 September 2020 corresponds with the increase in Covid-19 cases in Bong County. Case  
527 detection began in February 2020, with high numbers reported after completion of the integrated  
528 training in January; however, there were significantly fewer cases reported with the first  
529 lockdown between March and April. In contrast, NTDs cases reported between August to  
530 September 2020 also decreased when no new COVID-19 was reported in Bong county. This is  
531 due to the increased workload for CHWs as other health programs began to resume activities  
532 after the decrease in COVID-19 cases and the end of lockdown measures.

533  
534 Many CHWs mentioned how COVID-19 impacted their work in case identification and referrals.  
535 People feared attending health facilities for fears of contracting COVID-19. This resulted in  
536 reduced regular community visits, reduced referrals, limited health talks. To mitigate the impact  
537 of COVID-19 on the case detection activities, infection prevention and control measures were  
538 implemented. Hand sanitizers and face masks were provided to CHWs to reduce their fear of  
539 interacting with community members during case detection when possible.

540 Furthermore, many CHWs were recruited and trained to conduct COVID-19 surveillance, which  
541 was done simultaneously with NTDs case detection and awareness activities. COVID-19 also  
542 affected the work of peer advocates. While peer advocates' significant role in case finding was  
543 recognized, the duration of their case-finding activities was far less than that of CHWs. As a  
544 result, the impact of their work cannot be quantitatively measured for case-finding activities.

## 545 **Recommendations for Health System Strengthening**

546 Implementation of the optimal model did come with challenges, which are to be anticipated  
547 when integrating interventions into emerging health systems. Despite the global impetus

548 towards integration, restrictions on donor funding limited the scope of some activities, e.g.,  
549 length of training and supervision., greater flexibility concerning the scope and integration  
550 within the training of multi diseases would enable more effective case finding for NTDs.  
551 Similarly, a significant limitation was the logistical support required to implement a robust  
552 supervision model. The importance of supportive supervision was repeatedly emphasized;  
553 however, the investment in this area was insufficient. Based on these challenges as documented  
554 in the results, it is recommended that the time allocated for training should be increased. For  
555 example, a three-day training was suggested to allow more time to practice the referral and  
556 reporting forms. Supervisors need logistical support to provide on-spot mentoring for CHWs on  
557 the supervision forms. Provision of adequate copies of the referral forms to health facilities is  
558 suggested to improve documentation and reporting of the referral process. The remuneration  
559 system of cash, certificates, and other rewards should be strengthened to avoid delays and  
560 encourage CHWs in their work.

561 The challenges documented in the study also highlight the financial constraints and barriers  
562 faced by NTDs programs in implementing such an integrated model that requires skilled human  
563 resources and logistics for routine activities. Funding allocation will significantly impact the  
564 sustainability and scaling up of the optimal model. Implementing the optimal model is largely  
565 dependent on donor support; components of the optimal model, such as the incentive package,  
566 will be harder to sustain without financial support. With the documented success of integrating  
567 active case finding and collaboration at all health system levels through this study, national  
568 budgets should prioritize funding for integrated approaches to the case management of NTDs.  
569 Low reliance on donor funding will enable the NTDs program to make independent decisions  
570 and lead in the prioritization of critical interventions.

## 571 Investment in integration

572 This study has shown how integrated case detection of NTDs is pivotal to achieving the global  
573 target to leave no one behind(18). The second WHO NTDs roadmap 2021-2030 prioritizes  
574 integrated approaches to achieving new global targets(36). The increase in the number of cases  
575 detected and diagnosed at an early stage of the disease through the optimal model outlined in this  
576 study enables evidence-based advocacy with policymakers. Optimal models for case detection,  
577 referral, and confirmation of suspect NTD cases rely on integrated approaches to training,  
578 supervision, referral, and remuneration that are embedded within existing health systems  
579 infrastructure. Equity and inclusion are also critical, particularly the gender of health workers  
580 and the inclusion of affected persons. The global NTD target SDG 3:3 can only be achieved  
581 through optimal case detection, referral, and confirmation models. Together, these approaches  
582 improve access to health services, thus reducing morbidity associated with NTDs.

## 583 Acknowledgment

584 We would like to extend our gratitude to the following people, without whose support this  
585 project would not have been successful. Many thanks to the Ministry of Health Liberia, the data  
586 collection team comprised of Mohammad Dunbar and Otis Kpadeh, and field drivers for their  
587 commitment and endurance during the data collection process. Many thanks to the Ministry of  
588 Health NTD program monitoring and evaluation team and the UL-PIRE for the approval to  
589 conduct the research, and finally, our most tremendous thanks to the national stakeholders,  
590 clinical staff, CHAs, CHVs, CDDs, patients, and communities for their time to participate in this  
591 study. Finally, we appreciate Paul Saunderson and Stefanie Weiland for their review of earlier  
592 drafts of the manuscript, which contributed immensely to the final draft.

## 593 References

- 594 1. Engels D. Neglected tropical diseases in the Sustainable Development Goals. Vol. 387,  
595 The Lancet. Lancet Publishing Group; 2016. p. 223–4.
- 596 2. Ramchandra Atre S, Girish Rangan S, Prabhakar Shetty V, Gaikwad N, Furdoon Mistry  
597 N. Perceptions, health-seeking behavior and access to diagnosis and treatment initiation  
598 among previously undetected leprosy cases in rural Maharashtra, India. *Leprosy review*.  
599 2011;82(3):222.
- 600 3. Prochazka M, Timothy J, Pullan R, Kollie K, Rogers E, Wright A, et al. “Buruli ulcer and  
601 leprosy, they are intertwined”: Patient experiences of integrated case management of skin  
602 neglected tropical diseases in Liberia. *PLoS Neglected Tropical Diseases*. 2020 February  
603 5;14(2).
- 604 4. Hofstraat K, van Brakel WH. Social stigma towards neglected tropical diseases: a  
605 systematic review. *International Health*. 2016 Mar 3;8(suppl 1):53–70.
- 606 5. Nicholls PG, Chhina N, Bro AK, Barkataki P, Kumar R, Withington SG, et al. Factors  
607 contributing to delay in diagnosis and start of treatment of leprosy: analysis of help-  
608 seeking narratives in northern Bangladesh and in West Bengal, India. *Leprosy Review*.  
609 2005;76(1):35–47.
- 610 6. Weiss MG. Stigma and the social burden of neglected tropical diseases. *PLoS Neglected  
611 Tropical Diseases*. 2008 May 14;2(5):237.
- 612 7. Van GY, Onasanya A, van Engelen J, Oladepo O, Diehl JC. Improving access to  
613 diagnostics for schistosomiasis case management in Oyo state, Nigeria: Barriers and  
614 opportunities. *Diagnostics*. 2020 May 1;10(5):328.
- 615 8. World Health Organization. ACCELERATING WORK TO OVERCOME THE  
616 GLOBAL IMPACT OF NEGLECTED TROPICAL DISEASES: A ROADMAP FOR  
617 IMPLEMENTATION. 2012.
- 618 9. Mieras LF, Anand S, van Brakel WH, Hamilton HC, Martin Kollmann KH, Mackenzie C,  
619 et al. Neglected Tropical Diseases, Cross-Cutting Issues Workshop, 4–6 February 2015,  
620 Utrecht, the Netherlands: meeting report. *International Health*. 2016 March 3;8(suppl  
621 1):7–11.
- 622 10. Mitjà O, Marks M, Bertran L, Kollie K, Argaw D, Fahal AH, et al. Integrated Control and  
623 Management of Neglected Tropical Skin Diseases. *PLOS Neglected Tropical Diseases*.  
624 2017 January 19;11(1).
- 625 11. Ortu G, Williams O. Neglected tropical diseases: exploring long term practical approaches  
626 to achieve sustainable disease elimination and beyond. *Infectious Diseases of Poverty*.  
627 2017 Dec 27;6(1):1–2.
- 628 12. Rosenberg M, Utzinger J, Addiss DG. Preventive Chemotherapy Versus Innovative and  
629 Intensified Disease Management in Neglected Tropical Diseases: A Distinction Whose  
630 Shelf Life Has Expired. *PLOS Neglected Tropical Diseases*. 2016 April 14;10(4).
- 631 13. TRANSFORMING OUR WORLD: THE 2030 AGENDA FOR SUSTAINABLE  
632 DEVELOPMENT. New York, NY; 2015.
- 633 14. World Health Organization. Department of Control of Neglected Tropical Diseases.  
634 Integrating neglected tropical diseases into global health and development : fourth WHO  
635 report on neglected tropical diseases. Geneva; 2017.

- 636 15. Boisson S, Engels D, Gordon BA, Medlicott KO, Neira MP, Montresor A, et al. Water,  
637 sanitation and hygiene for accelerating and sustaining progress on neglected tropical  
638 diseases: a new Global Strategy 2015–20. *International Health*. 2016 March 3;8(suppl 1).
- 639 16. Yotsu R. Integrated Management of Skin NTDs—Lessons Learned from Existing Practice  
640 and Field Research. *Tropical Medicine and Infectious Disease*. 2018 Nov 14;3(4).
- 641 17. Simpson H, Quao B, van der Grinten E, Saunderson P, Ampadu E, Kwakye-Maclean C, et  
642 al. Routine Surveillance Data as a Resource for Planning Integration of NTD Case  
643 Management. *Leprosy Review*. 2018;89(3):178–96.
- 644 18. Fitzpatrick C, Engels D. Leaving no one behind: a neglected tropical disease indicator and  
645 tracers for the Sustainable Development Goals: Box 1. *International Health*. 2016 March  
646 3;8(suppl 1):15–8.
- 647 19. Essential Package of Health Services: Secondary & Tertiary Care: The District, County &  
648 National Health Systems, November 2011. Monrovia, Liberia; 2011.
- 649 20. Republic of Liberia Investment Plan for Building a Resilient Health System in Liberia.  
650 2015.
- 651 21. STRATEGIC PLAN FOR INTEGRATED CASE MANAGEMENT OF NEGLECTED  
652 TROPICAL DISEASES (NTDs). 2016.
- 653 22. Dean L, Tolhurst R, Nallo G, Kollie K, Bettee A, Theobald S. Neglected tropical disease  
654 as a 'biographical disruption': Listening to the narratives of affected persons to develop  
655 integrated people-centered care in Liberia. *PLOS Neglected Tropical Diseases*. 2019  
656 September 6;13(9).
- 657 23. Republic of Liberia 2008 Population and Housing Census. Monrovia, Liberia; 2008.
- 658 24. Collins SE, Clifasefi SL, Stanton J, The LEAP Advisory Board, Straits KJE, Gil-  
659 Kashiwabara E, et al. Community-based participatory research (CBPR): Towards  
660 equitable involvement of community in psychology research. *American Psychologist*.  
661 2018 Oct;73(7):884.
- 662 25. Zawolo G, Kollie K, Bettee A, Siakeh A, Thomson R, Theobald S, et al. Implementing  
663 and integrating NTD programmes in Liberia: Reflections from key stakeholders.  
664 Liverpool, UK; 2019 Jan.
- 665 26. Kasang C, Krishnan N, Menkor N, Puchner K. Impact of General Health Volunteers on  
666 Leprosy Control in Nimba County, Liberia: an experience from the Ganta Leprosy  
667 Rehabilitation Centre over 2013-2015. *Leprosy Review*. 2019;90(2):161–6.
- 668 27. Glennie M, Gardner K, Dowden M, Currie BJ. Active case detection methods for crusted  
669 scabies and leprosy: A systematic review. *PLOS Neglected Tropical Diseases*. 2021 July  
670 23;15(7).
- 671 28. Buyon L, Slaven R, Emerson PM, King J, Debrah O, Aboe A, et al. Achieving the  
672 endgame: Integrated NTD case searches. *PLOS Neglected Tropical Diseases*. 2018  
673 December 20;12(12).
- 674 29. Barogui YT, Sopoh GE, Johnson RC, de Zeeuw J, Dossou AD, Houezo JG, et al.  
675 Contribution of the Community Health Volunteers in the Control of Buruli Ulcer in Bénin.  
676 *PLoS Neglected Tropical Diseases*. 2014 October 2;8(10).
- 677 30. Vouking MZ, Tamo VC, Mbuagbaw L. The impact of community health workers (CHWs)  
678 on Buruli ulcer in sub-Saharan Africa: a systematic review. *Pan African Medical Journal*.  
679 2013;15.

- 680 31. Vouking MZ, Takougang I, Mbam LM, Mbuagbaw L, Tadenfok CN, Tamo CV. The  
681 contribution of community health workers to the control of Buruli ulcer in the Ngoantet  
682 area, Cameroon. *Pan African Medical Journal*. 2013;16.
- 683 32. Ukwaja KN, Alphonsus C, Eze CC, Lehman L, Ekeke N, Nwafor CC, et al. Investigating  
684 barriers and challenges to the integrated management of neglected tropical skin diseases in  
685 an endemic setting in Nigeria. *PLOS Neglected Tropical Diseases*. 2020 April 30;14(4).
- 686 33. Succhanda John A, Samuel P, Rao S, Das S. Assessment of needs and quality care issues  
687 of women with leprosy. 2010 Mar;81(1):34–40.
- 688 34. Ozano K, Dean L, Yoshimura M, MacPherson E, Linou N, Otmani del Barrio M, et al. A  
689 call to action for universal health coverage: Why we need to address gender inequities in  
690 the neglected tropical diseases community. *PLOS Neglected Tropical Diseases*. 2020  
691 March 12;14(3).
- 692 35. Casulli A. New global targets for NTDs in the WHO roadmap 2021–2030. *Plos NTDs*.  
693 2021;  
694

## 695 Supporting Information

696 S1 File. Detailed Optimal Model for case detection, referral, and confirmation

697

698 S2 File. Ethical approval

699

700 S3 File. An example of consent form administered

701

702 S1 Appendix. Supervision Tools

703

704 S2 Appendix. Certificate of Training

705

706 S3 Appendix. Training Manual

707

708 S4 Appendix. Referral forms

709

# Three models aimed at improving early case detection in 2017

**(1)** The Community Health Assistant Model (CHA Model) recruited semi-literate community volunteers to identify and refer people with clinical manifestations in the skin typical to NTDs as part of their routine work. CHAs operate in communities outside a 5 kilometers radius from the nearest health facility and receive monthly incentives of \$70.00USD.

**(2)** The Community Health Volunteer Motivation Model (CHV Model) encouraged community volunteers to identify and report suspected DMDI cases within their communities. The CHVs received USD 5 per confirmed positive case identified.

**(3)** The Integrated Mass Drug Administration and DMDI Active Case Search Model (MDA-CM model) trained Community Directed Distributors (who usually include CHVs and CHAs) to conduct active case searches for DMDI cases while distributing NTD medications during MDA campaigns.

## **TRAINING**

A cascaded model of training was developed for CHWs, peer advocates (people affected by NTDs), and health workers as supervisors. An integrated training manual and visual aids on disease signs and symptoms were collaboratively developed to include context specific language. CHWs participated in one-day refresher training to identify, refer and report suspected NTD cases using the visual aids. The CHVs who participated in MDA training used the integrated manual. A one-day peer advocate training focused on peer support, case detection, and referral. Certificates were provided to all trainees on completion of training.

## **SUPERVISION**

A comprehensive supervision plan was developed that included tools, personnel, and responsibilities at all levels (national, county, district, Health Facility, community). A cascaded supervision structure was created to monitor and support the skills transferred during the training. This supervision focused on ensuring effective implementation of the training at all levels. The national NTD program provided a one-day supervision training for Community Health Services Supervisors (CHSS), District Health Officers (DHO), and NTDs Focal Persons.

## **REFERRAL**

Tools were developed to document persons making the referral and patients referred. CHWs received a USD 5 performance-based incentive per confirmed case identified. Furthermore, a communication structure (feedback/referral form) was implemented for the CHWs to inform the HW receiving the referred patient at the facility. In return, the CHW received feedback about the outcome (diagnosis) of the patient referred. This structure promoted the follow-up of referred patients by CHWs to ensure they visit the health facility and receive proper diagnosis and care.

## **LOGISTICS**

All trained case detectors received a logistics toolkit. The toolkit included a flashlight, batteries, T. shirt, and backpacks.


## **GENDER BALANCE**

A balance in gender was considered when selecting CHWs and peer advocates to promote equity and increase patient accessibility and acceptability as many women and men preferred to see CHWs of the same gender. the burden of shame incurred by both women and men during case identification and referral. (Was there gender balance?)




Figure


# Identifying the optimal delivery model for the identification, confirmation and referral of NTD cases requiring MMDP services within an integrated health systems approach to NTD care in Liberia




Figure




Figure




Figure



Figure



Figure



Figure