

Alpha globin gene copy number and hypertension risk among Black Americans

A. Parker Ruhl^{1,2}, Neal Jeffries³, Yu Yang⁴, Orlando M. Gutierrez^{5,6}, Paul Muntner⁶, Rakhi P. Naik⁷, Lydia H. Pecker⁷, Bryan T. Mott⁸, Neil A. Zakai^{9,10}, Monika M. Safford¹¹, Leslie A. Lange¹², Cheryl A. Winkler¹³, Marguerite R. Irvin⁶, Mary Cushman^{9,10}, Hans C. Ackerman¹

¹Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland; ²Pulmonary Branch, National Heart, Lung, and Blood Institute, Bethesda, Maryland; ³Office of Biostatistics Research, National Heart, Lung, and Blood Institute, Bethesda, Maryland; ⁴Division of Blood Diseases and Resources, National Heart, Lung, and Blood Institute, Rockville, Maryland; ⁵Department of Medicine, University of Alabama at Birmingham, Birmingham, Alabama; ⁶Department of Epidemiology, University of Alabama at Birmingham, Birmingham, Alabama; ⁷Division of Hematology, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland; ⁸University of Alabama at Birmingham School of Medicine, Birmingham, Alabama; ⁹Department of Medicine, Larner College of Medicine at the University of Vermont, Burlington, Vermont; ¹⁰Department of Pathology & Laboratory Medicine, Larner College of Medicine at the University of Vermont, Burlington, Vermont; ¹¹Department of Medicine, Weill Cornell Medicine, New York, New York; ¹²Division of Biomedical Informatics and Personalized Medicine, Department of Medicine, University of Colorado, Denver, CO; Department of Epidemiology, ¹³Basic Research Laboratory, National Cancer Institute, Frederick National Laboratory for Cancer Research, Frederick, Maryland

Short title: Alpha globin and hypertension

Keywords: alpha globin, alpha thalassemia, hypertension, hemoglobin, nitric oxide

Word count: 3,764

Tables: 4

Figures: 1

Corresponding authors:

Dr. A. Parker Ruhl, MD, MHS
Building 10-Clinical Research Center, Rm. B3-4207
10 Center Dr., Bethesda MD, 20892-1684
parker.ruhl@nih.gov
Tel: 240-669-5776
Fax: 301-480-0590

Hans Ackerman, MD, DPhil
12735 Twinbrook Parkway, Room 3E-20
Rockville, Maryland 20852
Hans.ackerman@nih.gov

ABSTRACT (250/250 words)

BACKGROUND: Alpha globin is expressed in the endothelial cells of human resistance arteries where it binds to endothelial nitric oxide synthase and limits release of the vasodilator nitric oxide. Genomic deletion of the alpha globin gene (*HBA*) is common among Black Americans and could lead to increased endothelial nitric oxide signaling and reduced risk of hypertension.

METHODS: Community-dwelling US adults aged 45 years or older were enrolled and examined from 2003 to 2007, followed by telephone every 6 months, and reexamined from 2013 to 2016. At both visits, trained personnel performed standardized, in-home blood pressure measurements and pill bottle review. Prevalent hypertension was defined as systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg or anti-hypertensive medication use. Droplet digital PCR was used to determine *HBA* copy number. The associations of *HBA* copy number with prevalent hypertension, resistant hypertension, and incident hypertension were estimated using multivariable regression.

RESULTS:

Among 9,684 Black participants, 7,439 (77%) had hypertension at baseline and 1,044 of those had treatment-resistant hypertension. 1,000 participants were not hypertensive at baseline and participated in a follow up visit; 517 (52%) developed hypertension over median 9.2 years follow-up. Increased *HBA* copy number was not associated with prevalent hypertension (PR=1.00; 95%CI 0.98,1.02), resistant hypertension (PR=0.95; 95%CI 0.86,1.05), or incident hypertension (RR=0.96; 95%CI 0.86,1.07).

CONCLUSIONS: There were no associations between increased *HBA* copy number and risk of hypertension. These findings suggest that variation in alpha globin gene copy number does not modify the risk of hypertension among Black American adults.

INTRODUCTION

Black Americans have a higher prevalence of hypertension than other racial and ethnic groups such as White, Asian, and Hispanic Americans [1]. However, they and other minority racial and ethnic groups remain underrepresented in population studies of hypertensive disease [2,3]. A better understanding of genetic factors that influence blood pressure among Black Americans is needed to identify new targets for treatment of high blood pressure in this population.

The alpha subunit of hemoglobin is expressed in the endothelium of small arteries where it regulates the release of the vasodilator nitric oxide [4–6]. Specifically, alpha globin binds to endothelial nitric oxide synthase (eNOS) and reacts with nitric oxide to limit its diffusion into vascular smooth muscle. Silencing of the alpha globin gene, or targeting the alpha globin/eNOS complex pharmacologically, dilates isolated arteries and lowers blood pressure in mice [5,7,8]. In humans, alpha globin is expressed by the tandem duplicated genes *HBA1* and *HBA2* [9,10]. A common 3.7 kb deletion in the alpha globin gene cluster is common in people of African or Asian descent likely due to the protection it confers against severe malaria, [11–13] but how this deletion affects vasoregulation or vascular disease susceptibility is only beginning to emerge.

The 3.7 kb alpha globin gene deletion is associated with enhanced dynamic vasodilation in adults of African or Asian ancestry [14,15]; however, the same deletion is not associated with blood pressure in healthy Kenyan adolescents [16]. We recently discovered this deletion to be associated with protection from chronic kidney disease and end-stage kidney disease in the Reasons for Geographic and Racial Differences in Stroke (REGARDS) study cohort, a large population-based cohort study of older Black adults [17]; however, the question of whether *HBA* copy number is associated with hypertension risk or blood pressure in this cohort remains unaddressed. To fill this knowledge gap, we analyzed the association of *HBA* copy number with prevalent hypertension, resistant hypertension, or incident hypertension in approximately 10,000 Black participants from the national longitudinal REGARDS cohort using multivariable models

that included social, demographic, and biomedical risk factors for hypertension [18]. Among those not taking anti-hypertensive medications, we also examined the association of increasing *HBA* copy number with measured blood pressure.


METHODS

Study Design

REGARDS is a longitudinal cohort study designed to determine the reasons for racial disparities in stroke and cognitive decline in Black and White Americans aged ≥ 45 years.[18] REGARDS enrolled 30,239 participants from the 48 continental United States from 2003 to 2007. Exclusion criteria included self-reported race other than Black or White, residence in or on the waiting list for a nursing home, active cancer within the past year, or inability to communicate in English. A total of 12,514 (41%) REGARDS participants were Black, and 56% were from states considered to be the stroke belt where stroke incidence in the United States is highest. All self-reported Black participants consenting to genetic research were included in this study (Figure 1). All participants provided oral and written informed consent. The REGARDS study was approved by the Institutional Review Boards of the participating centers. This current study was determined to not be human subjects research by the NIH Office of Human Subjects Research Protection (OHSRP #13416) before this study began. This study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline. The analytic plan was prespecified and approved by the REGARDS Cohort Study Executive Committee.

Figure 1. Study flow diagram

REGARDS= REasons for Geographic and Racial Differences in Stroke Study; *HBB*= beta globin gene; *HBA*= alpha globin gene.



Genotyping

We determined *HBA* copy number using droplet-digital polymerase chain reaction (ddPCR) analysis of genomic DNA as previously described [17]; the method is briefly described here and further details can be found in the online supplement. The assay targeted a sequence within the common 3.7 kb insertion/deletion polymorphism to provide a quantitative measure of alpha globin gene copy number per genome. Two-dimensional clusters of positive and negative droplets for the target and reference gene (*EIF2C1*) were manually gated using Quantasoft (Bio-Rad) per the manufacturer's protocols. Droplet counts, copy number variant (CNV) values, and

95% CIs for CNV were extracted, visualized, and genotype was assigned using custom scripts in the R computing environment without user intervention. Copy number ranged from 2 to 6 gene copies, with 2 (-a/-a) and 3 (-a/aa) copies representing homozygosity and heterozygosity, respectively, for the 3.7 kb deletion, 4 (aa/aa) copies representing homozygosity for the reference allele, and 5 (aaa/aa) and 6 (aaa/aaa) copies representing heterozygosity and homozygosity for a known 3.7 kb insertion [19].

Data collection

Baseline REGARDS study data were collected via standardized telephone interview, self-administered questionnaire, and an in-home visit. Trained personnel administered computer-assisted telephone interviews to collect study participant age, sex, region of residence, insurance status, education level, income, self-reports of prior physician-diagnosed comorbid conditions, and verbal informed consent. Trained personnel performed in-home examinations during which height, weight, and blood pressure were recorded, medication reviewed (with pill bottle review), electrocardiograms (EKG), and collection of blood and spot urine samples were performed, and written informed consent was obtained. These measurements were repeated in subjects who were alive and consented during a second in-home visit conducted a median (25th, 75th percentile) of 9.5 (8.7, 9.9) years after their baseline study visit.

Blood pressure measurement

Systolic blood pressure and diastolic blood pressure were measured by trained personnel using a standardized protocol recommended by the Joint National Committee on Prevention, Detection, and Treatment of High Blood Pressure (JNC 7) [20] with an aneroid sphygmomanometer (American Diagnostic Corporation, Hauppauge, NY) [18]. Participants were in a fasting state and seated for five minutes prior to blood pressure measurements. Following a standardized protocol with an appropriate-sized cuff, an aneroid

sphygmomanometer was used to measure systolic and diastolic blood pressure twice, with a 30-second rest period between the two measurements. The average of both measurements was used for all analyses.

Outcome measures

Prevalent hypertension was defined as one or more of the following: 1) systolic blood pressure \geq 140 mmHg or diastolic \geq 90 mmHg; 2) self-reported current antihypertensive medication; or 3) two or more antihypertensive agents on in-home pill bottle review. Treatment-resistant hypertension was defined as taking medications from 4 or more antihypertensive classes or systolic blood pressure \geq 140 mmHg or diastolic blood pressure \geq 90 mmHg while taking medications from \geq 3 antihypertensive classes. The number of classes of antihypertensive medication being taken at baseline was determined by pill bottle review during the baseline in-home study visit. Pulse pressure was defined as systolic blood pressure minus diastolic blood pressure. Incident hypertension was defined as systolic blood pressure \geq 140 mmHg or diastolic blood pressure \geq 90 mmHg or antihypertensive medication use at the follow-up home visit among those who did not have hypertension at baseline.

Exposure variable

HBA copy number was evaluated as a numerical variable with values of 2, 3, 4, 5, or 6 as determined by ddPCR analysis of genomic DNA and modeled per increase of *HBA* copy number by one copy.

Covariates

Age, sex, race, health insurance (yes or no), highest education level obtained (less than high school, high school, some college, college or more), annual income (\leq \$20K, \$20-34K, \$35-74K, \geq \$75K), and smoking status (categorized by never, past, or current smoker) were self-reported.

Only those participants reporting Black race were included in the current analysis. Body mass index (BMI) was calculated from measured height and weight. Region was defined as three geographic areas: stroke belt buckle (coastal plains of North Carolina, South Carolina, and Georgia), stroke belt (the rest of North Carolina, South Carolina, and Georgia and the entire states of Tennessee, Alabama, Mississippi, Arkansas, and Louisiana), and stroke nonbelt (the remaining contiguous US) [21]. Chronic kidney disease was defined by the 2009 CKD-Epi equation and estimated glomerular filtration rate less than 60mL/min/1.73m², urine albumin to creatinine ratio ≥ 30 mg/g measured on urine collected during the baseline in-home examination, or end-stage kidney disease. Fasting glucose levels ≥ 126 mg/dL, random glucose ≥ 200 mg/dL, or self-reported use of glucose-lowering medication was used to define diabetes mellitus. Total cholesterol was measured from blood collected during the in-home examination. C-reactive protein (CRP), hemoglobin, mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), and red-cell distribution width-coefficient of variation (RDW-CV) values were measured or calculated from blood collected during the in-home examination [22]. CRP was natural log-transformed due to right skewness. The first 10 principal components of ancestry were calculated from Infinium Expanded Multi-Ethnic Genotyping Array data on 7,792 participants.

Descriptive analysis of patient characteristics by HBA copy number

For continuous measures, median and 25th and 75th percentiles were reported by *HBA* copy number. Covariates and outcome measures were examined for outliers.

Primary outcome

The primary outcome measure was prevalent hypertension. The association of *HBA* copy number with prevalent hypertension was evaluated first in an unadjusted regression model. A modified Poisson regression model with robust variance estimation for analyzing binary

dependent variables was employed [23]. This Poisson approach gives the estimated risk ratio (RR) of hypertension for each covariate. The RR is regarded as a more clinically meaningful and transparent assessment of risk than the odds ratio in cohort studies [23]. The use of Poisson regression with robust variance estimation allows for estimating the RR directly and is reported here as the prevalence ratio (PR) for prevalent hypertension, rather than the odds ratio that arises from logistic regression. The covariates age, sex, BMI, region, insurance status, education level, income, hemoglobin, chronic kidney disease, diabetes mellitus, total cholesterol, and smoking status were included in all adjusted models. Additional models were estimated in pre-specified sensitivity analyses that included principal components of ancestry and CRP added separately to the base model and a stricter definition of resistant hypertension that required at least one antihypertensive medication to be a diuretic. Several post-hoc sensitivity analyses were performed. We removed chronic kidney disease and hemoglobin each separately from the base model. In addition, we evaluated the outcome of prevalent hypertension using the thresholds of systolic blood pressure ≥ 130 or diastolic blood pressure ≥ 80 mm Hg per the 2017 ACC/AHA recommendation [24].

Secondary outcomes

The same modeling steps were used to analyze the following dichotomous outcomes: 1) the presence/absence of resistant hypertension at baseline reported as a PR, and 2) the presence/absence of incident hypertension (among those who had a second visit and were not hypertensive at baseline) reported as a RR (see Figure 1 illustrating participant follow up to the second visit). A similar set of models were estimated for the following continuous outcomes: systolic blood pressure, diastolic blood pressure, and pulse pressure. Multivariable linear regression was used to model these continuous outcomes. For these outcomes the *HBA* regression coefficients represent the estimated difference in the outcome measure associated with each additional *HBA* copy. The number of baseline anti-hypertensive medications in use

was analyzed using a zero-inflated Poisson model [25,26]. This approach models the number of medications as a mixture of two latent, unobserved subpopulations: a ‘susceptible’ group that may have zero or more medications that is described by a Poisson distribution and a ‘non-susceptible’ group that has exactly zero medications. The non-susceptible population is included as part of the model because a Poisson model by itself has too few parameters to capture the relatively large number of individuals with zero medications (3175 of 9684). For this outcome the exponentiated *HBA* allele regression coefficient represents the multiplicative effect of an additional *HBA* allele on the expected number of medications among the susceptible subpopulation. Other exponentiated coefficients were similarly interpreted.

Missing data considerations

Missing data for the primary outcome, secondary outcomes, and explanatory variables were typically rare (< 0.5%) with some exceptions, e.g., hemoglobin values (Table 1). Multiple imputation methods were employed in the multivariable models and methods as described in the online supplement [27,28].

Data availability

The data underlying the findings include potentially identifying participant information and cannot be made publicly available because of ethical/legal restrictions. However, data including statistical code from this article are available to researchers who meet the criteria for access to confidential data. Data can be obtained upon request through the University of Alabama at Birmingham at regardsadmin@uab.edu. Additional information on the REGARDS study is available at www.regardsstudy.org.

RESULTS

HBA copy number variation and prevalent hypertension

HBA copy number was variable among the 9,684 Black study participants. 6,507 (67%) had four *HBA* copies, 2,685 (28%) had three copies, and 387 (4%) had two copies (Table 1). In addition, 105 (1%) participants had five *HBA* copies and three participants had six *HBA* copies, consistent with heterozygosity and homozygosity for the known *HBA* gene triplication.[19] Among the 9,684 participants, 7,439 (77%) had hypertension at baseline, 1,044 (11%) had resistant hypertension, and 4,543 (47%) were taking two or more antihypertensive medications (Table 1). In multivariable regression analyses, *HBA* copy number was not associated with prevalent hypertension, (PR = 1.00; 95%CI 0.98, 1.02), resistant hypertension, or the number of antihypertensive medications being taken (Table 2).

Table 1. Clinical and demographic characteristics according to *HBA* copy number

	<i>HBA</i> copy number					<i>P</i> value[†]
	All Subjects	2	3	4	≥ 5*	
Subjects, N (%)	9,684 (100)	387 (4)	2685 (28)	6507 (67)	105 (1)	NA
Age, years	63 (57, 70)	64 (58, 70)	64 (57, 71)	63 (57, 70)	67 (61, 73)	0.004
Female sex, N (%)	5952 (61)	243 (63)	1638 (61)	4013 (62)	58 (55)	0.50
Body mass index, kg/m²	29.7 (26.1, 34.4)	30.3 (26.9, 34.9)	29.8 (26.1, 34.4)	29.7 (26.1, 34.4)	30.6 (27.5, 34.9)	0.19
Smoking status, N (%)						0.58
Never	4368 (45)	167 (43)	1225 (46)	2931 (45)	45 (43)	
Past	3588 (37)	156 (41)	986 (37)	2401 (37)	45 (43)	
Current	1683 (17)	61 (16)	460 (17)	1148 (18)	14 (13)	
Region, N (%)						0.06
Non-Belt	4816 (50)	169 (44)	1367 (51)	3220 (49)	60 (57)	
Belt	3197 (33)	138 (36)	850 (32)	2181 (34)	28 (27)	

Buckle	1671 (17)	80 (21)	468 (17)	1106 (17)	17 (16)	
Medically insured, N (%)	8716 (90)	350 (91)	2428 (91)	5842 (90)	99 (91)	0.61
Education level, N (%)						0.18
Less than high school	1896 (20)	70 (18)	532 (20)	1271 (20)	23 (21)	
High school graduate	2677 (28)	111 (29)	774 (29)	1762 (27)	30 (28)	
Some college	2572 (27)	99 (27)	738 (28)	1707 (26)	28 (28)	
College graduate or more	2527 (26)	106 (27)	639 (24)	758 (27)	24 (23)	
Income, N (%)						0.15
≤ \$20K	2600 (31)	105 (30)	721 (31)	1742 (30)	32 (33)	
\$20K - \$34K	2513 (30)	107 (31)	711 (30)	1681 (30)	32 (33)	
\$35K - \$74K	2489 (29)	103 (30)	708 (30)	1651 (29)	27 (28)	
≥ \$75K	894 (10)	32 (9)	210 (9)	670 (11)	6 (6)	
Chronic Kidney disease, N (%)	2498 (27)	83 (22)	688 (27)	1692 (27)	35 (35)	0.05
Diabetes mellitus, N (%)	2818 (29)	103 (27)	808 (30)	1878 (29)	29 (28)	0.47
Total Cholesterol, mg/dL	190 (165, 217)	188 (162, 215)	189 (164, 217)	191 (165, 218)	193 (172, 217)	0.15
C-reactive protein, mg/L units	2.9 (1.2, 6.5)	3.3 (1.3, 7.5)	2.8 (1.1, 6.3)	2.9 (1.2, 6.5)	2.9 (1.1, 6.0)	0.12
Hemoglobin, g/dL	13.1 (12.2, 14.0)	12.4 (11.6, 13.2)	12.9 (12.0, 13.8)	13.2 (12.4, 14.1)	13.0 (12.1, 14.1)	< 0.0001
MCV, fL	88 (84, 92)	74 (72, 77)	85 (82, 87)	90 (87, 93)	89 (87, 92)	< 0.0001
MCH, pg	29.6 (27.9, 30.9)	23.8 (22.9, 24.8)	27.9 (26.9, 29.0)	30.3 (29.1, 31.4)	29.9 (29.1, 31.1)	< 0.0001
MCHC, g/dL	33.4 (32.9, 33.9)	32.1 (31.6, 32.6)	33.0 (32.6, 33.5)	33.6 (33.2, 34.1)	33.8 (33.2, 34.0)	< 0.0001
RDW-CV, %	14.0 (13.3, 14.8)	15.0 (14.4, 15.9)	14.2 (13.5, 15.1)	13.8 (13.2, 14.6)	13.6 (13.2, 14.2)	< 0.0001

Prevalent hypertension[‡], N (%)	7439 (77)	304 (79)	2070 (77)	4978 (77)	87 (83)	0.20
Resistant hypertension, N (%)	1044 (11)	50 (13)	299 (11)	679 (10)	16 (15)	0.16
Anti-hypertensive medication, N (%)						0.20
None	3175 (33)	134 (35)	877 (33)	2135 (32)	29 (28)	
One	1966 (20)	67 (17)	525 (20)	1344 (21)	30 (29)	
Two or more	4543 (47)	186 (48)	1283 (48)	3028 (45)	46 (44)	
Systolic blood pressure, mm Hg	130 (120, 140)	130 (120, 140)	130 (120, 141)	130 (120, 140)	133 (120, 144)	0.42
Diastolic blood pressure, mm Hg	79 (72, 84)	79 (71, 84)	79 (72, 84)	79 (71, 84)	80 (71, 86)	0.82
Pulse pressure, mm Hg	51 (43, 60)	51 (43, 60)	51 (43, 60)	51 (43, 60)	53 (45, 64)	0.30

HBA= alpha globin gene; P= p value; N= number; K= thousand; RBC= red blood cell; MCV= mean corpuscular volume; MCH= mean corpuscular hemoglobin; MCHC= mean corpuscular hemoglobin concentration; RDW-CV= red cell distribution width-coefficient of variation; No.= number. Values are median (25th, 75th percentile) except where otherwise indicated.

*105 subjects had 5 *HBA* gene copies and 3 subjects had 6 *HBA* copies; [†]P values for tests of differences by alpha globin copy number using chi-square test for categorical variables and Kruskal-Wallis test for continuous variables; [‡]261 participants met the definition of prevalent hypertension solely based on having two or more anti-hypertensive medications on pill bottle review.

Missing data are as follows: medically insured (n=12, <1%); education (n=12, <1%); income (n=1,170 refused, 12%); kidney disease (n=389, 4%); diabetes mellitus (n=89, <1%); total cholesterol (n=57, <1%); body mass index (n=55, <1%); smoking status (n=45, <1%); c-

reactive protein (n=151, 1.6%); hemoglobin (n=3,089, 32%); MCV (n=3,095, 32%); MCH (n=3,089, 32%); MCHC (n=3,089, 32%); RDW-CV (n=3,102, 32%). All other variables in Table 1 had no missing values.

Table 2. Association of *HBA* copy number with prevalent hypertension, resistant hypertension, or anti-hypertensive use—fully adjusted analyses

	Prevalent hypertension*		Resistant hypertension†		Number of anti-hypertensive medications‡	
	PR	CI	PR	CI	$e^{\hat{\beta}}$	CI
<i>HBA</i> copy number	1.00	(0.98, 1.02)	0.95	(0.86, 1.05)	0.98	(0.95, 1.01)
Age, per year	1.01	(1.01, 1.01)	1.02	(1.01, 1.03)	1.01	(1.00, 1.01)
Sex						
Female (ref) §	1.00	(0.98, 1.03)	1.37	(1.18, 1.58)	1.10	(1.05, 1.15)
Male						
Body mass index 	1.10	(1.09, 1.11)	1.33	(1.26, 1.40)	1.09	(1.07, 1.12)
Region						
Non-Belt (ref)						
Belt	1.05	(1.02, 1.07)	1.03	(0.91, 1.17)	1.03	(0.99, 1.08)
Buckle	1.02	(0.99, 1.05)	1.05	(0.90, 1.22)	1.03	(0.98, 1.09)
Medically insured						
No (ref)						
Yes	1.07	(1.02, 1.12)	1.12	(0.90, 1.40)	1.06	(0.98, 1.14)
Education level						
< HS Grad (ref)						
HS Grad	1.00	(0.97, 1.02)	1.08	(0.92, 1.25)	1.02	(0.97, 1.08)
Some College	0.98	(0.95, 1.01)	0.81	(0.68, 0.96)	1.01	(0.95, 1.07)
≥ College Grad	0.95	(0.92, 0.98)	0.90	(0.75, 1.08)	0.99	(0.93, 1.05)
Income						
< \$20K (ref)						
\$20K - \$34K	0.99	(0.96, 1.01)	0.95	(0.82, 1.10)	0.98	(0.93, 1.03)
\$35K - \$74K	0.96	(0.93, 0.99)	0.95	(0.80, 1.14)	0.99	(0.93, 1.05)
≥ \$75K	0.94	(0.90, 1.00)	0.77	(0.58, 1.02)	0.93	(0.85, 1.02)
Hemoglobin, per 1 g/dL	0.99	(0.98, 1.00)	0.96	(0.91, 1.01)	0.98	(0.96, 0.99)
Chronic Kidney disease						
No (ref)						
Yes	1.14	(1.10, 1.16)	1.97	(1.74, 2.23)	1.20	(1.15, 1.25)

Diabetes mellitus							
No (ref)							
Yes	1.16	(1.14, 1.18)	1.64	(1.45, 1.85)	1.18	(1.13, 1.23)	
Total Cholesterol							
	0.97	(0.96, 0.98)	0.84	(0.78, 0.89)	0.93	(0.91, 0.95)	
Smoking status							
Never (ref)							
Past	1.04	(1.02, 1.06)	1.08	(0.95, 1.22)	1.04	(1.00, 1.09)	
Present	1.04	(1.01, 1.08)	0.93	(0.77, 1.12)	0.98	(0.92, 1.04)	

HBA= alpha globin gene; PR= estimated prevalence ratio; CI= confidence interval; $e^{\hat{\beta}}$ = zero-inflated Poisson regression; SE= standard error; K= thousand; HS= high school.

*Prevalent hypertension was defined as having one or more of the following: 1) systolic blood pressure \geq 140 mmHg or diastolic blood pressure \geq 90 mmHg; 2) self-reported use of medication to control blood pressure; or 3) two or more antihypertensive medications found on in-home pill bottle review. [†]Resistant hypertension was defined as taking medications from 4 antihypertensive medication classes or having systolic blood pressure \geq 140 mmHg or diastolic blood pressure \geq 90 mmHg while taking 3 or more antihypertensive medication classes. [‡]The number of antihypertensives being taken at baseline was determined by pill bottle review during the in-home baseline study visit. [§](ref) indicates reference category used for categorized explanatory variables. ^{||}Body mass index and total cholesterol were scaled by standard deviation. Multiple imputations were performed for missing data.

HBA copy number variation and incident hypertension

Among the 1,000 participants who were not hypertensive at baseline and who participated in the second visit, 517 (52%) developed hypertension over a median (25th, 75th percentile) of 9.2 (8.6, 9.9) years of follow-up. In multivariable regression analysis, *HBA* copy number was not associated with the risk of incident hypertension (RR = 0.96; 95%CI 0.86, 1.07; Table 3).

Table 3. Association of *HBA* copy number with incident hypertension – fully adjusted analysis

	Incident Hypertension*	
	(n=1,000) Modified Poisson	
	RR	CI
<i>HBA</i> Copy Number	0.96	(0.86, 1.07)
Age, per year	1.01	(1.00, 1.02)
Sex		
Female (ref) †		
Male	0.95	(0.82, 1.11)
Body mass index[‡]	1.14	(1.08, 1.20)
Region		
Non-Belt (ref)		
Belt	1.12	(0.98, 1.27)
Buckle	0.99	(0.84, 1.16)
Medically insured		
No (ref)		
Yes	0.97	(0.80, 1.17)
Education level		
< HS Grad (ref)		
HS Grad	1.00	(0.82, 1.23)
Some College	0.95	(0.77, 1.17)
≥ College Grad	0.83	(0.67, 1.03)
Income		
< \$20K (ref)		
\$20K - \$34K	1.01	(0.84, 1.22)
\$35K - \$74K	1.04	(0.87, 1.25)
≥ \$75K	0.92	(0.72, 1.18)
Hemoglobin, per 1 g/dL	1.00	(0.94, 1.06)
Chronic Kidney disease		
No (ref)		
Yes	1.10	(0.91, 1.32)

Diabetes mellitus		
No (ref)		
Yes	1.30	(1.12, 1.51)
Total Cholesterol[‡]		
	0.94	(0.88, 1.01)
Smoking status		
Never (ref)		
Past	0.98	(0.85, 1.12)
Present	1.15	(0.97, 1.36)

HBA= alpha globin gene; RR= risk ratio; CI= confidence interval; K= thousand; HS= high school.

*Incident hypertension was defined as systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg or reported medication use to control blood pressure during the REGARDS study cohort follow-up visit, among those who were not hypertensive at baseline (systolic blood pressure < 140 mmHg or diastolic blood pressure < 90 mmHg or no medication use to control blood pressure). The incident hypertension analysis was performed on 1,000 participants who were not hypertensive at baseline and had second visit data available. The median (25th, 75th percentile) time elapsed between the baseline and second visit was (9.2 [8.6, 9.9]) years. P values calculated by modified Poisson regression model for dichotomous outcome measures. [†](ref) indicates reference category used for categorized explanatory variables. [‡]Body mass index and total cholesterol were scaled by standard deviation. Multiple imputations were performed for missing data.

HBA copy number variation and measured blood pressure

The association between *HBA* copy number and measured blood pressure was assessed in 3,175 participants who were not taking antihypertensive medications at baseline. Each additional *HBA* copy was associated with lower diastolic blood pressure (-0.66 mmHg; 95%CI -1.25, -0.07; Table 4) but was not significantly associated with lower systolic blood pressure (-0.88 mmHg; 95%CI -1.89, 0.13; Table 4). There was no association between *HBA* copy number and pulse pressure (-0.22 mmHg; 95%CI -0.95, 0.52) as the direction of the effect of *HBA* copy number on systolic blood pressure and diastolic blood pressure was the same.

Table 4. Association of *HBA* copy number with measured blood pressure in subjects not taking anti-hypertensive medications—fully adjusted analyses

	Systolic blood pressure*		Diastolic blood pressure*		Pulse pressure* [†]	
	(n=3,175)		(n=3,175)		(n=3,175)	
	Multivariable linear		Multivariable linear		Multivariable linear	
	$\hat{\beta}$	CI	$\hat{\beta}$	CI	$\hat{\beta}$	CI
<i>HBA</i> copy number	-0.88	(-1.89, 0.13)	-0.66	(-1.25, -0.07)	-0.22	(-0.95, 0.52)
Age, by year	0.27	(0.20, 0.34)	-0.06	(-0.10, -0.02)	0.33	(0.28, 0.39)
Sex						
Female (ref)						
Male	4.53	(3.06, 5.99)	2.18	(1.33, 3.03)	2.34	(1.28, 3.40)
Body mass index[§]	3.27	(2.67, 3.88)	2.23	(1.87, 2.58)	1.04	(0.60, 1.48)
Region						
Non-Belt (ref)						
Belt	0.40	(-0.89, 1.69)	-0.10	(-0.86, 0.65)	0.50	(-0.44, 1.44)
Buckle	-2.24	(-3.84, -0.63)	-0.78	(1.72, 0.15)	-1.45	(-2.62, -0.28)
Medically insured						
No (ref)						
Yes	-1.83	(-3.61, -0.05)	-0.53	(-1.57, 0.51)	-1.30	(-2.61, -0.28)
Education level						
< HS Grad (ref)						

HS Grad	-3.10	(-4.99, -1.21)	-1.42	(-2.53, -0.32)	-1.68	(-3.06, -0.29)
Some College	-3.30	(-5.23, -1.38)	-1.05	(-2.18, 0.08)	-2.25	(-3.66, -0.85)
≥ College Grad	-4.17	(-6.19, -2.15)	-1.56	(-2.74, -0.37)	-2.61	(-4.09, -1.13)
Income						
< \$20K (ref)						
\$20K - \$34K	1.06	(-0.63, 2.76)	0.23	(-0.76, 1.22)	0.83	(-0.42, 2.08)
\$35K - \$74K	-0.18	(-1.96, 1.59)	-0.37	(-1.42, 0.68)	0.18	(-1.13, 1.49)
≥ \$75K	-0.99	(-3.40, 1.42)	-0.65	(-2.04, 0.73)	-0.34	(-2.11, 1.44)
Hemoglobin, per 1 g/dL						
	-0.15	(-0.70, 0.40)	0.62	(0.30, 0.94)	-0.77	(-1.17, -0.38)
Chronic Kidney disease						
No (ref)						
Yes	6.49	(4.78, 8.19)	2.96	(1.95, 3.96)	3.53	(2.30, 4.76)
Diabetes mellitus						
No (ref)						
Yes	0.58	(-1.12, 2.27)	-0.92	(-1.91, 0.08)	1.50	(0.25, 2.74)
Total Cholesterol [§]	1.34	(0.75, 1.92)	0.65	(0.30, 0.99)	0.69	(0.27, 1.12)
Smoking status						
Never (ref)						
Past	0.27	(-1.07, 1.60)	-0.72	(-1.50, 0.06)	0.98	(-0.01, 1.96)
Present	3.06	(1.48, 4.64)	0.52	(-0.41, 1.45)	2.54	(1.39, 3.70)

HBA= alpha globin gene; $\hat{\beta}$ = estimated linear regression coefficient; CI= confidence interval; K= thousand; HS= high school.

*The analyses of measured blood pressured outcomes were performed only on those subjects not currently taking antihypertensive medication (n=3,175). [†]Pulse pressure was defined as systolic blood pressure minus diastolic blood pressure. [‡](ref) indicates reference category used for categorized explanatory variables. [§]Body mass index and total cholesterol were scaled by standard deviation. Multiple imputations were performed for missing data

Sensitivity and interaction analyses

Pre-specified sensitivity analyses were performed to address possible population stratification, to explore the role of inflammation in the relationship between *HBA* copy number and prevalent hypertension, and to evaluate a stricter definition of resistant hypertension. The sensitivity analysis for population stratification was performed by adding the first 10 principal components of ancestry to the prevalent hypertension model; this did not change the association of *HBA* copy number with prevalent hypertension (Table S1). To explore the role of inflammation, c-reactive protein (CRP) was included as a covariate in the multivariable models. Log CRP was associated with increased risk of prevalent hypertension (PR = 1.03; 95% CI 1.02,1.04) but did not change the association of *HBA* copy number with prevalent hypertension (Table S2). We evaluated an alternative definition of resistant hypertension that required at least one of the anti-hypertensive medications to be a diuretic, and there was no change in the association of *HBA* copy number with resistant hypertension (Table S3). We conducted pre-specified tests for statistical interaction between age, sex, or chronic kidney disease with *HBA* copy number on the outcome of prevalent hypertension and found no significant interactions (all p values for interaction terms >0.20, Table S4). Post-hoc sensitivity analyses removing hemoglobin or chronic kidney disease from the model did not change the association of *HBA* copy number with prevalent hypertension (Tables S5 and S6). Defining hypertension with a lower blood pressure threshold also did not change the association of *HBA* copy number with prevalent hypertension (Table S7).

DISCUSSION

In light of the novel role of endothelial alpha globin as a restrictor of nitric oxide signaling in resistance arteries, we hypothesized that individuals who inherit a deletion of the alpha globin gene would be less likely to develop hypertension. To test this hypothesis, we used digital

droplet PCR to genotype a 3.7 kb insertion/deletion polymorphism that quantitatively alters alpha globin gene copy number in a large longitudinal cohort of Black Americans. We found no association between *HBA* copy number and the pre-specified primary outcome of prevalent hypertension. There was also no association between *HBA* copy number and the secondary outcomes of prevalent resistant hypertension or the number of anti-hypertensive medications being taken. *HBA* copy number was not associated with incident hypertension over a decade of observation.

There have been no studies reporting the association between *HBA* copy number and the clinical outcome of hypertension for direct comparison. A related study of Kenyan adolescents found no association between the 3.7 kb gene deletion and 24-hour ambulatory blood pressure [16]. Our study extends that observation into a population of older Black American adults with a range of social, economic, and medical risk factors where we find no association between *HBA* copy number and the clinical outcome of hypertension. One relevant co-morbidity is chronic kidney disease, which was recently found to be associated with *HBA* copy number in a separate study also performed in the REGARDS cohort [29]. Chronic kidney disease was included in the pre-specified model in the current study evaluating the association of *HBA* copy number with hypertension. We performed a pre-specified test of interaction with *HBA* copy number and chronic kidney disease (Table S4) and a post-hoc sensitivity analysis removing chronic kidney disease from the main model (Table S6) and neither demonstrated any impact on the finding that there was no association between *HBA* copy number and hypertension. Similarly, removing hemoglobin from the main model did not alter the negative findings of this study (Table S5).

In contrast to the study in Kenyan adolescents where there was no association between *HBA* and blood pressure, we observed that each additional copy of *HBA* was associated with lower diastolic blood pressure among adults. This observation implies that increased expression of

alpha globin is associated with lower blood pressure, rather than higher blood pressure as hypothesized. Keller et al. recently demonstrated that endothelial alpha globin can produce nitric oxide from the chemical reduction of nitrite under hypoxic conditions [30]. Thus, endothelial alpha globin may function not only as a nitric oxide scavenger, but also as a nitric oxide producer. Since NOS-dependent vascular nitric oxide production declines with age, the relative roles of alpha globin as a nitric oxide scavenger versus a nitric oxide producer may shift with age as well, potentially explaining the unexpected association of higher *HBA* copy number with lower diastolic blood pressure in this study of older Black Americans.

This study has strengths and limitations. The REGARDS Study is one of the largest cohorts of Black Americans, and the participants were well characterized using standardized measurements for blood pressure [20] and surveyed using instruments validated for collecting social and biomedical factors known to impact hypertension [31]. We used a robust and quantitative ddPCR approach to genotype the *HBA* insertion/deletion directly and avoided the potential for mis-assignment that can result from inferring genotype from sequence analysis or SNP genotyping. We focused on the most prevalent structural variant in Black Americans, the 3.7 kb deletion, but did not evaluate rare deletions or regulatory polymorphisms that could affect gene expression [32]. We utilized a pre-specified analysis plan to reduce the chance of identifying spurious associations, however, no statistical corrections were made for multiple testing.

While this study has strengths from an epidemiological perspective, the hypothesis tested relies on several implicit assumptions linking gene expression in the vascular endothelium to the phenotypes of blood pressure and hypertension. First, while *HBA* copy number is quantitatively associated with alpha globin expression in erythrocyte progenitors, the same relationship has not yet been established for endothelial cells in the wall of the resistance artery. In the future,

measuring gene expression directly in human arteries would establish the relationship between *HBA* copy number and gene expression in this important cellular context. Second, arteries may adapt to the loss of endothelial alpha globin by downregulating the expression or activity of nitric oxide synthase or its downstream receptors such as soluble guanylate cyclase. Thus, compensation may obscure the effect of reduced endothelial alpha globin gene expression on blood pressure. Third, while deletion of *HBA* has been associated with enhanced dynamic vasodilation [14,15], blood pressure is regulated by multiple pathways that are integrated across the renal, cardiovascular, and nervous systems. Thus changes in arterial nitric oxide signaling could be overcome by other regulatory systems to maintain a blood pressure set point [33,34]. Moreover, endothelial nitric oxide synthase has a diminishing role in the regulation of vasodilation as people age and specifically in hypertension [35–39]. In the older population studied here, endothelial nitric oxide synthase may no longer be as active and thus would not be subject to regulation by different levels of alpha globin expression. Therefore, the absence of association between *HBA* copy number and the clinical phenotype of hypertension does not exclude a role for alpha globin as a regulator of nitric oxide signaling in human resistance arteries [4,6–8]. Further investigation into the role of alpha globin in the dynamic vasoreactivity of human resistance arteries remains warranted.

PERSPECTIVES

Despite previous evidence that alpha globin regulates nitric oxide signaling in resistance arteries, we found no association between *HBA* copy number and risk of hypertension among older Black American adults.

ACKNOWLEDGEMENTS

The authors thank the investigators, staff, and participants of the REGARDS study for their valuable contributions. A full list of participating REGARDS investigators and institutions can be found at <http://www.regardsstudy.org>.

SOURCES OF FUNDING

This is an ancillary study supported by cooperative agreement U01 NS041588 co-funded by the National Institute of Neurological Disorders and Stroke (NINDS) and the National Institute on Aging (NIA), National Institutes of Health, Department of Health and Human Service. This research was supported in part by the Divisions of Intramural Research, National Institute of Allergy and Infectious Diseases project AI001150 (A.P.R., H.C.A) and National Heart, Lung, and Blood Institute (NHLBI) project HL006196 (A.P.R., Y.Y., H.C.A.). This work was also funded in part by the National Cancer Institute (NCI) Intramural Research Program under contract HHSN26120080001E (C.A.W), and the NHLBI grants K08HL12510 (R.P.N.) and K08HL096841 (N.A.Z.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NINDS, NIA, NIAID, NCI, or NHLBI. The content of this publication does not necessarily reflect the view or policy of the Department of Health and Human Services, nor does mention of trade names, commercial products or organizations imply endorsement by the government. The interpretation and reporting of these data are the responsibility of the author(s) and in no way should be seen as an official policy or interpretation of the U.S. government.

DISCLOSURES

Dr. Gutierrez discloses receiving grant funding and consulting fees from Akebia Therapeutics; grant funding and consulting fees from Amgen; grant funding from GlaxoSmithKline; consulting fees from Reata, AstraZeneca, and Ardelyx; and serving on the Data Monitoring Committee for QED Therapeutics.

REFERENCES

1. Hardy ST, Chen L, Cherrington AL, Moise N, Jaeger BC, Foti K, et al. Racial and Ethnic Differences in Blood Pressure Among US Adults, 1999–2018. *Hypertension*. 2021;78: 1730–1741. doi:10.1161/HYPERTENSIONAHA.121.18086
2. Rocca WA, Yawn BP, St Sauver JL, Grossardt BR, Melton LJ. History of the Rochester Epidemiology Project: half a century of medical records linkage in a US population. *Mayo Clin Proc*. 2012;87: 1202–1213. doi:10.1016/j.mayocp.2012.08.012
3. Kannel WB. The Framingham Study: ITS 50-year legacy and future promise. *J Atheroscler Thromb*. 2000;6: 60–66. doi:10.5551/jat1994.6.60
4. Straub AC, Lohman AW, Billaud M, Johnstone SR, Dwyer ST, Lee MY, et al. Endothelial cell expression of haemoglobin α regulates nitric oxide signalling. *Nature*. 2012;491: 473–477. doi:10.1038/nature11626
5. Brooks SD, Kamenyeva O, Ganesan S, Zeng X, Smith R, Ma D, et al. Hemoglobin Interacts with Endothelial Nitric Oxide Synthase to Regulate Vasodilation in Human Resistance Arteries. medRxiv. 2021; 2021.04.06.21255004. doi:10.1101/2021.04.06.21255004
6. Lechauve C, Butcher JT, Freiwan A, Biwer LA, Keith JM, Good ME, et al. Endothelial cell α -globin and its molecular chaperone α -hemoglobin-stabilizing protein regulate arteriolar contractility. *J Clin Invest*. 2018;128: 5073–5082. doi:10.1172/JCI99933
7. Straub AC, Butcher JT, Billaud M, Mutchler SM, Artamonov MV, Nguyen AT, et al. Hemoglobin α /eNOS Coupling at Myoendothelial Junctions Is Required for Nitric Oxide Scavenging During Vasoconstriction. *Arterioscler Thromb Vasc Biol*. 2014;34: 2594–2600. doi:10.1161/ATVBAHA.114.303974
8. Keller TCS, Butcher JT, Broseghini-Filho GB, Marziano C, DeLalio LJ, Rogers S, et al. Modulating Vascular Hemodynamics With an Alpha Globin Mimetic Peptide (Hb α X). *Hypertens Dallas Tex 1979*. 2016. doi:10.1161/HYPERTENSIONAHA.116.08171
9. Dozy AM, Kan YW, Embury SH, Mentzer WC, Wang WC, Lubin B, et al. alpha-Globin gene organisation in blacks precludes the severe form of alpha-thalassaemia. *Nature*. 1979;280: 605–607.
10. Higgs DR, Vickers MA, Wilkie AO, Pretorius IM, Jarman AP, Weatherall DJ. A review of the molecular genetics of the human alpha-globin gene cluster. *Blood*. 1989;73: 1081–1104.
11. Mockenhaupt FP, Ehrhardt S, Gellert S, Otchwemah RN, Dietz E, Anemana SD, et al. Alpha(+)-thalassemia protects African children from severe malaria. *Blood*. 2004;104: 2003–2006. doi:10.1182/blood-2003-11-4090
12. Wambua S, Mwangi TW, Kortok M, Uyoga SM, Macharia AW, Mwacharo JK, et al. The Effect of α +-Thalassaemia on the Incidence of Malaria and Other Diseases in Children

- Living on the Coast of Kenya. *PLoS Med.* 2006;3: e158.
doi:10.1371/journal.pmed.0030158
13. Ndila CM, Uyoga S, Macharia AW, Nyutu G, Peshu N, Ojal J, et al. Human candidate gene polymorphisms and risk of severe malaria in children in Kilifi, Kenya: a case-control association study. *Lancet Haematol.* 2018;5: e333–e345. doi:10.1016/S2352-3026(18)30107-8
 14. Denton CC, Shah P, Suriyany S, Liu H, Thuptimdang W, Sunwoo J, et al. Loss of alpha-globin genes in human subjects is associated with improved nitric oxide-mediated vascular perfusion. *Am J Hematol.* 2020. doi:10.1002/ajh.26058
 15. Romana M, Reminy K, Moeckesch B, Charlot K, Hardy-Dessources M-D, Doumido L, et al. Loss of alpha globin genes is associated with improved microvascular function in patients with sickle cell anemia. *Am J Hematol.* 2021;96: E165–E168. doi:10.1002/ajh.26126
 16. Etyang AO, Khayeka-Wandabwa C, Kapesa S, Muthumbi E, Odipo E, Wamukoya M, et al. Blood Pressure and Arterial Stiffness in Kenyan Adolescents With $\alpha(+)$ Thalassemia. *J Am Heart Assoc.* 2017;6. doi:10.1161/JAHA.117.005613
 17. Ruhl AP, Jeffries N, Yang Y, Naik RP, Patki A, Pecker LH, et al. Alpha Globin Gene Copy Number Is Associated with Prevalent Chronic Kidney Disease and Incident End-Stage Kidney Disease among Black Americans. *J Am Soc Nephrol.* 2022;33: 213–224. doi:10.1681/ASN.2021050653
 18. Howard VJ, Cushman M, Pulley L, Gomez CR, Go RC, Prineas RJ, et al. The reasons for geographic and racial differences in stroke study: objectives and design. *Neuroepidemiology.* 2005;25.
 19. Goossens M, Dozy AM, Embury SH, Zachariades Z, Hadjiminias MG, Stamatoyannopoulos G, et al. Triplicated alpha-globin loci in humans. *Proc Natl Acad Sci U S A.* 1980;77: 518–521. doi:10.1073/pnas.77.1.518
 20. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA.* 2003;289: 2560–2572. doi:10.1001/jama.289.19.2560
 21. Howard G, Howard VJ. Twenty Years of Progress Toward Understanding the Stroke Belt. *Stroke.* 2020;51: 742–750. doi:10.1161/STROKEAHA.119.024155
 22. Gillett SR, Boyle RH, Zakai NA, McClure LA, Jenny NS, Cushman M. Validating laboratory results in a national observational cohort study without field centers: the Reasons for Geographic and Racial Differences in Stroke cohort. *Clin Biochem.* 2014;47: 243–246. doi:10.1016/j.clinbiochem.2014.08.003

23. Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol.* 2004;159: 702–706.
24. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison Himmelfarb C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertens Dallas Tex* 1979. 2018;71: 1269–1324. doi:10.1161/HYP.0000000000000066
25. Lambert D. Zero-Inflated Poisson Regression, with an Application to Defects in Manufacturing. *Technometrics.* 1992;34: 1–14.
26. Preisser JS, Stamm JW, Long DL, Kincade ME. Review and recommendations for zero-inflated count regression modeling of dental caries indices in epidemiological studies. *Caries Res.* 2012;46: 413–423. doi:10.1159/000338992
27. Van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate Imputation by Chained Equations in R. *J Stat Softw.* 2011;45: 1–67.
28. Rubin D. Multiple Imputation for Nonresponse in Surveys. New York: John Wiley & Sons, Ltd;
29. Ruhl A, Jeffries N, Yang Y, Naik R, Patki A, Pecker L, et al. Alpha Globin Gene Copy Number is Associated with Prevalent Chronic Kidney Disease and Incident End-stage Kidney Disease among Black Americans. *J Am Soc Nephrol.* 2021 [cited 14 Dec 2021]. doi:10.1681/ASN.2021050653
30. Keller TCS, Keller AS, Broseghini-Filho GB, Butcher JT, Page HRA, Islam A, et al. Endothelial alpha globin is a nitrite reductase. *bioRxiv.* 2021; 2021.02.12.430954. doi:10.1101/2021.02.12.430954
31. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension.* 2020;75: 1334–1357. doi:10.1161/HYPERTENSIONAHA.120.15026
32. Raffield LM, Ulirsch JC, Naik RP, Lessard S, Handsaker RE, Jain D, et al. Common α -globin variants modify hematologic and other clinical phenotypes in sickle cell trait and disease. *PLOS Genet.* 2018;14: e1007293. doi:10.1371/journal.pgen.1007293
33. Raven PB, Chapleau MW. Blood Pressure Regulation XI: Overview and Future Research Directions. *Eur J Appl Physiol.* 2014;114: 579–586. doi:10.1007/s00421-014-2823-z
34. Ehret GB, Ferreira T, Chasman DI, Jackson AU, Schmidt EM, Johnson T, et al. The genetics of blood pressure regulation and its target organs from association studies in 342,415 individuals. *Nat Genet.* 2016;48: 1171–1184. doi:10.1038/ng.3667

35. Gerhard M, Roddy MA, Creager SJ, Creager MA. Aging progressively impairs endothelium-dependent vasodilation in forearm resistance vessels of humans. *Hypertens Dallas Tex* 1979. 1996;27: 849–853. doi:10.1161/01.hyp.27.4.849
36. Ungvari Z, Tarantini S, Kiss T, Wren JD, Giles CB, Griffin CT, et al. Endothelial dysfunction and angiogenesis impairment in the ageing vasculature. *Nat Rev Cardiol*. 2018;15: 555–565. doi:10.1038/s41569-018-0030-z
37. Hongo K, Nakagomi T, Kassell NF, Sasaki T, Lehman M, Vollmer DG, et al. Effects of aging and hypertension on endothelium-dependent vascular relaxation in rat carotid artery. *Stroke*. 1988;19: 892–897. doi:10.1161/01.str.19.7.892
38. Lang MG, Noll G, Lüscher TF. Effect of aging and hypertension on contractility of resistance arteries: modulation by endothelial factors. *Am J Physiol*. 1995;269: H837-844. doi:10.1152/ajpheart.1995.269.3.H837
39. Panza JA, García CE, Kilcoyne CM, Quyyumi AA, Cannon RO. Impaired Endothelium-Dependent Vasodilation in Patients With Essential Hypertension Evidence That Nitric Oxide Abnormality Is Not Localized to a Single Signal Transduction Pathway. *Circulation*. 1995;91: 1732–1738. doi:10.1161/01.CIR.91.6.1732