Single-cell genomics improves the discovery of risk variants and genes of Atrial Fibrillation

Alan Selewa[1*], Kaixuan Luo[2*], Michael Wasney[3+], Linsin Smith[4+], Xiaotong Sun[2+], Chenwei Tang[5], Heather Eckart[3], Ivan P. Moskowitz[2,6], Anindita Basu[3], Xin He[2#], Sebastian Pott[3#]

[1]Biophysical Sciences Graduate Program,
[2]Department of Human Genetics,
[3]Department of Medicine, Section of Genetic Medicine,
[4]Committee on Genetics, Genomics and Systems Biology,
[5]The College,
[6]Department of Pediatrics,
The University of Chicago, Chicago, IL 60637, USA;
*, +, # These authors contributed equally.

Correspondence: obasu@uchicago.edu, xinhe@uchicago.edu, spott@uchicago.edu

Abstract

Genome-wide association studies (GWAS) have linked hundreds of loci to cardiac diseases. However, in most loci the causal variants and their target genes remain unknown. We developed a combined experimental and analytical approach that integrates single cell epigenomics with GWAS to prioritize risk variants and genes. We profiled accessible chromatin in single cells obtained from human hearts and leveraged the data to study genetics of Atrial Fibrillation (AF), the most common cardiac arrhythmia. Enrichment analysis of AF risk variants using cell-type-resolved open chromatin regions (OCRs) implicated cardiomyocytes as the main mediator of AF risk. We then performed statistical fine-mapping, leveraging the information in OCRs, and identified putative causal variants in 122 AF-associated loci. Taking advantage of the fine-mapping results, our novel statistical procedure for gene discovery prioritized 46 high-confidence risk genes, highlighting transcription factors and signal transduction pathways important for heart development. In summary, our analysis provides a comprehensive map of AF risk variants and genes, and a general framework to integrate single-cell genomics with genetic studies of complex traits.

Introduction

Cardiac diseases are a leading cause of mortality across the world[1,2]. GWAS of cardiac traits have uncovered a large number of associations, including more than 100 loci linked to atrial fibrillation (AF)[3–7]. However, in most loci the disease-driving causal variants remain unknown. Given that

most trait-associated variants are located in non-coding regions[8], researchers often use regulatory

and epigenomic datasets to annotate possible effects of variants, and to prioritize putative causal

variants[8–10]. Existing datasets, however, were often collected from bulk tissue samples that

represent complex mixtures of cell types[11,12], while disease-causing variants often act in specific

cell types. Thus, lack of cell type-resolved epigenomic data in disease-related tissues limits our

ability of variant annotation and prioritization. Another challenge of post-GWAS analysis is that

long-range gene regulation is common, making it difficult to link non-coding variants with their

target genes.

Despite these challenges, researchers have made attempts to identify putative risk variants and

genes underlying the AF genetics. One study used epigenomic and gene expression data in the

human heart to nominate putative risk genes in 104 AF-associated loci. This study, however, did

not use rigorous statistical analysis to fine-map causal variants and instead used relatively lenient

cutoffs and an *ad hoc* scoring scheme to rank putative target genes[13]. This study nominated nearly

300 genes in these loci, many of which are likely not causal genes. Another study used STARR-

seq to map regulatory regions and variants to nominate risk variants in 12 AF-associated loci[14].

But the majority of AF-associated loci were not investigated in the study. A more recent study

collected single-cell RNA-seq and ATAC-seq data in the human heart, and performed fine-

mapping in AF-associated loci[15]. The study identified 38 putative risk variants in heart CREs.

Nevertheless, few nominated variants reach high confidence and in most loci the risk genes remain

unknown.

To address these challenges in the context of heart diseases, we developed an integrated

framework that unifies advances in single cell epigenomics, computational fine-mapping and a

novel procedure for risk gene discovery. Specifically, we performed single-cell chromatin

accessibility profiling to map CREs across major cell types in the heart. Our statistical fine-

mapping method utilizes the CRE maps to infer disease-relevant cell types and takes advantage of

such information to identify putative causal variants. Our novel gene-mapping approach then

aggregates information of all fine-mapped SNPs to predict the risk genes, considering multiple

sources of information such as distance and chromatin loops between enhancers and promoters.

Application of this framework to AF revealed a number of putative risk variants and genes,

highlighting biological processes important to the genetics of AF.

An unexpected finding from our study is that the majority of risk variants of AF we discovered

did not colocalize with heart eQTLs. Taking advantage of our cell-type resolved epigenomic data,

we found that this was largely due to the lack of power of bulk eQTL studies to identify regulatory

variants with cell-type specific effects. This finding thus sheds light on the common strategy of

annotating GWAS results using eQTLs.

Results

Overview of the experimental and computational approach. Our approach combines single-

cell genomics with novel computational procedures to study genetics of cardiac traits (Fig. 1).

Using single nucleus RNA-sequencing[16–18] (snRNA-seq) and single cell ATAC-seq (scATAC-

seq)[19,20], we obtained transcriptome and open chromatin regions (OCRs) across all major cell types

in the adult human heart (Fig. 1, step 1). These OCR profiles allow us to discover cell types

enriched with the genetic risks of traits of interest. To identify specific causal variants in trait-

associated loci, we performed Bayesian statistical fine-mapping. Fine-mapping is a technique that

aims to identify one or few causal variants that explain all the associations in a locus. It avoids the

use of arbitrary LD cutoffs in selecting candidate variants and is able to quantify the uncertainty

of each nominated variant. Recent fine-mapping techniques are also able to incorporate functional

information of variants, such as regulatory activities in trait-related cell types[9,21,22]. Because of

these benefits, fine-mapping techniques have been successfully applied to many common traits

such as Type 2 Diabetes[23], Schizophrenia[24] and autoimmune disorders[25]. Our fine-mapping

method takes advantage of the cell-type-resolved chromatin data to favor variants located in OCRs

of enriched cell types (Fig. 1, step 2). After fine-mapping, the candidate SNPs and their associated

cell-type information allow us to assign the cell type(s) through which the causal variants are likely

to act.

Finally, we developed a procedure to infer causal genes at each locus (Fig. 1, step 3), addressing

some common challenges. In "gene association tests" researchers test if the set of SNPs near a

gene collectively show disease association[26,27]. These types of methods, however, cannot

distinguish between multiple genes close to disease-associated variants. Alternatively, researchers

may perform fine-mapping first, then link the high-confidence SNPs to target genes using

additional information. However, fine-mapping alone rarely leads to a single, or even a few, high

confidence SNPs at associated loci [23], therefore this approach also has limited utility. In contrast,

our procedure aggregates information of all fine-mapped variants in a locus to nominate risk genes.

To see its benefit, suppose fine-mapping in a locus implicates 10 putative causal variants without

any single one reaching high confidence; however, if all 10 SNPs likely target the same gene, we

can be confident of the causal gene. We developed a statistical procedure to implement this idea,

and estimated a score, called gene PIP (Posterior Inclusion Probability), for each gene. Under

certain assumptions, we showed that the gene PIP estimates the probability of a gene being causal.

The details are described below (Fig. 5a) and in Methods.

Single-cell transcriptome and chromatin accessibility profiling reveals multiple cell types in

the human heart. We performed snRNA-seq and scATAC-seq using the Chromium platform

(10x Genomics) (Fig. 1, step 1). The heart samples were obtained from the left and right ventricles

(LV and RV), the interventricular septum, and the apex of three adult male donors (Supplementary

Table 1). After quality control, we retained data of 49,359 cells in snRNA-seq and 26,714 cells in

scATAC-seq, respectively (Extended Data Fig. 1 and 2).

We characterized cell populations with clustering analysis in both snRNA-seq and scATAC-seq

datasets. From snRNA-seq[28], we identified eight major cell types based on marker genes and

comparison to published single-cell heart atlas data[17] (Fig. 2a, left, Extended Data Fig. 3a), with

~70% of cells from cardiomyocytes (CMs), fibroblasts, and endothelial cells. Clustering based on

scATAC-seq data[29] revealed similar cell populations (Fig. 2a, right). We computationally

transferred cluster labels from snRNA-seq onto scATAC-seq clusters[28] (Methods) and

unambiguously identified matching cell types (Extended Data Fig. 3b, c). Indeed, expression and

chromatin accessibility near marker genes showed high cell-type specificity (Fig. 2b, c). Across

the eight clusters, gene scores inferred from scATAC-seq, a metric that summarizes the chromatin

accessibility near a gene[29] (Methods), were highly correlated with transcript levels in the matched

clusters (Extended Data Fig. 3d). We also found good agreement between cell types identified in

our scATAC-seq data and a recent study (Hocker *et al.*[15]), the only differences in annotation

between these two studies was that we did not detect atrial cardiomyocytes, owing to our use of

ventricular samples, and that we detected separate pericytes and smooth muscle clusters (Fig. 2a),

whereas Hocker *et al.* annotated a single large cluster as "smooth muscle". These results supported

our cell-type assignments in both modalities.

Analysis of scATAC-seq data identifies cell-type-specific regulatory elements and their regulators. We pooled cells of the same cell type and identified OCRs separately in each cell type. Combining samples of the same cell type (Extended Data Fig. 4a, b), we detected 45,000-150,000 OCRs per cell type (Extended Data Fig. 4c) yielding a union set of 352,904 OCRs. *K*-means clustering of these regions based on their accessibility suggested that most OCRs are active in specific cell types (Fig. 3a). Using differential accessibility (DA) analysis, we identified 173,782 (49%) OCRs with cell-type-specific accessibility (Methods). We divided the remaining 179,122 (51%) OCRs into three categories based on their detection across cell types: shared in 2-3 cell types, shared in >=4 cell types (denoted as Shared 2-3 and Shared 4), and remaining ones, denoted as "non-DA OCRs", which mostly comprise peaks with low read counts (Methods). In agreement with previous observations, shared OCRs were enriched in promoter regions[30] (Fig. 3b, c).

We compared our OCRs to regulatory regions identified in multiple tissues in ENCODE[12]. As expected, a large fraction of OCRs from major heart cell types (e.g., CMs, endothelial, fibroblasts) overlapped with DNase Hypersensitive sites (DHS) from the ventricles (Fig. 3d, top, Extended Data Fig. 6). In contrast, the proportions of OCRs from rare cell types (e.g., myeloid) overlapping with bulk DHS were significantly smaller, suggesting that scATAC-seq is more sensitive and detects more regulatory elements specific to rare cell types compared to bulk DHS (Fig. 3d, top, Extended Data Fig. 5d). This can be seen in several cell-type-specific OCRs near marker genes of rare cell types, which were largely undetected in the pseudo-bulk sample (Extended Data Fig. 4). Additionally, 60-80% of OCRs from major cell types overlapped with H3K27ac regions from LV and RV, suggesting enhancer activities (Fig. 3d, bottom, Extended Data Fig. 6). Together, these results showed that scATAC-seq identified cell-type-specific regulatory elements. We also compared cell-type-specific OCRs identified in our study to peaks identified by an earlier single-cell study in heart[15] and found that OCRs showed good agreement across studies (Extended Data Fig. 7). Importantly, more than 75% of OCRs detected in our CMs were also detected in ventricular CMs by Hocker *et al.* (Extended Data Fig. 7).

Chromatin accessibility is largely controlled by lineage-specific transcription factors (TFs)[31]. To identify these TFs, we assessed the enrichment of TF motifs in OCRs specific to each cell type and identified 260 significantly enriched motifs (Methods). Because TFs of the same family may share similar motifs, we performed additional analysis to infer the exact TFs driving the enrichment, assuming that for these TFs, their motif enrichment should correlate with gene

expression across cells. To test this, we correlated motif accessibility scores of TFs calculated by

chromVar[32] with their accessibility-derived gene scores, a proxy for gene expression[29] (Methods).

This analysis yielded 76 TFs with enriched motifs and correlation > 0.5 (Fig. 3e, Supplementary

Table 2). Many of these TFs are cell type-specific (Fig. 3e) and include known CM regulators,

such as TBX5, GATA4, and MEF2A[33] (Fig. 3f). These results provided a compendium of putative

transcriptional regulators across major cell types in the human heart.

Open chromatin regions in CMs are enriched with risk variants of heart diseases and inform

statistical fine-mapping. Using our cell-type-resolved OCRs, we assessed the contributions of

different cell types to genetics of heart-related traits[34]. Risk variants of AF were almost exclusively

enriched (>10-fold) in OCRs from CMs (Fig. 4a). Similar findings were reported in an earlier

study[15]. Interestingly, the variants associated with the PR interval showed a similar enrichment

pattern, suggesting a genomic link between PR interval and AF risk for future investigation[35] (Fig.

4a). In contrast, risk variants of cardiovascular traits and diseases, and blood pressure, were

enriched across multiple cell types (Fig. 4a). As control, non-cardiovascular traits showed little or

no enrichment in heart cell types (Fig. 4a). We also checked the enrichment of genetic risk of AF

at open chromatin regions at individual cells, using the method SCAVENGE[36]. This analysis

confirms that the vast majority of cells enriched with AF risk are CMs (Extended Data Figure 8).

Together, these results suggest distinct cell type origins of different heart-related traits,

highlighting CMs as the main cell type underlying AF.

This observation motivated us to take advantage of the epigenomic data to statistically fine-map

causal variants in 122 approximately independent AF-associated loci[37]. We first used TORUS[21] to

estimate how putative risk variants are enriched in multiple functional annotations, including

protein-coding regions, conserved sequences, and OCRs in CMs (Extended Data Fig. 9a,

Methods)[21]. This information was used to set prior probabilities of variants being causal. We then

performed fine-mapping analysis of all AF-associated loci with SuSiE[38]. Compared to fine-

mapping that treats all variants equally (uniform prior), this procedure increased the number of

high-confidence risk variants. In total, we identified 68 variants whose probabilities of being causal

variants, denoted as Posterior Inclusion Probabilities (PIP), are 0.5 or higher, compared with 44 at

PIP >= 0.5 under the uniform prior (Fig. 4bc, Supplementary Table 3, Extended Data Fig. 9bc).

Across 122 loci, our procedure narrowed down putative causal variants to 5 or fewer SNPs in 51
loci (Fig. 4d).

We highlighted the advantage of our functionally informed fine-mapping with some examples.
In the locus containing HCN4, several SNPs are in high LD with similar GWAS association
statistics (Fig. 4e, top). Our procedure identified a single SNP, rs7172038, as the most likely causal
variant (PIP = 0.99) in the locus (Fig. 4e, middle). This SNP is inside a CM-specific OCR, and
H3K27ac region in the heart. Interestingly, the Activity-by-contact (ABC) method [39] predicted a
loop between the SNP and the HCN4 gene. HCN4 is an ion channel and has been implicated in
the genetics of AF[14]. In another example, we nominated a likely causal variant (PIP = 0.96) among
several high LD variants in the locus containing GATA4 (Extended Data Fig. 9d), an important
TF for AF[40].

A recent study nominated putative causal variants in 12 AF-associated loci by detecting
regulatory variants using STARR-seq[14]. We compared these variants with our fine-mapped
variants (Supplementary Table 4). Among the 9 loci where fine-mapping identifies one variant at
PIP > 0.5, the fine-mapped variants agree with allele-specific variants from STARR-seq in two
loci. In the remaining cases, the disagreement is driven by two sources. Most of the allele-specific
variants from STARR-seq have much lower GWAS association than fine-mapped variants,
suggesting that statistically they are unlikely to be causal variants (see examples in Extended Data
Fig. 10). Also STARR-seq tested variant functions *in vitro,* and a few of allelic variants have no
regulatory annotations *in vivo* (Supplementary Table 4). These results together added to the
emerging picture that in a trait-associated locus, multiple variants may show regulatory effects *in*
vitro[41]. But to identify true causal variants, we believe one should consider both regulatory
information and the strength of GWAS evidence.

The fine-mapping results inform how the risk variants are partitioned into various functional
categories, such as exons and OCRs in different cell types. The sum of PIPs of all SNPs assigned
to a category can be interpreted as the expected number of causal variants in that category. We
found that >50% of causal signals are from OCRs and 30% of signals from CM-specific OCRs,
highlighting the key role of CMs in AF (Fig. 4f). As expected, exons and UTRs explain only 3%
of causal signals.

The same PIP summation approach can also be applied to each locus, with the PIP sum of a
functional category, e.g., OCRs or exons, now interpreted as the probability that the causal variant

in that locus falls into that category. Using this approach, we estimate that at more than half of all

loci, causal variants have >50% probability to localize to OCRs (Fig. 4g). Further partitioning of

OCRs into cell-type-level categories (Fig. 3b), we identified 37 loci where the causal signals

almost entirely (>90%) come from CM-OCRs (Fig. 4h). With the only exceptions of two loci, CM

OCRs explain the causal signals in most of the loci, based on OCR annotations (Fig. 4h,

Supplementary Table 5). Together these results highlighted that our approach can identify cell type

contexts of individual loci.

Fine-mapped variants are supported by regulatory annotations and experimental validation.

We characterized the regulatory functions of 68 specific variants at PIP >= 0.5. The majority

(42/68) were located in CM-OCRs (Fig. 4i, Supplementary Table 3). 60% (41/68) of all variants

and 86% (36/42) of variants in CM-OCRs overlapped H3K27ac marks in the heart, suggesting

enhancer activities (Fig. 4i). 40% of variants (27/68) overlapped with fetal DHS[12], suggesting that

these variants may act across fetal and adult stages (Fig. 4i). Additionally, 22% of variants were

linked to promoters through chromatin loops in Promoter-capture HiC (PC-HiC) from iPSC

derived CMs[42] (Fig. 4i). Using mouse ChIP-seq datasets of three key cardiac TFs (*GATA4, TBX5,*

NKX2-5)[33], we found that five candidate variants are located in human orthologous regions of TF

binding sites, representing 4-fold enrichment over expectation by chance (Extended Data Fig. 9e).

We also found that 22% (15/68) SNPs alter binding motifs (Fig. 4i) of one of the 76 TFs we

identified as likely transcriptional regulators in heart cell types (Fig. 3e). Together, these results

supported regulatory functions of many fine-mapped variants.

We experimentally tested six non-exonic variants with PIP > 0.95 that were located inside CM-

OCRs and overlapped with putative enhancers marked by H3K27ac or H3K4me1/3 (Fig. 4j,

Supplementary Table 6). Four out of six variant-containing OCRs induced reporter gene

expression in mouse cardiac cells (HL-1 cell line)[43,44] (Extended Data Fig. 11a, Methods), but not

in a fibroblast line (3T3), suggesting cell-type-specific activity of the four OCRs (Extended Data

Fig. 11b). Three out of these four variants showed allelic changes of reporter activities in cardiac

cells, for at least one alternative allele (Fig. 4j). The most striking effect was observed for

rs7172038. Two alternative alleles of this SNP (A and G) strongly reduced activation. The

enhancer containing this SNP interacts with the promoter of *HCN4* located about 5 kb away,

according to Activity-by-Contact (ABC) score[39] (Supplementary Table 3). *HCN4* is a well-known

AF risk gene and is physiologically implicated in cardiac rhythm control[45]. Consistent with these
results, deletion of a syntenic 20 kb region in mice containing this enhancer significantly reduced
the expression of *HCN4*[14]. Notably, in two out of three SNPs with allelic effects, the use of
functional information in fine-mapping significantly boosted their PIPs to >= 0.95 (PIP = 0.40 for
rs7172038 and 0.41 for rs1152591 under the uniform prior). These experimental results supported
regulatory functions of our high confidence variants.

In principle, we expect regulatory variants to affect transcript levels of target genes. Using GTEx
eQTL data from the left ventricle (LV), we found that only 31% (21/68) variants are eQTLs
(Supplementary Table 7). And only in four cases, the eQTLs showed plausible evidence of
colocalization (PP4 > 0.5 using coloc[46]) with the AF risk (Supplementary Table 7). The small
overlap of fine-mapped variants with heart eQTLs suggests a limitation of bulk eQTL data to
identify regulatory variants, an issue we will address in more detail below.

A novel computational procedure utilizes fine-mapping results to identify AF risk genes.
Despite our fine-mapping efforts, there remained considerable uncertainty of causal variants in
most loci (Fig. 4d). Even if the causal variants are known, assigning target genes can be difficult
due to long-range regulation of enhancers[47]. We developed a novel procedure, called Mapgen, to
address these problems (Fig. 5a, top): **(1)** For every putative causal SNP, we assign a weight to
each nearby gene, considering multiple ways the SNP may affect a gene (see below). The weight
of a gene can be viewed as the probability that the SNP affects that gene. **(2)** For each gene, we
then aggregate the causal evidence of all SNPs likely targeting this gene, expressed as the weighted
sum of the PIPs of all these SNPs. To ensure that the causal evidence of a variant is not counted
multiple times when it targets multiple genes, we normalize the SNP-to-gene weights in this
calculation. The resulting "gene PIP" approximates the probability of a gene being causal
(Methods). Similar to variant-level fine-mapping, we also define a "credible gene set", the set of
genes that capture the causal signal at a locus with high probability (Methods).

The weights of SNP-gene pairs reflect the strength of biological evidence linking SNPs to genes
(Fig. 5a, bottom). For a SNP in an exon or in a regulatory region linked to a particular gene, we
assign a weight of 1 to that gene. When a SNP cannot be linked to any gene in these ways, its
target genes are assigned using a distance weighted function so that nearby genes receive higher
weights (Methods).

We identified 46 genes with gene PIP >= 0.8, and 87 with gene PIP >= 0.5 (Fig. 5b,

Supplementary Table 8, and Table 1 for top prioritized genes). At each locus, we obtained credible

gene sets that captured at least 80% of the causal signal. These credible gene sets contained a single

gene in 42 out of 122 blocks, and two genes in 34 blocks (Fig. 5c, Supplementary Table 9). The

genes at PIP >= 0.8 included many known AF risk genes such as TFs involved in cardiac

development and atrial rhythm control (e.g. *TBX5*[40] and *PITX2*[48]), ion channels (e.g. *KCNN3*[50]),

and genes involved in muscle contraction (e.g. *TTN*).

We note that a key benefit of Mapgen is that even in the absence of high-confidence causal

variants, it may still identify putative risk genes. In 14 out of 46 genes at PIP >= 0.8, the SNP

level PIPs were diffused, i.e., no single SNP reached PIP >= 0.5 (Supplementary Table 8). As an

example, GJA5, a known AF risk gene[51], was supported by seven SNPs (highest PIP = 0.22), five

of which were linked to GJA5 by PC-HiC loops. This led to the gene PIP of 0.80 for GJA5 (Fig.

5d, Supplementary Table 3). NKX2-5, a well known transcription factor important for heart

development[52], was supported by four SNPs (highest PIP = 0.37), all likely targeting NKX2-5.

This led to a gene level PIP = 0.99 (Extended Data Fig. 12, Supplementary Table 3). These

examples highlighted the advantage of aggregating information from all putative causal variants.

We benchmarked the performance of Mapgen to nominate risk genes against alternative

methods. Given the absence of a comprehensive list of known AF risk genes, we used, as a proxy,

a set of Gene Ontology (GO) terms previously linked to AF[5]. A gene annotated with one or more

of these terms would be considered as a "true" gene in our evaluation, and otherwise a "false"

gene. We considered several methods: nearest gene to GWAS lead SNPs (nearest), Activity-by-

contact (ABC) score linking enhancers to promoters (ABC-max), a gene association test method

(MAGMA), and heart eQTLs linking variants to genes. Additionally, we included a recent study

that nominated risk genes in AF-associated loci based on functional genomic data in heart (denoted

as van Ouwerkerk[13]). We found that all these alternative methods, except ABC, have precision

below or near 50% (Fig. 5e). ABC score has a precision at 60%, but its sensitivity is low, detecting

only a few genes. Mapgen, at the threshold of gene PIP > 0.8, reaches a precision of 76%, while

detecting nearly 50 genes. These results thus demonstrated the advantages of Mapgen for risk gene

discovery.

We next examined specific loci in detail to gain insights of the weaknesses of existing methods,

and how Mapgen addresses them. We focused on the three commonly used methods, nearest gene

method, the use of chromatin interaction data, and eQTL. Among the 46 genes at PIP >= 0.8, eight

(17%) were not the nearest genes, by distance to TSS, to the top GWAS SNPs. Some of these

genes have been implicated in AF and related phenotypes, including KCNN3, TTN and HCN4.

Most of the other genes have plausible functions such as CALU[53], SSPN, and PKP2

(Supplementary Table 11). Most of the nearest genes in these loci, in contrast, showed limited or

no functional relevance (Supplementary Table 11). As an example, in the locus containing CALU,

the nearest gene of the top SNP, rs55985730 (PIP 0.91) is OPN1SW, an opsin gene with function

in color vision, but no clear relevance to AF. This SNP is looped to a distal gene CALU in PC-

HiC data (Fig. 5f), allowing Mapgen to identify CALU as the likely risk gene. CALU is a calcium-

binding protein and involved in alleviation of endoplasmic reticulum (ER) stress in

cardiomyocytes[54]. ER stress has a critical role in the pathophysiology of AF[55]. These results

suggest that by using chromatin loop information, Mapgen is able to identify distal risk genes.

We also considered the use of chromatin conformation in resolving target genes of high PIP

SNPs. We found that while chromatin looping data were useful, as shown in the *CALU* example

above, using such information alone may miss many potential risk genes. Among 68 SNPs at

PIP >= 0.5, only five showed chromatin interactions with promoters based on ABC scores[39], and

18 if we included both ABC and PC-HiC data. Additionally, it is common to observe multiple

chromatin loops at a single SNP. Among the 18 SNPs with chromatin interactions, 50% (9/18)

contact more than one promoter (Supplementary Table 3), highlighting the uncertainty of target

genes from chromatin looping data.

Use of expression QTLs is another common strategy for linking SNPs to genes. However, as

reported above, few fine-mapped variants colocalized with eQTLs. Even if a GWAS SNP is also

an eQTL, it may not identify the correct target gene. For example, in the *TTN* locus, the top SNP

(rs3731746) is an eQTL of *FKBP7*, but the true risk gene is very likely *TTN*[56,57].

Altogether, these results demonstrated the improved ability of Mapgen to nominate plausible

candidate genes compared to alternative approaches linking SNPs to genes.

Putative AF risk genes are supported by multiple lines of evidence. We evaluated our candidate

genes using multiple sources of data. Consistent with enrichment of AF variants in CM-OCRs,

candidate genes (PIP >= 0.8) tended to have higher expression in CMs, compared with other genes

in the AF-associated loci (Fig. 6a). Among 46 loci with PIP proportion >= 50% in cardiomyocyte

OCR, the likely target genes (gene PIP >= 0.5) were highly enriched (nearly 10-fold) in
cardiomyocyte differentially expressed genes (Extended Data Fig. 13a). Additionally, high PIP
genes were enriched in AF-related Mendelian disorders (Supplementary Table 12) (Fig. 6b). We
also compared our genes with those prioritized by earlier work that used additional functional data
such as AF-related gene ontology and heart gene expression[5,13]. While such functional data was
not used in our analysis, the genes at PIP >= 0.8 scored on average substantially higher in two
earlier studies than low PIP genes (Extended Data Fig. 13b), and 32 of them (71%) were supported
by at least one study (Supplementary Table 8).

We next assessed the functions of candidate genes using Gene Ontology (GO) and gene
networks[58]. GO analysis showed enrichment of Biological Processes related to heart development
and cardiac function, and of Molecular Functions such as ion channels, hormone binding and
protein tyrosine kinase (Fig. 6c, Supplementary Table 13). For network analysis, we used the
STRING gene network built with genes at a relaxed PIP threshold of 0.5 (87 genes) to increase the
number of interactions. This analysis highlighted some well-known processes in AF, such as ion
channels, and structure components of heart muscle (Fig. 6d). A prominent subnetwork consisted
of key TFs, including *GATA4, TBX5, NKX2-5* and *HAND2*, implicated previously in AF genetics
and/or heart development[40,59–61] (Fig. 6d). Two other TFs in the network, *PITX2* and *ZFHX3*, are
also well-known AF genes[40]. Combined with the fact that putative causal variants were enriched
in binding sites of TBX5, NKX2-5 and GATA4 (Fig. 4i, Extended Data Fig. 9e), these results
suggested that perturbation of transcriptional regulatory networks consisting of TFs and their
targets, plays a critical role in the genetics of AF. Additionally, the interaction network highlighted
signal transduction pathways, including MAPK signaling and Ephrin signaling (Fig. 6d). Both
processes are important in heart development[62–65]. Indeed, 19 out of 87 genes at PIP >= 0.5 were
annotated by the GO term "regulation of intracellular signal transduction" (FDR < 0.02)
(Supplementary Table 14).

Finally, we found additional literature support for the candidate genes. 39 out of 46 (85%) genes
at PIP >= 0.8 have reported roles in cardiac processes and/or diseases from literature
(Supplementary Table 10). The subset of genes at PIP >= 0.95 with literature support, as well as
their supporting SNPs, were shown in Table 1. The majority of these genes have not been
established as AF risk genes through functional studies, representing novel yet biologically
plausible risk genes.

Cell-type-specific epigenomes reveal insights to lack of colocalization of GWAS signals to heart eQTLs. While a large fraction of fine-mapped AF SNPs fell inside CM-specific OCRs (Fig. 4f), most of them did not colocalize with heart eQTLs (Supplementary Table 7). This result adds to the growing evidence that eQTLs may explain a relatively small percent of GWAS signals of complex traits[66,67]. It is unclear, however, why this is the case[68]. We took advantage of our cell-type resolved transcriptomic and epigenomic data to investigate this issue. We hypothesized that the heart eQTLs missed a large fraction of regulatory variants specific to CMs, thus were depleted of AF risk variants. To assess this hypothesis, we focused on 1,216 heart (LV) eQTLs from GTEx where the causal variants (known as eQTNs) were fine-mapped with high confidence (PIP >= 0.8) (Supplementary Table 15).

Given that the heart eQTL study was performed on bulk tissues, the cell types where these eQTLs act are unknown. We used eQTL information in other tissues to infer whether the eQTLs are likely CM-specific. Our reasoning is that eQTLs that were found across many tissues are likely to be functional in cell types shared across tissues, thus not specific to CMs. We found that the majority of eQTLs were highly shared, i.e., found in >30 tissues in GTEx (Fig. 7a). Less than 10% of heart eQTLs were found in 5 or fewer tissues. These results thus suggest that the detected heart eQTLs are highly biased towards variants with effects in cell types shared across tissues.

This finding thus confirms our hypothesis that the detected heart eQTLs are generally not specific to CMs, even though CMs constitute a relatively large fraction of heart cells (20-30%, Fig. 2a). To understand these results, we divided the heart eQTLs into different categories based on the location of the eQTNs, including exons, UTRs, introns, OCRs in specific cell types, and OCRs shared with varying numbers of cell types. We analyzed the tissue-sharing pattern of each category separately. The overall sharing pattern of all eQTLs would depend on the sharing pattern of each category, and the percent of eQTLs in each category (see Methods). This analysis would thus allow us to understand what drives the high degree of observed eQTL sharing across tissues.

As expected, eQTLs falling into OCRs shared in multiple cell types were extensively shared across tissues (Fig. 7b). On the other hand, eQTLs in cell-type-specific OCRs showed variable levels of

sharing. Fibroblast-eQTLs (eQTLs in fibroblast-specific OCRs) and myeloid-eQTLs were highly

shared (median 31 and 43 tissues, respectively), but most CM-eQTLs were found in <10 tissues

(Fig. 7b). We believed this variability reflected different degrees of cell type sharing between the

heart and other tissues. To test this, we compared heart eQTLs with those from the brain and whole

blood. As expected, heart eQTLs from immune cell OCRs had the highest sharing with whole

blood, while eQTLs of all heart cell types have low sharing with the brain (Fig. 7c). Together,

these results highlighted considerable variability of tissue sharing patterns of heart eQTLs,

depending on their likely cell-type origins.

We next assessed the proportions of heart eQTLs in functional categories, focusing on eQTLs in

OCRs, whose cell type origins could be inferred. A large proportion of those eQTLs were from

OCRs shared in multiple cell types, with eQTLs in CM-specific OCRs only explain <10% of heart

eQTLs (Fig. 7d). Given that different categories of OCRs have different genome sizes, we

compared the proportions of eQTLs in each category with random expectation (Methods). While

eQTLs in OCRs from single cell types showed 2-9 fold enrichment, those shared with 4 or more

cell types showed 26-fold enrichment (Fig. 7d). Indeed, the enrichment is highly correlated with

the number of cell types in which an OCR is detected (Fig. 7e). We thus concluded that discovered

eQTLs are biased towards those with broad effects across multiple cell types.

Altogether, our results suggest that eQTLs that are likely CM-specific are under-represented in the

data, constituting <10% of all heart eQTLs. Most of the remaining eQTLs have effects across

multiple cell types; or have effects in cell types shared across other tissues. As a result, the overall

level of tissue sharing of heart eQTLs is very high. Given that AF risk variants are specifically

enriched in CM-specific OCRs (Fig. 4a), the depletion of CM-specific eQTLs explains why heart

eQTL data fail to explain most GWAS signals.

We reason that this depletion of cell-type-specific regulatory variants in eQTLs can be explained

by the nature of bulk eQTL studies. When the effect of an eQTL on a gene is limited to a single

cell type, but the gene is expressed in other cell types, the effect of the variant on the bulk gene

expression would be diluted, leading to lower power of detecting this eQTL. This argument was

supported by the observation that gene expression was less cell-type-specific than accessibility of

regulatory elements. In heart eQTLs localized to CM-specific OCRs, the expression of

corresponding genes in CMs were only modestly higher than their expression in other cell types

(Extended Data Fig. 14a). We performed simulations to investigate the power loss in detecting

cell-type-specific eQTLs. When the cell type of interest is 20% of the bulk tissue, we estimate that

the power of detecting eQTL specific in this cell type is only about 8-40% of the power of detecting

eQTLs with shared effects across cell types (Supplementary Notes, Extended Data Fig. 14b). This

analysis demonstrated that the low power of detecting cell-type specific eQTLs is a key limitation

of bulk eQTL studies.

Discussion

While GWAS have been successful in a range of complex traits, the causal variants, their target

genes, and their mechanisms in disease-related cell types have been elucidated in few cases[47]. In

this work, we established a cell-type-resolved atlas of chromatin accessibility and transcription of

the human heart to study the genetics of heart-related traits, focusing on AF[3–5]. We statistically

fine-mapped AF-associated loci, and experimentally validated some of the candidate variants.

Using a novel computational procedure, we identified 46 high confidence genes, implicating key

biological processes, in particular TFs and signaling pathways important for heart development.

Motivated by our observation that the putative AF variants often were not colocalized with eQTLs,

we investigated how heart eQTLs are shared across tissue types. Our analysis suggests that eQTLs

with cell-type-specific effects are under-detected and that this is likely a factor explaining both

high tissue-sharing of eQTLs and the lack of eQTLs in GWAS variants.

Compared with several recent studies that aimed to identify risk variants and genes in AF[13,15],

our study has a few key advantages. Hocker *et al*. intersected fine-mapped variants with cell-type-

resolved OCRs to nominate putative regulatory variants. Their work and related studies[69]

demonstrated the utility of single cell ATAC-seq data for interpretation of non-coding variants

from GWAS. Our work extends these studies by using a computational procedure that leverages

the strong enrichment of genetic signals in CM-specific OCRs to fine-map causal variants,

identifying 68 high confidence SNPs at PIP > 0.5, including 42 in CM OCRs, compared to five

nominated by Hocker *et al.*[15] (Fig. 4b). Our gene-mapping procedure effectively leverages fine-

mapping results and multiple sources of information linking SNPs to putative targets. This avoids

the bias of previous work that only considers one metric, e.g., distance, to link SNPs to genes, and

increases the sensitivity of detecting risk genes. As a result, we found high confidence genes
(PIP >= 0.8) in more than 1/3 of known AF-associated loci.

Our set of 46 candidate genes shed light on the genetics of AF. Earlier linkage studies implicated
ion channels and structural proteins, as well as a few TFs[70]. Our results confirmed these earlier
findings and showed an even larger role of regulatory genes, including TFs and signaling proteins.
In total, we identified 13 TFs with PIP >= 0.8 (Supplementary Table 10), and 18 at PIP >= 0.5.
These included known AF genes, TBX5 (PIP 0.99), NKX2-5 (0.99), PITX2 (0.99), ZFHX3 (0.85)
and GATA4 (0.57), as well as TFs with roles in heart development such as HAND2 (0.87), ZEB2
(0.98), and PRRX1 (0.72). Our results also highlighted signal transduction pathways, including
MAPK signaling[62], Ephrin signaling[63–65] (Fig. 6d), G-protein coupled receptor signaling[71], Wnt
signaling[72] (Supplementary Table 13) and FGF signaling[73,74] (FGF9, PIP = 0.94 and FGF5 PIP =
0.53), all previously implicated in heart development.

Despite the advances described above, our study has a few limitations. Our experimental data
were limited to four anatomical locations of the ventricles, while some AF risk variants might act
through atrial-specific CMs. However, it is worth noting that a recent study, using scRNA-seq
based cellular atlas of the heart including all anatomic locations, found that AF candidate genes
were strongly enriched in ventricular CMs[17]. Additionally, our fine-mapping leveraged the almost
exclusive enrichment in CM-specific OCRs (Fig. 4a), and thus may miss variants acting on the AF
risk through other cell types. This possibility is suggested by a small number of candidate variants
showing accessibility specific to fibroblasts (Fig. 4h), known contributors to AF etiology[75].
Finally, some disease variants potentially act transiently during development and might be missed
using adult heart samples.

In conclusion, by combining novel experimental and computational approaches, our study
identified a number of risk variants and genes and revealed key insights of the genetics of AF.
These data provide a rich resource for future functional studies. Importantly, our analytic
framework, including the software for fine-mapping and risk gene identification, may provide a
general model for the study of other complex phenotypes.

Methods

Data collection. *Nuclei isolation from adult heart tissue.* Heart tissue samples were obtained from National Disease Research Interchange (NDRI) and were stored at -80ºC and kept on dry ice whenever outside of the freezer. We included samples from 4 regions (left and right ventricles, interventricular septum, apex) from 3 male individuals (Supplementary Table 1). Aliquots of each heart sample were prepared from frozen heart tissue using a tissue pulverizer, which was cooled prior to pulverization for 20 minutes over dry ice. Aliquots assayed in this study ranged from 86.7 mg to 141.6 mg. Prior to library preparation, we purified nuclei using fluorescence-activated cell sorting (FACS) to remove debris and minimize contamination from ambient RNA.

Single nuclei isolation was performed on the heart tissue aliquots as described in Litvinukova *et al.* 2020[17], with some modifications. Single heart aliquots were kept on dry ice until being transferred into a precooled 2 mL dounce homogenizer (Sigma) with 2 mL homogenization buffer (250 mM sucrose, 25 mM KCl, 5 mM MgCl2, 10 mM Tris-HCl, 1 mM dithiothreitol (DTT), 1x protease inhibitor, 0.4 U/µl, RNaseIn, 0.2 U/µl SUPERaseIn, 0.1% Triton X-100 in nuclease-free water). Samples were dounced 25 times with pestle A (loose) and 15 times with pestle B (tight), filtered through a 40-µm cell strainer, and centrifuged (500g, 5 minutes, 4ºC). Supernatant was discarded and the nuclei pellet was suspended in nuclei resuspension buffer (1x PBS, 1% BSA, 0.2 U/µL RnaseIn) and stained with NucBlue Live ReadyProbes Reagents (ThermoFisher). Hoechst-positive nuclei were enriched using fluorescence-activated cell sorting (FACS) on the FACSAria (BD Biosciences), obtaining between 172,500 and 350,000 nuclei while targeting a maximum of 350,000. Nuclei were sorted into 0.75 ml of resuspension buffer. Flow-sorted nuclei were counted in a C-Chip Disposable Hemocytometer, Neubauer Improved (INCYTO) before commencing with library preparation.

snRNA-seq library preparation and sequencing. A portion of the sorted nuclei suspension was removed and brought to a concentration of between 700 and 1,200 nuclei per microliter. An appropriate number of nuclei were loaded on the Chromium controller (10X genomics) in order to target between 6,000-8,000 nuclei, according to V3 of the manufacturer's instructions for the Chromium Next GEM Single Cell 3′ Reagent Kits (10X Genomics)[76]. 3' gene expression libraries were amplified with 15 cycles during sample index PCR. QC was performed on 3' gene expression cDNA and final libraries using a Qubit Fluorometer (ThermoFisher) and an Agilent 2100

Bioanalyzer (Agilent). Libraries were sequenced on the NovaSeq 6000 (Illumina) or the NextSeq

500 (Illumina) at the University of Chicago's Genomics Facility using paired-end sequencing.

scATAC-seq library preparation and sequencing. scATAC-seq libraries were prepared according

to v1 of the manufacturer's guidelines for the Chromium Next GEM Single Cell ATAC Reagent

Kits (10X Genomics), with the modification that we started from nuclei that were isolated as

described above. Between 9,300 and 25,000 nuclei were tagmented using Transposition Mix (10X

Genomics) at 37°C for 1 h and loaded on the Chromium controller. We targeted between 6,000

and 10,000 nuclei for library preparation. QC was performed on final ATAC-seq libraries using a

Qubit Fluorometer and an Agilent 2100 Bioanalyzer. Libraries were sequenced on the

NovaSeq6000 or the NextSeq500 at the University of Chicago's Genomics Facility using paired-

end sequencing.

Single-cell genomic data analysis. *snRNA-seq pre-processing*. FastQ files from 12 sequencing

experiments were individually processed using an in-house scRNA-seq pipeline dropRunner[77].

Briefly, dropRunner utilizes FastQC[78,79] to obtain quality control metrics followed by fast and

efficient alignment to human reference genome hg38 using `STARsolo 2.6.1`[80] in `GeneFull`

mode with other parameters set to default. STARsolo performs alignment and quantification of

gene expression in one package. We quantified expression at the gene level using `Gencode v29`

gene annotations[81] utilizing both intronic and exonic reads to improve clustering and downstream

analyses of the snRNA-seq data. We extracted the raw gene-by-barcode expression matrices output

by STARsolo for downstream analyses. We used `Seurat 3.2.1`[82] in R to analyze the snRNA-

seq data. We combined all 12 expression matrices into a single Seurat object together with the

corresponding metadata such as donor and anatomical region. To filter low-quality nuclei, we

removed barcodes that contained less than 1000 UMI. We also used `DoubletFinder 2.0.3`[83]

with `pN = 0.015` and `pK = 0.005` to account for doublets, which works by generating in-

silico doublets and performs clustering to identify nuclei that fall in the neighborhood of the

generated doublets. After quality control, we retained a total of 49,359 nuclei.

scATAC-seq pre-processing. FastQ files from 12 sequencing experiments were individually

processed using 10x Genomics `CellRanger-atac 1.2.0`[84]. We used the command

`cellranger-atac count` to align the fastq files to human reference genome hg38, followed by
marking and removing duplicate reads, and producing a fragment file containing the mapped
location of each unique fragment in each nucleus. We used `ArchR 0.9.5`[29] to further pre-process
the data and perform downstream analyses of the scATAC-seq data. Using ArchR, we converted
the fragments file into a tile matrix, which is a bin-by-barcode Tn5 insertion count matrix, using a
bin-size of 500 bp. We also generated a gene score count matrix using the "model 42" from ArchR,
which aggregates Tn5 insertion signals from the entire gene body, scales signals with bi-directional
exponential decays from the TSS (extended upstream by 5 kb) and the transcription termination
site, and accounts for neighboring gene boundaries. Gene annotations were obtained from
`Gencode v29`. To filter low quality nuclei, we kept nuclei with at least 5,000 unique fragments
and a TSS enrichment score of 6. We also used ArchR's doublet removal approach with default
parameters, which is based on in-silico doublet generation. We removed nuclei with a doublet
enrichment score greater than 1. After quality control, we retained a total of 26,714 nuclei.
574
Cell-type identification from snRNA-seq and scATAC-seq. We performed normalization,
dimensionality reduction, and unsupervised clustering on snRNA-seq and scATAC-seq data in
order to identify cell-types. For snRNA-seq, we used Seurat's workflow which begins with
converting counts to log2 TP10k values using the `NormalizeData` function. Next, we found the
top 2000 variable genes using `FindVariableGenes` and used these genes as input features for
Principal Component Analysis (PCA). We computed the top 30 principal components (PCs) for
each cell and used these for downstream analyses. We observed batch effects due to different
donors, and corrected this batch effect. This was done using the `RunHarmony` function from the
`Harmony 1.0`[85] package with default parameters to regress out the donor variable from the PCs.
Next, we used the `FindClusters` in Seurat with a resolution of 0.2 on the harmony-corrected
PCs to define clusters. We also computed the corresponding UMAP to visualize the harmony-
corrected PCs in two dimensions. We used previously established cell-type markers in order to
map clusters to cell types[17,18].
We performed cell-type mapping for scATAC-seq using the `ArchR` package. We performed
dimensionality reduction on the tile matrix using the top 20,000 bins in terms of count across all
cells. We used the function `addIterativeLSI` with 2 iterations in order to perform latent
semantic indexing (LSI) on the scATAC-seq tile matrix and retained the top 50 LSI vectors.

Similar to snRNA-seq, we observed batch effects across different donors, and removed this effect

using the `RunHarmony` function. We used `addClusters` with `resolution = 0.2` in order to

cluster nuclei based on the harmony-corrected LSI vectors. `addUMAP` with `min.dist = 0.4` was

used to compute a 2-dimensional representation of the harmony-corrected LSI vectors. We

visualized gene activity scores, as defined in ArchR, using the same marker genes as in snRNA-

seq to assign clusters to cell-types.

Defining and classifying open chromatin regions. Insertion read counts were aggregated across

all cells in each cell-type to form a cell-type pseudo-bulk and peak calling was performed on

pseudo-bulk data of each cell-type. Using the function `addReproduciblePeakSet` in ArchR in

conjunction with MACS2[86], a union set of 352,900 peaks were called in total across all cell-types

at FDR < 0.1. This set of peaks, called union set, were used for all downstream analyses.

In order to discover cell-type specific regulatory elements, a single-cell insertion count matrix

was created using the function `addPeakMatrix` in ArchR. Cells were grouped into their

respective cell-types and differential accessibility (DA) analysis was performed in a one-vs-all

fashion, i.e., one cell type vs. all other ones. To perform DA, we used `getMarkerFeatures` in

ArchR with default parameters, which uses the Wilcoxon rank-sum test on the log-normalized

insertion count matrix. To control for technical variation, cells from the cell-type group and the

group of remaining cell types are matched in terms of TSS enrichment and number of fragments.

Using FDR < 10% and $\log_2$ fold-change > 1, we found about 47% of the union set to be cell-type

specific.

For OCRs that were not differentially accessible, we reasoned that these are more likely to be

shared. To further stratify these OCRs into different classes, based on sharing among different cell

types, we used a simple quantile-based method. First, we aggregated the ATAC-seq counts across

all cells within each cell-type for each non-DA peak and normalized the counts by the total sum

of counts in each cell-type. Next, we binarized the peaks within each cell-type based on whether

they are in the top 25% or not in terms of their normalized counts. In this way, we identify the top

25% accessible peaks in each cell-type. Finally, we count how many times a peak is 1, or highly

accessible, across cell-types. Through this strategy, we defined three disjoint sets: shared in 2-3

cell types, shared in 4+ cell types and the remaining peaks denoted as "non-DA". The last category

corresponds to peaks that are only highly accessible (top 25%) in one cell type but are not found

to be differentially accessible based on our criteria above.

Comparing snRNA-seq and scATAC-seq. We calculated correlation scores of gene expressions

from snRNA-seq and gene activities from scATAC-seq in the following manner: First, we selected

genes that were up-regulated in each cell type according to differential expression analysis of

snRNA-seq data. Approximately 3000 genes were identified in this manner. For each gene, the

ATAC-seq gene scores and RNA-seq transcript counts, respectively, were aggregated across all

cells in each cell-type cluster, followed by a log transformation. We then used the log-transformed

pseudo-bulk gene scores and normalized expression levels to calculate Pearson correlation

between gene scores and expression across cell types.

Comparing the cell labels in our study with Litviňuková's et al. Label transfer was performed

using Seurat to compare the labels in our study with Litviňuková's *et al.* [17]. The processed scRNA-

seq data from Litviňuková *et al.* were downloaded from

<u>https://www.heartcellatlas.org/#DataSources</u>. `LoadH5Seurat` from `Seurat` R

package was used to convert the h5ad format into a Seraut object. Next, anchors were identified

using the `FindIntegrationAnchors` function and used as reference, which takes the earlier

Seurat object as input. Then, we predicted the labels of our cells with the `TransferData` function,

which used the anchors and our scRNA-seq data (also a Seurat object) as inputs and returned the

predicted labels for each cell in our dataset. We summarized the number of matched cells with a

heatmap, showing the proportion of matched cells in each cluster.

Comparing the OCRs in our study with Hocker et al. We compared the OCRs from our dataset

with Hocker *et al.*[15] using peak sets called on individual cell type clusters. Cell-type level peaks

identified by Hocker *et al.* were obtained from their CARE portal

(`http://cepigenomics.org/CARE_portal/Cell_Type_Diversity.html`). We only

included their peak set from ventricular CMs to match our cell types. For each cell type, we

computed a simple overlap between peaks from both datasets using `GenomicRanges`

`findOverlaps`[87] in R. Overlaps for each cell type were represented as Venn diagrams generated

with the `eulerr` R package.

Identifying putative TFs regulating chromatin accessibility. We used a set of 870 human motif
sequence instances from `CisBP`[88]. These motif annotations were added onto the ArchR object
using the `addMotifAnnotations` function. Next, enrichment analysis was performed for each
motif in each cell-type-specific set of peaks, using the `peakAnnoEnrichment` function in ArchR.
The function uses the hypergeometric test to assess the enrichment of the number of times a motif
overlaps with a given set of peaks, compared to random expectation. After correcting for multiple
testing within each cell-type, we used FDR < 1% to ascertain a set of motifs and their enrichment.

Motif enrichment analysis may find multiple TFs with similar motifs. To reduce the redundancy
and identify true TFs that drive gene regulation, we correlated the motif accessibility with gene
score activity of each TF, expecting that for true TFs, their expression levels should be positively
correlated with accessibility of their motifs across cells. We obtained motif accessibility scores
from `chromVAR`[32] (using the `addDeviationsMatrix` function in ArchR) for each TF across all
cells. We obtained the corresponding TF gene activity scores using the "model 42" by ArchR (see
"scATAC-seq pre-processing"). These single-cell-level motif accessibility scores and gene scores,
however, are noisy given the sparsity of data at individual cells. We thus used a strategy similar to
Cicero[89], by aggregating cells into "metacells" based on similarity using a k-nearest neighbor
approach. Specifically, we found the k nearest neighbors to each cell using the LSI vectors of the
single-cell ATAC-seq data. We only retained sets of metacells that shared a maximum of 25% of
constituting cells. Metacells that shared more than 25% of cells were removed at random. Using k
= 100, we created about 200 non-redundant meta-cells based on these criteria and averaged the
motif accessibility scores and gene scores across cells within each meta-cell. We then computed
Pearson's correlation between the gene scores and the motif accessibility scores across meta-cell.
We selected all TFs with a Pearson's correlation greater than 0.5.

SCAVENGE analysis. SCAVENGE[36] was used to calculate for each cell a trait relevance score
(TRS) for Atrial fibrillation. SCAVENGE was run under default settings, with ATAC-seq peak
matrix and fine-mapping results (under the uniform prior) as inputs.

Testing enrichment of GWAS risk variants in functional annotations. We obtained
harmonized GWAS summary statistics for cardiovascular and some non-cardiovascular traits from

the IEU OpenGWAS project. We removed SNPs with missing values, SNPs on non-autosomal

chromosomes, and indels. Utilizing approximately independent Linkage Disequilibrium (LD)

blocks generated by ldetect[37], we assigned each SNP to one of 1700 LD blocks.

We used TORUS[34] to estimate the genome-wide enrichment of risk variants of GWAS traits in

various functional annotations, including cell-type specific OCRs obtained from DA testing, and

some generic annotations including coding, retrieved from UCSC Genome Browser database, and

conserved sequences from Lindblad-Toh, K. *et al.* 2011[90]. We ran TORUS on each annotation,

one at a time, to get the marginal enrichment reported in Fig. 4a. P-values for enrichment were

estimated from the 95% confidence intervals returned by TORUS and were adjusted for multiple

testing across all traits/cell-types using the Benjamini-Hochberg approach.

Fine-mapping causal variants in AF-associated loci. We start with a general description of

statistical fine-mapping analysis. We assume the trait of interest, Y, is related to the genotypes of

all variants in a locus by a linear model. Let X_j be the genotype of the j-th variant, we have: $Y =$

$\sum_j X_j \beta_j + \epsilon$, where β_j is the effect size of the j-th variant. Because causal variants in a locus are

generally "sparse", it is often assumed that most β_j's would be zero. It is easy to see that, under

this model, even if a single variant has $\beta_j \neq 0$, other variants in LD with this variant would appear

associated with the trait in the standard single-variant association analysis. But in the joint

regression model here, once we choose the correct causal variant(s), conditioned on them, the non-

causal variants in LD would no longer be associated with the trait. The goal of fine-mapping is

then to select as few variants with $\beta_j \neq 0$ as possible to explain all associations in the locus. This

"variable selection" step is often accomplished using a Bayesian spike-and-slab prior, which

assumes that β_j follows a mixture distribution of point mass at 0, and a normal distribution. The

mixture proportion of the point mass is typically very large (close to 1), ensuring that at most a

few variants would have non-zero effects. Inference of this model is computationally difficult. We

used SuSiE in our analysis [38]. SuSiE uses an efficient variational Bayes procedure, and generally

outperforms other fine-mapping tools. The main output of SuSiE is the posterior probability that

$\beta_j \neq 0$, denoted as Posterior Inclusion Probability (PIP).

To run SuSiE on our GWAS summary statistics, we first partitioned the genotype into LD blocks

using LDetect[37,38]. Then, we ran the `susie_rss()` function on each LD block. The input of this

function includes GWAS z-scores and the LD matrix for the SNPs in a block. The GWAS

summary statistics were available publicly. For the LD matrix, we used out-of-sample genotype

information from 1000 Genome Project[91]. We ran SuSiE with `L = 1`, which allows a single causal

signal for each LD block and is robust to mismatching LD patterns. We fine-mapped a total of 122

LD blocks in the AF GWAS, each containing at least 1 SNP at genome-wide significance ($P < 5$

$\times 10^{-8}$).

To incorporate functional information of variants in fine-mapping, we allowed SNPs to have

different prior probabilities in SuSiE. Specifically, each SNP has a different prior distribution of

β_j, with the prior probability that $\beta_j \neq 0$, denoted as π_j, dependent on the functional information of

of that SNP. These prior probabilities are estimated using TORUS[21]. Briefly, TORUS assumes that

π_j is related to the annotations of the SNP through a logistic regression model. These annotations

may include, for example, whether a SNP is located in an OCR in CMs, or in an evolutionarily

conserved region. The parameter of an annotation in the model encodes the extent to which causal

variants are enriched in this annotation. TORUS uses the entire GWAS data of all variants in the

genome to estimate these parameters. We included the following annotations in TORUS: CM

specific OCRs, CM shared OCRs, CM non-DA OCRs, non-CM OCRs, UCSC conserved, coding,

or finemapped eQTLs.

Annotating putative AF causal variants with additional functional data. Fetal DHS and heart

H3K27ac data were obtained from ENCODE. PC-HiC interactions were obtained from an earlier

study conducted in iPSC derived CMs[42]. Only interactions found in at least 2 out of 3 replicates

were included. Motif analysis was performed using R `motifbreak` package[92] . Only "strong"

effects on motif scores, according to the package, were considered.

Assessing regulatory effects of candidate variants by Luciferase assay. Candidate regulatory

elements were designed from CM-specific accessibility in hg38 and synthesized by IDT, with

either the reference allele or SNP allele(s). Sequence was verified and then cloned into the

pGL4.23 enhancer luciferase response vector with a minimal promoter. HL-1 cardiomyocytes

were co-transfected with luciferase response vector and a pRL control using Lipofectamine 3000,

cultured for 48 hr after transfection, then lysed and assayed using the Dual-Luciferase Reporter

Assay system (Promega). For each construct reporter gene activity was assayed in 5 replicates.

Gene mapping procedure with Mapgen. We used the posterior inclusion probabilities (PIPs)

generated by SuSiE to calculate a gene-level PIP, reflecting the probability that a gene is a risk

gene. We assume there is a single causal gene per disease associated locus. Let Z_g be an indicator

variable describing whether gene g is causal ($Z_g = 1$) or not ($Z_g = 0$) for the trait. Assuming a single

causal SNP per locus, the probability that the gene is causal, which is denoted as "gene PIP", can

be then related to the probabilities of SNPs being causal variants:

754
$$P(Z_g = 1|\, D) = \sum_i P(Z_g = 1|\gamma_i = 1)P(\gamma_i = 1|D),$$

where γ_i is the indicator variable for whether SNP i is causal or not, and D is the GWAS summary

statistics. The term $P(Z_g = 1|\gamma_i = 1)$ is the probability that g is the causal gene if the causal SNP

is SNP i, and the term $P(\gamma_i = 1|D)$ is simply the PIP of SNP i, or PIP_i. So the gene PIP of a gene

is a weighted sum of PIPs of all SNPs, weighted by how much that gene is supported by each SNP

(see below). Since the PIPs of all SNPs in a block sum to 1, the gene PIP has an upper-bound of

1. In the rare cases where a gene spans two nearby blocks - e.g. when a gene has large introns, the

gene PIP may exceed 1, which can be interpreted as the expected number of causal variants

targeting the gene g.

To calculate the term $P(Z_g = 1|\gamma_i = 1)$, we consider the location of the SNP i with relation to

the gene g, as well as functional genomic data linking SNP i with gene g. These data were used to

assign the weights, denoted as w_{ig}, between SNP i *and* gene g, reflecting how likely the SNP i

affects gene g. For example, if a SNP is inside an exon of a gene, then the SNP-gene will have

weight 1. We note that w_{ig} and $P(Z_g = 1|\gamma_i = 1)$ have different semantics: it is possible that a

SNP affects multiple genes with weights all equal to 1, but there is only a single causal gene

supported by any SNP. In other words, for a causal SNP i, the conditional probabilities $P(Z_g =$

$1|\gamma_i = 1)$ should sum to 1 across all nearby genes g. So we need to normalize w_{ig} with:

771
$$P(\gamma_i = 1) = \frac{w_{ig}}{\sum_g w_{ig}}$$

To assign the weight terms, w_{ig}, we follow these four steps capturing several scenarios where a
SNP may affect a gene: **1)** If a SNP is in an exon or active promoter (promoter overlapping with
OCR) of a gene, we assign the SNP to that gene with weight $w_{ig} = 1$. **2)** If a SNP can be linked
to a gene's promoter via "enhancer loops", we assign the linked gene with weight $w_{ig} = 1$. Here,
"enhancer loops" are defined based on Activity-By-Contact (ABC) scores (constructed from heart
ventricle data with ABC scores >= 0.015)[39] and promoter-capture HiC data (from iPSC-CMs)[42].
Considering the fact that Hi-C and PC-HiC may miss contacts between close regions due to
technical reasons, we also consider a SNP in OCR within 20 kb of an active promoter as an
"enhancer loop". **3)** If a SNP is in a UTR but not in OCRs, suggesting that the SNP likely regulates
the containing gene through RNA processing mechanisms, e.g. RNA stability or alternative
polyadenylation, we will assign the SNP to the UTR-containing gene with weight $w_{ig} = 1$. **4)** If
a SNP is not linked to any gene via the criteria above, we use a distance-based weighting to assign
it to all genes within 1Mb. The weights follow an exponential decay function as below, where
d_{ig} is the SNP-gene distance:

$$w_{ig} = e^{-d_{ig}/5 \times 10^4} \ .$$

The parameter of this weight function, 50 kb, was chosen based on the fact that most enhancers,
estimated to be 84% using CRISPR deletion experiments[93], are located within 100 kb of the target
promoters. Using a weight of 50 kb here would lead to 87% of weights within 100 kb, with a
simple area-under-curve calculation of the weight function above.

At any locus, having PIPs for all the genes in the locus allows us to define the "credible gene
set" of the locus, much like the use of the term for SNPs[38]. Simply speaking, the credible set at the
80% level means the minimum set of genes in the locus whose sum of PIPs is greater than or equal
to 80%. One complication is that some of the genes in the locus may span another nearby locus,
as described above. In this case, while the final reported gene PIP is computed from both loci, we
only use the PIP of the gene from the locus of interest to define the credible gene set of that locus.
798
Benchmarking performance of different methods for risk gene identification. We compared
the accuracy of Mapgen (gene PIP >= 0.8), and several other commonly used methods that
nominate risk genes from GWAS (see below). Given that we do not have a gold standard list of
known AF genes, we used a set of Gene Ontology (GO) terms that have been associated with AF

genetics (using DEPICT method) from an earlier study (Nielsen *et al.*[5], Supplementary Table 7).

We used FDR < 5% and required three or more genes in a gene set, to select 173 GO terms. We

call a candidate gene "plausible", if the gene is annotated with any of those GO terms. Then we

compared the precision of the methods, calculated as the number of plausible genes divided by the

total number of nominated genes.

We included the following methods in the comparison: (1) Nearest gene to the top GWAS SNP

(based on distance to gene TSS). (2) eQTL, linking gene to the top GWAS SNP in each locus using

GTEx eQTL from the left ventricle (LV). (3) Activity-by-Contact (ABC) scores, linking promoters

with enhancers based on chromatin-looping data. Following Nasser *et al* [39], we used the ABC-

max approach, linking each top SNP to the gene with the maximum ABC score. (4) Multi-marker

Analysis of GenoMic Annotation (MAGMA), gene association test method[27]. We ran MAGMA

gene analysis and identified genes with Bonferroni adjusted p-value < 0.05. (5) In addition to the

above methods, we also included the nominated genes (gene score >= 11) from van Ouwerkerk *et*

al.[13].

Gene interaction network analysis. We used the STRING database (`STRING 11.5`)[94] to

construct gene network. The analysis was done using `Cytoscape 3.8.2`[95]. The input genes are

those at PIP >= 0.5 from our gene-mapping analysis. To create the gene network (Fig. 5h), we use

all default settings except that we use the recommended threshold for high-confidence interactions

(0.700) for interaction scores. Singletons, i.e., genes not having any interactions with other ones,

were not shown from the output network. We also used STRING to run functional enrichment

analysis based on sources including Gene Ontology[96,97], Reactome Pathways[98] and KEGG[99].

eQTL tissue sharing analysis. We started with the rationale of our eQTL tissue sharing analysis.

For simplicity, consider eQTLs found in one tissue (heart in our case), and we study the sharing

of these eQTLs in a second tissue. Let p denote the probability of eQTLs in the first tissue being

shared in the second tissue. Assuming we have several functional categories of eQTLs, e.g.

regulatory elements specific in a cell type, or shared across cell types, we can then break down p

into several categories with the simple relation:

833
$$p = \sum_c p_c w_c ,$$

where c denotes a category, p_c is the probability of tissue sharing in eQTLs from category c, and

w_c is the proportion of eQTLs in category c. We hypothesize that different eQTLs categories have

distinct molecular mechanisms of modulating transcript levels, and thus different tissue sharing

patterns. This simple analysis thus suggests that both w_c and p_c are important for our understanding

of tissue sharing. For instance, some categories may have a highly tissue-specific pattern (low p_c),

but may constitute a small proportion of all eQTLs (low w_c), thus these categories would have

limited contribution to the overall level of tissue sharing among eQTLs.

Summary statistics of GTEx heart eQTLs. Summary statistics of eQTLs from the left ventricle

were obtained from the GTEX v8 release[100]. We also obtained fine-mapping results using DAP-

G[21]. The variants with posterior inclusion probability (PIP) greater than 0.8 were kept for

downstream analyses. We refer to these putative causal variants as eQTLs henceforth. The total

number of eQTL-gene pairs that passed the threshold is 1,216. Tissue sharing data on the same

eQTLs were also obtained from GTEx[100]. These data provide information of whether these heart

eQTLs are also associated with gene expression in the other tissues in GTEx.

Defining functional categories of heart eQTLs. eQTLs were intersected with genomic features. To

obtain a set of disjoint genomic features, we used a combination of the union peak set and generic

annotations. For generic annotations, the longest transcript was chosen for each gene body, and its

corresponding exons, UTRs, and introns were obtained for all protein coding genes. We partitioned

the union peak set into cell-type-specific categories based on the differential accessibility (DA)

analysis, as well as the shared categories defined using the quantile approach, as described earlier.

We note that DA analysis does not guarantee disjoint sets of features. Indeed, we find that cell

types such as lymphoid and myeloid share about 6% of their DA peaks, while CMs share at most

1% with the other cell-types. To make these cell-type DA sets disjoint, we moved any DA peaks

that occurred in multiple cell types from DA analysis, to the "Shared 2-3" and "Shared 4+"

categories (see "Defining and classifying OCRs") depending on the number of cell types in which

it occurred. A small percentage of peaks ($< 1\%$) were affected by this step. The eQTLs in OCRs

that overlap with exons or UTRs, or eQTLs in non-DA OCRs, are ambiguous to assign, so they

were filtered from our analysis. The eQTLs in intronic OCRs were assigned based on the OCR

categories. Those eQTLs not intersecting with any functional category were designated in an

"unassigned" category.

Estimating extent of tissue sharing in different categories of heart eQTLs. GTEx has performed

eQTL mapping jointly across all tissues. Using these results, we call a SNP an eQTL in a given

tissue, if it passes the local false sign rate (LFSR) threshold of 1%. For any eQTL, we can thus

determine the number of tissues where it is active.

Estimating eQTL enrichment in functional categories. All the fine-mapped heart eQTLs are

assigned to our set of categories. The proportion of eQTLs in each category is then compared with

the expected proportion by chance to obtain enrichment reported in Fig. 6d and 6e. We used

SNPsnap[101] to create a set of random control SNPs that match our eQTLs in LD and minor allele

frequency. The LD data is obtained from the European population genotypes from 1000 Genomes.

We generated 1000 random SNPs which is roughly how many high-confidence eQTLs were used.

The proportion of random SNPs in each category is then used as our estimated proportion by

chance.

880
881 **Data availability**
882 Our snRNA-seq and scATAC-seq data has been deposited in GEO repository: accession
883 GSE224997.

884
885 **Code availability**
886 Mapgen R package is available from https://github.com/xinhe-lab/Mapgen. Code for data
887 processing and analyses are available at https://github.com/xinhe-lab/heart_atlas

888
889
890

Acknowledgements:
This work was funded by National Institutes of Health (NIH) grants, R01MH110531 and
R01HG010773 (to X.H.), R01HL163523 (to X.H., S.P., and I.P.M.), and R21 AI144417-02 (to
O.B). This project has been made possible in part by grant number CZF2019-002431 from the
Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation.
This work was completed in part with resources provided by the University of Chicago Research
Computing Center. We thank Xuanyao Liu for helpful comments on the manuscript.

899
900 **Main figures:**
901
902

903
Fig. 1 | Overview of our experimental and computational framework. Left: SnRNA-seq and
scATAC-seq profiling to cluster cells and obtain open chromatin regions (OCRs) in each cell type.
Middle: Using OCRs and GWAS summary statistics to assess variant enrichment in cell-type-
resolved OCRs. The enrichment results then provide prior for Bayesian statistical fine-mapping.
The resulting Posterior Inclusion Probabilities (PIPs) represent the probabilities of variants being
causal. The likely cell types through which the causal signals at each locus act can be identified by
considering cell type information of likely causal variants. We may not always be able to identify
a single cell type per locus, so we assign probabilities to cell types. **Right:** Computational gene-
mapping using PIPs from SNP fine-mapping and SNP-to-gene links to obtain gene level PIPs.
Note that the PIP of a SNP is partitioned into nearby genes in a weighted fashion, with more likely
target genes receiving higher weights (as indicated by thicker arrows). Prioritized genes can be
further assessed through external evidence such as gene networks and expression.
916
917

918
Fig. 2 | Mapping cell types in the human heart. a, UMAP projection of individual cells from
snRNA-seq and scATAC-seq colored by cell types. Stacked barplots on the right represent the
proportions of cell-types from each of the three donors. **b,** UCSC genome browser track plots of
chromatin accessibility at selected marker genes across cell types. The bottom part shows the gene
track (RefSeq annotation). Shown are two marker genes, TNNT2: cardiomyocyte marker; DCN:
fibroblast marker. **c,** Stacked violin plots of marker gene expression (log-normalized expression
values) in each cell type.
926

927
Fig. 3 | Discovery of OCRs and transcriptional regulators in the human heart. a, Row-
normalized accessibility of OCRs across all cell types. **b,** Number of cell-type-specific and shared
OCRs and their genomic distributions. **c,** Density plot of the $\log_{10}$ distance to nearest gene for all
cell-type-specific and shared OCRs. Colors of the lines for cell-type-specific OCRs follow the
same convention as in Figure 2a. Gray and black lines represent shared 2-3 and shared 4 OCRs. **d,**
Proportions of cell-type specific OCRs that overlap with DHS (upper panel). Bar graph (lower
panel) shows the proportions of cell-type specific OCRs that overlap with H3K27ac regions (LV
= left ventricle, RV = right ventricle). Smooth muscle cells and neuronal cells are not shown due
to the small numbers of peaks in these cell types. **e,** Enrichment of TF motifs in the OCRs specific
to each cell type. Shown are 76 TFs with FDR < 1% from motif enrichment analysis in at least one
cell-type, and correlation between motif enrichment and gene activity > 0.5. **f,** Gene scores (from
ArchR) and motif accessibility scores calculated with chromVar in OCRs for MEF2A (top) and
TBX5 (bottom) across all cells. Abbreviations for cell-types: CM = Cardiomyocyte, S.M = Smooth
Muscle, Peri = Pericyte, Endo = Endothelial, Fibro = Fibroblast, Neuro = Neuronal, Mye =
Myeloid, Lymph = Lymphoid.
943

944
Fig. 4 | Statistical fine-mapping of loci associated with the AF risk. a, log₂ fold enrichment
(from the tool TORUS) of risk variants of various traits in cell-type-specific OCRs. **b,**
Comparison of AF fine-mapping results under the informative prior using OCRs (Y-axis) vs. the
results under the uniform prior (X-axis). Each dot is a SNP, and color represents the annotation
of SNPs. Dashed line has a slope of 1. **c,** Summary of PIPs of variants. **d,** Summary of credible
set sizes from fine-mapping of AF. **e,** Trackplot at the HCN4 locus and the finemapped variant
rs7172038 (PIP = 0.99). The top two tracks represent the -log₁₀ p-value of SNPs from AF GWAS
(with color representing LD with the lead SNP) and their PIPs from SNP-level fine-mapping.
Middle three tracks represent cell-type aggregated ATAC-seq signals (CM: red, endothelial:
green; fibroblast: purple), followed by heart H3K27ac and fetal DHS peak calls. The bottom
track represents ABC scores from the heart ventricle. Abbreviations for cell-types: CM =
Cardiomyocyte, Endo = Endothelial, Fibro = Fibroblast. **f,** Proportions of summed PIPs in
disjoint functional annotation categories among all the loci. **g,** Proportion of summed PIPs in
disjoint functional annotation categories at each individual locus. **h,** Proportion of summed PIPs
into cell type-specific OCRs at each individual locus, for loci with summed PIPs in OCR >=
0.25. Highlighted are two loci with high proportions in non-CM cells: fibroblast, lymphoid
specific OCRs. **i,** Chromatin accessibility and additional functional genomic annotations of all
SNPs with PIP >= 50%. **j,** Reporter activities in cardiac cells (HL-1) of regions containing
selected SNPs, with both reference and alternative alleles. Data are from 5 replicates for each
construct. P-values were calculated using a paired two-sided t-test.

965
Fig. 5 | Mapping putative risk genes of AF. a. Schematic demonstrating the calculation of gene-
level PIPs. g_1 and g_2 represent genes, and s_1, s_2 represent SNPs. Vertical bars show the PIPs of the
SNPs from fine-mapping. Dashed arrows show the linking of SNPs to genes. $w_{s,g}$ represents the
weight of a gene (g) with respect to a SNP (S). The PIP of a gene is the weighted sum of PIPs of
SNPs that are linked to that gene. The weight of a SNP-gene pair is set according to the locations
of the SNP relative to the gene. See Methods for details. **b,** Manhattan plot of gene PIPs. Genes at
PIP >= 0.8 at labeled. **c,** Summary of the sizes of 80% credible gene sets from gene mapping. **d,**
GJA5 locus: similar to Fig. 4e, except that the links here represent interactions identified from
promoter-capture HiC data in iPSC-derived CMs. Vertical bars show the locations of the four fine-
mapped SNPs supporting GJA5 as the risk gene in this locus. The red links in the PC-HiC track
show interactions linked to these four fine-mapped SNPs. **e,** The performance of Mapgen and other
methods in nominating AF risk genes. Precision is measured as the proportion of likely risk genes,
those annotated with AF-related GO terms, in the set of nominated genes by each method. See
Methods for the details of each method. **f,** CALU locus, tracks shown are similar to the panel **d.**
The vertical bar shows the location of the top fine-mapped SNP. Abbreviations for cell-types: CM
= Cardiomyocyte, Endo = Endothelial, Fibro = Fibroblast.
982

983
984

985
986

Fig. 6 | Functional support of putative AF risk genes. a, Log-normalized CM expression of genes at PIP >= 80% vs. other genes from the AF loci. **b,** Percentage of Mendelian disease genes from OMIM in each gene PIP bin. **c,** Top 5 Biological Processes (BP) and Molecular Functions (MF) GO terms from gene-set enrichment analysis of the 46 genes with PIP >= 80%. **d,** Gene interaction network of candidate AF genes (PIP >= 0.5) using STRING. Only genes with interactions are shown. Interactions are defined using a confidence threshold of 0.7 by STRING. Node sizes represent gene PIPs. Colors of genes indicate their shared molecular functions.

994
995

Fig. 7 | Tissue-sharing patterns of heart (LV) eQTLs from GTEx. a, Number of tissues where LV-eQTLs are detected at local false sign rate (LFSR) < 1%. **b,** Violin plot showing the number of tissues in which a specific eQTL is detected. Each row represents a different class of eQTLs, assigned based on their overlap with OCRs categories and other genomic locations. Unassigned: eQTLs that cannot be assigned to any functional class. **c,** Proportion of LV-eQTLs located in OCRs of selected cell types (Cardiomyocytes, Endothelial cells, Fibroblast, and Immune cells) that were also detected as eQTLs in a second tissue. **d,** Proportion of LV-eQTLs (n = 1216) in each functional class. For comparison, the proportions of random SNPs in all the classes are also shown. The numbers near the bars represent the fold enrichment in heart eQTLs compared to random SNPs. **e,** Enrichment of GTEx heart eQTLs in OCRs vary with the number of cell types where the OCRs are active. Lower panel shows the proportion of eQTLs (light blue) and control SNPs (dark blue, chosen to match eQTLs in LD and MAF) overlapping OCRs. The OCRs are divided into 4 categories, based on the degree of sharing across cell types in heart: 1= not shared, 4+= shared in >=4 cell types. The upper panel shows the enrichment of eQTLs in each OCR class compared to expectation based on control SNPs. Abbreviations for cell-types: CM = Cardiomyocyte, S.M = Smooth Muscle, Peri = Pericyte, Endo = Endothelial, Fibro = Fibroblast, Neuro = Neuronal, Mye = Myeloid, Lymph = Lymphoid.

1015
1016 **Table 1 | Top prioritized genes (gene PIP >= 0.95).**
1017
1018

Gene	Gene PIP	Supporting SNPs	SNP PIP	Link Method*	OMIM	CM-specific expression	Known AF risk gene	Reference
SYNPO2L	1.113	rs60632610	0.971	exon			✓	[20215401, 33768119]
HCN4	1.095	rs7172038	0.989	ABC	✓		✓	[29987112]
ASAH1	1.061	rs7508	1	exon		✓		[32015399]
ATXN1	1.000	rs59430691	0.809	PC-HiC				[21475249, 22306179]
ERBB4	1.000	rs6738011	0.12	distance		✓		[19632177]
KCNN2	1.000	rs337705 rs337708	0.528 0.113	distance distance			✓	[19139040]
RPL3L	1.000	rs140185678	1	exon				[32870709, 32514796]
TUBA8	1.000	rs464901 rs361834	0.886 0.114	nearby OCR nearby OCR				[31398994]
EPHA3	0.999	rs35124509 rs6771054 rs2117137	0.345 0.1720.117	exons distance distance				[17046737]
THRB	0.999	rs73041705 rs73032363 rs9841040 rs1865712	0.177 0.139 0.130 0.119	distance distance distance distance		✓		[28740583]
ETV1	0.998	rs55734480 rs12154315 rs12112152	0.403 0.338 0.218	distance distance distance			✓	[27775552, 29930145]
BEND5	0.997	rs11590635	0.973	distance				
PITX2	0.997	rs1906615 rs7689774	0.798 0.15	distance distance			✓	[28217939, 29367545, 32309338]
TBX5	0.997	rs7312625 rs883079 rs7955405	0.511 0.194 0.126	distance exon PC-HiC		✓	✓	[28057264]

Gene	PIP	Supporting SNPs	frac PIP	Connection				Reference
PKP2	0.992	rs12809354 rs2045172	0.771 0.211	PC-HiC PC-HiC		✓		[28740174]
CAMK2D	0.991	rs11938486 rs17446418	0.649 0.17	distance PC-HiC		✓	✓	[24030498]
NKX2-5	0.989	rs6882776 rs6891790 rs2277923 rs10071514	0.37 0.318 0.157 0.133	active promoter ABC/nearby OCR exon ABC/nearby OCR	✓			[26805889]
GYPC	0.985	rs28387148 rs28387129 rs28387153 rs28387149	0.191 0.153 0.144 0.103	distance nearby OCR distance distance				
LRIG1	0.982	rs34080181 rs900171 rs12633819	0.329 0.119 0.106	distance exon distance				[23558895, 19632177]
ZEB2	0.979	rs10496971	0.896	distance				[33398012]
NR3C1	0.975	rs72804738	0.208	distance				[23595884]
MYO18B	0.974	rs133902 rs133885	0.685 0.185	distance exon		✓		[27858739]
DPF3	0.972	rs3814866 rs3814864	0.744 0.176	active promoter active promoter		✓	✓	[26582913, 30240284]
WIPF1	0.971	rs35215597	0.874	Nearby OCR				

1019
1020
* In the Supporting SNPs column, only SNPs that contribute a fractional PIP (SNP PIP multiplied by the weight of
the SNP to that gene) of 0.1 or more are shown. *EFNA5* (gene PIP =1) is not included because it does not have any
SNPs with fractional PIP >= 0.1.
* Nearby OCR is defined as OCR within 20 kb of active promoter of the gene.
* Reference column shows the PMIDs of the relevant papers supporting the connections of the genes to AF or heart
physiology.
1027

References

1029

1030 1. Benjamin, E. J. *et al.* Independent Risk Factors for Atrial Fibrillation in a Population-Based

Cohort: The Framingham Heart Study. *JAMA* **271**, 840–844 (1994).

2. Kornej, J., Börschel, C. S., Benjamin, E. J. & Schnabel, R. B. Epidemiology of Atrial Fibrillation in the 21st Century. *Circ. Res.* **127**, 4–20 (2020).

3. Roselli, C. et al. Multi-ethnic genome-wide association study for atrial fibrillation. *Nat. Genet.* **50**, 1225–1233 (2018).

4. Roselli, C., Rienstra, M. & Ellinor, P. T. Genetics of Atrial Fibrillation in 2020. Circ. Res. 127, 21–33 (2020).

5. Nielsen, J. B. et al. Biobank-driven genomic discovery yields new insight into atrial fibrillation biology. Nat. Genet. 50, 1234–1239 (2018).

6. Arking, D. E. et al. Genetic association study of QT interval highlights role for calcium signaling pathways in myocardial repolarization. Nat. Genet. **46**, 826–836 (2014).

7. Pfeufer, A. et al. Genome-wide association study of PR interval. Nat. Genet. **42**, 153–159 (2010).

8. Maurano, M. T. et al. Systematic Localization of Common Disease-Associated Variation in Regulatory DNA. *Science* **337**, 1190–1195 (2012).

9. Pickrell, J. K. Joint Analysis of Functional Genomic Data and Genome-wide Association Studies of 18 Human Traits. *Am. J. Hum. Genet.* **94**, 559–573 (2014).

10. Finucane, H. K. et al. Partitioning heritability by functional annotation using genome-wide association summary statistics. Nat. Genet. **47**, 1228–1235 (2015).

11. Kundaje, A. et al. Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–330 (2015).

12. Consortium, E. P. et al. Expanded encyclopaedias of DNA elements in the human and mouse genomes. Nature **583**, 699–710 (2020).

13. Ouwerkerk, A. F. van et al. Identification of atrial fibrillation associated genes and functional non-coding variants. *Nat. Commun.* **10**, 4755 (2019).

14. Ouwerkerk, A. F. van et al. Identification of Functional Variant Enhancers Associated With Atrial Fibrillation. Circ. Res. 127, 229–243 (2020).

15. Hocker, J. D. et al. Cardiac cell type–specific gene regulatory programs and disease risk association. *Sci. Adv.* **7**, eabf1444 (2021).

16. Habib, N. et al. Massively-parallel single nucleus RNA-seq with DroNc-seq. *Nat. Methods* **14**, 955–958 (2017).

17. Litviňuková, M. et al. *Cell*s of the adult human heart. Nature **588**, 466–472 (2020).

18. Tucker, N. R. et al. Transcriptional and Cellular Diversity of the Human Heart. *Circulation* **142**, 466–482 (2020).

19. Buenrostro, J. D. et al. Single-cell chromatin accessibility reveals principles of regulatory variation. Nature **523**, 486–490 (2015).

20. Cusanovich, D. A. et al. Multiplex single cell profiling of chromatin accessibility by combinatorial cellular indexing. Science **348**, 910–914 (2015).

21. Wen, X., Lee, Y., Luca, F. & Pique-Regi, R. Efficient Integrative Multi-SNP Association Analysis via Deterministic Approximation of Posteriors. Am. J. Hum. Genet. **98**, 1114–1129

(2016).

22. Weissbrod, O. et al. Functionally informed fine-mapping and polygenic localization of
complex trait heritability. Nat. Genet. 52, 1355–1363 (2020).

23. Mahajan, A. et al. Fine-mapping type 2 diabetes loci to single-variant resolution using high-
density imputation and islet-specific epigenome maps. Nat. Genet. 50, 1505–1513 (2018).

24. Trubetskoy, V. et al. Mapping genomic loci implicates genes and synaptic biology in
schizophrenia. Nature **604**, 502–508 (2022).

25. Huang, H. et al. Fine-mapping inflammatory bowel disease loci to single-variant resolution.
Nature **547**, **173**–178 (2017).

26. Segrè, A. V. et al. Common Inherited Variation in Mitochondrial Genes Is Not Enriched for
Associations with Type 2 Diabetes or Related Glycemic Traits. *Plos Genet* **6**, e1001058 (2010).

27. Leeuw, C. A. de, Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: Generalized Gene-Set
Analysis of GWAS Data. *Plos Comput Biol* **11**, e1004219 (2015).

28. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat. Biotechnol.* **36**,
411–420 (2018).

29. Granja, J. M. et al. ArchR is a scalable software package for integrative single-cell chromatin
accessibility analysis. Nat. Genet. 53, 403–411 (2021).

30. Thurman, R. E. et al. The accessible chromatin landscape of the human genome. Nature **489**,
75–82 (2012).

31. Buenrostro, J. D. et al. Integrated Single-Cell Analysis Maps the Continuous Regulatory
Landscape of Human Hematopoietic Differentiation. Cell 173, 1535-1548.e16 (2018).

32. Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat Methods* **14**,
975–978 (2017).

33. Olson, E. N. Gene regulatory networks in the evolution and development of the heart.
Science **313**, 1922–7 (2006).

34. Wen, X. Molecular QTL discovery incorporating genomic annotations using Bayesian false
discovery rate control. *Ann. Appl. Statistics* **10**, **16**19–1638 (2016).

35. Cheng, S. et al. Long-term Outcomes in Individuals With Prolonged PR Interval or First-
Degree Atrioventricular Block. JAMA **301**, 2571–2**577** (2009).

36. Yu, F. et al. Variant to function mapping at single-cell resolution through network
propagation. *Nat Biotechnol* 1–10 (2022) doi:10.1038/s41**58**7-022-01341-y.

37. Berisa, T. & Pickrell, J. K. Approximately independent linkage disequilibrium blocks in
human populations. *Bioinformatics* **32**, 283–285 (2016).

38. Wang, G., Sarkar, A., Carbonetto, P. & Stephens, M. A simple new approach to variable
selection in regression, with application to genetic fine mapping. *J. R. Stat. Soc. Ser. B.*
Statistical Methodol. **82**, 1273–1300 (2020).

39. Nasser, J. et al. Genome-wide enhancer maps link risk variants to disease genes. Nature **593**,
238–243 (2021).

40. Ouwerkerk, A. F. van et al. Epigenetic and Transcriptional Networks Underlying Atrial Fibrillation. Circ. Res. 127, 34–50 (2020).

41. Sobreira, D. R. et al. Extensive pleiotropism and allelic heterogeneity mediate metabolic effects of IRX3 and IRX5. Science **372**, 1085–1091 (2021).

42. Montefiori, L. E. et al. A promoter interaction map for cardiovascular disease genetics. *Elife* **7**, e35788 (2018).

43. Kapoor, A. et al. Multiple SCN5A variant enhancers modulate its cardiac gene expression and the QT interval. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 201808734 (2019).

44. Claycomb, W. C. et al. HL-1 cells: A cardiac muscle cell line that contracts and retains phenotypic characteristics of the adult cardiomyocyte. Proc. Natl. Acad. Sci. U.S.A. **95**, 2979–2984 (1998).

45. DiFrancesco, D. HCN4, Sinus Bradycardia and Atrial Fibrillation. *Arrhythmia Electrophysiol Rev.* **4**, 9 (2015).

46. Giambartolomei, C. et al. Bayesian Test for Colocalisation between Pairs of Genetic Association Studies Using Summary Statistics. *PLoS Genet.* **10**, e1004383 (2014).

47. Claussnitzer, M. et al. A brief history of human disease genetics. Nature 577, 179–189 (2020).

48. Nadadur, R. D. et al. Pitx2 modulates a Tbx5 -dependent gene regulatory network to maintain atrial rhythm. *Sci Transl Med* **8**, 354ra115 (2016).

49. Drabkin, M. et al. Nocturnal Atrial Fibrillation Caused by Mutation in KCND2 , Encoding Pore-Forming (α) Subunit of the Cardiac Kv4.2 Potassium Channel. *Circulation Genom Precis Medicine* **11**, e002293 (2018).

50. Mahida, S. et al. Overexpression of KCNN3 results in sudden cardiac death. *Cardiovasc Res* **101**, 326–334 (2013).

51. H., G. M. et al. Somatic Mutations in the Connexin 40 Gene (GJA5) in Atrial Fibrillation. *New Engl J Med* **354**, 2677–2688 (2006).

52. Chung, I.-M. & Rajakumar, G. Genetics of Congenital Heart Defects: The NKX2-5 Gene, a Key Player. *Genes (Basel).* **7**, 6 (2016).

53. Sahoo, S. K. & Kim, D. H. Characterization of calumenin in mouse heart. *BMB Rep.* **43**, 158–63 (2010).

54. Lee, J. H., Kwon, E. J. & Kim, D. H. Calumenin has a role in the alleviation of ER stress in neonatal rat cardiomyocytes. *Biochem Bioph Res Co* **439**, 327–332 (2013).

55. Sirish, P., Diloretto, D. A., Thai, P. N. & Chiamvimonvat, N. The Critical Roles of Proteostasis and Endoplasmic Reticulum Stress in Atrial Fibrillation. *Front Physiol* **12**, 793171 (2022).

56. Ahlberg, G. et al. Rare truncating variants in the sarcomeric protein titin associate with familial and early-onset atrial fibrillation. Nat. Commun. 9, **4**316 (2018).

57. Choi, S. H. et al. Monogenic and Polygenic Contributions to Atrial Fibrillation Risk. *Circ Res* **126**, 200–209 (2020).

58. Szklarczyk, D. et al. STRING v11: protein–protein association networks with increased

coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* **47**, gky1131 (2018).

59. Benaglio, P. et al. Allele-specific NKX2-5 binding underlies multiple genetic associations with human electrocardiographic traits. Nat. Genet. 51, 1506–1517 (2019).

60. Schindler, Y. L. et al. Hand2 elevates cardiomyocyte production during zebrafish heart development and regeneration. *Development* **141**, 3112–3122 (2014).

61. Cohen, A. S. A. et al. Haploinsufficiency of the basic helix–loop–helix transcription factor HAND2 causes congenital heart defects. *Am. J. Med. Genet.* **182**, 1263–1267 (2020).

62. Romero-Becerra, R., Santamans, A. M., Folgueira, C. & Sabio, G. p38 MAPK Pathway in the Heart: New Insights in Health and Disease. *Int. J. Mol. Sci.* **21**, 7412 (2020).

63. O'Neal, W. T. et al. Ephrin–Eph signaling as a potential therapeutic target for the treatment of myocardial infarction. *Med. Hypotheses* **80**, 738–744 (2013).

64. Su, S. et al. Essential roles of EphrinB2 in mammalian heart: from development to diseases. *Cell Commun. Signal.* **17**, 29 (2019).

65. Chen, K. et al. EphB4 Forward-Signaling Regulates Cardiac Progenitor Development in Mouse ES Cells. *J Cell Biochem* **116**, 467–475 (2015).

66. Chun, S. et al. Limited statistical evidence for shared genetic effects of eQTLs and autoimmune disease-associated loci in three major immune cell types. *Nat Genet* **49**, 600–605 (2017).

67. Yao, D. W., O'Connor, L. J., Price, A. L. & Gusev, A. Quantifying genetic effects on disease mediated by assayed gene expression levels. Nat. Genet. 52, 626–633 (2020).

68. Umans, B. D., Battle, A. & Gilad, Y. Where Are the Disease-Associated eQTLs? *Trends Genet* **37**, 109–124 (2021).

69. Chiou, J. et al. Single-cell chromatin accessibility identifies pancreatic islet cell type– and state-specific regulatory programs of diabetes risk. Nat Genet **53**, 455–466 (2021).

70. Lozano-Velasco, E., Franco, D., Aranega, A. & Daimi, H. Genetics and Epigenetics of Atrial Fibrillation. *Int J Mol Sci* **21**, 5717 (2020).

71. Wang, J., Gareri, C. & Rockman, H. A. G-Protein–Coupled Receptors in Heart Disease. Circ. Res. 123, 716–735 (2018).

72. Foulquier, S. et al. WNT Signaling in Cardiac and Vascular Disease. *Pharmacol Rev* **70**, 68–141 (2018).

73. Huang, J. et al. Fibroblast growth factor 9 (FGF9) inhibits myogenic differentiation of C2C12 and human muscle cells. *Cell Cycle* **18**, 1–19 (2019).

74. Itoh, N. & Ohta, H. Pathophysiological roles of FGF signaling in the heart. Front Physiol 4, 247 (2013).

75. Nattel, S. Molecular and Cellular Mechanisms of Atrial Fibrosis in Atrial Fibrillation. *JACC Clin. Electrophysiol.* **3**, 425–435 (2017).

76. Chromium Single Cell 3' Reagent Kits User Guide (v3.1 Chemistry). https://support.10xgenomics.com/single-cell-gene-expression/library-prep/doc/user-guide-chromium-single-cell-3-reagent-kits-user-guide-v31-chemistry (2019).

77. Selewa, A. et al. Systematic Comparison of High-throughput Single-Cell and Single-Nucleus
Transcriptomes during Cardiomyocyte Differentiation. *Sci. Rep.* **10**, 1535 (2020).
78. FastQC. https://qubeshub.org/resources/fastqc (2015).
79. Andrews, S. FastQC: A Quality Control Tool for High Throughput Sequence Data [Online].
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (2010).
80. Kaminow, B., Yunusov, D. & Dobin, A. STARsolo: accurate, fast and versatile
mapping/quantification of single-cell and single-nucleus RNA-seq data. *Biorxiv*
2021.05.05.442755 (2021) doi:10.1101/2021.05.05.442755.
81. Frankish, A. et al. GENCODE reference annotation for the human and mouse genomes.
Nucleic Acids Res 47, gky955- (2018).
82. Stuart, T. et al. Comprehensive Integration of Single-Cell Data. Cell **177**, 1888-1902.e21
(2019).
83. McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in
Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-337.e4
(2019).
84. Satpathy, A. T. et al. Massively parallel single-cell chromatin landscapes of human immune
cell development and intratumoral T cell exhaustion. Nat. Biotechnol. 37, 925–936 (2019).
85. Korsunsky, I. et al. Fast, sensitive and accurate integration of single-cell data with Harmony.
Nat Methods 16, 1**28**9–1296 (2019).
86. Zhang, Y. et al. Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137–R137
(2008).
87. Lawrence, M. et al. Software for computing and annotating genomic ranges. *PLoS Comput.*
Biol. **9**, e1003118 (2013).
88. Weirauch, M. T. et al. Determination and Inference of Eukaryotic Transcription Factor
Sequence Specificity. Cell 158, 1431–1443 (2014).
89. Pliner, H. A. et al. Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell
Chromatin Accessibility Data. *Mol. Cell.* **71**, 858-871.e8 (2018).
90. Lindblad-Toh, K. et al. A high-resolution map of human evolutionary constraint using 29
mammals. Nature **478**, 476–82 (2011).
91. Clarke, L. et al. The international Genome sample resource (IGSR): A worldwide collection
of genome variation incorporating the 1000 Genomes Project data. Nucleic Acids Res 45, D854–
D859 (2017).
92. Coetzee, S. G., Coetzee, G. A. & Hazelett, D. J. motifbreakR: an R/Bioconductor package for
predicting variant effects at transcription factor binding sites. Bioinformatics 31, 3847–3849
(2015).
93. Fulco, C. P. et al. Activity-by-contact model of enhancer–promoter regulation from
thousands of CRISPR perturbations. Nat Genet **51**, 1664–1669 (2019).
94. Szklarczyk, D. et al. The STRING database in 2021: customizable protein–protein networks,
and functional characterization of user-uploaded gene/measurement sets. Nucleic Acids Res. 49,
gkaa1074- (2020).

95. Shannon, P. et al. Cytoscape: A Software Environment for Integrated Models of
Biomolecular Interaction Networks. *Genome Res.* **13**, 2498–2504 (2003).
96. Ashburner, M. et al. Gene Ontology: tool for the unification of biology. Nat. Genet. 25, 25–
29 (2000).
97. Consortium, T. G. O. et al. The Gene Ontology resource: enriching a GOld mine. Nucleic
Acids Res 49, D325–D334 (2020).
98. Jassal, B. et al. The reactome pathway knowledgebase. Nucleic Acids Res **48**, D498–D503
(2019).
99. Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids
Res 28, 27–30 (2000).
100. Consortium, T. Gte. The GTEx Consortium atlas of genetic regulatory effects across human
tissues. Science **369**, 1318–1330 (2020).
101. Pers, T. H., Timshel, P. & Hirschhorn, J. N. SNPsnap: a Web-based tool for identification
and annotation of matched SNPs. Bioinformatics 31, 418–420 (2015).