Early guidance for Sars-Cov-2 health policies in India: Social Distancing amidst Vaccination and Virus Variants. *

Yi Zhang and Sanjiv Kapoor [†]
Computational Decision Science Laboratory
Illinois Institute of Technology, Chicago, USA.

Abstract

Policy decisions during the SARS-COV-2 pandemic were complicated due to virus variants and the impacts of societal restrictions. Accurate predictive models were required in this context. In this paper we report results from a model that helped in predicting the impact of SARS-CoV-2 virus transmission in India over a period of a number of months from June, 2021 to March 2022. These models were applied in the context of enabling policy decisions to tackle the impact of the pandemic in India culminating in early warning projections for the Omicron variant and used for advise on preemptive policy actions.

Methods: Our model utilizes a deterministic compartment models incorporating a dynamic transmission factor, dependent on the population's behavior as a function of the reported confirmed cases of virus transmission as well as methods for estimation of the increase in susceptible population when social distancing mandates are relaxed. The model used to predict viral growth incorporates the state of vaccination and the virus variants that form part of the transmission dynamics as well the lockdown state of the population. NPI actions were used in India to contain the spread of infections during the period of study, especially during the surge of the Omicron variant of the virus. Further we present the impact of lockdown policies and illustrate the advantage of adopting partial lockdown policies in the early period of 2022. Based on the models, our predictive analysis, when applied to the Omicron variant, illustrated substantial improvement even when partial lockdown is planned.

Findings: This report presents models and results that incorporated the impact of vaccination rates and the Omicron variant and were used to establish projections on the growth of Sars-Cov2 infections in India for the period from July 2021 till March 2022. The growth rate of the Omicron virus was deduced from data that originated from South Africa in November 2021. These projections were submitted to a pivotal government organization involved in developing a national public health strategy to address the pandemic and, as per personal communication, were considered when formulating national policy. The pandemic had a subdued impact in India during the period from July 2021 till date as evident from the deaths reported by the government. The projections were made every month and cases were projected over the next 4-16 weeks. The projections of cumulative cases during the Omicron wave had low errors when measured using RMSE per capita and had a MAPE error of 17.8% when measured 15 days after start of the projection on December 5th, 2021.

Discussion: The composed model was found to be useful in providing predictive and data based analytic input to inform early warning approaches in the context of policy based interventions to

---

[*]Research sponsored in part by NSF. grant No. 2028274.

[†]Corresponding author, kapoor@iit.edu. Part of this work was performed when S. Kapoor was a visiting Professor at IIT-Delhi.

control the pandemic in India. The model provided monthly early prediction of the spread and impact of the SARS-COV-2 virus in India, state-wise, during the phase of removal of government lockdown in the second half of 2021. The early warning system incorporated the impact of the Omicron variant to provide predictions for Indian states and the country.

1 Introduction

The SARS-COV-2 virus has been plaguing the world since January 2020, compounding the original problem with multiple variants. Containment measures like social distancing are key, even in the presence of vaccinations, to preventing the spread and also to contain variants. Accuracy in models that predict the growth in infections are very important. While short term analysis of a few weeks is typically more accurate, long range models are important in planning. Our motivation was the impact of the virus in India during March to May 2021, in this paper we consider a model for virus infection spread that accounts for the impact of lockdown policies, vaccinations and the impact of vaccine-resistant virus variants.

Effective public health policy requires timely social distancing guidelines; as such it is critical to develop early-warning systems that would predict the rise of the virus, especially as lockdown or social distancing policies are relaxed. These warning systems would include parameters that reflect the relaxation of NPI practices related to masking, enhanced mobility etc. that would be indicative of behavior patterns that create an influx of susceptible into the population. Since the incubation period of the disease and the spread in asymptomatic population, especially the vaccinated population, results in delayed impact on the progression of the disease, long term modeling of the spread of the virus is critical.

This report deals with an analysis of the spread of infection as more population emerges from lockdown and adopts lax behavior with respect to social distancing. Impact of relaxed norms are apparent in the multiple waves that have afflicted countries around the world. We construct a model that reflects the vaccination status of the population, accounting for the time-dependent efficacy of the vaccines, and also incorporates virus variants. The analysis is based on epidemic spread models that utilize compartment models which have along history (see [6, 9] and [10, 11, 12]). A key differentiation in our work is the use of dynamic social distancing factor in the model, termed the SIR-SD model introduced in [25]. The role of reducing contacts to reduce transmission is well studied [24, 7, 13] as more recent modeling efforts have focused also on mobility [5], amongst other factors [15, 21, 14, 22, 3, 18, 19, 16, 23]. Our study incorporates virus variants. The compartment models we utilize includes compartments reflecting the various states of the population including vaccinated population, population under lockdown etc. As mentioned above, we also use a social distancing factor to modify the virus transmission rate that reflects the behavior of the population as the number of infection cases rises and reporting of the infections, as well as government mandates, becomes more prevalent. As introduced earlier in [25] the transmission of the virus incorporates a factor that is inversely dependent on the number of cumulative infections incorporated by a time-dependent susceptible population size.

This model had been utilized for multiple months, starting from June 2021, in order to provide predictive scenarios months in advance. We calibrated resurgence of the virus in our model using a release of population into the susceptible pool during the second wave of infections in India in April 2021. The wave had come as a surprise to policy makers and we designed a predictive long range model since it was important to ensure that the infection growth was contained as lockdown was removed and vaccination progressed. Our model incorporates a population that transfers from a lockdown state to a susceptible

state. To quantify the growth of the susceptible population in our model, we assessed the growth of susceptible population, H, that led to the second wave and based future projections on this baseline behavior estimate, using this growth to determine the rate at which the population entered the susceptible set, as a result of relaxed government policies and social distancing behavior. We report modeling two rates of relaxation of lockdown procedures after the second wave that direct the rate of policy changes after June 2021: a fast rate of release where the susceptible population H was released in 30 days; a moderate or medium rate of release where the susceptible population H was released in 45 days. We illustrate the succession of projections for the fast release rate (30 day release) in Figure 1. The projections range is bounded by two scenarios, an optimistic and a pessimistic scenario. The two scenarios differ in the estimates of the addition to the susceptible population resulting from the relaxation of the lockdown that was initiated in June 2021. Over the period we studied, projections showed a reduction in cases and then an increase during the winter months ending in a sharp rise in predicted cases when Omicron started and became prevalent. Our initial lower projections of the growth of infections during August, September and October could possibly be attributed to government actions including slow release from lockdown state and enhanced vaccination rates. Our projections in December, 2021 appear higher than the actual confirmed cases as likely the rise in cases was tempered by government policies, that included night curfews as well as business and school closings during January. The dynamic progression of infection case projections is illustrated in the accompanying video[2]. Detailed estimates provided in later sections illustrate the mixed impact of the two variants, with the Omicron variant rapidly dominating the currently existing Delta variant. In later sections in this paper we also consider the impact of lockdown on reduction in viral spread.

This work was motivated by the problem of determining policy decisions as vaccinations progressed and we used this model to provide monthly updates on projections to a pivotal government organization. The projections focused on the onset of the third wave and covered the period from June 2021 till December 2021. The third wave in India started in December 2021. As per personal communication these projections were considered when formulating national policy. The policies implemented at state levels included mask mandates, physical school closings, partial curfews and office closings, an example provided by Delhi state government policies [1]. The projections were accompanied by data on social distancing and mask usage from IHME. However we did not use these social distance metrics as part of our model as this data was typically delayed.

Accuracy of our model was established during the initial part of study, from April to May 2021. The changing governmental policies on non-pharmaceutical interventions, as and when rise in infection cases were evident, made accuracy analysis for the projections difficult. However, during the Omicron variant wave the confirmed cases of viral infection were substantial and an accuracy analysis of our projections versus the actual confirmed cases, showed that the cumulative case counts had very low RMSE per capita (0.0013 per capita, and had MAPE errors ranging from 17.8% to 41.6% for projections 15 to 45 days from December 15, 2021. As time progressed the deviation from our projections grew due to the NPI policies that were implemented.

While the cases incidences were kept limited till November, relaxed social distancing norms and waning vaccine efficacy likely had an impact on the cases incidences in December 2021, along with increased transmissibility of the Omicron variant.

Figure 1: Monthly sequence of projections for the case of fast release rate. The peak case load varied according to the current infection trends and successively delayed by almost a month.

2 SEIAR-SD-L Model

We use a mechanistic model as illustrated in Figure 2. Apart from standard compartments representing the Susceptible S, Exposed E, Infected (confirmed via testing) I, Undetected Symptomatic and Asymptomatic A, we include hospitalized I_H and home treated I_N, along with dead D and recovered R. There was no data on deaths due to undetected symptomatic cases and so we omit the corresponding transition.

Each of these compartments are replicated for various types of virus variants to model the impact of each type of virus variant. In the context of the Omicron virus in addition to the base variant, the Delta variant, these compartments are indexed by superscript O. Additionally we have a population that is on lockdown, grouped by different vaccination states, H_0: the set of unvaccinated population under lockdown; H_1: the population that has received the first shot of vaccination (based on the rate of vaccination of the population); H_2: population after two weeks of receiving the first dose(first dose effective two weeks after the dose) ; H_3: population after two weeks of receiving the second dose of vaccination given after a fixed delay from the first dose); H_4: vaccinated population after 4 months of the second dose. Vaccination efficacy has been known to drop by 50% after 4 months and H_4 models the population in this state [4]. Similarly the vaccination status of the susceptible population is modeled by S_0, S_1, S_2, S_3 and S_4.

A release from lockdown transfers population from the corresponding lockdown compartments to the susceptible population, while lockdown policies transfer susceptible back to the H compartments.

2.1 The Compartment Model

We let

$$\frac{dH_i}{dt} = v_i H_{i-1} - v_{i+1} H_i - \kappa^t \cdot \frac{H_i}{\hat{H}} \ , \ i = 0 \ldots 4;$$

$$\frac{dS_i}{dt} = v_i S_{i-1} - v_{i+1} S_i + \kappa^t \cdot \frac{H_i}{\hat{H}} - \sum_C \frac{\beta_i^C S_i \hat{I}^C}{N} \ , \ i = 0 \ldots 4;$$

where $\hat{H} = H_0 + H_1 + H_2 + H_3 + H_4$, $v_0 = 0, v_5 = 0$, and v_1, v_2, v_3, v_4 represent transition rates between the population groups as represented in Figure 2. And for all strains, C, the following conditions hold, where we omit the notation for the C for ease of understanding.

$$\frac{dE}{dt} = \frac{\overline{\beta S} \cdot \hat{I}}{N} - \gamma_E E; \quad \frac{dI}{dt} = (1 - \alpha)\gamma_E E - (\gamma + \gamma_2)I;$$

$$\frac{dA}{dt} = \alpha\gamma_E E - \gamma_3 A; \quad \frac{dI_H}{dt} = \gamma I - (a_1 + a_2)I_H;$$

$$\frac{dI_N}{dt} = \gamma_2 I - a_3 I_N; \quad \frac{dD}{dt} = a_2 I_H;$$

$$\frac{dR}{dt} = a_1 I_H + a_3 I_N;$$

where $\overline{\beta S} = \beta_0 S_0 + \beta_1 S_1 + \beta_2 S_2 + \beta_3 S_3 + \beta_4 S_4$ and $\hat{I} = I + A$. The model parameters, $\gamma_E, \gamma, \gamma_2, \alpha, a_1, a_2, a_3$ are all with respect to the strain C.

To model the impact of the virus on the population's behavior we utilize dynamic transmission factor β [25]. The value of β is based on the following generic transmission function that models the dynamics

Figure 2: The SEIAR-SD-L Compartment Model with vaccination and two strains of the virus; delta and omicron (red)

of population behavior (for further discussion see [25]):

$$\beta = \beta_0 \frac{1 - c_1}{1 - c_1 \cdot \frac{S}{S_0}}$$

where S is the number of the current susceptible population and S_0 the original set of susceptible population.

The function κ^t varies dependent on whether the policies are in a release phase or in a lockdown phase. We use a linear function in our model:

- Release Phase: In the release phase, between time t_1 and t_2, the function $\kappa^t = h^R, t_1 \leq t \leq t_2$ where h^R is a positive constant and where t_1 and t_2 is the range of time over which the population enters into a susceptible state. The value of h^R is determined by the release policies and adherence.

- Lockdown Phase: In the lockdown phase, between time t_3 and t_4, the function $\kappa^t = h^L, t_3 \leq t \leq t_4$ where h^L is a negative constant, represents the rate at which the population withdraws into a state of lockdown, over the range of time $[t_3, t_4]$. The value of h^L is determined by the strength of the lockdown and adherence.

The theoretical value of R_0 in this model is analyzed by the method provided in the supplementary material and is:

$$R_0 = \max_C \{ \sum_i \beta_i^C \eta_i (1 + \rho)(\frac{(1 - \alpha^C)}{\gamma^C + \gamma_2^C} + \frac{\alpha^C}{\gamma_3^C}) \}$$

where $\rho = \kappa^t / \sum_i S_i$ is the rate at which population enters the susceptible set S_i, η_i is the fraction of population in susceptible set S_i.

3 Model based Projections

The SEIAR-SD-L model was used to create projections in order to help guide policy advice.

While successive projections illustrated in Figure 1 had been communicated periodically to Niti Ayog, we provide details here of the projections that include both the delta and the highly transmissible Omicron variant. Given that substantial time had elapsed since the start of the vaccination program, we utilized vaccination rate efficacy to include breakthrough cases.

These projections on the Sars-Cov2 crisis were based on infection case data until December 15th, 2021. We estimated the transmissibility of Omicron variant from the South African data of the virus and comparing the transmissible factor with the previous variant to compute a beta_factor that was used to determine the transmission factor that is used for Omicron. The fitting for South Africa is detailed in the supplementary material (Figure S3). In the absence of data on incubation period and infectious period we utilized the parameters of the delta variant as estimated from our model during the phase where the delta variant was prevalent. The vaccine efficacy is adjusted for and is considered less effective w.r.t. Omicron. We present results that are parameterized on the vaccine efficiency and listed below.

Baseline Behavior and Assumptions

We established projections that were based on the initial growth in population that were part of transmission dynamics in the first half of 2021 between February and June (first phase), and utilize that to create a baseline for the release of population from lockdown during February-March 2021 when restrictions were removed in late January. This estimate of the release with $\kappa^t = h^L$, i.e. the slope of release, is established during model fitting during the first phase and is critical to establish a rise in cases. We utilized a best-fit model that optimized a loss function. After lockdown removal in June, a release of population that adopts the behaviour of the baseline is estimated. termed the second release. Using the baseline scenario we quantified the second release with respect to the release of population (given by the rate κ^t) into the susceptible population during the first phase. We list the key features and parameters of our methodology below:

- Baseline rate of population increase (rate κ_t) was established during February to March, 2021. An optimization method was used to achieve a best-fit model for the rise in confirmed cases and deaths.

- Average Vaccination rates: 3.0M/day or approximately 0.225%/day till August 16th.

- Subsequent rates averaged to 5M vaccinations/day. Optimistic projected vaccination during releases from lockdown is averaged to approximately 8M vaccinations/day for first dose.

- Additional lockdown removal date is estimated in late November due to local cultural events and its impact estimated to be starting on December 16th, 2021. Lockdown release rate are 100% of baseline over 30 days (high release) and 60 days (moderate release) given the lack of social distancing during the months of November and December.

- We modeled the loss of efficacy of vaccines as reduced linearly over 4 months by 50%. This loss of efficiency results in breakthrough infections.

- The efficacy of vaccines against Omicron was incorporated, considering two case with 50% and 75% of the original efficacy.

- We assume that there is a seeding of 50 cases per state on December 15th, 2021.

We provide two scenarios, one where there will be (i) no subsequent lockdowns and (ii) there will be mild lockdowns, impacting the social distancing and the susceptible population by a factor of 15, 20 and 25 %.

Discussion

The modeling shows projections (Figure 3) that were concerning, especially due to reduction in social distancing countered by reported milder symptoms of the new variant. We modeled this release in two scenarios where the susceptible population is increasing at a substantial rate, the rate being represented by a release of the estimated base population in 30 and 45 days. The first corresponds to a fast release and the second to a moderate release. The estimates illustrate mixed impact of the two variants, with the Omicron variant rapidly dominating the currently existing Delta variant.

We determined estimates from our parameterized model when there was no lockdown as compared to the situation when there was partial lockdown initiated.

No Lockdown assumption:

The two scenarios considered are provided below and evaluated in December, 2021. The indicated peaks for daily infections are substantial.

Scenario 1 (pessimistic scenario–red curve): The pessimistic scenario is when the numer of future release into susceptible pool is adopted to be equivalent to the release during the established baseline estimate. The population that has already become part of the susceptible pool after the relaxation from lockdown is ignored.

This results indicated a substantial rise in infections when rules regarding lockdown are relaxed. Our results projected that there would be increasing high daily case loads peaking at the end of January, 2022 and the beginning of February 2022, extending into March and April months. This indicated that utmost care be taken to ensure social distancing and mask usage, reduced capacity of interior locations and marketplaces, especially in restaurants and bars [25]. Our model indicated infection cases peaking at around 1.3 Million cases per day for 50% efficacy loss and 1.0 Million cases for 25% efficacy loss.

Further peaks were estimated to be progressively delayed and limited in size if social distancing measure are strictly followed. Note that this does not account for any variants that can breakthrough the vaccine defense. The second wave of infections in March to June 2021 illustrated the impact of lack of continued emphasis on these measures.

Figure 3: **50% Vaccine efficacy against Omicron, No lockdown:** Comparison of two scenarios: releasing population having occurred from lockdown illustrating two release rates, high (30 days) and moderate (60 days) Red indicates the worst case behavior (Scenario 1) and Orange indicate the optimistic Scenario 2. The leftmost graph in each row is for the projected cases of the omicron variant whereas the third graph is for the projected confirmed cases of the two variants together.

Figure 4: **75% Vaccine efficacy against Omicron, No lockdown:** Comparison of two scenarios: releasing population having occurred from lockdown with high release (30 days) and low release (60 days). Red indicates the worst case behavior (Scenario 1) and Orange indicate the optimistic Scenario 2. The leftmost graph in each row is for the projected cases of the omicron variant whereas the third graph is for the projected confirmed cases of the two variants together.

Scenario 2 (optimistic scenario-orange curve): This scenario assumes that the susceptible that are added after relaxation of lockdown, but before 15th December, 2021, as estimated from the release function $\kappa^t = h^R$ are discounted from future release of population into the susceptible pool (supplementary table S7). This is an optimistic scenario which models reduced susceptible size due to spreading infection. Even then this indicated a substantial rise in daily case numbers peaking at above 400 thousand cases.

These two scenarios are illustrated in Figure 3 and Figure 4.

Impact of Lockdown:

We also consider the situation when a mild lockdown is implemented after the first week of January, when substantial cases will be visible. January 8th was chosen as the lockdown date in our simulation. This lockdown is assumed to reduce the susceptible population by 15, 20 and 25% for the high release rate scenario (30 day release) prior to the date of lockdown. It is assumed that the release of population stops, and is actually reversed, as the lockdown is implemented.

We observe substantial benefits due to partial lockdown, being limited to less than half of the cases at peak when no lockdown is enforced (Figures 5 and 6), which would have resulted in substantial cases and the resulting hospitalization could have impacted the health system.

The benefits of partial lockdown are further illustrated in figure 7 where the peak number of confirmed cases shows an almost linear decrease with respect to the increase in the percentage of people in partial lockdown. The results indicate that partial lockdown of 25% is reasonably effective in bringing the case incidence down by about 50% even with low vaccine efficacy (Efficacy=0.2). Furthermore, figure 8 shows that delaying partial lockdown increases the possible peak daily infections. It was indicated to be critical that partial lockdown be implemented as soon as feasible.

4 Methodology

Model parameters were determined by an optimization method to minimize a loss function that is dependent on cumulative cases and deaths.

We let G, D denote the simulated cumulative cases and deaths, $\overline{G}, \overline{D}$ are the corresponding reported data. Our optimization methodology utilizes a random initial choices of the parameters over defined ranges as well as a search over the space of release dates to achieve the best fit. The ranges were chosen to be wide enough so that the optimum point was not constrained. The optimization method used is L-BGFS that converges to a best-fit estimate of the model parameters, which due to the non-convex nature of the problem could possibly be a local optima. Sampling multiple starting points to find the best fit leads to reasonably accurate fitting as measured by R^2 values.

4.1 First-Phase fitting

In the first phase we determined the parameters related to the first variant of the virus. We consider the period from February up till June 10th, 2021 when the removal of lockdown began, as an effective release phase where the population behavior trended towards low mask usage and high mobility. While around mid April, 2021 there was a lockdown, we consider the entire period as an effective release period. The release of susceptible population was assumed to be a linear function with a starting date in a range of 1

Figure 5: **50% Vaccine efficacy against Omicron with Lockdown proportion 25,20 and 15 % starting January 8th:** Comparison of three scenarios based on the proportion of population under lockdown. The leftmost graph in each row is for the projected cases of the omicron variant whereas the third graph is for the projected confirmed cases of the two variants together.

Figure 6: **75% Vaccine efficacy against Omicron with Lockdown proportion 25,20 and 15 % starting January 8th:** Comparison of three scenarios based on the proportion of population under lockdown. The leftmost graph in each row is for the projected cases of the omicron variant whereas the third graph is for the projected confirmed cases of the two variants together.

Figure 7: Predicted Peak of confirmed cases as dependent on efficacy of vaccine against the Omicron variant and percentage of population under lockdown. These illustrations are for the high release rate scenario. The release stops at the time of lockdown.

Figure 8: Predicted Peak of confirmed cases as dependent on percentage of population under lockdown and delay of lockdown. These illustrations are for the high release rate scenario. The release stops at the time of lockdown.

week beginning on 15th March. These dates were sampled to compute the release function that resulted in the best fit.

The parameters we considered for fitting are $\beta, \gamma_E, \alpha, \gamma, \gamma_2, \gamma_3, a_1, a_2, a_3, h_R, t_1, t_2$ that are used to generate the series G and D according to the differential equations governing the model.

We use the following loss function.

$$L_1(G, D, r_{ab}) =$$
$$- \left[\theta_1 R^2(G, \overline{G}) + \theta_2 R^2(dG, d\overline{G}) + \theta_3 R^2(D, \overline{D}) + \theta_4 R^2(dD, d\overline{D}) + \theta_5(1 - \frac{|r_{ab} - \overline{r_{ab}}|}{r_{ab}}) \right]$$

where the series G, D and $\overline{G}, \overline{D}$ are weighted geometrically emphasizing the latest data points in each phase, R^2 is the coefficient of determination, $r_{ab}, \overline{r_{ab}}$ are the antibody ratio at the end of the first phase of the simulation and reported, respectively. We obtained $\overline{r_{ab}}$ from sero-surveys carried out by ICMR [17, 20] Our simulation is on a daily basis with $dG, dD, d\overline{G}, d\overline{D}$ being calculated daily. Relative weighting of the parameters $\theta_i, i = 1, 2, 3, 4, 5$ was used to fit the multi-objective function, the choices being provided in the supplementary material.

4.2 Second-Phase Fitting for reopen

In the second phase when removal of lockdown started we considered release of people from lockdown via a linear function with a starting date in a range after 1 week of release of lockdown, i.e. starting on the 17th of June. The release of population into the susceptible set is assumed to be a linear function again and the rate of release dictates the speed at which the policy restrictions are being relaxed and the behavior of the population.

The growth of the infection curve is strongly dependent on this rate, as we will illustrate in our projections. During this phase we estimate the rate of release, the start of release and the size of the release.

$$L_2(h_R) = - \left[\theta_1 R^2(G, \overline{G}) + \theta_2 R^2(dG, d\overline{G}) + \theta_3 R^2(D, \overline{D}) + \theta_4 R^2(dD, d\overline{D}) \right]$$

4.3 Fitting parameters and Projection accuracy

We illustrate our fitting for a combined first and second phase in Figure 9. The goodness of fit for all Indian states is illustrated in the supplementary material. The R^2 values illustrating the goodness of fit was determined and listed in Table S4 in the supplementary material. The box plot in Figure 9 illustrates that the per-capita RMSE is bounded to be less than a small fraction with maximum at less than 0.1% indicating a high degree of accuracy, with a corresponding range of R^2 scores close to 1.

To determine the accuracy of our projections, before the start of the projections, we estimated parameters for our model using data until May 11th, 2021. We fit parameters for the first phase of our model using 4 sets of experiments where we used training data until 4/11/2021, 4/18/2021,4/24/2021, 4/30/2021 and estimated the error in projections of daily cases of infections and deaths beyond these dates for 1-4

Figure 9: Fitting results for India, obtained by accumulating fitting for each state within India, illustrates model performance for both confirmed cases and deaths during the period from February 1st to 15th December, 2021. Data indicating cumulative numbers for each state was used and the model's performance in terms of the per capita RMSE and R^2 scores for each of the states shows the accuracy of the model. Outliers were removed from the Box-plots. Oulier data is in the supplementary material **??**

weeks. The error in projections of daily cases beyond the 4 chosen training dates was below 5% as detailed in the supplementary material S2. Since the vaccination rates were still low, the projections used for testing the accuracy of the model did not account for vaccinations.

Projection accuracy for the Omicron Wave The projection accuracy was measured by evaluating RMSE, MAPE and R^2 values comparing the projections and the actual confirmed cases. We compared the error in our pessimistic estimates versus the actual data for cumulative and daily confirmed cases and deaths at three time intervals, corresponding to 15, 30 and 45 days starting from 15th December, 2021. Cumulative cases were counted from December 15th, 2021. For cumulative cases, the RMSE error per capita was extremely small for projections 15-45 days forward with the maximum error being 0.0013 per capita. The R^2 values were 0.744,0.36,0.203 for 15, 30 and 45 day projections, respectively. And MAPE measure was 17.8%, 27.2% and 41.6% for 15,30 and 45 day intervals respectively. The corresponding numbers for cumulative deaths were: R^2 values of 0.74,0.803 and 0.936 for the 15, 30 and 45 day interval projections. The MAPE values were 20.9%, 21% and 16.9%, while RMSE error/per capita was negligible. For daily cases, there was more variability and the details of the errors may be found in Table S3.

Our projections were based on initial estimate of 50 Omicron cases on December 15th, 2021, in each state, chosen due to the absence of confirmed numbers as accurate Omicron variant case numbers were unknown at that stage.

5   Conclusions and Limitations

Early warning systems which project long range behavior more than a month in advance, while not able to pinpoint exact peaks and accurate case projections, are very useful for policy decisions. Our results were made available to a nodal body (at the central government) for policy decisions. Partial lockdowns, based on inputs from multiple sources, were implemented by various state governments in India.

As indicated from the results, which re-evaluated projections monthly, long term projection of the disease is impacted by governmental policy decisions and adherence to social distancing policies by the population. While it appears that policy decisions, including vaccinations, induced monthly shifts delaying the third wave until December 2021, further investigation of the impact of the policies is required. While we have accounted for a measured reduction of lockdown as well as investigated implementation of levels of lockdown strategies, the impact of a variety of factors including adherence to policy declarations remains an important aspect of further study.

We would like to thank Dr. V.K. Paul (Niti Aayog, Government of India) and Professor S.N. Maheshwari (Honorary Professor, IIT Delhi) for their suggestions and help during the development of these projections.

References and Notes

[1] http://it.delhigovt.nic.in/writereaddata/Odr2022711235.pdf. Policy document for Delhi State.

[2] Forecast video showing projections. https://drive.google.com/file/d/1ohzgOveRDM5Ncgh85mN42Metsxee1Zbo/view?usp=sharing, 2022.

[3] Cleo Anastassopoulou, Lucia Russo, Athanasios Tsakris, and Constantinos Siettos. Data-based analysis, modelling and forecasting of the COVID-19 outbreak. *PloS One*, 15(3):e0230405, March 2020.

[4] Nick Andrews, Julia Stowe, Freja Kirsebom, Samuel Toffa, Tim Rickeard, Eileen Gallagher, Charlotte Gower, Meaghan Kall, Natalie Groves, Anne-Marie O'Connell, et al. Effectiveness of covid-19 vaccines against the omicron (b. 1.1. 529) variant of concern. *MedRxiv*, 2021.

[5] Sercan Arik, Chun-Liang Li, Jinsung Yoon, Rajarishi Sinha, Arkady Epshteyn, Long T. Le, Vikas Menon, and Shashank Singh. *Interpretable Sequence Learning for COVID-19 Forecasting*. Available at https://storage.googleapis.com/covid-external/COVID-19ForecastWhitePaper.pdf.

[6] David W Berger, Kyle F Herkenhoff, and Simon Mongey. An SEIR Infectious Disease Model with Testing and Conditional Quarantine. Working Paper 26901, National Bureau of Economic Research, March 2020.

[7] I De Falco, A Della Cioppa, U Scafuri, and E Tarantino. Coronavirus covid-19 spreading in italy: optimizing an epidemiological model with dynamic social distancing through differential evolution. *arXiv preprint arXiv:2004.00553*, 2020.

[8] Odo Diekmann, JAP Heesterbeek, and Michael G Roberts. The construction of next-generation matrices for compartmental epidemic models. *Journal of the Royal Society Interface*, 7(47):873–885, 2010.

[9] Herbert W Hethcote. The basic epidemiology models: models, expressions for R0, parameter estimation, and applications. In *Mathematical Understanding of Infectious Disease Dynamics*, pages 1–61. World Scientific, 2009.

[10] William Ogilvy Kermack and Anderson G McKendrick. A contribution to the mathematical theory of epidemics. *Proceedings of the Royal Society of London. Series A, Containing papers of a mathematical and physical character*, 115(772):700–721, 1927.

[11] William Ogilvy Kermack and Anderson G McKendrick. "contributions to the mathematical theory of epidemics. ii.—the problem of endemicity". *Proceedings of the Royal Society of London. Series A, Containing papers of a mathematical and physical character*, 138(834):55–83, 1932.

[12] William Ogilvy Kermack and Anderson G McKendrick. Contributions to the Mathematical Theory of Epidemics. III.—Further studies of the problem of endemicity. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 141(843):94–122, 1933.

[13] Stephen M Kissler, Christine Tedijanto, Edward Goldstein, Yonatan H Grad, and Marc Lipsitch. Projecting the transmission dynamics of sars-cov-2 through the postpandemic period. *Science*, 368(6493):860–868, 2020.

[14] Adam J Kucharski, Timothy W Russell, Charlie Diamond, Yang Liu, John Edmunds, Sebastian Funk, Rosalind M Eggo, Fiona Sun, Mark Jit, James D Munday, et al. Early dynamics of transmission and control of COVID-19: a mathematical modelling study. *The Lancet Infectious Diseases*, 20:553–558, May 2020.

[15] Li, Ruiyun and Pei, Sen and Chen, Bin and Song, Yimeng and Zhang, Tao and Yang, Wan and Shaman, Jeffrey. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science*, 368(6490):489–493, 2020.

[16] Clément Massonnaud, Jonathan Roux, and Pascal Crépey. COVID-19: Forecasting short term hospital needs in France. *medRxiv, doi: https://doi.org/10.1101/2020.03.16.20036939*, March 2020.

[17] Manoj V Murhekar, Tarun Bhatnagar, Sriram Selvaraju, V Saravanakumar, Jeromie Wesley Vivian Thangaraj, Naman Shah, Muthusamy Santhosh Kumar, Kiran Rade, R Sabarinathan, Smita Asthana, et al. Sars-cov-2 antibody seroprevalence in india, august–september, 2020: findings from the second nationwide household serosurvey. *The Lancet Global Health*, 9(3):e257–e266, 2021.

[18] K Roosa, Y Lee, R Luo, A Kirpich, R Rothenberg, JM Hyman, P Yan, and G Chowell. Real-time forecasts of the COVID-19 epidemic in China from February 5th to February 24th, 2020. *Infectious Disease Modelling*, 5:256–263, 2020.

[19] Kimberlyn Roosa, Yiseul Lee, Ruiyan Luo, Alexander Kirpich, Richard Rothenberg, James M Hyman, Ping Yan, and Gerardo Chowell. Short-term forecasts of the COVID-19 epidemic in Guangdong and Zhejiang, China: February 13–23, 2020. *Journal of Clinical Medicine*, 9(2):596, 2020.

[20] Lok Sabha. *Questions*. Available at http://loksabhaph.nic.in/Questions/QResult15.aspx?qref=24949lsno=17.

[21] Thomas C Tsai, Benjamin H Jacobson, and Ashish K Jha. American hospital capacity and projected need for COVID-19 patient care. *Health Affairs Blog, doi: 10.1377/hblog20200317.457910*, March 2020.

[22] Joseph T Wu, Kathy Leung, and Gabriel M Leung. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *The Lancet*, 395(10225):689–697, 2020.

[23] Bo Xu, Bernardo Gutierrez, Sumiko Mekaru, Kara Sewalk, Lauren Goodwin, Alyssa Loskill, Emily L Cohn, Yulin Hswen, Sarah C Hill, Maria M Cobo, et al. Epidemiological data from the COVID-19 outbreak, real-time case information. *Scientific data*, 7(1):1–6, 2020.

[24] Juanjuan Zhang, Maria Litvinova, Yuxia Liang, Yan Wang, Wei Wang, Shanlu Zhao, Qianhui Wu, Stefano Merler, Cécile Viboud, Alessandro Vespignani, Marco Ajelli, and Hongjie Yu. Changes in contact patterns shape the dynamics of the covid-19 outbreak in china. *Science*, 368(6498):1481–1486, 2020.

[25] Yi Zhang and Sanjiv Kapoor. Hidden Parameters Impacting Resurgence of SARS-CoV-2 Pandemic. *medRxiv*, 2021.

Supplementary Materials

S1 $\mathcal{R}_0$ with multiple variants

We determine the value of R_0 for a SEIR model with multiple virus variants comprising the set C. The equations governing the behavior of this system are as follows:

$$\frac{dS_i}{dt} = -\sum_C \frac{\beta_i^C(1+\rho)S_i\hat{I}^C}{N}; \qquad \frac{dE^C}{dt} = \frac{\overline{\beta S}^C \cdot \hat{I}^C}{N} - \gamma_E^C E^C;$$

$$\frac{dI^C}{dt} = (1-\alpha)\gamma_E E^C - (\gamma+\gamma_2)I^C; \quad \frac{dA^C}{dt} = \alpha\gamma_E E^C - \gamma_3 A^C;$$

$$\frac{dI_H^C}{dt} = \gamma I^C - (a_1+a_2)I_H^C; \qquad \frac{dI_N^C}{dt} = \gamma_2 I^C - a_3 I_N^C;$$

$$\frac{dD^C}{dt} = a_2 I_H^C; \qquad \frac{dR^C}{dt} = a_1 I_H^C + a_3 I_N^C;$$

where ρ is the rate at which the susceptible set changes due to the population during relaxation of lockdown or during lockdown. $\overline{\beta S}^C = (1+\rho)(\beta_0^C S_0 + \beta_1^C S_1 + \beta_2^C S_2 + \beta_3^C S_3 + \beta_4^C S_4)$, $\hat{I}^C = I^C + A^C$. The parameter ρ is negative in a lockdown state and positive when lockdown is released. The growth R_0 can be obtained from the simplified linear subsystem, where $S = S_0 + S_1 + S_2 + S_3 + S_4$ is assumed to be N:

$$\frac{dE^C}{dt} = \hat{\beta}^C \cdot (I^C + A^C) - \gamma_E^C E^C$$

$$\frac{dI^C}{dt} = (1-\alpha)\gamma_E E^C - (\gamma+\gamma_2)I^C$$

$$\frac{dA^C}{dt} = \alpha\gamma_E E^C - \gamma_3 A^C$$

where $\hat{\beta}^C = \sum_{i=0,1,2,3,4} \beta_i^C(1+\rho)\eta_i$ for the types of the variant. The fractions η_i represent the distribution of susceptibles into the various groups S_0, S_1, S_2, S_3 and S_4. Using the method of [8] we can estimate the value of R_0 from the eigenvalues of $T\Sigma^{-1}$ where T and Σ are block diagonal matrices: $\begin{bmatrix} T_1 & 0 \\ 0 & T_2 \end{bmatrix}$ and $\begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix}$ with

$$T_C = \begin{bmatrix} 0 & \hat{\beta}^C & \hat{\beta}^C \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

and

$$\Sigma_C = \begin{bmatrix} -\gamma_E^C & 0 & 0 \\ \alpha'\gamma_E^C & -(\gamma+\gamma_2) & 0 \\ \alpha\gamma_E^C & 0 & -\gamma_3 \end{bmatrix}$$

where $\alpha' = 1 - \alpha$. Thus

$$\Sigma_C^{-1} = \begin{bmatrix} -1/\gamma_E^C & 0 & 0 \\ -\alpha'/(\gamma+\gamma_2) & -1/(\gamma+\gamma_2) & 0 \\ -\alpha/\gamma_3 & 0 & -1/\gamma_3 \end{bmatrix}$$

The dominant eigenvalue of $T\Sigma^{-1}$ is $\max_C\{\hat{\beta}^C(\alpha'/(\gamma+\gamma_2)+\alpha/\gamma_3)\}$

S2   Model and Projection Accuracy

To establish the accuracy of our model we estimated parameters for our model using training data and projected until May 11th, 2021. We trained our our model using data until 4/11/2021, 4/18/2021,4/24/2021, 4/30/2021 and estimated the error in projections of daily cases of infections and deaths. The error for projections beyond the 4 chosen dates is illustrated in Table S1 and Table S2. Since death projections had a time lag, projections were limited as the training data dates progressed.

Fitting Date	1 week	2 weeks	3 weeks	4 weeks
4/11/21	4.54%	4.11%	3.85%	4.13%
4/18/21	3.67%	3.51%	3.99%	
4/24/21	3.41%	4.21%	4.4%	
4/30/21	4.37%	4.7%	5.0%	

Supplementary Table S1: Error in daily new case projections (all India)

Fitting Date	1 week	2 weeks	3 weeks	4 weeks
4/11/21	11.15%	7.48%	7.73%	6.83%
4/18/21	3.80%	6.02%	5.39%	
4/24/21	8.64%	6.42%		
4/30/21	5.33%			

Supplementary Table S2: Error in daily new death projections (all India)

Omicron Projection Accuracy: We measured the accuracy of our long range projections during the Omicron phase. We measured the cumulative and daily case count errors at three intervals, 15, 30 and 45 days after December 15th, 2021. The results are shown in the Table S3.

metric	15 days	30 days	45 days
RMSE per capita(daily cases)	0.00000250	0.00003963	0.00027739
R2(daily cases)	-12.0704	0.5361	-6.1591
MAPE(daily cases)	38.115%	40.541%	80.549%
RMSE per capita(daily death)	0.00000010	0.00000011	0.00000015
R2(daily death)	-0.8340	-0.1965	-0.5837
MAPE(daily death)	32.621%	37.510%	39.884%
RMSE per capita(cumulative cases)	0.00001232	0.00028131	0.00137607
R2(cumulative cases)	0.7440	0.3763	0.2032
MAPE(cumulative cases)	17.857%	27.268%	41.611%
RMSE per capita(cumulative deaths)	0.00000059	0.00000095	0.00000093
R2(cumulative deaths)	0.7405	0.8036	0.9360
MAPE(cumulative deaths)	20.989%	21.112%	16.869%

Supplementary Table S3: Error in Case Projection of Omicron wave starting from 12/15/2021

S3 Results of model fitting for States in India

The results for the fitting model fitting are illustrated in Figure S2. The fitting was over two phase, the first phase ending on June 10th, the lockdown phase and the second phase was the release phase from June 10th on-wards till December 15th.

The R^2 values for the combined fitting, i.e. first and second phase, of the confirmed infection case numbers is contained in Table S4 in the first column while the R^2 values for the combined fitting of the number of deaths is in the 4th column. The R^2 values were consistently high over all the states. R^2 values for individual states are inconsistent in the second phase for both the infected cases and the deaths since each individual state had applied varying policies, some including multiple attempts at restrictive policies including lockdown during this phase. The parameters obtained from the fitting are detailed in Table S6.

State	R^2(confirmed)	R^2(confirmed phase1)	R^2(confirmed phase2)	R^2(death)	R^2(death phase1)	R^2(death phase2)
India	0.9961	0.9997	0.7951	0.9993	0.9998	0.9750
Andaman and Nicobar	0.9967	0.9962	0.5327	0.1906	0.9662	-1831.9827
Andhra Pradesh	0.9970	0.9999	0.7953	0.9504	0.9899	-0.3936
Arunachal Pradesh	0.9952	0.9980	0.9640	0.9925	0.9725	0.9446
Assam	0.9379	0.9998	-0.4587	0.9603	0.9977	0.2740
Bihar	0.9997	0.9996	-0.6258	0.0798	0.9389	-63.8096
Chandigarh	0.9987	0.9982	0.8754	0.9540	0.9930	-0.4292
Chhattisgarh	0.9996	0.9997	-0.3986	0.9994	0.9995	-1.3777
Daman and Diu	0.9996	0.9996	-0.0732	0.6843	0.6575	0.0000
Delhi	0.9996	0.9996	-4.5287	0.9996	0.9991	-0.4389
Goa	0.9908	0.9988	-0.6514	0.9859	0.9991	-0.3780
Gujarat	0.9998	0.9997	-1.8440	0.9979	0.9934	0.6432
Haryana	0.9994	0.9981	-0.0964	0.9795	0.9993	-2.6716
Himachal Pradesh	0.9886	0.9931	0.1555	0.9966	0.9910	0.7871
Jammu and Kashmir	0.9941	0.9991	-0.1017	0.9954	0.9943	-0.4672
Jharkhand	0.9998	0.9995	0.7872	0.9927	0.9984	-187.2522
Karnataka	0.9991	0.9999	0.7588	0.9989	0.9995	0.8907
Kerala	0.9633	0.9997	0.8058	0.9825	0.9821	0.9538
Ladakh	0.9941	0.9955	-0.6089	0.7665	0.9558	-55.2289
Lakshdweep	0.9223	0.9985	-8.3217	0.7206	0.9570	-35.6842
Madhya Pradesh	0.9989	0.9997	-40.8878	0.7687	0.9852	-6.0478
Maharastra	0.9919	0.9991	0.2586	0.9848	0.9975	0.5334
Manipur	0.9967	0.9972	0.9811	0.9472	0.9714	0.6193
Meghalaya	0.8741	0.9937	0.0778	0.9854	0.9900	0.8856
Mizoram	0.5215	0.9979	0.0192	0.9473	0.9899	0.8839
Nagaland	0.9974	0.9988	0.9628	0.9327	0.9913	-0.0927
Odisha	0.9966	0.9998	0.8474	0.5614	0.9976	-0.8083
Puducherry	0.9920	0.9977	0.2387	0.9932	0.9957	-0.0703
Punjab	0.9989	0.9981	-0.7284	0.9992	0.9989	0.7297
Rajesthan	0.9992	0.9988	-33.4340	0.9990	0.9996	-9.8509
Sikkim	0.9987	0.9989	0.9882	0.8127	0.9726	-1.4990
Tamil Nadu	0.9992	0.9976	0.9520	0.9431	0.9974	-2.1433
Telangana	0.9969	0.9995	0.7377	0.9963	0.9982	0.7271
Tripura	0.9936	0.9985	0.9132	-0.5002	0.9864	-23.2177
Uttar Pradesh	0.9980	0.9998	-158.6117	0.9873	0.9934	-8.8216
Uttarakhand	0.9993	0.9984	-0.4632	0.9972	0.9993	-0.8500
West Bengal	0.9995	0.9996	0.9377	0.9911	0.9980	0.5844

Supplementary Table S4: R^2 score for India and all states resulting from fitting the model

State	RMSE(confirmed)	RMSE(confirmed phase1)	RMSE(confirmed phase2)	RMSE(death)	RMSE(death phase1)	RMSE(death phase2)
India	569636.48	113749.61	733661.93	3304.07	829.50	4235.04
Andaman and Nicobar	62.17	46.85	70.80	26.96	3.41	34.89
Andhra Pradesh	26735.58	2645.13	34647.63	663.33	133.60	854.22
Arunachal Pradesh	1134.67	161.87	1467.28	8.26	3.41	10.35
Assam	41156.92	921.37	53437.89	411.82	37.26	533.87
Bihar	3233.97	3575.20	2977.29	3586.26	319.19	4649.34
Chandigarh	618.92	620.56	617.78	43.61	11.96	55.75
Chhattisgarh	5748.32	4717.63	6359.64	96.49	77.87	107.41
Daman and Diu	59.11	54.87	61.85	0.48	0.58	0.39
Delhi	6291.99	6123.40	6405.11	116.01	150.88	84.10
Goa	4887.45	1308.37	6253.25	127.64	19.90	164.93
Gujarat	3355.40	3865.25	2955.10	110.69	172.75	13.43
Haryana	5322.70	8136.19	1532.45	419.71	51.70	543.31
Himachal Pradesh	7170.09	4028.09	8692.12	67.01	71.24	63.95
Jammu and Kashmir	6617.75	1802.87	8462.55	73.26	51.79	84.91
Jharkhand	1369.79	1974.87	698.32	148.73	60.57	186.50
Karnataka	25090.19	6161.03	32178.07	371.76	126.90	471.16
Kerala	289886.35	9990.76	376334.03	1545.60	212.57	1999.26
Ladakh	327.22	178.66	398.30	10.84	2.35	13.94
Lakshdweep	1270.53	127.25	1646.45	18.09	4.33	23.22
Madhya Pradesh	7091.03	3401.33	8766.24	1334.04	189.01	1725.20
Maharastra	151604.85	42609.65	193672.89	4485.81	768.40	5790.07
Manipur	2287.77	395.55	2952.61	150.22	25.71	193.89
Meghalaya	10225.00	585.58	13268.57	65.42	15.00	84.04
Mizoram	33030.03	130.26	42890.22	39.14	1.17	50.82
Nagaland	411.79	118.01	525.72	64.18	9.12	83.00
Odisha	17036.52	2188.91	22047.94	1826.18	14.72	2371.31
Puducherry	3265.99	1109.18	4140.25	42.26	19.74	52.38
Punjab	5511.12	6142.31	5032.42	124.38	99.69	138.81
Rajesthan	7773.64	8272.69	7411.79	84.38	42.84	103.66
Sikkim	389.91	108.16	498.31	46.74	7.00	60.41
Tamil Nadu	21991.04	21318.63	22440.78	2425.46	207.48	3144.83
Telangana	8388.12	2435.60	10703.71	57.69	26.79	71.55
Tripura	1757.24	245.57	2272.74	226.48	5.30	294.06
Uttar Pradesh	20620.56	7021.12	26137.08	682.43	355.82	835.70
Uttarakhand	2876.79	3558.88	2294.29	131.02	44.43	166.11
West Bengal	9129.80	5651.29	10891.90	351.23	83.79	450.77

Supplementary Table S5: RMSE score for India and all states resulting from fitting the model

Supplementary Figure S1: RMSE per Capita and R2 scores

Supplementary Figure S2: Fitting of Indian States

Supplementary Figure S3: Fitting of South Africa. The ratio of β in the 2nd period versus the 1st is 1.8.

State	β	γ_E	α	γ	γ_2	γ_3	a_1	a_2	a_3
Andaman and Nicobar	49.99956	0.3	0.275821	0.159024	0.060573	0.04	0.056469	0.003398	0.2
Andhra Pradesh	7.882238	0.295294	0.177333	0.040099	0.197636	0.199207	0.211944	0.007038	0.15097
Arunachal Pradesh	3.685567	0.2	0.779355	0.2	0.2	0.134107	0.5	0.005904	0.094896
Assam	4.621663	0.210916	0.815336	0.2	0.2	0.128884	0.5	0.013201	0.011474
Bihar	50	0.3	0.469605	0.063626	0.2	0.2	0.01	0.001736	0.01
Chandigarh	10.22508	0.3	0.1	0.2	0.2	0.2	0.114719	0.002874	0.013961
Chhattisgarh	7.971897	0.3	0.189233	0.154885	0.184541	0.2	0.33631	0.010948	0.021935
Daman and Diu	10.82086	0.3	0.870973	0.04	0.2	0.170437	0.5	0.001	0.165116
Delhi	10.7955	0.3	0.1	0.110269	0.2	0.2	0.137628	0.007119	0.01
Goa	5.666019	0.3	0.1	0.2	0.2	0.106537	0.222253	0.009409	0.015194
Gujarat	42.40642	0.3	0.1	0.2	0.2	0.2	0.5	0.010229	0.010791
Haryana	23.97823	0.3	0.1	0.2	0.2	0.2	0.118534	0.003017	0.011975
Himachal Pradesh	22.01681	0.3	0.1	0.2	0.2	0.2	0.235399	0.008519	0.013447
Jammu and Kashmir	14.32116	0.3	0.130402	0.2	0.199394	0.2	0.440694	0.011909	0.12049
Jharkhand	50	0.3	0.1	0.2	0.2	0.2	0.5	0.019294	0.015639
Karnataka	11.06724	0.210959	0.100545	0.045264	0.120618	0.199422	0.137694	0.006871	0.021345
Kerala	2.483146	0.21049	0.788614	0.04	0.2	0.112912	0.014261	0.001	0.099696
Ladakh	2.306169	0.295377	0.533661	0.06354	0.173535	0.199869	0.098133	0.002325	0.182724
Lakshdweep	1	0.2	0.369919	0.2	0.2	0.04	0.119597	0.001973	0.199669
Madhya Pradesh	49.99987	0.290687	0.107274	0.172474	0.2	0.2	0.086072	0.001673	0.013247
Maharastra	4.339277	0.247925	0.156601	0.09567	0.142352	0.167112	0.027022	0.001429	0.2
Manipur	10.16117	0.2	0.370882	0.2	0.2	0.04	0.5	0.020669	0.2
Meghalaya	31.28439	0.296347	0.1	0.2	0.2	0.2	0.493665	0.023404	0.162305
Mizoram	1	0.2	0.898921	0.04	0.124766	0.064663	0.061328	0.002265	0.188491
Nagaland	28.61501	0.3	0.272491	0.056024	0.185681	0.061247	0.247969	0.046448	0.161904
Odisha	15.43327	0.206192	0.1	0.04	0.2	0.060363	0.03037	0.001	0.15063
Puducherry	3.725994	0.3	0.1	0.192101	0.194099	0.2	0.478134	0.015057	0.2
Punjab	17.46567	0.3	0.1	0.2	0.2	0.2	0.139514	0.007409	0.05231
Rajesthan	43.54541	0.3	0.1	0.2	0.199045	0.2	0.173863	0.003455	0.012366
Sikkim	4.184508	0.2	0.46698	0.2	0.2	0.04	0.5	0.01308	0.2
Tamil Nadu	1.751252	0.283865	0.764172	0.154537	0.040295	0.2	0.12411	0.002323	0.156475
Telangana	50	0.3	0.117522	0.2	0.2	0.04	0.5	0.00634	0.090637
Tripura	3.200484	0.202165	0.899591	0.199954	0.199863	0.069605	0.046824	0.0016	0.06073
Uttar Pradesh	50	0.3	0.365293	0.040047	0.2	0.2	0.099243	0.007514	0.01
Uttarakhand	17.52281	0.3	0.1	0.2	0.2	0.2	0.199483	0.009511	0.015712
West Bengal	8.37165	0.287528	0.551159	0.158685	0.154366	0.2	0.133173	0.002176	0.088322

Supplementary Table S6: Parameters determined for all the states. The model was fit from February, 2021 to June 10, 2021.

State	Ratio	State	Ratio
India	**109.3%**		
Andaman and Nicobar	100.0%	Lakshdweep	105.6%
Andhra Pradesh	131.7%	Madhya Pradesh	159.5%
Arunachal Pradesh	92.4%	Maharastra	175.2%
Assam	76.7%	Manipur	69.0%
Bihar	204.5%	Meghalaya	93.5%
Chandigarh	81.1%	Mizoram	245.7%
Chhattisgarh	146.3%	Nagaland	55.0%
Daman and Diu	63.6%	Odisha	69.1%
Delhi	111.7%	Puducherry	99.1%
Goa	76.7%	Punjab	124.6%
Gujarat	148.0%	Rajesthan	128.0%
Haryana	109.6%	Sikkim	34.7%
Himachal Pradesh	83.9%	Tamil Nadu	89.6%
Jammu and Kashmir	161.7%	Telangana	84.4%
Jharkhand	101.5%	Tripura	38.3%
Karnataka	76.2%	Uttar Pradesh	107.4%
Kerala	113.6%	Uttarakhand	100.0%
Ladakh	109.9%	West Bengal	92.2%

Supplementary Table S7: Release ratios for susceptible population: Population release in second lockdown relaxation as a fraction of release in lockdown relaxation in February and March. Release of vaccinated population may be more.