ARTICLE SUMMARY LINE: Regional spread of SARS-CoV-2 is dependent on weather,

socio-economic and mandatory lockdowns, but the effectiveness of the latter varies with

virus lineage.

RUNNING TITLE: SARS-CoV-2 lineages, socio-economics and lockdowns

KEYWORDS: SARS-CoV-2, lineages, spatio-temporal models, deprivation

TITLE: Spatio-temporal epidemiology of SARS-CoV-2 virus lineages in Teesside, UK, in

2020: effects of social deprivation, weather and lockdown on lineage dynamics

AUTHORS: Rushton, S.P.[1], Moss, E.D.[1], Baker, P.[2], Bashton, M.[3], Crown, M.R.[3], dos

Santos, R.N.[4], Nelson, A.[4], O'Brien, S.J.[1], Richards, Z.[4], Sanderson, R.A.[1]*, Yew, W.C.[4],

Young, G.R.[3], Brooks, E.[5,6]** and Smith, D.L.[3]

AFFILIATIONS:

[1] School of Natural and Environmental Sciences, Newcastle University, Newcastle upon

Tyne, UK

[2] South Tees Hospitals NHS Foundation Trust

[3] Hub for Biotechnology in the Built Environment, Northumbria University, Newcastle upon

Tyne, UK

[4] Department of Applied Sciences, Northumbria University, Newcastle upon Tyne, UK

[5] Department of Medicine, University of Cambridge, UK.

[6] Covid-19 Genomics UK Consortium (COG-UK) https://www.cogconsortium.uk

*Correspondence Author roy.sanderson@newcastle.ac.uk

** See Supplementary Information for full list of Covid-19 Genomics UK Consortium names

and affiliations.

ABSTRACT (150 words)

SARS-CoV-2 emerged in the UK in January 2020 and the Government introduced national

'lockdowns' and regional 'tiers' to control virus transmission. As the outbreak continued,

new variants were detected. We analysed spatio-temporal dynamics of positive tests for

COVID-19 on Teesside, UK throughout 2020, in relation to: socio-economic deprivation,

weather, and Government interventions. We used a combination of disease mapping and

mixed-effect modelling to investigate the dynamics of positive tests from two sampling

strategies and the spread of particular variants of the virus as they emerged on Teesside.

SARS-CoV-2 spread was related to the extent of social deprivation, lockdown interventions

and weather. SARS-CoV-2 spread faster in some lineages than others, with positive tests

related to levels of socio-economic deprivation. The interventions appeared to have different

effects in the two waves of disease, and were associated with reduced numbers of records in

the first wave, but having no effect during the second.

INTRODUCTION

In 2019 a new coronavirus respiratory infection (COVID-19) caused by the SARS-CoV-2
virus (1) emerged in China and rapidly spread globally, with first UK patients in January
2020. COVID-19 is highly infectious with transmission primarily via airborne droplets (2).
Initial UK COVID-19 control measures aimed to break transmission through home
quarantine of cases and suspected cases, social distancing, and disease tracking. Disease
cases increased despite these controls and hence a complete social and economic lockdown
was introduced on March 23rd 2020 (3). Understanding of the epidemiology of the disease in
the early phases of the pandemic was low, for example it was not clear that at least 30% of
infected individuals could be asymptomatic (4). However, there was early evidence that both
the severity of disease, and mortality, were age-related, with older individuals (5) and those
with comorbidities at substantially higher risk of developing severe illness requiring
respiratory support (6). Whilst the first lockdown led to a rapid reduction in cases and
pressures on secondary care, it was insufficient to stop the epidemic in the UK and the
number of cases increased rapidly in autumn 2020 (7), leading to the introduction of further
interventions to reduce social contact and spread (including a short national lockdown in
November 2020). This failed to reduce hospital admissions and deaths, and a further
lockdown was introduced in January 2021 (8).

In contrast with previous pandemics, that of COVID-19 occurred at a time when molecular
techniques were available for rapid diagnosis of disease. This meant that from early in the
epidemic, polymerase chain reaction (PCR) analyses were available not only to diagnose
COVID-19 disease, but also for genomic sequencing (9) to characterise genetic lineages of
SARS-CoV-2 in individual cases. The UK government introduced two 'pillars' of testing for
case ascertainment. Initial pillar 1 testing was undertaken in Public Health England (PHE)
laboratories and NHS hospitals for patients identified with a clinical need (10). It also in-

cluded health and care workers who were likely to encounter infected individuals or transmit

it to people vulnerable to disease. Later pillar 2 testing assessed community-level incidence

of infection. Inevitably, pillar 1 case ascertainment omitted asymptomatic individuals or

those who had not sought help from primary care. It therefore represents an underestimate of

the spatio-temporal disposition of disease and a biased estimate of the burden of disease

across different age demographics (11).

SARS-CoV-2 has had high mutation rates throughout the pandemic. Pathan *et al.* (12) re-

ported high levels of three types of RNA substitution mutations: missense mutations, where

the amino acid being coded is changed; nonsense mutations, which stop gene translation; and

silent mutations, where the functional nature of the protein produced is unchanged (13). Mis-

sense, nonsense and silent mutation rates were 34%, 6.7% and 0.8% respectively (12). High

rates of mutation led to the development of several virus lineages, often with different epide-

miological characteristics. Lineage B.1.1.7 (the Alpha variant, first identified in Kent, UK)

emerged in autumn 2020 and had high transmission rates (14,15). SARS-CoV-2 spike pro-

teins initiate infection and host-cell colonisation via binding the ACE2 receptor (16,17) and

mutations that change the spike protein occur in lineages with higher infectivity (18). Muta-

tion rates provide a tool for forensic analysis of spread in communities. Here we use longitu-

dinal records of COVID-19 cases to determine the spatio-temporal dynamics of disease

spread in the Teesside region of North East England during 2020. We use records of individ-

ual lineages from pillar 1 and 2 tests in conditional autoregression and mixed-effects models

to quantify the impacts of social deprivation, demography, weather, and UK Government in-

terventions on disease spread in this region.

METHODS

Data collation

Records of positive tests were collated for individual postcode districts (TS1 to TS29) in the Teesside area. The 29 separate postcode districts have a range of population sizes and different socio-demographic conditions (Table S1).

COVID-19 lineage types were derived from the Pangolin (19) and Civet (20) analyses of genomes for both pillar 1 and pillar 2 tests, analysed as part of the COG-UK initiative. Available data provided the dates and postcodes of all positive tests for SARS-CoV-2 from January 2020 to January 2021. Pillar 1 samples were collected 7 hospital trusts across the North East and Cumbria on a weekly basis and random sampling was employed to select a representative set of positive cases with CT values below 30 across these trusts. Pilar 2 samples were sequenced at the Wellcome Sanger Institute using a random sampling procedure again for samples with a CT below 30 from UK Lighthouse Laboratories. Lineages were defined as those recognised as of 31[st] January 2021, restricting analyses to pillar 1 and pillar 2 tests, and summarised for subsequent analyses by week of test. The extent to which the different sampling methods used by pillar 1 versus pillar 2 may have biased detection of lineages amongst cases was investigated by correlating the total numbers of each lineage recorded by each pillar during the period when sampling was run contemporaneously for both pillars (from week of year 30 to 53).

We collated covariates hypothesised as important in determining spread of disease. These were socio-demographics of the exposed populations, weather and interventions introduced by UK government to mitigate spread. Recent demographic data on socio-economic deprivation, Index of Multiple Deprivation IMD (21), was available for each postcode district, as SARS-CoV-2 may be higher in areas of social deprivation (22). Mean weekly

temperature and rainfall were obtained from Teesside Airport to provide weather covariates

that might affect spread. We collated timings for Government interventions for the UK

COVID-19 epidemic in two forms: first, complete national lockdowns; and second, more

heterogeneous interventions that were applied at regional scales, sometimes referred to as a

"Tier" system. These interventions were included as additional covariates into the models to

estimate the extent to which the observed patterns of disease could be associated with

attempts to mitigate spread.

Disease mapping

118 We used disease mapping (23) with postcode as the areal unit. We calculated expected risk of

119 disease in each area as a function of the proportion of positive tests that might be expected

120 given the population of that area. Excess or reduced risk in each area was then determined by

121 comparison with the risk for the whole Teesside region.

Mixed-effect modelling

We used mixed-effect models to investigate the role of social deprivation, weather

(temperature and rainfall), and UK Government interventions (lockdowns) on the dynamics

of positive PCR test cases recorded in each postcode district for all positive tests. We used a

negative binomial mixed-effect model with postcode as the random effect because data in

postcodes represent repeated measures and there would be unmeasured variation in the

postcodes that might affect recorded cases. Furthermore, we anticipated that cases in post-

codes would not be normally distributed but aggregated as recorded in other large scale

studies (24). To account for variation in date of infection to being tested, we included a time

lag of two weeks on the introduction of lockdown interventions.

There were insufficient data to model the trends in each individual lineage across all

postcodes and many postcodes had zero counts. To investigate spread of lineages in time and

space we therefore used a zero-inflated negative binomial mixed-effect model with lineage

type as a random effect, to relate the number of positive tests of each lineage against time,

using the same covariates above. The model had random intercepts and random gradients for

time for each lineage to enable comparison of the number of positive tests for each lineage as

well as differences in their rates of increases with time.

For both the standard negative binomial and zero-inflated models all covariates were initially

fitted (full model), and then covariates were eliminated as models were simplified based on

Akaike Information Criterion (AIC) scores to find the best minimal model. All models were

fitted in R with the glmmTMB package (25)

RESULTS

The data and trends in positive tests for all tests and individual lineages

The Teesside postal area has a population of 599,600 people across 29 postcode districts,

with considerable variation in the proportion of the total population in the top 10^{th} IMD decile

(Table S1). The populations of urban postcode districts of central Middlesbrough (TS1, TS4

and TS3) all had high proportions of their populations in the top 10^{th} decile for IMD, whereas

many of the suburban and rural postcode districts had little to none of their population in this

decile. There were 2,328 positive COVID-19 tests for all lineages of which 1,073 were pillar

1 and 1,255 were pillar 2. Temperature was highest in summer, but there were also periods of

heavy rain in late spring and early autumn (Figure 1a).

The overall temporal trend in positive tests was bimodal with a first peak in week 12 (March

2020) followed by a decline through the summer, before the onset of a minor peak and

decline in week 40 and a rapid increase in numbers of positive tests from week 42 (Figure

1b). This reflects the use of pillar 1 testing in clinical settings and the introduction of pillar 2

testing later in the epidemic.

Pillar 1 cases were detected from week 10 (2nd March 2020), reached a peak of 100 cases in

week 13, before declining to 0 in week 28 after a UK Government lockdown (Figure 1c).

Pillar 1 cases rose again from week 37 and stayed high to the end of 2020. In contrast, pillar 2

cases were first detected in week 18 (6th May 2020) as community testing was introduced.

There was a strong positive correlation between the total number of cases recorded by the

two different pillars (r=0.94, t=18.87, df=47, P<0.001), during the period when both pillars

were in operation (weeks 30 to 53 inclusive), suggesting that both pillars were reflecting the

similar patterns of infection across Teesside.

Out of 86 distinct lineages only 8 were recorded more than 60 times: B.1.1.1, B.1.1.119,

B.1.1.309, B.1.1.315, B.1.1.37, B.1.1.7, B.1.177 and B.1.177.10 (Figure 1d). These 8

lineages were recorded elsewhere in the UK before being recorded on Teesside. There were

two dominant lineages in the first wave of the epidemic, B.1.1.1 and B.1.1.119, records for

which declined rapidly during the first lockdown and to zero by week 27. Patterns for other

lineages were more complex in the second and third waves. B.1.177 and B.1.1.315 both

increased in frequency during weeks 32 to 40 before declining in the second lockdown.

However, after the end of this lockdown, B.1.177 increased substantially whilst B.1.1.315 did

not. A new, highly transmissible SARS-CoV-2 variant, B.1.1.7 (Alpha variant; VoC

202012/1) was first detected in the UK in December 2020 (26) and increased hugely after the

end of the second lockdown. The first lockdown appeared successful in reducing the most

prevalent lineages during that part of the epidemic, whereas the second lockdown was less

effective.

Disease mapping

The relative risk of cases in each Teesside postcode for the 8 commonest lineages varied greatly amongst lineages and between the two waves of the epidemic (Figure 2). The spatial pattern of risk of a test being positive for COVID-19 of *any* lineage was highest in western Middlesbrough (TS5). In the first wave of the pandemic lineages B.1.1.1 and B.1.1.119 were predominant; both were associated with urban postcodes of Middlesbrough although B.1.1.1 was more widespread and was also relatively high in parts of Stockton-on-Tees (TS18), Redcar (TS10), and Guisborough (TS14) as well as some south western suburban and rural areas. Cases during the second wave of the epidemic were more widespread and represented more lineages (B.1.1.119, B.1.1.309, B1.1.7, B.1.1.37, B.1.177, B.1.177.10). The risks for B.1.177 and B.1.1.315 were widespread in urban and suburban areas across the central part of Teesside. The risk of B.1.1.37 was also widespread but had a patchier distribution. The remaining three lineages had more concentrated distributions, with the highest risks in either Thornaby in TS17 (B.1.1.309), southern Redcar in TS25 (B.1.1.7), or both (B.1.177.10).

Mixed-effect modelling

There was a significant negative relationship between total positive tests for COVID-19 and mean weekly temperature, rainfall, and the time since imposition of the second lockdown (Table 1a). Cases were positively related to week of the year, the time since imposition of the first lockdown, the total population, and the population resident in areas in the top decile of social deprivation for England. There was broad agreement in the observed and predicted distributions of cases (Figure 3).

For individual lineages, parameter estimates for the best zero-inflated model with postcode and lineage as random intercepts and with a random gradient for week for lineages are shown in Table 1b. The more urban and suburban postcodes of Middlesbrough (TS1 to TS8) and

some of the rural postcodes in the south east (TS9, TS12 and TS14) and north west (TS21

and TS22) had positive random effects (Figure 4a). This indicates that records of positive

cases in these postcodes were higher even after adjusting for total population density.

Random effects for week, showing relative rate of change in number of positive cases of each

lineage in sequenced samples, indicated that B.1.1.7 (Alpha variant) increased substantially

faster than the others (Figure 4b).

DISCUSSION

Our analyses indicate considerable spatial and temporal variation in occurrence of SARS-

CoV-2 lineages in Teesside in 2020 and that the patterns were related to several hypothesised

covariates. These results must be interpreted with caution because of the way in which

positive test results were ascertained and also due to relatively large sizes of geographical

areas used. Positive tests were derived from two collection strategies: first, pillar 1, that

focussed on patients and individuals showing illness in primary or secondary care; and

second, pillar 2, from general sampling in the community. As the first wave declined the

number of pillar 1 samples declined, whereas pillar 2 sampling accelerated, and this may

have biased the results. Pillar 1 samples may have been biased away from asymptomatic

carriage, which will have been detected in pillar 2 samples. Furthermore, lineages may differ

in the extent to which they lead to serious illness, which may have impacted on what was

detected in pillar 1 (27,28). However, the relative abundance of each lineage in the

overlapping period for pillar 1 and pillar 2 was highly correlated, suggesting that both pillars

provided similar representations of lineages in general circulation amongst the population on

Teesside. The geographies (postcodes) used to define the areas for the mixed-effect

modelling may not capture spatial variation in the underlying epidemiological process of

transmission or other social and environmental drivers. Postcodes ensured anonymity of

records, but many covariates associated with disease transmission, including proximity to

infectious individuals and social contacts, could not be measured. The covariates used in

these models were therefore surrogates for the underlying mechanisms associated with

disease transmission and spread.

Notwithstanding these caveats, four main conclusions can be drawn from our analyses. First,

there was considerable spatio-temporal variation in positive SARS-CoV-2 tests across the

region. Middlesbrough has the most ethnically diverse communities on Teesside with an

ethnic minority population of 27.2% (29). There is a well-established link between risk of

serious disease from SARS-CoV-2 with both ethnicity (30) and deprivation (22). However,

separating the possible contributions of ethnicity and deprivation on Teesside is difficult as

we do not have information on the ethnic status of the individuals subject to PCR testing.

Furthermore, whilst deprivation, population density, and 'urbanness' are strongly related,

analysis of COVID-19 in Norfolk indicates that they can have independent contributions to

levels of disease (31).

Second, there was variation in the extent to which cases appeared spatially and temporally.

Many of the more urban postcodes recorded cases earlier and records tended to expand in

different directions from these foci with time. The lineages had all been recorded elsewhere

in the UK before they appeared emerged on Teesside suggesting that lineages were

introduced rather than arising locally. Indeed, B.1.177 emerged in Spain during summer

2020 before rapidly spreading across Europe (32). Teesside urban centres (TS1 to TS6),

which function as a local transport and population hubs, may have played a role in the

initiation and local spread of cases.

Third, Government interventions introduced to manage the disease during 2020 had mixed

impacts on the spatial and temporal dynamics of records in Teesside.   Numbers of positive

tests were, counter-intuitively, correlated positively with the imposition of three of the

interventions and only negatively associated with the introduction of Tier 2 which was a local

intervention. This probably reflects the fact that the imposition of early interventions on

Teesside was not driven by the state of disease in the region (only 100 pillar 1 cases out of a

regional population of nearly 600,000) but rather spread elsewhere in the UK.

Fourth, the mixed-effect models demonstrated distinct differences in both the spatial and

temporal patterns amongst the numbers of positive tests for different lineages in Teesside. For

example, lineage B.1.1.7 increased rapidly across the region towards the latter part of 2020,

despite initially having a relatively restricted geographical distribution in Teesside (Figures 2

and 4). B.1.1.7 emerged in the South East of the UK in December 2020 after a temporary

relaxation of social restrictions relaxed prior to Christmas.

These analyses have demonstrated low effectiveness of national-level interventions for

COVID-19 on disease patterns in Teesside, particularly in late 2020. This contrasts with other

studies undertaken at larger national and trans-national scales (24) where timings of

interventions were demonstrated to have impacts on spread of disease. However, our results

suggest that disease management should be more orientated toward local epidemiological

conditions at the time point of delivery, indicating a need for more temporal precision. In

addition, our analyses also support the hypothesis that socio-economic deprivation played a

role in the spread of the disease. Since this is to a large extent spatially defined it is likely that

interventions could also have been targeted with greater spatial precision. Our results also

suggest that the rate of increase in cases, as measured by the pillar testing system, was similar

for most lineages, except for the alpha variant (B.1.1.7) which subsequently spread over

much of the UK in 2021. We conclude that the spatio-temporal spread of COVID-19 on

Teesside in 2020 was to a large extent opportunistic, but mediated locally by socio-economic

deprivation and reactive rather than proactive interventions.

ACKNOWLEDGEMENTS

The authors thank colleagues within COG-UK for constructive comments on earlier versions

of this manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

FUNDING

This research was funded under COG-UK; this is supported by funding from the Medical

Research Council (MRC) part of UK Research & Innovation (UKRI), the National Institute

of Health Research (NIHR) [grant code: MC_PC_19027], and Genome Research Limited,

operating as the Wellcome Sanger Institute.

RESEARCH ETHICS APPROVAL

The surveillance activities within which this study was conducted are part of Public Health

England's responsibility to monitor COVID-19 during the current pandemic. Public Health

England has legal permission, provided by Regulation 3 of The Health Service (Control of

Patient Information) Regulations 2002 to process confidential patient information under

Sections 3(i) a–c, 3(i) d(i and ii), and 3(iii) as part of its outbreak response activities. This

study falls within the research activities approved by the Public Health England Research

Ethics and Governance of Public Health Practice Group.

AUTHOR BIO

Steve Rushton is Professor of Biological Modelling at Newcastle University. His research

focuses on prediction of disease risks in humans and wildlife.

REFERENCES

1. WHO. Naming the coronavirus disease (COVID-19) and the virus that causes it [Internet]. 2021 [cited 2021 May 19]. Available from: https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it

2. Wilson N, Corbett S, Tovey E. Airborne transmission of covid-19. BMJ. 2020 Aug 20;370:m3206.

3. UK Government. Prime Minister's statement on coronavirus (COVID-19): 23 March 2020 [Internet]. www.gov.uk. 2020 [cited 2021 May 19]. Available from: https://www.gov.uk/government/speeches/pm-address-to-the-nation-on-coronavirus-23-march-2020

4. Qiu J. Covert coronavirus infections could be seeding new outbreaks. Nature [Internet]. 2020 Mar 20 [cited 2021 May 19]; Available from: http://www.nature.com/articles/d41586-020-00822-x

5. Yanez ND, Weiss NS, Romand J-A, Treggiari MM. COVID-19 mortality risk for older men and women. BMC Public Health. 2020 Nov 19;20(1):1742.

6. Atkins JL, Masoli JAH, Delgado J, Pilling LC, Kuo C-L, Kuchel GA, et al. Preexisting Comorbidities Predicting COVID-19 and Mortality in the UK Biobank Community Cohort. The Journals of Gerontology: Series A. 2020 Oct 15;75(11):2224–30.

7. UK Government. Coronavirus in the UK [Internet]. 2021 [cited 2021 May 19]. Available from: https://coronavirus.data.gov.uk

8.   UK Government. Prime Minister's address to the nation: 4 January 2021 [Internet].

www.gov.uk. 2021 [cited 2021 May 19]. Available from:

https://www.gov.uk/government/speeches/prime-ministers-address-to-the-nation-4-

january-2021

9.   COVID-19 Genomics UK (COG-UK) Consortium. An integrated national scale SARS-

CoV-2 genomic surveillance network. The Lancet Microbe. 2020 Jul 1;1(3):e99–100.

10.   Dept. Health & Social Care. COVID-19 testing data: methodology note [Internet].

GOV.UK. 2021 [cited 2021 May 19]. Available from:

https://www.gov.uk/government/publications/coronavirus-covid-19-testing-data-

methodology/covid-19-testing-data-methodology-note

11.   Clare T, Twohig KA, O'Connell A-M, Dabrera G. Timeliness and completeness of

laboratory-based surveillance of COVID-19 cases in England. Public Health. 2021 May

1;194:163–6.

12.   Pathan RK, Biswas M, Khandaker MU. Time series prediction of COVID-19 by

mutation rate analysis using recurrent neural network-based LSTM model. Chaos,

Solitons & Fractals. 2020 Sep 1;138:110018.

13.   Lodish H, Berk A, Zipursky SL, Matsudaira P, Baltimore D, Darnell J. Mutations:

Types and Causes. In: Molecular Cell Biology 4th edition [Internet]. W. H. Freeman;

2000 [cited 2021 May 19]. Available from:

https://www.ncbi.nlm.nih.gov/books/NBK21578/

14.   Davies NG, Abbott S, Barnard RC, Jarvis CI, Kucharski AJ, Munday JD, et al.

Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England.

Science [Internet]. 2021 Apr 9 [cited 2021 May 19];372(6538). Available from:

https://science.sciencemag.org/content/372/6538/eabg3055

15. Kirby T. New variant of SARS-CoV-2 in UK causes surge of COVID-19. The Lancet

Respiratory Medicine. 2021 Feb 1;9(2):e20–1.

16. Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, et al. Structure of the SARS-CoV-2 spike

receptor-binding domain bound to the ACE2 receptor. Nature. 2020

May;581(7807):215–20.

17. Shang J, Ye G, Shi K, Wan Y, Luo C, Aihara H, et al. Structural basis of receptor

recognition by SARS-CoV-2. Nature. 2020 May;581(7807):221–4.

18. Teruel N, Mailhot O, Najmanovich RJ. Modelling conformational state dynamics and its

role on infection for SARS-CoV-2 Spike protein variants. bioRxiv. 2021 Jan

11;2020.12.16.423118.

19. Rambaut A, Holmes EC, O'Toole Á, Hill V, McCrone JT, Ruis C, et al. A dynamic

nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. Nat

Microbiol. 2020 Nov;5(11):1403–7.

20. O'Toole A, Hill V. Civet - Cluster Investigation and Virus Epidemiology Tool

[Internet]. Civet - Cluster Investigation and Virus Epidemiology Tool. 2021 [cited 2021

May 19]. Available from: https://artic-network.github.io/civet/

21. UK Government. English indices of deprivation 2019 [Internet]. GOV.UK. 2019 [cited

2021 Nov 5]. Available from: https://www.gov.uk/government/statistics/english-

indices-of-deprivation-2019

22.   Woodward M, Peters SAE, Harris K. Social deprivation as a risk factor for COVID-19

mortality among women and men in the UK Biobank: nature of risk and context

suggests that social interventions are essential to mitigate the effects of future

pandemics. J Epidemiol Community Health [Internet]. 2021 Apr 27 [cited 2021 May

19]; Available from: https://jech.bmj.com/content/early/2021/04/27/jech-2020-215810

23.   Lawson AB. Bayesian Disease Mapping: Hierarchical Modeling in Spatial

Epidemiology. New York: Chapman and Hall/CRC; 2008. 368 p.

24.   Hunter PR, Colón-González FJ, Brainard J, Rushton S. Impact of non-pharmaceutical

interventions against COVID-19 in Europe: A quasi-experimental study. medRxiv. 2020

Jul 17;2020.05.01.20088260.

25.   Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al.

glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated

Generalized Linear Mixed Modeling. The R Journal. 2017;9(2):378–400.

26.   Public Health England. Variant Of Concern 202012/01: Technical Briefing 2.

Investigation of novel SARS-CoV-2 variant [Internet]. 2021. Available from:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment

_data/file/959361/Technical_Briefing_VOC202012-2_Briefing_2.pdf

27.   Burki T. Understanding variants of SARS-CoV-2. The Lancet. 2021 Feb

6;397(10273):462.

28.   Dumonteil E, Fusco D, Drouin A, Herrera C. Genomic Signatures of SARS-CoV-2

Associated with Patient Mortality. Viruses. 2021 Feb;13(2):227.

29.   UK Government. Regional ethnic diversity [Internet]. 2021 [cited 2021 Nov 5].

384   Available from: https://www.ethnicity-facts-figures.service.gov.uk/uk-population-by-

385   ethnicity/national-and-regional-populations/regional-ethnic-diversity/latest#areas-of-

386   england-and-wales-by-ethnicity

387   30.   Niedzwiedz CL, O'Donnell CA, Jani BD, Demou E, Ho FK, Celis-Morales C, et al.

388   Ethnic and socioeconomic differences in SARS-CoV-2 infection: prospective cohort

389   study using UK Biobank. BMC Medicine. 2020 May 29;18(1):160.

390   31.   Brainard J, Rushton S, Winters T, Hunter PR. Spatial risk factors for Pillar 1 COVID-19

391   case counts and mortality in rural eastern England, UK. medRxiv. 2020 Dec

392   4;2020.12.03.20239681.

393   32.   Hodcroft EB, Zuber M, Nadeau S, Vaughan TG, Crawford KHD, Althaus CL, et al.

394   Spread of a SARS-CoV-2 variant through Europe in the summer of 2020. Nature. 2021

395   Jul;595(7869):707–12.

ADDRESS FOR CORRESPONDENCE: **Roy Sanderson, School of Natural and Envi-**

ronmental Science, Newcastle University, Newcastle upon Tyne, NE1 7RU, United

Kingdom; email: roy.sanderson@newcastle.ac.uk

Table 1. Summary of results of mixed-effects models for a) all-lineages combined with a

negative binomial error with postcode district as a random effect and b) zero-inflated

negative binomial errors with random intercepts for postcode district and lineages, and week

of year as a random gradient for lineages. Both models relate observed number of positive

tests of COVID-19 in Teesside postcode districts to demographics of postcode, weather, and

imposition of interventions to restrict disease spread. Temperature and rainfall are the mean

values per week, calculated from daily means and daily totals respectively. The interventions

include a two-week time lag to account for the delay in symptom onset (and testing) after

infection.

a) Negative binomial error model

Variable	Estimate	Std. Error	z value	p value
(Intercept)	-2.879	0.226	-12.712	<0.001
Temperature	-0.041	0.009	-4.759	<0.001
Rainfall	-0.098	0.021	-4.571	<0.001
First Lockdown (+2 weeks)	0.052	0.015	3.503	0.001
Second lockdown (+2 weeks)	-0.132	0.036	-3.686	<0.001
Week of year	0.073	0.003	22.352	<0.001
Tier 2 (+2 weeks)	0.100	0.040	2.511	0.012
Deprived population (IMD 10^{th} decile)	0.021	0.009	2.248	0.025
Total population	0.047	0.006	7.834	<0.001

411   b) Zero-inflated negative binomial error model

Variable	Estimate	Std. Error	z value	p value
(Intercept)	-7.935	1.950	-4.068	<0.001
Temperature	0.061	0.014	4.248	<0.001
Rainfall	-0.060	0.025	-2.426	0.015
First Lockdown (+ 2 weeks)	-0.040	0.021	-1.877	0.061
Week of year	0.165	0.097	1.705	0.088
Tier 2 (+ 2 weeks)	0.254	0.036	7.094	<0.001
Deprived population (IMD 10[th] decile)	0.025	0.011	2.207	0.027
Total population	0.051	0.007	7.054	<0.001
Zero-inflated model				
(Intercept)	-32.475	9.066	-3.582	<0.001
Week of year	-2.432	0.662	-3.672	<0.001

FIGURE LEGENDS

Figure 1. Trends in weather and positive tests of COVID-19 over the course of 2020, in

Teesside. (a) Mean temperature and rainfall per week (calculated from daily means and daily

totals respectively). (b) Total positive tests per week (across all lineages), coloured areas

represent the periods of different interventions: complete national lockdown (National

lockdown), weak regional restrictions (Regional tier 2), stricter regional restrictions

(Regional tier 3), and the 'Eat out to help out' government scheme to promote eating within

restaurants via financial incentive (Hospitality promotion). (c) Positive tests detected via the

two testing regimes per week. (d) Positives tests of the eight most commonly detected

lineages in Teesside per week.

Figure 2. Relative risk of the 8 most common SARS-CoV-2 lineages (and all cases) being

detected in each Teesside postcode district during 2020 (from the spatial CAR models).

Lineages B.1.1.1 and B.1.1.119 occurred mainly in the first wave in spring 2020. Lineages

B.1.1.309, B.1.1.315, B.1.1.37, and B.1.177 occurred mainly in the second wave beginning in

late August 2020, while lineages B.1.1.7 and B.1.177.10 occurred mainly at the end of the

year. The central Middlesbrough postcode districts predominated in the first wave and the

remaining districts (including Hartlepool) more in the second wave.

Figure 3. Predicted and observed total number of positive cases of COVID-19 (across all

lineages) in Teesside during 2020. The predicted estimates (red) are fitted values from a

negative binomial mixed effect model of the total number of positive test results in different

Teesside postcodes.

Figure 4. Random effects and associated 95% confidence intervals for the intercepts from the

zero-inflated negative binomial mixed effect model relating counts of positive test results for

different lineages of SARS-CoV-2 in a) different Teesside postcodes and b) amongst

lineages. Blue values indicate random effects that were greater than the mean and an earlier

increase in positive cases, while red indicates the converse.

Random effects of Postcode (Intercept)

Random effects of Week