

Factors associated with contraceptive failure in Uganda: Analysis of the 2016 Uganda Demographic and Health Survey

Short title

Contraceptive failure in Uganda

Authors

Ruth Ketty Kisuza¹, Saviour Kicaber¹, Derrick Abila Bary¹, Felix Bongomin², Christopher Orach Garimoi¹.

Affiliation

1) College of Health Sciences, Makerere University, Kampala Uganda.

2) Department of Medical Microbiology and Immunology, Faculty of Medicine, Gulu University, Gulu, Uganda

Corresponding author

Ruth Ketty Kisuza.

College of Health Sciences,

Makerere University, Kampala Uganda

P.O Box 7072

Kampala, Uganda

1 **Abstract**

2 **Introduction**

3 Sustained motivation is essential for effective use of contraceptive methods by women in low- and

4 middle-income countries as many women are likely to abandon use of contraceptives especially

5 when they continually experience episodes of failure. We aimed to determine contraceptive failure
6 rates and associated factors among Ugandan women using data from the 2016 Uganda
7 Demographic Health Survey (UDHS).

8 **Methods**

9 We analyzed data collected by the UDHS conducted in Uganda 2016. All eligible women aged 15
10 to 49 years at the time of the survey were enrolled. Discontinuation of contraceptive use due to
11 failure within a 5-year period preceding the survey was the dependent variable.

12 **Results**

13 A total of 18,505 women were included in this study, 70.8% (n=5153) lived in rural areas while
14 56.9% (n=5153) owned a mobile phone. The mean age of the women was 29.6years (SD 7.6). The
15 overall prevalence of contraceptive failure was 5.6%, and was higher (7.8%) among women aged
16 25-29 years or had completed secondary education (7.1%). The odds of contraceptive failure was
17 38% lower in women who had an informed choice on contraceptives compared to those who didn't
18 [Adjusted Odd ratio, 0.62; 95% confidence interval, 0.50 – 0.77; p< 0.001].

19 **Conclusion**

20 The burden of contraceptive failure among women of reproductive age in Uganda is substantial
21 and significantly varied by socio-demographic characteristics.

22 **Key words:** Contraceptive failure, Uganda, Demographic Health Survey (DHS) and Socio-
23 demographic Characteristics.

24

25 **Introduction**

26 Improving access to family planning (FP) services is fundamental to achieving the Sustainable
27 Development Goals (SDGs) because it is strongly related with women's and children's health,

28 poverty reduction, education, gender equality, and human rights. Access to family planning
29 contributes up to a 44% reduction in maternal deaths [1]. Since most unplanned pregnancies and
30 abortions occur in women who were either not using contraception or not using it consistently.
31 Greater access to contraception and more consistent use of contraception are crucial in the
32 reduction of unplanned pregnancies and abortions.

33
34 The reasons for discontinuation of contraceptives can be grouped by whether they represent
35 discontinuations due to reduced need for contraception (not in need) or discontinuations while
36 women were presumably still exposed to the risk of pregnancy and did not want to become
37 pregnant (in need). Among discontinuations that are not in need, the most common reason given
38 is wanting to become pregnant. Among in-need discontinuations, the most common reasons given
39 are women that become pregnant while using the method (contraceptive failure) or those that
40 stopped using the method because of side effects[2].

41
42 High rates of discontinuation for reasons other than the desire for pregnancy are problematic
43 because of their association with several negative reproductive health outcomes. In countries with
44 moderate-high contraceptive prevalence most of the unintended pregnancy is the result of
45 contraceptive discontinuation or failure. Several studies have associated contraceptive
46 discontinuation for reasons other than the desire to become pregnant with unmet need for
47 contraception and induced abortion ([2–4]. Unintended pregnancy has been associated with
48 increased risk of maternal morbidity, health behaviors during pregnancy that are associated with
49 adverse maternal health, and adverse fetal, infant and child health outcomes [5] Additionally,

50 unintended pregnancy has been associated with negative psychological effects on women and their
51 children[6].

52
53 In the developing world, 74 million unintended pregnancies occur annually, and nearly a third
54 30%, are due to contraceptive failure among women using some type of contraceptive method
55 (whether traditional or modern). This includes both method-related failures (i.e., failure of a
56 method to work as expected) and user-related failures (i.e., failure due to incorrect or inconsistent
57 use of a method)[7].

58
59 Sustained motivation is essential for effective use of contraceptive methods by women in low- and
60 middle-income countries as many women are likely to abandon use of contraceptives especially
61 when they continually experience episodes of failure. Assessing failure rates among demographic
62 and socioeconomic groups is important to inform efforts to improve contraceptive information,
63 services and use, and to minimize contraceptive failures. Additionally, detailed information on
64 contraceptive failure rates is critical to inform improvements in provision of contraceptive
65 information, supplies and services, which can help women and couples to use methods correctly
66 and consistently. This study aimed to determine the contraceptive failure rates and the factors
67 associated with contraceptive failure among Ugandan women using the 2016 Uganda
68 Demographic Health Survey (UDHS) data.

69
70

71 **Methods**

72 **Study design and setting:**

73 We analyzed data collected during the UDHS conducted in 2016. “The Demographic and Health
74 Surveys (DHS) are internationally comparable household surveys that collect information on
75 demographic, socioeconomic, and health-related variables among nationally representative
76 samples of households in developing countries. Details of the DHS sampling design and strategies
77 are described elsewhere”[8].

78 **Study population**

79 In this study, we used the women recode file which included women aged 15 to 49 years at the
80 time of the survey. We included all the women in the dataset into our analysis.

81 **Study variables**

82 Discontinuation of contraceptive use due to failure within a five-year period preceding the survey
83 was the dependent variable. The DHS does not have a variable that records discontinuation of
84 contraceptive use due to failure. This was derived from the variable v360 that records the various
85 reasons for discontinuation such as pregnancy, wanting to become pregnant, side effects and
86 wanting a more effective method. In this study, a woman becoming pregnant while on
87 contraceptives was termed as a method failure (exposure) while discontinuation due to another
88 reason was termed as non-exposure.

89 The independent variables used in the study included (1) age, (2) type of place of residence (rural
90 vs urban), (3) wealth index, (4) education attainment, (5) phone ownership, (6) internet use, (7)
91 number of children delivered, (8) Women's participation in decision-making for using
92 contraceptives, and (9) work status. We also included the variable on the method of contraception
93 that was discontinued over the 5-year period preceding the survey.

94 **Data management**

95 The data used were obtained after receiving permission from the DHS program website. The
96 dataset for the Uganda DHS 2016 was then downloaded from the DHS program website. The
97 women's recode file readable by STATA version 13 was selected for use.

98 To calculate failure rate, we created a new variable for which the denominator would be the total
99 of the number of women who were currently using contraceptives at time of survey interview and
100 the number of women who were not using at the time of survey because they had discontinued.

101 The number of women who were currently using contraceptives at time of survey were obtained
102 by converting variable v312 in the DHS into a binary variable (currently use vs. non-currently
103 using). The number of women who were not using contraceptives at the time of the survey because
104 they discontinued was obtained in the following way. First, we created a new variable for only
105 women who were not currently using contraceptives at the time of the survey. This was recorded
106 into a binary variable with 0 if a woman had not discontinued contraceptives and 1 if a woman
107 discontinued contraceptives. The latter was used to obtain the number of women who were not
108 using contraceptives at the time of the survey because they discontinued.

109 The numerator for calculating contraceptive failure rate was obtained by converting DHS variable
110 v360 (reason for discontinuation) into a binary variable with yes (1) if reason was
111 failure/pregnancy and no (0) if other reason for discontinuation. The numerator considered was
112 the number of women who discontinued due failure.

113 **Data analysis**

114 All the analysis in this study was performed in STATA13 and each survey dataset was analyzed
115 separately (StataCorp LP 2013). Weighting was performed for all the descriptive statistics using
116 the weight variable (v005) after dividing it by 1,000,000.

117
$$\text{weight} = \frac{V005}{1,000,000}$$

118 For the regression analysis in STATA13, weighting was also performed using the Primary
119 Sampling Unit (PSU) and Strata as the variables v021 and v022 respectively in the DHS survey
120 datasets for the surveys and weight (wgt) calculated previously. The STATA13 code below was
121 used to apply weights before calculating means, standard deviation and 95% confidence intervals
122 of continuous variables and regression analysis.

123 `svyset v021 [pw = wgt], strata(v022) single unit(centered)`

124 where pw is the probability weight (sampling weight), psu is primary sampling unit, v021 is
125 variable for the primary sampling unit, and v022 is the variable in the DHS that indicates the strata
126 used in the DHS surveys.

127 **Summary statistics**

128 For the categorical variables, we summarized them as weighted proportions. Variables that were
129 numeric like age were summarized as weighted means reporting standard deviations and 95%
130 confidence intervals.

131 **Regression analysis**

132 We used bivariate and multivariate logistic regression to determine the association between
133 demographic characteristics and contraceptive failure among women. We reported crude odds
134 ratios, adjusted odds ratios, p-values, and 95% confidence intervals. First, bivariate logistic
135 regression was used to determine the association between contraceptive failure and the women's
136 demographic characteristics. Then, multivariate logistic regression was used including the
137 variables that had a p-value of less than 0.200 i.e. (1) age group (2) number of children a woman
138 has, and (3) Frequency of using the internet last month.

139

140

141 **Quality control**

142 To ensure that we are working with the correct DHS dataset for a given year, a test analysis was
143 performed to try to replicate the tables in the respective DHS survey reports for the various years.
144 We created a frequency distribution table for the variable of type of place of residence for the
145 different years in STATA13 and compared the result with what was reported in the respective
146 DHS survey report. All results from this test analysis matched those in the DHS survey reports.

147 **Results**

148 **Characteristics of the women**

149 **Table 1** shows the socio-demographic characteristics of the sample of women aged 15-49 who
150 had used and never used a contraceptive method within the five years before the 2016 UDHS.
151 Among contraceptive users (n=8000), 23.4% were between age 25-29 years, only 7.6% (n=691)
152 were adolescents (age 15-19 years) and 4.4% (n=401) were above age 45-49 years. Only 7.8% had
153 no formal education. Majority (80.3%) of the women were working and about 70.7% lived in rural
154 areas of Uganda. About 28.9% (n=2573) of the women belonged to the highest wealth quintile.
155 Majority of women who used contraceptives had never been exposed to the internet in the last 12
156 months (89.0%), 9.6% (n= 873) had used the internet in the last 12 months. Additionally, 56.8%
157 (n= 5153) of the women who used contraceptives owned a mobile phone. Among cultural factors,
158 about two-thirds of the women made a joint decision with their partners on contraceptive use.
159 41.8% (n=3788) of the women using contraceptives made an informed choice while 58.2%
160 (n=5273) of the women did not make an informed choice. The most common methods used and
161 discontinued in the past 5 years were injectable (54.8%), followed by implants (10.5%).

162

163 **Table 1: Distribution of method related, demographic, socio economic and cultural**
 164 **characteristics of women by status of contraceptive use.**

Characteristics	Current users or ever used contraceptives in past 5 years		Neither current user nor ever used contraceptive in past 5 years	
	Number	Percentage	Number	Percentage
Total	9060		9445	
Demographic factors				
Age at the start of the episode				
Mean (SD)	29.64 (7.68)		26.3 (10.60)	
95% CI	29.44 – 29.85		26.06 – 26.54	
5-year age group				
15-19	691	7.62	3574	37.83
20-24	2081	22.97	1740	18.43
25-29	2122	23.42	929	9.84
30-34	1704	18.81	839	8.88
35-39	1220	13.46	791	8.38
40-44	841	9.28	767	8.12
45-49	401	4.43	805	8.53

Parity (living children at the end of the episode)				
Mean (SD)	3.78 (2.66)		2.41 (3.11)	
95% CI	3.7 – 3.87		2.33 – 2.49	
Socioeconomic factors				
Internet use in the past year				
Never	8067	89.03	8624	91.31
Yes, last 12 months	873	9.64	723	7.66
Yes, before last 12 months	120	1.33	98	1.04
Frequency of using internet last month				
Not at all	8776	92.70	8230	90.83
Less than once a week	127	1.35	119	1.32
At least once a week	187	1.98	285	3.14
Almost every day	376	3.98	426	4.71
Mobile phone ownership				
No	3907	43.12	6170	65.32
Yes	5153	56.88	3275	34.68
Women's education attainment				
No education	678	7.48	1103	11.68
Incomplete primary	3783	41.76	4474	47.36
Complete Primary	1291	14.26	1081	11.45
Incomplete secondary	2251	24.85	2093	22.16
Complete secondary	169	1.87	125	1.32

Higher	887	9.79	569	6.02
Currently/formerly/never in union				
Never in union	863	9.52	3920	41.50
Currently in union/living with a man	6939	76.58	4284	45.36
Formerly in union/living with a man	1259	13.89	1241	13.14
Partners/Husbands education attainment (n = 11,223)				
No formal education	287	4.13	425	9.93
Incomplete primary	2191	31.58	1609	37.55
Complete Primary	1233	17.77	798	18.62
Incomplete secondary	1744	25.15	861	20.11
Complete secondary	304	4.39	101	2.35
Higher	976	14.07	382	8.91
Do not know	202	2.91	109	2.54
Husband's/partner's age				
15-19	18	0.26	56	1.31
20-24	503	7.25	621	14.49
25-29	1220	17.59	685	15.98
30-34	1433	20.65	650	15.16
35-39	1256	18.11	581	13.56
40-44	1055	15.20	488	11.39
45-49	759	10.94	490	11.44
50-54	423	6.10	380	8.88

55-59	135	1.95	140	3.27
60 years or more	136	1.96	194	4.52
Work status				
Not currently employed	1784	19.69	3201	33.90
Currently employed	7276	80.31	6243	66.10
Place of residence				
Urban	2650	29.24	2293	24.28
Rural	6411	70.76	7152	75.72
Household wealth index quintile				
Lowest	1205	13.30	2041	21.61
Second	1600	17.67	1796	19.01
Middle	1725	19.04	1734	18.36
Fourth	1956	21.59	1727	18.28
Highest	2573	28.39	2147	22.73
Cultural factors				
Women participation in decision-making for using contraceptives				
Mainly woman	1340	30.64	-	-
Mainly husband/partner	312	7.13	-	-
Joint decision	2710	61.98	-	-
Other	11	0.25	-	-
Informed choice				
No	5273	58.20	-	-

Yes	3788	41.80	-	-
Method-related factors				
Contraceptive method discontinued				
Pill	553	9.64	-	-
IUD	161	2.80	-	-
Injectable	3150	54.85	-	-
Male condom	444	7.72	-	-
Male sterilization	1	0.01	-	-
Periodic abstinence	188	3.27	-	-
Withdraw	371	6.46	-	-
Other traditional	42	0.73	-	-
Implants/Norplant	605	10.54	-	-
Lactational amenorrhea (lam)	155	2.71	-	-
Female condom	2	0.03	-	-
Emergency contraception	22	0.38	-	-
Other modern method	15	0.26	-	-
Standard days method (sdm)	35	0.61	-	-

165 IUD= intrauterine contraceptive device

166 **Prevalence of contraceptive failure**

167 As shown in **Table 2**, the prevalence of contraceptive failure among Ugandan women aged 15-49
 168 years during the 5-year period by demographic, socio economic and cultural factors was 5.6%

169 (n=506). Among the demographic factors, failure rate 7.83% (n=157) was prevalent in women
 170 aged 25-29 years and having five or more children (6.35%). For socioeconomic factors, the highest
 171 prevalence rates were seen in women who used internet almost every day in the one month (7.78%,
 172 n=426), owned a mobile phone (5.72%, n=294), completed higher education (7.09%, n=63),
 173 unemployed (5.83%, n=104) and lived in a rural area (5.70%, n=364).
 174 Among cultural factors, women who had a joint decision for contraceptive use with their partner
 175 had the highest failure of the 4.95% (n=134). Women who made an informed choice on
 176 contraceptive use had low contraceptive failure rates (4.43%, n=168) as compared to those who
 177 didn't make an informed choice (6.41%, n=338).

178

179 **Table 2: Prevalence of contraceptive failure among Ugandan women aged 15-49 years**
 180 **during a five-year period by demographic, socioeconomic, and cultural factors.**

		5-year contraceptive failure rate	
Demographic characteristics	Number using contraceptives	No. disc.	Percentage
Total	9061	506	5.58
Demographic factors			
5-year age group			
15-19	690	19	2.82

20-24	2081	105	5.02
25-29	2122	157	7.38
30-34	1704	105	6.19
35-39	1220	66	5.40
40-44	841	41	4.84
45-49	401	13	3.22
Parity (total children ever born)			
One	1298	58	4.46
Two	1489	73	4.88
Three	1312	77	5.88
Four	1164	73	6.25
Five or more	3119	198	6.35
Socioeconomic factors			
Internet use in the past year			
Never	8067	440	5.45
Yes, last 12 months	873	57	6.58
Yes, before last 12 months	120	8	6.79
Frequency of using internet last month			
Not at all	8230	450	5.47
Less than once a week	119	6	4.67
At least once a week	285	17	5.91
Almost every day	426	33	7.78

Mobile phone ownership			
No	3907	211	5.40
Yes	5153	295	5.72
Education			
No education	678	35	5.10
Incomplete primary	3783	194	5.12
Complete Primary	1292	66	5.10
Incomplete secondary	2251	136	6.05
Complete secondary	169	12	7.14
Higher	887	63	7.09
Currently/formerly/never in union			
Never in union	863	24	2.74
Currently in union/living with a man	6939	431	6.21
Formerly in union/living with a man	1259	51	4.03
Partners/Husbands education attainment (n = 11,223)			
No education	287	23	8.12
Incomplete primary	2191	122	5.56
Complete Primary	1233	76	61.6
Incomplete secondary	1745	104	5.94
Complete secondary	304	27	8.98
Higher	976	63	6.41
Do not know	202	16	8.15

Husband's/partner's age			
15-19	18	0	0
20-24	503	32	6.28
25-29	1220	67	5.50
30-34	1433	106	7.38
35-39	1256	91	7.27
40-44	1055	46	4.38
45-49	759	56	7.39
50-54	423	21	4.97
55-59	135	2	1.73
60 years or more	136	10	7.09
Work status			
Not currently employed	1784	104	5.83
Currently employed	7277	401	5.52
Place of residence			
Urban	2650	140	5.30
Rural	6411	365	5.70
Household wealth index quintile			
Lowest	1205	64	5.31
Second	1601	85	5.29
Middle	1725	97	5.60
Fourth	1956	116	5.93
Highest	2573	144	5.60

Cultural factors			
Women participation in decision-making for using contraceptives			
Mainly woman	1340	64	4.81
Mainly husband/partner	311	9	3.00
Joint decision	2710	134	4.95
Other	-	-	-
Informed choice			
No	5273	338	6.41
Yes	3788	168	4.43

181

182 **Table 3** shows the contraceptive failure rates in Uganda by method used. Generally traditional
 183 methods had the highest failure rates. Of this, withdrawal (36.30%, n=134) was the single most
 184 common reason for failure. For modern contraceptive methods, short term methods like emergency
 185 contraception (20.85%, n=5) and pills (10.68%, n=59) had higher failure rates when compared to
 186 long term modern contraceptives like IUD (1.62%, n=3) and implants (2.60%, n=15) which had
 187 very low failure rates.

188 **Table 3: Contraceptive failure rates in Uganda by methods used.**

Method-related factors			
Contraceptive method discontinued	Number who used the method	Number failed	Percentage (%)

Pill	553	59	10.68
IUD	161	3	1.62
Injectable	3150	140	4.43
Male condom	444	31	6.93
Male sterilization	1	0	0
Periodic abstinence	188	65	34.87
Withdraw	371	134	36.30
Other traditional	41	19	44.78
Implants/Norplant	605	15	2.60
Lactational amenorrhea (lam)	155	20	12.69
Female condom	2	1	50.00
Emergency contraception	22	5	20.85
Other modern method	15	2	13.43
Standard days method (sdm)	35	12	34.17

189

190

191 **Factors associated with contraceptive failure.**

192 From the multivariate analysis, women aged 45 to 49 years [Adjusted Odds ratio (aOR), 0.33; 95% confidence interval (95CI), 0.12 –
 193 0.94; p value = 0.038] were less likely to experience a contraceptive failure compared to those aged 15 to 19 years. Women who had an
 194 informed choice on contraceptives were less likely to experience a failure [aOR, 0.62; 95CI, 0.50 – 0.77; p-value < 0.001] compared to
 195 those who did not have informed choice (**Table 4**).

196

197 **Table 4: Factors associated with contraceptive failure among women in Uganda.**

Demographic characteristics	Bivariate analysis		Multivariate analysis	
	Crude Odds Ratio (95% CI)	P - value	Adjusted Odds Ratio (95% CI)	P - value
Demographic factors				
5-year age group				
15-19	1		1	
20-24	1.73 (1.04 – 2.86)	0.034	1.28 (0.66 – 2.49)	0.464
25-29	2.61 (1.59 – 4.27)	< 0.001	1.56 (0.75 – 3.25)	0.234

30-34	2.19 (1.32 – 3.63)	0.002	0.98 (0.45 – 2.16)	0.967
35-39	2.05 (1.22 – 3.46)	0.007	0.83 (0.36 – 1.90)	0.664
40-44	1.56 (0.88 – 2.76)	0.127	0.53 (0.22 – 1.3)	0.166
45-49	1.05 (0.496 – 2.211)	0.903	0.33 (0.12 – 0.94)	0.038
Parity (total children ever born)				
No child	1		1	
One child	1.58 (0.93 – 2.66)	0.089	0.68 (0.34 – 1.36)	0.277
Two children	1.45 (0.87 – 2.44)	0.157	0.66 (0.33 – 1.32)	0.239
Three children	1.80 (1.08 – 3.01)	0.024	0.79 (0.39 – 1.62)	0.522
Four children	1.895 (1.131 – 3.175)	0.015	0.92 (0.43 – 1.93)	0.818
Five or more children	2.16 (1.35 – 3.47)	0.001	1.70 (0.80 – 3.61)	0.166
Socioeconomic factors				
Internet use in the past year				
Never	1		-	
Yes, last 12 months	1.08 (0.78 – 1.49)	0.637	-	-
Yes, before last 12 months	1.42 (0.70 – 2.88)	0.329	-	-

Frequency of using internet last month				
Not at all	1		-	
Less than once a week	0.73 (0.28 – 1.88)	0.512	-	-
At least once a week	1.46 (0.89 – 2.398)	0.136	-	-
Almost every day	1.01 (0.64 – 1.59)	0.971	-	-
Mobile phone ownership				
No	1		-	
Yes	0.99 (0.82 – 1.19)	0.918	-	-
Education				
No education	1		-	
Incomplete primary	1.01 (0.698 – 1.454)	0.969	-	-
Complete Primary	1.09 (0.72 – 1.65)	0.678	-	-
Incomplete secondary	1.18 (0.799 – 1.73)	0.410	-	-
Complete secondary	1.32 (0.63 – 2.77)	0.461	-	-
Higher	1.20 (0.77 – 1.89)	0.416	-	-
Currently/formerly/never in union				

Never in union	1			
Currently in union/living with a man	2.15 (1.41 – 3.28)	< 0.001	-	
Formerly in union/living with a man	1.39 (0.84 – 2.30)	0.205	-	
Partners/Husbands education attainment (n = 11,223)				
No education	1			
Incomplete primary	0.66 (0.42 – 1.04)	0.074	-	-
Complete Primary	0.76 (0.47 – 1.22)	0.253	-	-
Incomplete secondary	0.68 (0.42 – 1.09)	0.110	-	-
Complete secondary	1.06 (0.58 – 1.94)	0.843	-	-
Higher	0.76 (0.46 – 1.25)	0.278	-	-
Do not know	0.90 (0.45 – 1.81)	0.772	-	-
Husband's/partner's age				
15-19	1		1	
20-24	2.93 (0.17 – 49.43)	0.456	2.75 (0.16 – 47.27)	0.486
25-29	2.69 (0.16 – 44.71)	0.491	1.99 (0.11 – 34.47)	0.638

30-34	3.81 (0.23 – 63.21)	0.350	2.43 (0.14 – 42.32)	0.543
35-39	3.39 (0.20 – 56.34)	0.394	2.26 (0.13 – 39.68)	0.577
40-44	2.49 (0.15 – 41.64)	0.524	1.75 (0.098 – 31.00)	0.704
45-49	3.42 (0.21 – 57.14)	0.392	2.78 (0.16 – 49.49)	0.487
50-54	2.66 (0.16 – 45.18)	0.499	2.5 (0.14 – 45.56)	0.537
55-59	1.4 (0.07 – 27.98)	0.826	1.46 (0.07 – 31.20)	0.810
60 years or more	3.34 (0.19 – 59.89)	0.412	3.02 (0.16 – 57.77)	0.463
Work status				
Not currently employed	1		-	
Currently employed	1.02 (0.80 – 1.31)	0.849	-	-
Place of residence				
Urban	1		1	
Rural	1.03 (0.83 – 1.27)	0.803	1.04 (0.8 – 1.4)	0.789
Household wealth index quintile				
Lowest	1		-	
Second	1.06 (0.78 – 1.45)	0.712	-	-

Middle	0.98 (0.71 – 1.34)	0.901	-	-
Fourth	1.05 (0.77 – 1.43)	0.747	-	-
Highest	0.95 (0.70 – 1.29)	0.760	-	-
Cultural factors				
Women participation in decision-making for using contraceptives				
Mainly woman.	1		-	
Mainly husband/partner	0.75 (0.38 – 1.46)	0.399	-	-
Joint decision	1.01 (0.73 – 1.399)	0.933	-	-
Other	1.27 (0.07 – 22.20)	0.872	-	-
Informed choice				
No	1		1	
Yes	0.65 (0.53 – 0.79)	< 0.001	0.62 (0.50 – 0.77)	< 0.001

199 **Discussion**

200 In this study, we aimed to establish and describe the prevalence of contraceptive failure amongst
201 women of reproductive age in Uganda. We used data from the demographic and health survey
202 conducted in Uganda in 2016. From this study, it is evident that the prevalence of contraceptive
203 failure varies by age, type of place of residence, phone ownership, education attainment and wealth
204 index. The highest failure rates were seen in women who use periodic abstinence and withdraw as
205 methods of contraception. Finally, women who had delivered four or more children were more
206 likely to experience contraceptive failure when compared with those who had not delivered any
207 child.

208
209 The study shows that women with access to the internet were more likely to experience
210 contraceptive failure compared to women with no access to the internet. This could be because the
211 women with access to the internet can easily access information on the different contraceptive
212 methods and are therefore more likely to stop or switch to better contraceptive methods when they
213 experience side effects [9]. Additionally, women who were provided with adequate information
214 about a particular contraceptive method and given an opportunity to make an informed choice were
215 less likely to experience contraceptive failure than women who did not make an informed choice.
216 This could be because a woman who made an informed choice had adequate information about
217 the contraceptive method of her choice and was able to effectively deal with its side effects without
218 necessarily halting contraceptive use [10,11]. This finding suggests that more time should be
219 dedicated by clinicians to providing information on contraceptives to women before use.

220 In this study, failure rates by contraceptive methods were highest in women who used traditional
221 methods than in those who used modern contraceptive methods. For modern contraceptive

222 methods, women using short term methods like pills and Injectables were more likely to experience
223 failure than those using long term contraceptive methods. This could be because of inconsistency
224 when using short term contraceptive methods which exposes women to increased failure rates.
225 Similar findings were reported in [2,12,13]. Thus, information and counselling should be
226 emphasized for short term method users to prevent failure.

227
228 At multivariate analysis, the most important factor associated with contraceptive failure was a
229 woman's age. Women above 45 years of age had lower contraceptive failure rates than women
230 below 40 years. This could be because women above 40 years are more likely than younger women
231 to desire a permanent form of contraception[14,15]. Additionally, older women of reproductive
232 age have lower rates of contraceptive failure than younger women and this could be explained by
233 lower fecundity, less frequent sexual intercourse and higher compliance with contraceptive
234 regimens as evidenced from previous studies [16].

235
236 This study used retrospective data from the DHS. The DHS follows a rigorous methodology
237 employing a two-stage cluster randomized sampling. We used sample weights to provide estimates
238 reported in this study. However, we acknowledge that contraceptive failure could best be studied
239 using cohort studies. But this study provides further information about the current state of
240 contraceptive failure among women of reproductive age in Uganda.

241 242 **Conclusion**

243 Our study found that there was a high burden of contraceptive failure among women of
244 reproductive age in Uganda that and varies by socio-demographic characteristics. Traditional

245 contraceptive methods including periodic abstinence and withdrawal are associated with higher
246 failure rates.

247 **Declarations**

248 **Ethics approval and consent to participate**

249 Ethical approval for the Demographic and Health Surveys is obtained from relevant research
250 ethical approval committees and research regulatory bodies before data collection. We obtained
251 permission to use the DHS survey datasets from the DHS program website. No personally
252 identifiable information is available in the used datasets.

253 **Consent for publication**

254 Not Applicable

255 **Availability of data and materials**

256 The datasets generated and/or analyzed during the current study are available in the Demographic
257 and Health Survey program website, <https://dhsprogram.com/data/available-datasets.cfm> and can
258 be accessed after obtaining approval.

259 **Competing interests**

260 The authors declare that they do not have any competing interests.

261 **Funding**

262 The authors did not receive any funding for this work.

263 **Authors' contributions**

264 DBA and RKK conceived the study. DBA and RKK requested data from the DHS program
265 website, performed statistical analysis, interpretation of data and spearheaded the writing of the
266 manuscript. RKK and DBA discussed the relevant results. DBA, RKK, SK, FB and CGO
267 contributed to the writing of the manuscript, read, and approved the final manuscript.

268 **Acknowledgements**

269 We thank the Demographic and Health Survey (DHS) program for granting us access to the DHS
270 datasets used in this study

271

272 **References**

- 273 1. Starbird E, Norton M, Marcus R. Investing in family planning: Key to achieving the
274 sustainable development goals. *Glob Heal Sci Pract*. 2016;4: 191–210.
275 doi:10.9745/GHSP-D-15-00374
- 276 2. Ali MM, Cleland JG, Shah IH, Organization WH. Causes and consequences of
277 contraceptive discontinuation: evidence from 60 demographic and health surveys. 2012.
- 278 3. Black KI, Gupta S, Rassi A, Kubba A. Why do women experience untimed pregnancies?
279 A review of contraceptive failure rates. *Best Pract Res Clin Obstet Gynaecol*. 2010;24:
280 443–455.
- 281 4. Polis CB, Bradley SEK, Bankole A, Onda T, Croft T, Singh S. Contraceptive Failure
282 Rates in the Developing World: An Analysis of Demographic and Health Survey Data in
283 43 Countries. *Contraception*. 2016;94: 11–17. Available:
284 <https://linkinghub.elsevier.com/retrieve/pii/S0010782416001037>
- 285 5. Yazdkhasti M, Pourreza A, Pirak A, Abdi F. Unintended pregnancy and its adverse social
286 and economic consequences on health system: a narrative review article. *Iran J Public*
287 *Health*. 2015;44: 12.
- 288 6. Gipson JD, Koenig MA, Hindin MJ. The effects of unintended pregnancy on infant, child,
289 and parental health: a review of the literature. *Stud Fam Plann*. 2008;39: 18–38.
- 290 7. Polis C, Bradley SEK, Bankole A, Onda T, Croft TN, Singh S. Contraceptive failure rates

- 291 in the developing world: an analysis of demographic and health survey data in 43
292 countries. 2016.
- 293 8. Corsi DJ, Neuman M, Finlay JE, Subramanian S V. Demographic and health surveys: a
294 profile. *Int J Epidemiol.* 2012;41: 1602–1613.
- 295 9. Samosir OB, Kiting AS, Aninditya F. Role of information and communication technology
296 and women’s empowerment in contraceptive discontinuation in Indonesia. *J Prev Med
297 Public Heal.* 2020;53: 117.
- 298 10. Pradhan MR, Patel SK, Saraf AA. Informed choice in modern contraceptive method use:
299 pattern and predictors among young women in India. *J Biosoc Sci.* 2020;52: 846–859.
- 300 11. Hilger DJ, Raviele KM, Hilgers TA. Hormonal contraception and the informed consent.
301 *Linacre Q.* 2018;85: 375–384.
- 302 12. Bradley SEK, Schwandt H, Khan S. Levels, trends, and reasons for contraceptive
303 discontinuation. *DHS Anal Stud.* 2009;20.
- 304 13. Westhoff CL, Heartwell S, Edwards S, Ziemann M, Stuart G, Cwiak C, et al. Oral
305 contraceptive discontinuation: do side effects matter? *Am J Obstet Gynecol.* 2007;196:
306 412-e1.
- 307 14. Alemayehu M, Belachew T, Tilahun T. Factors associated with utilization of long acting
308 and permanent contraceptive methods among married women of reproductive age in
309 Mekelle town, Tigray region, north Ethiopia. *BMC Pregnancy Childbirth.* 2012;12: 1–9.
- 310 15. Azmoude E, Behnam H, Barati-Far S, Aradmehr M. Factors affecting the use of long-
311 acting and permanent contraceptive methods among married women of reproductive age
312 in East of Iran. *Women’s Heal Bull.* 2017;4: 1–8.
- 313 16. Allen RH, Cwiak CA, Kaunitz AM. Contraception in women over 40 years of age. *CMAJ.*

314 2013;185: 565–573.

315