

1 **The impact of changes in mental health services in response to**
2 **COVID-19 on people with mental health conditions: protocol for a**
3 **rapid review**

4
5 G Yu^{*1,2}, D Craig^{1,2}, Y Fu^{1,2}

- 6
7 1. NIHR Applied Research Collaborative North East and North Cumbria, Cumbria,
8 Northumberland and Tyne and Wear NHS Foundation Trust, The Clocktower Building, St
9 Nicholas Hospital, Gosforth, Newcastle NE3 3XT, UK
10 2. Population Health Sciences Institute, Faculty of Medical Sciences, Newcastle University,
11 Sir James Spence Building, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP, UK

12
13
14 *Correspondence

15 **Dr Ge Yu**

16 Baddiley-Clark Building
17 Richardson Road
18 Population Health Sciences Institute
19 Newcastle University
20 Newcastle
21 NE2 4AX

22 ge.yu@newcastle.ac.uk

23 01912 085307

24
25 Keywords: mental health, mental health conditions, psychotic disorders, mental health
26 service, COVID-19, rapid review

27 Word count: 1782

28
29
30

1 **ABSTRACT**

2 **Introduction**

3 The COVID-19 pandemic has caused disruptions to mental health services, forcing the rapid
4 implementation of alternative ways of delivering services alongside a greater immediate,
5 and continuously growing, demand across those services. The care and level of mental
6 health service provided are felt to be inadequate to respond to the increasing demand for
7 mental health conditions in the time of the pandemic, leading to an urgent need to learn
8 from service change and consequences to inform solutions and plans to support the NHS
9 post-pandemic plan in the UK. This rapid review aims to understand the changes in mental
10 health services during the pandemic and summarise the impact of these changes on the
11 health outcomes of people with mental health conditions.

12 **Methods and analysis**

13 Cochrane CENTRAL, MEDLINE, EMBASE, and PsycInfo will be searched for eligible studies
14 with key terms indicating mental health AND COVID-19 AND health services. Studies will be
15 included if objective and subjective effects of changes to mental health services in response
16 to COVID-19 are reported on adults with mental health conditions, peer-reviewed and
17 published in the English language. Study selection and data extraction will be undertaken
18 dependently by two reviewers. Evidence will be summarised narratively and in a logic
19 model.

20 **Ethics and dissemination**

21 Ethics approval is not required for this review. A list of interventions/services/models of
22 care delivered to people with mental health conditions will be grouped as “Do”, “Don’t” and
23 “Don’t know” based on the evidence on effectiveness and acceptability. The results will be
24 written for publication in an open-access peer-reviewed journal and disseminated to the
25 public and patients, clinicians, commissioners, funders, and academic conferences.

26 **PROSPERO registration number**

27 CRD42022306923

1 **Strengths and limitations of this study**

- 2 • This is a rapid review with a systematic search of literature
- 3 • Evidence on changes to mental health services and associated health outcomes will be
- 4 examined to make practice recommendations
- 5 • This review will provide the evidence base to inform solutions and plans to support the
- 6 mental health service provision for the post-pandemic period
- 7 • Some limitations to the study design include limited to OECD studies, exclusion of non-
- 8 English studies, publication bias, quality of data, selection bias, and no quality
- 9 assessment in the rapid evidence review.

1 INTRODUCTION

2 In the UK, up to December 2021, 1 in 7 people have had a positive COVID-19 test result. It is
3 recognised that as well as the physical effects, a COVID-19 diagnosis has adverse mental
4 health effects both in previously healthy people and those with pre-existing mental health
5 conditions.¹ About 1 in 5 people with COVID-19 have experienced poorer mental health
6 within 90 days of diagnosis.² It is estimated that the pandemic will lead to new or additional
7 mental health support for up to 10 million people in England (around 20% of the
8 population).³ Evidence from previous studies illustrated that the virus and periods of
9 lockdown have deteriorated population mental health and disproportionately worsened the
10 mental health burden for certain population subgroups including those with pre-existing
11 mental health conditions.^{4 5} People with existing psychiatric diagnoses have reported increased
12 symptoms and poorer access to services and support leading to relapse and suicidal behaviour.¹ At
13 the same time, they are more likely to contract COVID-19 than people with no history of poor
14 mental health. The adaptations required to enable the delivery of mental healthcare during
15 this period of extended infection-control measures could have been disproportionately
16 detrimental to those now living with mental health conditions that have arisen since the
17 first UK-wide lockdown began in March 2020. Difficulties attending review appointments in
18 person and closure of support services are likely to have impacted all those in, or in need of,
19 active treatment.⁶ The unequal impact of the pandemic is likely to further entrench and
20 exacerbate the existing structural inequalities in mental health among people with pre-
21 existing mental health conditions before COVID-19, and the mental health services provided
22 have failed to meet their increasing demand for mental health conditions during the time of
23 the COVID-19 pandemic.⁷

24 The NHS has set up a long term plan to improve mental healthcare services that are widely
25 regarded as being under-resourced.⁸ However, for people with mental health conditions,
26 there is an incomplete picture of the impact of the pandemic on the pattern of mental
27 health services. Despite bringing current service inadequacies to the forefront, the
28 pandemic could provide an opportunity to rethink conventional approaches to mental
29 health services planning to meet patients' needs. For example, remote community
30 treatment and support has long been suggested but has not previously been implemented
31 widely because of barriers and challenges from both healthcare staff and service users.
32 Since the onset of the pandemic, the situation has changed.⁹ Similarly, the threshold for
33 hospital admission for mental illness varies between individuals and requires continuous
34 adaptation over time. Therefore, learning from health service changes throughout the
35 pandemic, and their consequences for people's physical and mental health is vital to inform
36 practical policy solutions for integrated service recovery and effectively plan services that
37 reach those with the greatest need.

38 The WHO recommends rapid review methods as an efficient approach to provide rapid but
39 relevant and contextualised evidence to the health decision-makers when there are time,
40 resources or other logistical constraints.¹⁰⁻¹⁵ We will conduct a rapid review to synthesise
41 the evidence on changes in mental health services because of the COVID-19 pandemic and
42 their impacts in high-income countries.^{16 17}

1 **METHODS AND ANALYSIS**

2 A rapid review will be undertaken to provide an evidence base supporting the
3 recommendation of mental health services and identify areas where the evidence base is
4 lacking, and future research is required. The review will be guided by the Cochrane guidance
5 for rapid reviews¹⁸. Preferred Reporting Items for Systematic reviews and Meta-Analyses
6 extension for Rapid Reviews guidance¹⁹ will be followed for reporting.

7 This protocol has been developed in advance of the review to improve the transparency and
8 quality of the methods to help reduce bias and enhance the reproducibility of the results.
9 This has been registered with the PROSPECT CRD42022306923.

10 **Eligibility**

11 **Type of studies**

12 Peer-reviewed quantitative or qualitative empirical studies describing the setting, problems
13 addressed, resource requirements, aim, service components, provider, method of delivery,
14 objective and subjective effects of changes to mental health services in response to COVID-
15 19 will be included. Included studies will be those undertaken in an Organisation for
16 Economic Co-operation and Development (OECD) country²⁰, to ensure a degree of
17 commonality in the health system and socioeconomic and demographic context.

18 **Type of participants**

19 People aged 18 or over experiencing mental health conditions as described by NHS²¹ who
20 were in need of mental health support during the pandemic.

21 **Type of health services**

22 Interventions, services and models of care delivered in response to COVID-19 to provide
23 support for adults with mental health conditions will be included.

24 **Type of outcome measures**

25 Primary outcomes are objective measures and subjective effects of changes, efficacy or use
26 of a service by mental health patients. Secondary outcomes are changes in knowledge,
27 attitudes or satisfaction of service users and/or professionals and health inequalities.

28 **Search methods**

29 Cochrane CENTRAL, MEDLINE, Embase and PsycInfo will be searched for from 2019 to the
30 present. A search strategy has been developed for MEDLINE with support from an
31 independent information specialist, using a range of keywords and subject headings
32 representing COVID-19, mental health and low- and middle-income countries
33 (supplementary material 1). This will be used to inform the detailed search strategy for
34 other databases. Reference lists and citation indexes of relevant studies will also be
35 examined. Only OECD studies published in or after 2019 and in the English language (no
36 resource available for translation) will be searched.

37 **Selection of studies**

38 Studies identified from databases will be exported to EndNote X9²² for deduplication. Study
39 titles of abstracts will be screened independently according to the selection criteria. Any
40 results that are inconclusive at the initial screen will be included and considered at full-text

1 screening. All full-text papers will be screened independently by two researchers (GY and
2 YF). Any discrepancies will be resolved by discussion and consensus. Where there is a
3 disagreement between two reviewers, a third researcher (DC) will be consulted to reach a
4 consensus.

5 **Data extraction**

6 A data extraction sheet will be designed to capture information including author's first
7 name, publication date, setting, study design, sample size, mental health conditions,
8 characteristics of participants, service components, service provider, method of delivery,
9 resources required, outcome measures and main study results. GY will extract all the data.
10 YF will check for accuracy and completeness through random double-extraction of 10% of
11 included studies. Where a study appears to have multiple citations, original authors will be
12 contacted for clarification. All information from multiple citations will be used if no replies
13 are received.

14 **Quality assessment and quality control**

15 This is a rapid review, so no quality assessment will be conducted. This can be done
16 retrospectively if time and resources allow.

17 The following steps will be taken to ensure quality control for the searching, screening, data
18 extraction, and coding process. GY will conduct screening and data extraction following pre-
19 determined inclusion criteria and data extraction framework. For articles that are retrieved
20 and full text saved, YF will check 10% of the coding to ensure they meet the screening
21 criteria. Where there is a disagreement between two reviewers, a third researcher (DC) will
22 be consulted to reach a consensus. Synthesis of each outcome will be conducted by GY and
23 independently revised by YF.

24 **Data synthesis and analysis**

25 A tabulated and narrative synthesis of the results will be undertaken following current best
26 practice²³⁻²⁵ to conduct synthesis systematically and transparently. It will focus on the
27 mental health services, mechanisms and their impact on health outcomes. A logic model will
28 be produced to present context, service provision and outcomes. Possible unintended
29 adverse outcomes will also be reported. Also, a list of interventions/services/models of care
30 delivered to people with mental health conditions will be grouped as "Do", "Don't" and
31 "Don't know" based on the strength of the evidence on effectiveness and acceptability.

32 If data is available, outcomes of studies will be synthesised according to characteristics of
33 study participants, for example, deprived communities, ethnic minorities, to produce
34 evidence on health inequalities that is likely to have been exacerbated during the pandemic.

35 **PATIENT AND PUBLIC INVOLVEMENT**

36 This study has been designed and developed in consultation with two public members (one
37 with lived experience), to ensure their input on the study design. They both read and
38 commented on the review summary, search strategies, eligibility and plans to synthesise
39 data and dissemination strategies. They valued the potential impact of this review on NHS
40 plans for mental health post-pandemic. It has been agreed that the process of this rapid
41 review will be presented to both members for their further comments.

1 **ETHICS AND DISSEMINATION**

2 As this rapid review will only consider published literature, no ethics approval is needed.
3 Dissemination will be led by the research team and supported by the public member and
4 the wider project advisory group. Results of this review will contribute to reports which will
5 be produced and shared with the National Institute for Health Research (NIHR) Three
6 Research Schools and NIHR Applied Research Collaboration (ARC) North East and North
7 Cumbria (NENC). The findings will be published in peer-reviewed journals and a plain study
8 summary will be disseminated to people receiving mental health care, groups and forum
9 that the project public members are connected, practitioners and commissioners. An
10 abstract will be prepared for academic conferences such as the Society for Academic
11 Primary Care Annual Conference.

12 **CONCLUSION**

13 This rapid review protocol proposes methods for rapid evidence synthesis. A list of
14 interventions/services/models of care delivered to people with mental health problems
15 during the pandemic will be grouped as “Do”, “Don’t” and “Don’t know” based on the
16 strength of the evidence on effectiveness and acceptability. The proposed rapid review is
17 expected to have an immediate impact on the care of patients by increasing awareness of
18 the impacts associated with service changes and factors enabling high-quality services, in
19 both affected individuals and their service providers. Evidence and practise
20 recommendations produced will be useful in supporting the NHS strategy for planning
21 mental health services in a post-pandemic future, and also in highlighting the needs of
22 service users, potentially the needs of underserved populations. In addition, it will identify
23 important knowledge gaps to help guide research direction.

24 Accepting the potentially increased risk of biases introduced by rapid review methods is a
25 compromise for many rapid review protocols.²⁶ The decision to restrict the evidence review
26 to OECD studies, research published in English and a limited number of databases, and not
27 to undertake a formal quality appraisal exercise will increase the risk of bias and the
28 likelihood that relevant research may be missed or excluded. To help mitigate the
29 limitations of the proposed study design, strategies such as regular team meetings
30 throughout and publishing the protocol in advance of conducting the review have been
31 employed. Furthermore, a quality appraisal may be undertaken retrospectively at a later
32 date for a more comprehensive assessment of the evidence.

1 **AUTHORS' CONTRIBUTIONS**

2 GY and YF conceived the study idea and design. GY drafted the initial manuscript and all
3 authors reviewed the manuscript and provided input to the final version.

4 **FUNDING STATEMENT**

5 This project/research/work is supported by the National Institute of Health Research (NIHR)
6 Three Research Schools (award number: MHF018) and the NIHR Applied Research
7 Collaboration (ARC) for the North East and North Cumbria (NENC) (grant number not
8 applicable). The NIHR is the nation's largest funder of health and care research and provides
9 the people, facilities and technology that enables research to thrive. [Find out more about](#)
10 [the NIHR](#).

11 **COMPETING INTERESTS STATEMENT**

12 The authors declare that they have no competing interests.

1 REFERENCES

- 2 1. Moreno C, Wykes T, Galderisi S, et al. How mental health care should change as a consequence of
3 the COVID-19 pandemic. *The Lancet Psychiatry* 2020;7(9):813-24. doi: 10.1016/S2215-
4 0366(20)30307-2
- 5 2. Rogers JP, Chesney E, Oliver D, et al. Psychiatric and neuropsychiatric presentations associated
6 with severe coronavirus infections: a systematic review and meta-analysis with comparison
7 to the COVID-19 pandemic. *The Lancet Psychiatry* 2020;7(7):611-27. doi:
8 [https://doi.org/10.1016/S2215-0366\(20\)30203-0](https://doi.org/10.1016/S2215-0366(20)30203-0)
- 9 3. O'Shea N. Covid-19 and the nation's mental health: Forecasting needs and risks in the UK. 2020
- 10 4. Pierce M, Hope H, Ford T, et al. Mental health before and during the COVID-19 pandemic: a
11 longitudinal probability sample survey of the UK population. *The Lancet Psychiatry*
12 2020;7(10):883-92.
- 13 5. Iob E, Steptoe A, Fancourt D. Abuse, self-harm and suicidal ideation in the UK during the COVID-19
14 pandemic. *The British Journal of Psychiatry* 2020;217(4):543-46.
- 15 6. Anderson G, Frank JW, Naylor CD, et al. Using socioeconomic factors to counter health disparities arising
16 from the covid-19 pandemic. *Bmj* 2020;369
- 17 7. Bambra C, Riordan R, Ford J, et al. The COVID-19 pandemic and health inequalities. *J Epidemiol
18 Community Health* 2020;74(11):964-68.
- 19 8. Alderwick H, Dixon J. The NHS long term plan: British Medical Journal Publishing Group, 2019.
- 20 9. Shore JH, Schneck CD, Mishkind MC. Telepsychiatry and the coronavirus disease 2019 pandemic—
21 current and future outcomes of the rapid virtualization of psychiatric care. *JAMA psychiatry*
22 2020;77(12):1211-12.
- 23 10. Tricco AC, Langlois E, Straus SE, et al. Rapid reviews to strengthen health policy and systems: a
24 practical guide: World Health Organization 2017.
- 25 11. McIntosh HM, Calvert J, Macpherson KJ, et al. The healthcare improvement Scotland evidence
26 note rapid review process: providing timely, reliable evidence to inform imperative decisions
27 on healthcare. *JBI Evidence Implementation* 2016;14(2):95-101.
- 28 12. Tricco AC, Antony J, Zarin W, et al. A scoping review of rapid review methods. *BMC medicine*
29 2015;13(1):1-15.
- 30 13. Polisena J, Garritty C, Kamel C, et al. Rapid review programs to support health care and policy
31 decision making: a descriptive analysis of processes and methods. *Systematic reviews*
32 2015;4(1):1-7.
- 33 14. Featherstone RM, Dryden DM, Foisy M, et al. Advancing knowledge of rapid reviews: an analysis
34 of results, conclusions and recommendations from published review articles examining rapid
35 reviews. *Systematic reviews* 2015;4(1):1-8.
- 36 15. Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and associated
37 methodologies. *Health information & libraries journal* 2009;26(2):91-108.
- 38 16. Brunier A, Drysdale C. COVID-19 disrupting mental health services in most countries, WHO
39 survey. *World Heal Organ* 2020
- 40 17. Byrne A, Barber R, Lim CH. Impact of the COVID-19 pandemic—a mental health service
41 perspective. *Progress in Neurology and Psychiatry* 2021;25(2):27-33b.
- 42 18. Garritty C, Gartlehner G, Kamel C, et al. Cochrane Rapid Reviews Interim Guidance from the
43 Cochrane Rapid Reviews Methods Group. March 2020, 2020.
- 44 19. Page MJ, McKenzie JE, Bossuyt PM, et al. Updating guidance for reporting systematic reviews:
45 development of the PRISMA 2020 statement. *Journal of Clinical Epidemiology* 2021;134:103-
46 12.
- 47 20. Reibling N, Ariaans M, Wendt C. Worlds of Healthcare: A Healthcare System Typology of OECD
48 Countries. *Health Policy* 2019;123(7):611-20. doi:
49 <https://doi.org/10.1016/j.healthpol.2019.05.001>
- 50 21. NHS. Mental health conditions: NHS; 2022 [Available from: [https://www.nhs.uk/mental-
51 health/conditions/](https://www.nhs.uk/mental-health/conditions/) accessed 08 Feb 2022.

- 1 22. EndNote [program]. EndNote X9 version. Philadelphia, PA: Clarivate, 2013.
- 2 23. Campbell M, Katikireddi SV, Sowden A, et al. Improving Conduct and Reporting of Narrative
- 3 Synthesis of Quantitative Data (ICONS-Quant): protocol for a mixed methods study to
- 4 develop a reporting guideline. *BMJ open* 2018;8(2):e020064.
- 5 24. Popay J, Roberts H, Sowden A, et al. Guidance on the conduct of narrative synthesis in systematic
- 6 reviews. *A product from the ESRC methods programme Version 2006*;1:b92.
- 7 25. Campbell M, McKenzie JE, Sowden A, et al. Synthesis without meta-analysis (SWiM) in systematic
- 8 reviews: reporting guideline. *bmj* 2020;368
- 9 26. Ganann R, Ciliska D, Thomas H. Expediting systematic reviews: methods and implications of rapid
- 10 reviews. *Implementation Science* 2010;5(1):1-10.