

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

Mesoamerican Nephropathy in Central Panama

Karen Courville MD^{1,2}, Norman Bustamante MD^{1,2}, Bárbara Hurtado MD³, Maydelin Pecchio MD^{1,4}, Clarissa Rodríguez MD⁵, Virginia Núñez-Samudio MD, MPH, PhD^{1,6} and Iván Landires MD, PhD^{1,7,}*

- ¹ Instituto de Ciencias Médicas, Las Tablas, Los Santos, Panamá
- ² Unidad de Hemodiálisis, Departamento de Nefrología, Hospital Dr. Gustavo N. Collado, Caja de Seguro Social, Chitré, Herrera, Panamá
- ³ Departamento de Epidemiología, Hospital Dr. Gustavo N. Collado, Caja de Seguro Social, Chitré, Herrera, Panamá
- ⁴ Unidad de Infectología, Hospital Dr. Gustavo Nelson Collado, Caja de Seguro Social, Chitré, Herrera, Panamá
- ⁵ Departamento de Infecciones Nosocomiales, Hospital Dr. Gustavo N. Collado, Caja de Seguro Social, Chitré, Herrera, Panamá
- ⁶ Sección de Epidemiología, Departamento de Salud Pública, Región de Salud de Herrera, Ministerio de Salud, Chitré, Herrera, Panamá
- ⁷ Hospital Joaquín Pablo Franco Sayas, Ministerio de Salud, Las Tablas, Los Santos, Panamá

Correspondence to

Iván Landires, MD, PhD, Instituto de Ciencias Médicas, Las Tablas, Los Santos, Panama. PO Box 0710-00043. Phone: +507 65937727
Email address: ilandires@institutodecienciasmedicas.org
ivanlandires@yahoo.es

Running headline: Mesoamerican Nephropathy in Panama

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86

Abstract.

Background. In the last three decades, there has been an increase in the frequency of patients diagnosed with chronic kidney disease of nontraditional causes (CKDnt) in the Mesoamerican region. A region with an increased frequency of patients with chronic kidney disease (CKD) has been identified in central Panama. The present study aims to characterize the clinical presentation of patients with CKDnt in an understudied population of the central region of Panama and compare them with patients with traditional CKD (CKDt).

Methods. A retrospective descriptive study was conducted in a nephrology reference hospital in the central provinces of Herrera and Los Santos, comparing a group of 15 patients with CKDnt to 91 patients with CKDt. Sociodemographic variables, personal history, laboratory parameters, and of renal ultrasound were compared.

Results. CKDnt is more common among relatively younger male patients who engage in strenuous work activities at high temperatures. CKDnt is not associated with type 2 diabetes mellitus or chronic hypertension, as it is the case in patients with CKDt. Renal atrophy and hyperuricemia are significant clinical markers of CKDnt.

Conclusion. To our knowledge, this is the first study in Panama and one of the few in Central America and the world to address the clinical presentation of patients with CKDnt compared to patients with CKDt. Because CKDnt remains asymptomatic for a long time, early detection is important, and efforts should be directed at halting disease progression at an early stage. Current evidence can also inform policies addressing occupational and environmental risk factors associated with CKDnt.

Keywords: chronic kidney disease; Panama; nontraditional; traditional; mesoamerican nephropathy; clinical presentation

87

88

89 **Background**

90 Chronic kidney disease (CKD) belongs to the group of noncommunicable diseases and, according to
91 estimates from the Global Burden of Disease group, produced a 19.6% increase in the disability-adjusted
92 life years (DALYs) and has been associated with 4% of deaths worldwide from 2005 to 2015,
93 representing 2.2 million deaths per year [1,2].

94 In 60% of patients, the cause of CKD is identified. Among the traditional causes of CKD are diabetes
95 mellitus, essential hypertension, and obesity, followed by a minority secondary to immunological
96 diseases, nephrolithiasis, and genetic conditions [3]. Approximately 25% of adult patients with CKD may
97 have a family history of this pathology. In 10% of patients, the cause of CKD cannot be identified, which
98 is classified as unknown or nontraditional cause (CKDnt) [4]. For several years, an increase in the
99 frequency of diagnoses of CKDnt from agricultural areas of Central America (El Salvador, Nicaragua,
100 Guatemala, and Costa Rica) has been identified, which is why it has been denominated Mesoamerican
101 nephropathy. This pathology has been presented mainly in young male patients with irreversible
102 impairment of kidney function. It has been proposed that among the probable causes of CKDnt would be
103 chronic exposure to pesticides, working conditions with exposure to high temperatures and dehydration,
104 chronic use of anti-inflammatory drugs, high consumption of alcohol and tobacco. These risk factors also
105 contribute with the high morbidity and mortality of CKD, since exposure increases with the diagnoses
106 advanced-stage CKD [5,6].

107 The Pan-American Health Organization defined CKDnt as impaired kidney function with a glomerular
108 filtration rate (GFR) below 60 mL/min/m² [7]. in absence of predisposing factors for traditional CKD
109 (namely, type 2 diabetes mellitus, essential hypertension, heart disease, urinary tract malformations,
110 immunological and congenital diseases. In addition, the diagnosis of CKDnt includes kidney damage
111 defined by structural abnormalities (i.e., renal atrophy without obstructive pattern) or abnormality in the
112 urinary sediment as a marker of kidney damage. There may be exposure to occupational risk factors or
113 living in a risk area [8].

114 In Panama, since 2014, an increase in patients who meet the criteria for CKDnt have been reported in the
115 provinces of Coclé, Herrera, and Los Santos [9]. In 2017, Panamanian health authorities started the
116 Information System for the Epidemiological Surveillance of CKD to establish mandatory notification of
117 all patients diagnosed with CKD in all its stages and to identify associated risk factors [10]. The system
118 aims to diagnose and establish the real prevalence of CKD in Panama to inform public policies for
119 promotion, prevention, and treatment of CKD. The 2017 Preventive Health Census of the Ministry of
120 Health estimated the national prevalence of CKD at 3.24% [11].

121 The objective of this study is to characterize the risk factors and clinical phenotype of patients diagnosed
122 with CKDnt from the provinces of Herrera and Los Santos in the central region of Panama and to
123 compare them with patients with CKDt from the same geographic area.

124 **Methods**

125 We conducted a retrospective, descriptive study with the patients records at the Nephrology Department
126 of the Dr. Gustavo N. Collado Hospital. This hospital is the nephrology reference center for Central
127 Panama (provinces of Herrera and Los Santos, serving a population estimated in 2012 at 214,539
128 inhabitants) [12].

129 Records between January 1 and December 31, 2018 and a minimum of two evaluations by Nephrology at
130 least 3 months apart during the study period were included. Records also reported at least two urinalyses
131 and blood samples. The patient record was also the data source for sociodemographic variables, personal
132 history, laboratory results, and renal ultrasound evaluation.

133 Data was extracted into MS Excel (The Microsoft Corporation; Redmond, WA) and the clean dataset was
134 exported to Stata v. 11.0 (StataCorp, LLC; College Station, TX) for data analyses. Analyses included
135 descriptive statistics. Fisher's exact test was used to compare proportions and the Mann-Whitney U test to
136 compare medians, setting alpha at 0.05 for statistical significance when comparing the frequencies of the
137 CKDt and CKDnt groups.

138 **Results**

139 The chart review identified 224 records of patients diagnosed with stage 3 and 4 CKD during 2018. After
140 two reviews of the inclusion criteria, 106 patient files were included (Figure 1).

141 Of the 106 patient charts included, 66% (70) were male and 34% (36) were female, with a median age of
142 68.8 years. Forty-five percent of patients were 70 years or older. The median weight was 73 kg, the
143 median height was 1.61 m and the median body mass index (BMI) was 27.7 kg/m².

144 In the occupation evaluation, 22% of the patients were retired, followed by 20% agricultural workers,
145 19% household administrators, 14% unemployed, and 13% practiced other professions. Eighty percent of
146 the patients had a history of essential hypertension, 30% type 2 diabetes mellitus, 19% cardiovascular
147 disease, 15% hyperuricemia, and 10% cerebrovascular disease.

148 Of the total of 106 patients included with a diagnosis of CKD, 14% ($n=15$) met the definition of CKDnt,
149 while 86% ($n=91$) were diagnosed with CKDt. **Table 1** summarizes the relevant clinical, history,
150 ultrasound, and laboratory data for both groups.

151 A statistically significant predominance of young men was observed in the group of patients with CKDnt
152 compared to the group with CKDt. Additionally, workers in agriculture or transportation presented with
153 significantly ($P<0.001$) higher frequencies of CKDnt than with CKDt. Although not statistically
154 significant ($P=0.09$), patients with CKDnt tend to exhibit a lower BMI than patients with CKDt. The
155 personal history of type 2 diabetes mellitus, essential hypertension, and cardiovascular disease were
156 significantly less frequent among patients with CKDnt than among those with CKDt. Of note, ultrasound-
157 confirmed that renal atrophy was significantly more frequent ($P<0.001$) among patients with CKDnt than
158 among patients with CKDt. Patients with CKDnt show significantly lower levels of glucose and higher
159 levels of uric acid in the blood.

160 **Discussion**

161 In Central American countries, the cases of CKDnt have been increasing during the last decades.
162 However, it has not yet been possible to establish a single causal factor, but rather different agents that
163 can produce chronic kidney damage leading to CKDnt. Various causes have been proposed, including
164 chronic occupational exposure to high temperatures and repeated dehydration due to frequent episodes of
165 heat stress, exposure to pesticides with nephrotoxic effect, contamination of water with heavy metals,
166 lifestyle risk factors (e.g., alcoholism, smoking, and diet), all of which could have a significant impact on
167 the pathophysiology of CKD [13-15].
168

169 In 2015 in El Salvador, a country where the epidemiology of CKDnt has been extensively studied, the
170 regional prevalence ranged from 7.1% in the Paracentral region; 5.2% in the Eastern region; and 2% in
171 the Central region [16], and that depending on the community and the work activity carried out by the
172 inhabitants, prevalence peaked at 18% in regions near the coast [17]. In Nicaragua, regional prevalence
173 ranged between 8% and 10% [18]. In Panama, a 2014 study estimated the prevalence of CDK requiring
174 dialysis in the province of Coclé at 40 per 100,000 population towards the north of the province, but up to

175 200 per 100,000 population towards the south of the province. The southern part of Coclé's economy is
176 characterized by sugar cane and rice agriculture. In this study, the prevalence of CKDnt could not be
177 determined, but it was the first study that reported an increase in CKD prevalence in a specific region of
178 the country [9].

179
180 For greater robustness of the data and to confirm chronicity of the renal function, patient charts should
181 have included at least two measurements of serum creatinine instead of only one [19]. Most previously
182 published studies have measured creatinine only once. The diagnosis of CKD required the calculation of
183 the GFR estimate after achieving a stable creatinine value, at least 3 months apart, to establish the degree
184 of chronicity and be able to rule out acute renal failure [20].

185
186 In the group of patients with CKDnt, a male predominance was observed compared to the group with
187 CKDt. This has also been observed in other countries, such as El Salvador (78%) [21], Nicaragua (77%)
188 [22] and Costa Rica (70%) [5]. The median age was lower in the CKDnt group (55 years), compared with
189 a median age of 68 years in the CKDt group. The latter is similar to what was observed in the study group
190 of agricultural communities in El Salvador, where the age of the patients was an average of 45 years [21].
191 The increased frequencies of patients with diagnoses of CKDnt have been observed mainly in some
192 defined areas in Central America. Particularly, areas heavy on agricultural employment are involved,
193 where young labor is hired to perform difficult tasks under the tropical sun. These factors may explain, at
194 least in part, the younger age, lower BMI, and higher prevalence in males presenting with CKDnt. In turn,
195 these patients did not present with chronic diseases typically associated with CKDt, such as type 2
196 diabetes mellitus and essential hypertension [23].

197
198 This study showed that a significantly higher proportion of patients with CKDnt presented with a history
199 of agricultural work, compared to patients in the CKDt group (60% vs. 15%, respectively; $P < 0.001$). The
200 central region of Panama has the largest area with crops of corn in the country, with over 26,000 hectares
201 of land per year. The province of Herrera is the fourth in sugarcane production nationwide [24]. The
202 patients in this study belong to the central provinces of Herrera and Los Santos, mostly land at sea level
203 dedicated to agriculture, in a peninsular area that frequently exhibits the highest temperatures in the
204 country of up to 35°C, especially in the dry season between December and April [25]. It has also been
205 described that in the region of Central America and Panama there has been an increase in temperature of
206 up to 1°C between 2010 and 2015 and that this has been important in the central provinces of Herrera and
207 de Los Santos [25]. Predictive models of temperature increase have been proposed, up to 15% in the next
208 decades in the central region of Panama [25]. This could contribute to the increase in cases of CKDnt
209 compared to the CKDt causes in recent years, as a consequence of continuous dehydration due to
210 exposure to high temperatures during strenuous working hours, as has been studied in other groups of
211 sugarcane farmers in Central America [26,27].

212
213 Twenty percent of patients with CKDnt worked as drivers of road transport vehicles or heavy equipment.
214 Studies have associated poor diet, continuous hours of work, lack of adequate hydration, and exposure to
215 high temperatures during the sunniest and hottest hours of the day while driving in freight transport
216 drivers [28]. This phenomenon has also been described for workers in a block factory [29]. Thus, this
217 pathology would not be exclusive to agriculture and transportation, but in occupations where there are
218 sustained exposure to high temperatures with risk of dehydration.

219
220 Glycosuria without hyperglycemia and HbA1c levels greater than 6.5% can be found in less than 3% of
221 patients with CKDnt [23], probably secondary to the pathophysiological damage caused by CKDnt and
222 not to the presence of type 2 diabetes mellitus [18]. In the present study, no patient in the CKD group
223 presented with type 2 diabetes mellitus, compared to 34% in the CKDt group. A trend towards increased
224 glucose associated with the higher frequency of type 2 diabetes mellitus was observed in the CKDt group,
225 in which poor control could help the progression to worsening of CKDt.

226

227 A higher frequency of essential hypertension was evidenced in the group of patients with CKDt compared
228 to the group of patients with CKDnt. Previous studies have found that arterial hypertension is rare in
229 patients with CKDnt and it is highly prevalent in CKDt [30]. Hypertensive nephrosclerosis is a chronic
230 disease that gradually and progressively produces chronic kidney involvement [30,31]. The finding of
231 mild hypertension in some patients with CKDnt (33%) in our study could be due to the fact that it is a
232 consequence of CKD and not the cause, as it has been previously suggested [19]. The presence of mild
233 hypertension of less than 5 years of evolution, associated with advanced chronic kidney disease, in the
234 absence of other risk factors, should lead us to suspect CKDnt.

235
236 In this study, uric acid values showed a significant elevation above the normal value in the group of
237 patients with CKDnt compared to CKDt. In dehydration states, fructose is produced endogenously from
238 glucose, leading to uric acid generation, inflammation, and fibrosis in the kidney [32]. Furthermore,
239 asymptomatic hyperuricemia has been associated with the development of mild hypertension, which,
240 when treated and normalized, improves blood pressure control [33]. The present study supports the concept
241 that hyperuricemia is a biomarker of CKDnt. More studies are necessary to determine its role in the
242 diagnosis, evolution and prognosis of CKDnt.

243
244 In the evaluation of the imaging studies, by renal ultrasound, 87% of the patients diagnosed with CKDnt
245 presented with a decrease in the cortico-medullary relationship with increased echogenicity and renal
246 atrophy. This translates into renal morphological damage observed in early stages of asymptomatic
247 patients. Renal ultrasound studies in El Salvador have found an increase in echogenicity of up to 95% and
248 a decrease in the cortex-medulla ratio in up to 82% of patients with CKDnt [21]. The present study
249 supports the concept that renal atrophy is a clinical marker of CKDnt. More studies are necessary to
250 determine its role in the diagnosis, evolution and prognosis of CKDnt.

251
252 Patients with CKD frequently present proteinuria mostly due to glomerular involvement, hypertension, or
253 diabetic nephropathy. In the present study, there were no significant differences proteinuria levels
254 between patients with CKDnt and CKDt. This may be due to the fact that patients included in this study
255 had moderate stages (3 and 4), and not advanced stages of CKD. In patients with CKDnt, low proteinuria
256 (<1 g/dL) was observed. Renal biopsy studies in patients with CKDnt have shown that the
257 histopathological damage is not due to damage to the glomeruli but rather to tubulointerstitial
258 nephropathy [34,35]. This could explain the difference in the presentation of proteinuria that they have
259 these patients in advanced stages of CKD.

260
261 There are various studies with patients affected by CKDnt, but there are few prevalence studies with
262 progression to end-stage CKD [23,36]. In the present study, in the group of patients with CKDt, five
263 patients progressed to stage 5 CKD, requiring initiation renal replacement therapy. At follow-up, no
264 patient in the CKDnt group progressed to terminal illness. In early stages of kidney involvement by an
265 external agent, recovery depends on exposure time, as has been observed in drug-caused interstitial
266 nephropathies. When the causative agent is withdrawn early, the patient can recover due to the fact that
267 the damage does not involve the glomeruli. When evaluating histopathological studies, it is important to
268 point out that damage in CKDnt mainly affects the tubules and interstices and not the glomeruli [37,38],
269 so that early identification could delay the progression to terminal disease. Due to the lack of knowledge
270 on prevalence, the information on the frequency of patients diagnosed with CKDnt who progress to renal
271 replacement therapy is still uncertain in Central America [39]. This entity has different names depending
272 on the region, either as CKDnt, CKD of unknown cause, or Mesoamerican nephropathy, which further
273 complicates its epidemiological traceability [15]. Thus, it is essential not only to diagnose it, but also
274 consistently code it as the same diagnosis.

275
276 There are environmental exposures that interact with gene function. Several occupational-environmental
277 risk factors predisposing to CKDnt were already discussed, with a possible summative effect and
278 sustained for a certain time that would lead to kidney involvement. However, not all people who live in
279 the same geographic region with the same gender and occupational risk factors will develop kidney

280 involvement leading to CKDnt. Thus, so it is also believed that genetic and epigenetic factors could play
281 an important role in the etiology of CKDnt [40]. It is possible that individual genetic susceptibilities in a
282 triggering environment could induce development and/or worsening of renal involvement [41].
283 Longitudinal follow-up studies of patients in early stages that could include the genetic and epigenetic
284 component that can also contribute to the identification of biomarkers in patients at risk and help prevent
285 kidney involvement [42].

286
287 A limitation of this study is the small sample size in the case of patients with CKDnt. Thus, the
288 interpretation of its implications, including considerations on risk factors, should be prudent. Another
289 limitation is that not all patients had renal ultrasound available to compare the results. Despite its
290 limitations, this study makes important contributions to the understanding of CKDnt since to our
291 knowledge it is the first published study in Panama and one of the few in Central America to address the
292 clinical presentation of patients with early stage CKDnt compared to patients with CKDt.
293

294 **Conclusions**

295 CKDnt is more common in younger-aged male patients than in patients with CKDt. CKDnt is diagnosed
296 in areas where agricultural activity predominates; however, it is not the only work activity to which this
297 condition has been associated, but more generally it is related to long-term strenuous work and high
298 temperatures that could lead to dehydration.

299
300 Because CKDnt remains asymptomatic for a long time, early detection is important in patients with
301 elevated creatinine values, without comorbidities such as diabetes and hypertension, in which it is
302 important to identify whether there are other risk factors for occupational and environmental exposure and
303 efforts should be directed at halting the progression of the disease at an early stage. The early detection of
304 patients with CKDnt could have an important effect in preventing progression to terminal CKD.

305
306 It is important to provide information and education to the population that works in chronic exposure to
307 high temperatures. Likewise, work should be done on the generation of public policies so that employers
308 are aware of the risk factors and allow the necessary rest and hydration times during the working hours of
309 their employees at risk.

310
311 New prospective follow-up studies are important to allow the study of genetics, epigenetics, the
312 identification of biomarkers and a better understanding of the pathophysiology, as well as the better
313 therapeutic management of patients at risk or diagnosed with CKDnt.

314 315 **Abbreviations**

316
317 CKD: chronic kidney disease; CKDnt: nontraditional chronic kidney disease; CKDt: traditional chronic
318 kidney disease; DALYs: Disability-adjusted life years; GFR: glomerular filtration rate; BMI: body mass
319 index.

320
321 **Acknowledgments:** The authors would like to acknowledge Humberto López Castillo, MD, PhD for his
322 review of the manuscript. Iván Landires is a member of the Sistema Nacional de Investigación (SNI),
323 which is supported by Panama's Secretaría Nacional de Ciencia, Tecnología e Innovación (SENACYT).

324
325 **Authors' contributions:** Conceptualization, K.C., N.B., V. N-S., and I.L.; methodology, K.C., N.B.,
326 B.H., M.P., C. R., V. N-S., and I.L.; software, K.C., and I.L.; validation, K.C., N.B., V. N-S., and I.L.;
327 formal analysis, K.C., N.B., V. N-S., and I.L.; investigation, K.C., N.B., B.H., M.P., C. R., V. N-S., and
328 I.L.; resources, K.C., V. N-S., and I.L.; data curation, K.C., N.B., B.H., M.P., C. R., V. N-S., and I.L.;

329 writing—original draft preparation, K.C., N.B., V. N-S., and I.L.; writing—review and editing, I.L.;
330 visualization, K.C., N.B., V. N-S., and I.L.; supervision, K.C., and I.L.; project administration, K.C., and
331 I.L.; funding acquisition, K.C., V. N-S., and I.L.; All authors have read and agreed to the published
332 version of the manuscript.

333 **Funding:** No funding was received for this study.

334 **Availability of data and materials:** All data generated or analysed during this study are included in this
335 published article.

336 **Ethics approval and consent to participate:** The study was conducted according to the guidelines of the
337 Declaration of Helsinki, and approved by the Interinstitutional Ethics Committee of the Social Security
338 Fund and the National Directorate for Teaching and Research (DENADOI-SIBI-008-2020). The need for
339 informed consent was waived by the Interinstitutional Ethics Committee of the Social Security Fund and
340 the National Directorate for Teaching and Research (DENADOI-SIBI-008-2020). No further
341 administrative permissions were needed to access the raw data used in this study. The data used in this
342 study were anonymized before use.

343 **Consent for publication:** Not applicable.

344 **Competing interests:** The authors declare that they have no competing interests.

345

346 **References**

347

- 348 1. GBD 2015 DALYs and HALE Collaborators (2016). Global, regional, and national disability-
349 adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE),
350 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* (London,
351 England), 388(10053), 1603–1658. [https://doi.org/10.1016/S0140-6736\(16\)31460-X](https://doi.org/10.1016/S0140-6736(16)31460-X)
- 352 2. Thomas, B., Matsushita, K., Abate, K. H., Al-Aly, Z., Ärnlöv, J., Asayama, K., Atkins, R., Badawi,
353 A., Ballew, S. H., Banerjee, A., Barregård, L., Barrett-Connor, E., Basu, S., Bello, A. K., Bensenor,
354 I., Bergstrom, J., Bikbov, B., Blosser, C., Brenner, H., Carrero, J. J., ... Global Burden of Disease
355 Genitourinary Expert Group (2017). Global Cardiovascular and Renal Outcomes of Reduced GFR.
356 *Journal of the American Society of Nephrology : JASN*, 28(7), 2167–2179.
357 <https://doi.org/10.1681/ASN.2016050562>
- 358 3. Go, A. S., Yang, J., Tan, T. C., Cabrera, C. S., Stefansson, B. V., Greasley, P. J., Ordonez, J. D., &
359 Kaiser Permanente Northern California CKD Outcomes Study (2018). Contemporary rates and
360 predictors of fast progression of chronic kidney disease in adults with and without diabetes mellitus.
361 *BMC nephrology*, 19(1), 146. <https://doi.org/10.1186/s12882-018-0942-1>
- 362 4. Groopman, E. E., Marasa, M., Cameron-Christie, S., Petrovski, S., Aggarwal, V. S., Milo-Rasouly,
363 H., Li, Y., Zhang, J., Nestor, J., Krithivasan, P., Lam, W. Y., Mitrotti, A., Piva, S., Kil, B. H.,
364 Chatterjee, D., Reingold, R., Bradbury, D., DiVecchia, M., Snyder, H., Mu, X., ... Gharavi, A. G.
365 (2019). Diagnostic Utility of Exome Sequencing for Kidney Disease. *The New England journal of*
366 *medicine*, 380(2), 142–151. <https://doi.org/10.1056/NEJMoa1806891>
- 367 5. Luyckx, V. A., Tuttle, K. R., Garcia-Garcia, G., Gharbi, M. B., Heerspink, H., Johnson, D. W., Liu,
368 Z. H., Massy, Z. A., Moe, O., Nelson, R. G., Sola, L., Wheeler, D. C., & White, S. L. (2017).
369 Reducing major risk factors for chronic kidney disease. *Kidney international supplements*, 7(2), 71–
370 87. <https://doi.org/10.1016/j.kisu.2017.07.003>
- 371 6. Wesseling, C., Crowe, J., Hogstedt, C., Jakobsson, K., Lucas, R., & Wegman, D. (2013).
372 Mesoamerican nephropathy: report from the first international research workshop on Mesoamerican
373 nephropathy. San Jose. SALTRA. International Research Workshop.
374 <http://hdl.handle.net/11056/8584>
- 375 7. Arreola-Guerra, J. M., Rincón-Pedrero, R., Cruz-Rivera, C., Belmont-Pérez, T., Correa-Rotter, R., &
376 Niño-Cruz, J. A. (2014). Performance of MDRD-IDMS and CKD-EPI equations in Mexican

- 377 individuals with normal renal function. *Nefrología : publicación oficial de la Sociedad Española*
378 *Nefrología*, 34(5), 591–598. <https://doi.org/10.3265/Nefrologia.pre2014.Jun.12538>
- 379 8. Lozier, M., Turcios-Ruiz, R. M., Noonan, G., & Ordunez, P. (2016). Chronic kidney disease of
380 nontraditional etiology in Central America: a provisional epidemiologic case definition for
381 surveillance and epidemiologic studies. *Revista Panamericana de Salud Pública*. 2016; 40, 294-300
- 382 9. Rodríguez F, Cuero C, Delgado E, Camargo I, Tuñon R. Diagnóstico de la Enfermedad Renal
383 Crónica y Factores de Riesgo Asociados en Áreas Seleccionadas de la Provincia de Coclé, Panamá.
384 . *Rev Med Panamá*. 2014; 34(1): 31-38. <https://doi.org/10.37980/im.journal.rmdp.2019810>
- 385 10. Ministerio de Salud MINSA. (16 de abril de 2018). *Gaceta Oficial*. Recuperado el 29 de enero de
386 2019: Disponible en http://www.gacetaofical.gob.pa/pdfTemp/28505_A
- 387 11. Ministerio de Salud MINSA. (4 de septiembre de 2017). Available in:
388 <http://www.css.gob.pa/cnsp.html>
- 389 12. Instituto Nacional de Estadística y Censo, XI Censo de Población y VII de Vivienda de Panamá: Año
390 2020. Contraloría General de la República de Panamá.
391 [https://www.inec.gob.pa/publicaciones/Default3.aspx?ID_PUBLICACION=499&ID_CATEGORIA](https://www.inec.gob.pa/publicaciones/Default3.aspx?ID_PUBLICACION=499&ID_CATEGORIA=3&ID_SUBCATEGORIA=10)
392 [=3&ID_SUBCATEGORIA=10](https://www.inec.gob.pa/publicaciones/Default3.aspx?ID_PUBLICACION=499&ID_CATEGORIA=3&ID_SUBCATEGORIA=10)
- 393 13. Correa-Rotter, R., Wesseling, C., & Johnson, R. J. (2014). CKD of unknown origin in Central
394 America: the case for a Mesoamerican nephropathy. *American journal of kidney diseases : the*
395 *official journal of the National Kidney Foundation*, 63(3), 506–520.
396 <https://doi.org/10.1053/j.ajkd.2013.10.062>
- 397 14. Correa Rotter, R. & García Trabanino, R. (2018). Nefropatía mesoamericana: Una nueva enfermedad
398 renal crónica de alta relevancia regional. *Acta Médica Grupo Ángeles*, 16(S1), 16-22
- 399 15. González-Quiroz, M., Pearce, N., Caplin, B., & Nitsch, D. (2018). What do epidemiological studies
400 tell us about chronic kidney disease of undetermined cause in Meso-America? A systematic review
401 and meta-analysis. *Clinical kidney journal*, 11(4), 496–506. <https://doi.org/10.1093/ckj/sfx136>
- 402 16. Orantes, C. M., Herrera, R., Almaguer, M., Brizuela, E. G., Hernández, C. E., Bayarre, H., Amaya, J.
403 C., Calero, D. J., Orellana, P., Colindres, R. M., Velázquez, M. E., Núñez, S. G., Contreras, V. M., &
404 Castro, B. E. (2011). Chronic kidney disease and associated risk factors in the Bajo Lempa region of
405 El Salvador: Nefrolempa study, 2009. *MEDICC review*, 13(4), 14–22
- 406 17. Ferreiro, A., Álvarez-Estévez, G., Cerdas-Calderón, M., Cruz-Trujillo, Z., Mena, E., Reyes, M.,
407 Sandoval-Díaz, M., Sánchez-Polo, V., Valdés, R., & Ordúñez, P. (2016). Confirmed clinical case of
408 chronic kidney disease of nontraditional causes in agricultural communities in Central America: a
409 case definition for surveillance. *Revista panamericana de salud publica = Pan American journal of*
410 *public health*, 40(5), 301–308
- 411 18. Fischer, R., Vangala, C., Mandayam, S., Chavarria, D., García-Trabanino, R., Garcia, F., Garcia, L.
412 L., & Murray, K. O. (2018). Clinical markers to predict progression from acute to chronic kidney
413 disease in Mesoamerican nephropathy. *Kidney international*, 94(6), 1205–1216.
414 <https://doi.org/10.1016/j.kint.2018.08.020>
- 415 19. Pearce, N., Caplin, B., Gunawardena, N., Kaur, P., O'Callaghan-Gordo, C., & Ruwanpathirana, T.
416 (2018). CKD of Unknown Cause: A Global Epidemic?. *Kidney international reports*, 4(3), 367–369.
417 <https://doi.org/10.1016/j.ekir.2018.11.019>
- 418 20. KDIGO (2013). 2012 Clinical practice guideline for the evaluation and management of chronic
419 kidney disease. *Kidney Int Suppl*, 3, 1-150
- 420 21. Herrera, R., Orantes, C. M., Almaguer, M., Alfonso, P., Bayarre, H. D., Leiva, I. M., Smith, M. J.,
421 Cubias, R. A., Torres, C. G., Almendárez, W. O., Cubias, F. R., Morales, F. E., Magaña, S., Amaya,
422 J. C., Perdomo, E., Ventura, M. C., Villatoro, J. F., Vela, X. F., Zelaya, S. M., Granados, D. V., ...
423 Aparicio, C. E. (2014). Clinical characteristics of chronic kidney disease of nontraditional causes in
424 Salvadoran farming communities. *MEDICC review*, 16(2), 39–48
- 425 22. Raines, N., González, M., Wyatt, C., Kurzrok, M., Pool, C., Lemma, T., Weiss, I., Marín, C., Prado,
426 V., Marcas, E., Mayorga, K., Morales, J. F., Aragón, A., & Sheffield, P. (2014). Risk factors for
427 reduced glomerular filtration rate in a Nicaraguan community affected by Mesoamerican
428 nephropathy. *MEDICC review*, 16(2), 16–22

- 429 23. Lunyera, J., Mohottige, D., Von Isenburg, M., Jeuland, M., Patel, U. D., & Stanifer, J. W. (2016).
430 CKD of Uncertain Etiology: A Systematic Review. *Clinical journal of the American Society of*
431 *Nephrology : CJASN*, 11(3), 379–385. <https://doi.org/10.2215/CJN.07500715>
- 432 24. Contraloría General de la República. Instituto Nacional de Estadística y Censo Panamá en cifras:
433 Años 2012-2016, INEC 2018.
434 <http://www.contraloria.gob.pa/inec/archivos/P8551PanamaCifrasCompleto.pdf>
- 435 25. Candanedo, M., Villarreal, D. G., & Bernal, S. (2020). Uso de registros de temperatura máxima
436 promedio de las estaciones meteorológicas de ETESA, para la creación de mapas de temperatura
437 mediante el uso de programa ArcGIS. *Revista de Iniciación Científica*, 6(2), 9-14.
438 <https://doi.org/10.33412/rev-ric.v6.2.2878>
- 439 26. Peraza, S., Wesseling, C., Aragon, A., Leiva, R., García-Trabanino, R. A., Torres, C., Jakobsson, K.,
440 Elinder, C. G., & Hogstedt, C. (2012). Decreased kidney function among agricultural workers in El
441 Salvador. *American journal of kidney diseases : the official journal of the National Kidney*
442 *Foundation*, 59(4), 531–540. <https://doi.org/10.1053/j.ajkd.2011.11.039>
- 443 27. García-Trabanino R, Jarquín E, Wesseling C, Johnson R et al. Estrés térmico, deshidratación y
444 función renal en cortadores de caña de azúcar: estudio pre y post jornada.
445 https://www.researchgate.net/publication/280312210_Estres_termico_deshidratacion_y_funcion_renal_en_cortadores_de_cana_de_azucar_Estudio_pre_y_post_jornada_de_trabajadores_en_riesgo_de_Nefropatia_Mesoamericana
446 Nefropatia Mesoamericana
- 448 28. Berrones Sanz, L. D., Cano Olivios, P., Sánchez Partida, D., Martínez Flores, J. L. (2018). Lesiones,
449 enfermedades y accidentes de trabajo de los conductores del autotransporte de carga en México. *Acta*
450 *Univ*, 28(3), 47-55. <https://doi.org/10.15174/au.2018.1946>
- 451 29. Gallo-Ruiz, L., Sennett, C. M., Sánchez-Delgado, M., García-Urbina, A., Gámez-Altamirano, T.,
452 Basra, K., Laws, R. L., Amador, J. J., Lopez-Pilarte, D., Tripodis, Y., Brooks, D. R., McClean, M.
453 D., Kupferman, J., Friedman, D., Aragón, A., González-Quiroz, M., & Scammell, M. K. (2019).
454 Prevalence and Risk Factors for CKD Among Brickmaking Workers in La Paz Centro,
455 Nicaragua. *American journal of kidney diseases : the official journal of the National Kidney*
456 *Foundation*, 74(2), 239–247. <https://doi.org/10.1053/j.ajkd.2019.01.017>
- 457 30. Freedman, B. I., & Sedor, J. R. (2008). Hypertension-associated kidney disease: perhaps no
458 more. *Journal of the American Society of Nephrology : JASN*, 19(11), 2047–2051.
459 <https://doi.org/10.1681/ASN.2008060621>
- 460 31. Mann, J., & Hilgers, K. (2017). Clinical features, diagnosis, and treatment of hypertensive
461 nephrosclerosis. *UpToDate*.
462 <https://somepomed.org/articulos/contents/mobipreview.htm?30/11/30911>
- 463 32. Nakagawa, T., Johnson, R. J., Andres-Hernando, A., Roncal-Jimenez, C., Sanchez-Lozada, L. G.,
464 Tolan, D. R., & Lanaspá, M. A. (2020). Fructose Production and Metabolism in the Kidney. *Journal*
465 *of the American Society of Nephrology : JASN*, 31(5), 898–906.
466 <https://doi.org/10.1681/ASN.2019101015>
- 467 33. Kanbay, M., Jensen, T., Solak, Y., Le, M., Roncal-Jimenez, C., Rivard, C., Lanaspá, M. A.,
468 Nakagawa, T., & Johnson, R. J. (2016). Uric acid in metabolic syndrome: From an innocent
469 bystander to a central player. *European journal of internal medicine*, 29, 3–8.
470 <https://doi.org/10.1016/j.ejim.2015.11.026>
- 471 34. López-Marín, L., Chávez, Y., García, X. A., Flores, W. M., García, Y. M., Herrera, R., Almaguer,
472 M., Orantes, C. M., Calero, D., Bayarre, H. D., Amaya, J. C., Magaña, S., Espinoza, P. A., & Serpas,
473 L. (2014). Histopathology of chronic kidney disease of unknown etiology in Salvadoran agricultural
474 communities. *MEDICC review*, 16(2), 49–54
- 475 35. Elinder, C. G. & Wernerson, A. (2020). Mesoamerican nephropathy. *UptoDate*.
476 <https://www.uptodate.com/contents/mesoamerican-nephropathy>
- 477 36. Gómez-Sánchez, V. (2018). Nefropatía mesoamericana. *Revista Brasileira de Medicina do*
478 *Trabalho*, 16, 35-36. <https://doi.org/10.5327/z16794435201816s1018>
- 479 37. Caravaca-Fontán F, González E, Fernández-Juárez G. Nefropatías Intersticiales. En: Lorenzo V,
480 López Gómez JM (Eds). *Nefrología al día*. <https://www.nefrologiaaldia.org/252>

- 481 38. Raghavan, R., & Shawar, S. (2017). Mechanisms of Drug-Induced Interstitial Nephritis. *Advances in*
482 *chronic kidney disease*, 24(2), 64–71. <https://doi.org/10.1053/j.ackd.2016.11.004>
483 39. Yaqub F. (2014). Kidney disease in farming communities remains a mystery. *Lancet (London,*
484 *England)*, 383(9931), 1794–1795. [https://doi.org/10.1016/s0140-6736\(14\)60867-9](https://doi.org/10.1016/s0140-6736(14)60867-9)
485 40. Perez-Gomez, M. V., Martin-Cleary, C., Fernandez-Fernandez, B., & Ortiz, A. (2018). Meso-
486 American nephropathy: what we have learned about the potential genetic influence on chronic
487 kidney disease development. *Clinical kidney journal*, 11(4), 491–495.
488 <https://doi.org/10.1093/ckj/sfy070>
489 41. Wuttke, M., & Köttgen, A. (2016). Insights into kidney diseases from genome-wide association
490 studies. *Nature reviews. Nephrology*, 12(9), 549–562. <https://doi.org/10.1038/nrneph.2016.107>
491 42. Friedman, D., & Luyckx, V. A. (2019). Genetic and Developmental Factors in Chronic Kidney
492 Disease Hotspots. *Seminars in nephrology*, 39(3), 244–255.
493 <https://doi.org/10.1016/j.semnephrol.2019.02.002>
494

495

Table 1. Relevant Clinical and Paraclinical Data of patients with CKDt and CKDt.

Characteristics	CKDt (n=91)	CKDt (n=15)	P value
Median age, years	68.8	55.4	<0.001
Median weight, kg	73.7	70.7	0.50
Median height, m	1.6	1.64	0.14
Median BMI, kg/m ²	28.03	25.85	0.09
Gender			0.016
Men	56 (61)	14 (93)	
Women	35 (38)	1 (7)	
Agricultural workers	14 (15)	9 (60)	<0.001
Transportation workers	0 (0)	3 (20)	<0.001
Personal history of:			
Type 2 diabetes mellitus	31 (34)	0 (0)	0.02
Hypertension	80 (88)	5 (33)	<0.001
Cardiovascular disease	20 (22)	0 (0)	0.04
Cerebrovascular disease	11 (12)	0 (0)	0.32
Obesity	1 (1)	0 (0)	0.84
CKD	17 (19)	2 (13)	0.80
Obstructive uropathy	9 (10)	0 (0)	0.40
Hyperuricemia	12 (13)	4 (27)	0.37
Renal ultrasound ¹			
Normal findings	28 (31)	2 (13)	0.16
Renal atrophy	36 (50)	13 (87)	<0.001
Hydronephrosis	7 (10)	0 (0)	0.27
Agenesis	1(1)	0 (0)	0.68
Urinalysis			
Erythrocyturia >3 RBCs/field	8 (9)	1 (7)	0.78
Proteinuria >3+	15 (16)	1 (7)	0.32
Median blood chemistry values			
Glucose 1, mg/dL	105	97.5	0.07
Glucose 2, mg/dL	111	96.5	0.04
Glycated hemoglobin A1c, %	6.71	5.005	0.11
Creatinin 1, mg/dL	1.85	2.14	0.10
Creatinin 2, mg/dL	1.89	2.17	0.13
Blood urea nitrogen, mg/dL	26.8	23.5	0.31
Uric acid 1, mg/dL	6.8	7.8	0.04
Uric acid 2, mg/dL	7	8.1	0.09
Sodium, mEq/L	139	138	0.60
Potassium, mEq/L	4.5	4.4	0.55
Calcium, mg/dL	9.75	9.7	0.72
Triglycerides, mg/dL	171	167	0.90
Total cholesterol, mg/dL	182	188	0.68


HDL cholesterol, mg/dL	51.4	45.7	0.70
LDL cholesterol, md/dL	108.8	101.7	0.59

496
497
498
499
500
501

¹ Only 72 of the CKDt files had renal ultrasound on file.

BMI: body mass index, CKDt: traditional chronic kidney disease, CKDnt: non-traditional chronic kidney disease, RBC, red blood cells; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

Figure 1. Patient chart selection flow diagram.



ARI: acute renal insufficiency; CKD: chronic kidney disease; GFR: glomerular filtration rate; KT: kidney transplant; PKD: polycystic kidney disease.