

A community-based ANP-led integrated oncology care model for adults receiving oral anticancer medication-protocol for a pilot study.

Richmond J.P., RN, RANP, RNP, BSc (Hons), PGDip, DNSc¹; Johnston A., RN, RNT, BSc (Hons) MSc, Kelly M.G., R.N, MSc¹, Murphy, P.J., BSc CS, HDip Psych, PG Dip Stats, PhD Psych²; Gillespie P., PhD Economics³. Murphy A.W., MB; FRCGP; MICGP; MD²

Affiliations

¹Letterkenny University Hospital, Donegal Ireland; ²HRB Primary Care Clinical Trials Network Ireland, Discipline of General Practice, National University of Ireland, Galway; ³Ireland Health Economics & Policy Analysis Centre (HEPAC), Institute for Lifecourse & Society (ILAS), National University of Ireland, Galway. Ireland.

Abstract

Oncology has been undergoing a profound transition in the last ten years or more with the increased usage in oral anti-cancer medication (OAM). Approximately 25% of all anti-cancer medication is now designed for oral use and this is likely to increase prospectively. Oral anti-cancer medications have the potential to alleviate capacity issues in cancer treating units as patients receive their treatment at home. There remains however a requirement for safe and efficient assessment and monitoring but this does not necessarily require them to repeatedly attend a hospital

day unit. Therefore the opportunity exists to transition this cohort to a community-based setting to be assessed by a specialist such as an Advanced Nurse Practitioner (ANP) in nurse-led clinics. Having an OAM assessment closer to their home would be more convenient to the patient. Furthermore, this could help alleviate hospital capacity issues which were brought into sharp focus with the onset of the COVID-19 pandemic and the use of nurse-led clinics are promoted in the aims of the current healthcare system reform process in Ireland.

Within the context of the Irish healthcare system reform and the COVID-19 pandemic this protocol will outline a collaboration between an Oncology Department in Letterkenny University Hospital in Ireland and the National University of Ireland, Galway aimed to develop and pilot a community-based Advanced Nurse Practitioner-led integrated oncology care model for adults receiving OAM. Phase 1 of this two-phase study commenced in September 2020 and comprised a scoping review, a benchmarking exercise and a qualitative analysis of relevant stakeholders. This protocol paper presents a pilot to be undertaken in phase 2 as OAM care is transitioned to an ANP-led community-based model, which is a radical shift for oncology care in Ireland. The pilot outlined will provide data that will identify potential refinements to the model and address specific uncertainties about a definitive trial.

Corresponding author: janicep.richmond@hse.ie

Keywords: cancer, care, management, oral anticancer medications, community, pilot, protocol

Introduction

The emergence of oral anti-cancer medication (OAM) has revolutionised oncology care, particularly within the last decade and these treatments are being approved at a record setting pace (Meier et al., 2018). Oral anticancer medications are a sub-set of systemic anti-cancer treatments (SACT) which are administered enterally with a narrow therapeutic window and a unique mechanism of action (National Cancer Control Programme (NCCP), 2018; National Cancer Institute, 2020). OAMs have the same benefits and risks as SACT given intravenously in terms of positive disease outcomes, treatment-related toxicities, and potential for serious medication errors leading to patient harm (NCCP, 2018). While these medications are convenient and often preferred or even requested by patients, it shifts the responsibility for medication management from the oncology healthcare professionals (HCPs) to the patient. Consequently, there are concerns regarding adherence and patients' management of toxicities or adverse effects (Greer et al., 2016; Hammond et al., 2012; NCCP, 2018; Paoletta et al., 2018; Wood, 2012).

Due to the associated safety challenges, ongoing specialised assessment and monitoring of patients receiving OAMs is essential; the organisation of this is usually not much different to parenteral anti-cancer treatment. In Ireland, historically the practice has been that the patient generally attends the Oncology or Haematology Day Unit for a dedicated health assessment, which includes review of recent laboratory tests and/or other investigative results (Hammond et al, 2012; Department of Health (DOH), 2017; NCCP, 2018). This practice, while necessary for the aforementioned reasons, when performed in the context of increasing incidence and

treatment of cancer, has contributed to hospital overcrowding (Sung et al, 2021; DOH, 2017).

The Impact of the COVID-19 Pandemic

Such pre-existing hospital capacity issues were exacerbated with the advent of the COVID-19 pandemic and the requirement for social distancing. Since March 2020, internationally, HCPs have been forced to reassess routine practices (Volger & Lightner, 2020) and to reduce unnecessary patient hospital visits (Cucinotta & Vanelli, 2020). This has been especially pertinent to individuals receiving cancer treatment due to potential immunosuppression and concomitant risk of infection (Leung et al, 2020).

At the Letterkenny University Hospital (LUH) Oncology Department (institution of authors JR, AJ, and MGK), the onset of the pandemic required the immediate transfer of the caseload of individuals receiving OAMs from the Oncology Day Unit to the Advanced Nurse Practitioner (ANP) Oncology Clinic for ongoing assessment. This was in order to maintain COVID restrictions and free up capacity for patients receiving intravenous SACT. This is an example of a successful nurse-led clinic. Nurse-led clinics have emerged internationally as an ideal means to achieve improved organisation and efficiency in health services (Randall et al., 2017, House of the Oireachtas, 2018; Torrens et al, 2020), with associated high levels of patient satisfaction (Molassiotis et al., 2020, Linedale et al., 2020), and potential cost-savings (Thompson & McNamara, 2021).

The Shift to Primary Care

While there is a requirement for patients receiving OAMs to have on-going assessment and monitoring, this does not necessarily require them to repeatedly attend a hospital day unit. The opportunity exists to transition this cohort to a community-based setting.

Transitioning to a community-based setting would align with the Government of Ireland's Sláintecare reform (House of the Oireachtas, 2018), which aims to shift healthcare from hospitals to the community. Indeed, internationally there is a transformative vision to shift patient care from acute to primary care, with a universal consensus regarding the crucial and central role for primary care (Calnan et al., 2006; Richmond et al., 2021).

The Current Study

With this background of Irish health service reform and the COVID-19 pandemic, a collaboration between the LUH Oncology Department and the National University of Ireland, Galway (NUIG) aimed to develop and pilot a community-based ANP-led integrated oncology care model for adults receiving OAM.

Phase 1 of this two-phase study commenced in September 2020 and comprised of three elements: a scoping review, a benchmarking exercise and a qualitative analysis of relevant stakeholders. The scoping review aimed to determine current clinical management practices for the ongoing assessment and monitoring of patients receiving OAM (Richmond, 2021a). The authors reported that in the studies reviewed that there was a unanimous endorsement of a dedicated OAM clinic as a

means to achieve improved care for this patient cohort. The scoping review also identified a range of best-practice recommendations for clinical practice, which were collated alongside existing national and international guidelines to develop a benchmarking tool. A member of the research team, who was not actively involved in the day-to-day OAM clinic, used this tool to retrospectively examine the standard of existing care of the ANP-led hospital-based OAM clinic. This benchmarking exercise demonstrated safe practice yet noted scope for improvement, especially with regard to documentation of patient education and standardisation of OAM prescription writing. The third piece of work in Phase 1 was a qualitative analysis of relevant stakeholders' (nurses, doctors, pharmacists, general practitioners, and service users) perceptions of a community-based ANP-led integrated model for OAM care as well as identification of the infrastructural and cultural supports required for this to happen.

The research team presented the results from the three elements of Phase 1 to an advisory panel of local and national experts in October 2021. The outcome was an agreement that a pilot study should proceed to examine the feasibility of a community-based ANP-led integrated oncology care model for adults receiving OAMs. The results of Phase 1 have directly informed the development of the protocol for Phase 2, presented in the remainder of this article.

Phase 2: Pilot Study Protocol

A pilot study is essential to assess the acceptability and feasibility of interventions and study protocols (Skivington et al, 2021), and can prevent the cost of a failed trial

(Arain et al, 2010). Richards & Halberg (2015) identified that commonly researchers fail to adequately develop and pilot phases of trials. As the authors propose transforming OAM care to an ANP-led community-based model, which is a radical shift for oncology care in Ireland, a pilot will provide data that will identify potential refinements to the model and address specific uncertainties about a definitive trial. We use the framework of 14 questions identified by Shanyinde et al (2011) that should be asked and answered in a pilot trial. These are presented in Table 1.

Study setting

The on-going management of patients who are established on OAMs will, for the pilot study, be performed at a medical practice in a primary care building in Letterkenny, Co. Donegal, Ireland. This venue is external to the LUH Oncology Department, and has availability for a maximum of two half-days per week to accommodate the OAM clinic. Once patients are stable on their OAM treatment (which generally is after two cycles of treatment) the Medical Oncology team formally refers the patient to the ANP oncology for on-going assessment and monitoring for the third cycle onwards. A cycle of OAM treatment is usually 3-4 weeks in duration. The approach of reviewing the patient in the hospital oncology department for their first two cycles (or more should they be unstable on treatment initially) facilitates any major toxicities to be managed within the acute hospital setting at their review and maximises patient safety. Virtual assessments can be facilitated at subsequent cycles when patients are stable on OAMs if this is clinically acceptable and preferable to the patient (see Figure 1, Appendix 1, and Appendix 5).

Recruitment

All participants will be recruited from the cohort of patients being cared for by the ANP in the hospital-based OAM clinic. Patient eligibility criteria for this existing clinic are:

- Over 16 years of age.
- Have an oncology diagnosis (i.e. solid tumour).
- Be under the care of a Consultant Medical Oncologist.
- Have been prescribed OAMs

The cyclic nature of OAMs requires that patients be assessed on certain days of the week, reflective of the day they commenced treatment. This is also necessary in order to adhere to the NCCP guidelines

(<https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/oral-anti-cancer-medicines>). Patients whose assessments are currently fall on the two half-days per week that the study setting is available will be approached to participate in-person by the ANP Oncology. They will be provided with a participant information sheet, which will be explained. Furthermore, they will be advised that should they not wish to participate, their care will continue with the ANP oncology and their review will be facilitated on designated assessment day in the usual hospital setting. If they are content to proceed, they will be given a consent form to read, sign at home and return in a stamped addressed envelope to another member of the research team.

There are no specific days for certain treatments or patient groups to attend the ANP Oncology in the hospital-based clinic, therefore the sample enrolled in this pilot study

should represent a cross-section of the entire OAM patient population within the locality.

Primary Outcomes

The primary outcomes of the pilot will be: patient safety, patient satisfaction and the cost of an ANP-led OAM clinic in a community setting. A secondary outcome will be staff satisfaction with this new clinic.

Sample size

The pilot duration will be 4 months (January to April 2022), a timeframe adequate to include those who are receiving OAMs on an 8 or 12 weekly assessment schedule. Considering the availability of the study setting for the pilot, approximately 7-8 patient visits to the ANP will take place weekly which equates to about one third of the entire OAM weekly patient workload. Over the period of the 4-month pilot, this equates to over 100 virtual and/or face-to-face visits.

Data Collection

Patient Safety

Patient safety in an ANP-led OAM clinic will be measured twofold. Firstly, there will be a repetition of, and comparison with, the benchmarking exercise for patient assessments & OAM prescriptions, which was initially performed in Phase 1 of this study (Figure 1 and Appendix 1). This consists of a Microsoft Excel tool with 82 items with the options of yes/no/not applicable. It will be completed by one of the authors (MGK or AJ) performing a retrospective chart review of 20 patient visits which

includes 20 OAM prescriptions written by the ANP. This data analysis and comparison to Phase 1 will be carried out locally by the LUH staff (AJ).

Secondly, a Microsoft® Excel® spreadsheet (Appendix 2) has been developed to capture data on the patient safety aspects including patient attendance, waiting times, timeliness of review, drug toxicities requiring medical review +/- admission, ad hoc queries from General Practitioner (GP)/community pharmacists or from patient/family since last review and clinical incidents/near misses/adverse events. This data will be captured retrospectively from a review of the patient records.

Patient Satisfaction

Patient satisfaction with the ANP-led OAM clinic will be measured using the 7-item EORTC-OUT-PATSAT7 instrument (Bredart et al, 2018), developed to capture the perceptions of patients with cancer regarding the service and organisation of care they receive. Ratings can range from 'fair' to 'excellent', providing scores ranging from 7 to 35, with a higher score indicating higher satisfaction (Appendix 3).

Cost

The economic analysis will consist of two components. Firstly, the economic burden of illness falling on patients will be estimated using the EuroQol EQ-5D-5L survey instrument, to capture health-related quality of life (Balestroni & Bertolotti, 2012 ; Schwenkgenks & Matter-Walstra, 2016) and a Health Economic Analysis questionnaire created specifically for this study (Appendix 4). This questionnaire includes a series of questions that will capture healthcare service usage, out-of-pocket expenses, and employment or education participation related impacts on

participants (Herdman et al, 2011). Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Quality of life research*, 20(10), 1727-1736. This questionnaire will be administered to individuals at the end of pilot OR at the patient's last visit using an allocated unique identification number (UID) to ensure participant confidentiality.

Secondly, a cost analysis will be undertaken to identify, measure and value the healthcare resources required to implement the proposed ANP-led OAM model of care in clinical practice. Resource items required for the OAM clinic will be identified by the study team. This process will be directly informed by the process flow diagram, which was developed, presented and endorsed by the advisory panel of local and national experts at the end of Phase 1 of this study (Appendix 5). Unit cost data will be identified and applied to calculate individual resource costs and the total cost per patient of implementing the proposed ANP-led OAM model of care will be estimated. This data will be extrapolated to present estimates of the healthcare budget impact to implementing the proposed model of care at local, regional and national levels.

A brief clinical workflow analysis will be carried out to identify the time required for certain aspects of care via self-reported activity tracking (Lopetegui et al, 2014).. The ANP Oncology will record the time to complete toxicity assessment, physical examination and care plan (to include OAM prescription writing). To minimise onerous timing of care in a busy clinical context, data will be collected 1 day per week for the duration of the pilot and data subsequently extrapolated from this.

Staff Satisfaction

Staff satisfaction will be measured through an anonymised online questionnaire with a single item: “How acceptable is the intervention to you”. Responses will be measured with a 5-point Likert scale ranging from ‘totally unacceptable’ to ‘perfectly acceptable’. Space will be given for “any further comments” and contact details of a non-clinical research team member will be provided should respondents want to arrange a confidential telephone interview to provide further feedback or comment. The staff satisfaction questionnaire link will be administered to all nursing, medical, pharmacy and administrative staff who currently work in the hospital Day Unit. This questionnaire will be administered in Month 4 of the 4-month pilot.

Data Analysis

Methodological issues will be the central focus of this pilot, using the list of 14 methodological issues that need to be examined in feasibility research (Shanyinde et al, 2011). How each of these questions will be specifically addressed in this pilot is outlined in table 2. To direct a future definitive trial, possible solutions to identified feasibility issues will be generated using a process for decision-making after pilot and feasibility trials (ADePT) (Bugge et al, 2013).

Data Management and Protection

All collected data will be treated confidentially in line with the seven key principles related to the processing of personal data General Data Protection Regulation requirements (Health Service Executive, 2019). All consented patients and staff will be allocated a UID and this will be used on all study data collection forms. Their

identification will be recorded on a password-protected document saved on a server accessed only by one of the members of the research team (JR).

At the end of the pilot, data from the questionnaires will be posted using registered mail to NUIG staff to be inputted to SPSS for statistical analysis. Similarly, at the end of the pilot, the Microsoft Excel data will be emailed to this same research team member for eventual uploading and analysis. Using the UID on all patient data obtained adheres the overriding principle of General Data Protection regulation (GDPR) in accordance to ethical approval obtained. All of the hard copy questionnaires will then be shredded by NUIG staff.

Study Status

At the time of submission of this study protocol, (14.02.2022) recruitment of patients has begun and consents are being obtained.

Discussion

Development of a protocol is an essential component in the process of performing a pilot trial and provides a detailed method for data collection that can be adhered to throughout. The objectives of this pilot study will be different from those of the future definitive study as it is not designed to consider the effectiveness of the intervention but rather determine whether a definitive trial can be done (Lancaster, 2015). Piloting of a new intervention for use in a definitive trial ensures that the methodological approach taken in any future research is robust and feasible and is an important step of the development process (Lancaster, 2015). Using the ADePT framework to guide

decision-making (Bugge et al, 2013), completion of this pilot will evaluate whether a definitive future trial is feasible.

Declarations

Ethical approval has been granted by Letterkenny University Hospital's Research Ethics Committee (17/11/2021). Informed consent was obtained before recruiting participants.

Competing interests: The authors declare that they have no competing interests.

Funding: This work was supported by The Irish Cancer Society (Grant number CNRA19RIC) in collaboration with the Health Research Board (HRB) the National Cancer Control Programme (NCCP) and the Office of the Nursing and Midwifery Services Director (ONMSD). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

J.R. Supervision, conceptualisation, methodology, project administration, writing—review & editing; M.G.K conceptualisation, methodology, writing—review & editing; A.J. methodology, writing—review & editing; P.J.M. conceptualisation, methodology,

review & editing; A.J. methodology, writing—review & editing; P.G. health economics methodology, review & editing, A.W.M. conceptualisation, methodology, review & editing.

ORCID

Janice P Richmond 0000-0002-1086-238X

Patrick J. Murphy 0000-0002-2299-1699

Andrew W. Murphy 0000-0001-5549-8246

Alison Johnston 0000-0001-8219-853X

Paddy Gillespie 0000-0001-6297-9478

Tables and figures


Table 1: Methodological framework

	Methodological issue	Relevant Pilot Data
1.	Did the feasibility/pilot study allow a sample size calculation for the main trial?	Pilot data for the primary outcomes will be used to estimate population-level variation for these outcomes, thus allowing a sample size calculation for a main trial.
2.	What factors influenced eligibility and what proportion of those approached were eligible?	Patient inclusion criteria have been specified above, and all patients attending the clinic should be eligible. Any reasons for non-eligibility will be recorded.
3.	Was recruitment successful?	Researchers will record the number of patients approached, the number eligible for inclusion, the number not eligible, and the number consented.
4.	Did eligible participants consent?	The number of eligible patients that consent will be recorded, as will the number of eligible patients that do not consent. Reasons for non-consent will be recorded.
5.	Were participants successfully randomized and did randomization yield equality in groups?	Not applicable.
6.	Were blinding procedures adequate?	Not applicable.
7.	Did participants adhere to the intervention?	Patient adherence will be measured as: <ul style="list-style-type: none"> • Clinical attendance, to include all cancellations/did not attend episodes/delay in attending. • Questionnaire returns at baseline and at end of treatment

		<p>Staff adherence will be measured as:</p> <ul style="list-style-type: none"> • Referral numbers to ANP Oncology for the duration of the pilot. • Rates of completion of the staff satisfaction questionnaire.
8.	Was the intervention acceptable to the participants?	<p>Acceptability for patients will be measured using the EORTC OUTPAT questionnaire (Appendix 3) at end of treatment.</p> <p>Acceptability for staff will be measured using the staff satisfaction questionnaire and optional interview as described above.</p>
9.	Was it possible to calculate intervention costs and duration?	<p>Relevant patient cost data will be gathered using the Health Economic Analysis Questionnaire developed for this study (see Appendix 4).</p> <p>For staff costs, a cost analysis will be undertaken to identify, measure and value the healthcare resources required to implement the proposed ANP-led OAM model of care in clinical practice. See above.</p>
10.	Were outcome assessments completed?	<p>For patients, questionnaire returns will be recorded at all time points, including reasons for non-returns (if available). Missing data within questionnaires will be recorded.</p> <p>For staff, the benchmarking exercise will identify levels of missing data for patient assessments and for OAM prescriptions.</p> <p>Researchers will also record the number of staff satisfaction questionnaires completed, and reasons for non-completion (if available). Repeat the benchmarking exercise to include retrospective review of the patient assessments and the High-tech OAM prescriptions written by the ANP Oncology (appendix 1).</p>

11.	Were outcomes measured those that were the most appropriate outcomes?	The research team will assess appropriateness of the outcomes measures based on patient adherence, completion rates, variance of the measures, and expert judgement.
12.	Was retention to the study good?	<p>For patients, clinical attendance will be measured to include all cancellations/did not attend episodes/delay in attending.</p> <p>The research staff will also measure the number of patients who had to be referred back to the hospital for medical assessment/personal preference and reason(s) for this.</p>
13	Were the logistics of running a multicenter trial assessed?	This will be a continuous element within the pilot and the tools used to collect data will be assessed as being fit for purpose (by analysing level of missing data or questionnaires not returned).
14	Did all components of the protocol work together?	This will be a continuous element within the pilot and the tools used to collect data will be assessed at the completion of the pilot.

Figure 1: OAM processes (NCCP, 2018)



Appendix 1: Best-practice standard used for bench-marking

Presented in the logical sequence of the Systemic Anti-Cancer Treatment Model of Care identified by NCCP [2018] {Figure 1}

1: Treatment decision

Standard 1.1: Patients who are to commence on an Oral Anti-Cancer Medicine have an assessment to determine their suitability for home management of treatment (NCCP (2018)).

Features of a service meeting this standard should include:

1.1.1 Baseline documented assessment of the patient regarding their capabilities and understanding of OAM.

Standard 1.2: Written consent should be obtained from the patient initially prior to the Medical team commencing treatment with OAM (ONS, 2016; Hall, 2016).

Features of a service meeting this standard should include:

- 1.2.1 Provision of pre-treatment timely and relevant information to patients regarding the proposed treatment to include reason for treatment, goal of therapy and potential side-effects.
- 1.2.2 MOATT© - MASCC tool (Teaching Tool for Patients Receiving Oral Agents for Cancer) used to assess and document suitability and understanding of OAM (NCCP Systemic Anti-Cancer Treatment assessment form)¹.
- 1.2.3 Written consent should be obtained from the patient prior to commencing and a copy available in the patient's notes.
- 1.2.4 Provision of information (1.2.1) and written consent (1.2.2) should be repeated should drug regime change.

¹ <https://www.hse.ie/eng/services/list/5/cancer/profinfo/medonc/sactguidance/extravasation.html>

2: Prescribing

Standard 2.1: Oral Anti-Cancer Medicines should be prescribed to the same safety standards as parenteral Systemic Anti-Cancer Therapy (NCCP (2018)).

Features of a service meeting this standard should include:

- 2.1.1 The first cycle of a course of an Oral Anti-Cancer Therapy should be written by a consultant or registrar based on the consultant's written therapy plan (NCCP (2018)).
- 2.1.2 Subsequent cycles should be written by a consultant, specialist registrar, registrar or Registered Nurse Prescriber following the consultant's written therapy plan (NCCP (2018)).
- 2.1.3 For any dose modification, a new prescription should be written by any of the Health Care Professionals listed in 2.1.2 (NCCP (2018)).
- 2.1.4 Oral Anti-Cancer Medicine prescriptions should be written for one cycle only except when a patient's scheduled review is longer than one cycle (NCCP (2018)).
- 2.1.5 All Oral Anti-Cancer Medicine prescriptions should be verified by an oncology pharmacist (NCCP 2018).

Standard 2.2: OAM prescriptions should include adequate information to facilitate prescription review by the community pharmacist (NCCP, 2018).

Features of a service meeting this standard should include:

- 2.2.1 An OAM prescription should include the following information (unless otherwise stated this is as per NCCP (2018))¹:
 - Patient name
 - Patient address
 - Date of Birth
 - Hospital number
 - Patients consultant
 - Height/weight/Body Surface Area (BSA) (if relevant)

¹ Email orders are considered written orders (ASCO, 2013).

- Diagnosis
- Diagnosis code
- Frequency of patient review
- Allergies/sensitivities/contraindications
- Protocol name
- Protocol code
- Deviations from protocol
- Course number
- Medication name (generic name but biosimilars require trade name also) (LUH¹)
- Medication dose (to include M2 if relevant)
 - Doses rounded to the nearest tablet size (ASCO, 2013)
 - Doses do not include trailing zeros (ASCO, 2013).
 - Use a leading zero for doses (ASCO, 2013).
- Medication administration frequency
- Number of treatment days
- Route (ONS, 2016)
- Total quantity to be dispensed (ONS, 2016).
 - Wording should read to not dispense more than x tablets/capsules (LUH).
- Planned treatment start date and/or cycle prescription commences
- Signature and printed name.
- Clinician registration number
- Date
- Contact details of prescriber
- GP name & address
- Nominated pharmacy & address

¹ LUH is Letterkenny University Hospital (author's place of work)

3: Dispensing

Standard 3.1: To facilitate appropriate counselling of the patient, the first cycle of Oral Anti-Cancer Medicines and the first cycle of a dose adjustment should be dispensed in a hospital setting where possible (NCCP, 2018) (exception mentioned by NCCP, 2018) .

Features of a service meeting this standard should include:

- 3.1.1 Face-to-face assessment of the patient by an Oncology health care professional pre first treatment with OAM.
- 3.1.2 Face-to-face assessment of the patient by an Oncology health care professional in the event of an OAM dose adjustment.

Standard 3.2: Only one cycle of an OAM should be dispensed at a time (NCCP, 2018)¹

Features of a service meeting this standard should include:

- 3.2.1 The prescription should identify the number of days treatment and/or the number of medications to be dispensed (NCCP, 2018).

Standard 3.3: The quantity dispensed should not exceed the number of doses required to complete the cycle (NCCP, 2018)².

Features of a service meeting this standard should include:

- 3.3.1 Days supply limitation rules should be identified on the prescription (Battis et al, 2016).

¹ Community pharmacy dispenses the medication so the treating hospital team have no control over this aspect of care.

² Community pharmacy dispenses the medication so the treating hospital team have no control over this aspect of care.

4: Medication administration

Standard 4.1: Patients prescribed Oral Anti-Cancer Medicines should have access to standardised education to support safe administration, safe handling, and management of side effects (NCCP, 2018)

Features of a service meeting this standard should include:

- 4.1.1 Dedicated initial pre-treatment face-to-face education session with an Oncology Health Care Professional (Muluneh et al, 2018) detailing storage, handling/preparation, administration, disposal of oral chemotherapy, possible drug/drug and drug/food interactions and plan for missed doses and management of side effects of OAM and documented appropriately (ASCO, 2013; NCCP, 2018).
- 4.1.2 MOATT© - MASCC tool (Teaching Tool for Patients Receiving Oral Agents for Cancer) used to frame, check understanding and document the pre-treatment education (NCCP Systemic Anti-Cancer Treatment assessment form)¹.
- 4.1.3 Patients should be informed of the required monitoring arrangements and have access to information in the written protocol and treatment plan from the hospital where treatment was initiated (UKONS, 2010).

¹ <https://www.hse.ie/eng/services/list/5/cancer/profinfo/medonc/sactguidance/extravasation.html>

5: Patient monitoring

Standard 5.1: Patients should not commence their treatment until the results of their monitoring tests are known (NCCP 2018)

Features of a service meeting this standard should include:

5.1.1 Documented review of monitoring tests and contact with the patient prior to commencing a treatment cycle (NCCP, 2018).

Standard 5.2 Patients on Oral Anti-Cancer Medicine treatment should be reviewed by specialist health care professional in an appropriate location at the predefined intervals (NCCP, 2018) prior to every treatment cycle (UKONS, 2010).

Features of a service meeting this standard should include:

5.2.1 Assessment of the patient¹ by an Oncology health care professional in an appropriate location at the predefined intervals (NCCP, 2018) as identified by the NCCP OAM protocols² prior to every treatment cycle (UKONS, 2010).

5.2.2 Completion of patient assessment template specific to an OAM can standardise medication monitoring and facilitate documentation of patient interactions and interventions (Battis et al, 2016).

5.2.3 Completion of patient telephone assessments in a template which allows flexibility in the conduction of the phone follow up and can be performed by a specialist oncology health care professional team member (May et al, 2017).

5.2.4 Patient assessment requires determining adherence (Muluneh et al, 2018)³.

5.2.2 Patient assessment includes but not limited to documentation of the following (laboratory tests required is as per NCCP protocol for specific regime) (unless otherwise stated this is per NCCP SACT assessment forms)⁴:

- Primary diagnosis

¹ Monitoring laboratory tests is not sufficient for patient management pts on OAMs. Close monitoring and follow up of patients on OAMs is crucial to achieve intended therapeutic outcome, improve drug safety and adherence and to reduce drug's adverse events and healthcare cost (Battis et al, 2016)

² <https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/oral-anti-cancer-medicines/>

³ No benefit noted to treatment calendars (Wong et al, 2014)

⁴ <https://www.hse.ie/eng/services/list/5/cancer/profinfo/medonc/sactguidance/extravasation.html>

- Allergies¹
- Frequency of review
- NCCP regime
- Cycle number
- Day number
- Translator present (changed to translator required)
- Has the patient been admitted to hospital or seen their GP since their last treatment?
 - Did the patient receive a discharge prescription?
- Any patient infection control alert/issues?
- Grading of toxicities
- Performance status documented
- Vital Signs/Early Warning Score (EWS) completed
- MST (Malnutrition assessment TOOL) Score completed
- Tumour markers as per medical instruction
- LMP (last menstrual period) documented (if applicable)
- Urinalysis completed (if applicable)
- MOATT© - MASCC tool completed (Teaching Tool for Patients Receiving Oral Agents for Cancer).
 - Assessment of ability to manage treatment at home
- Weight in metrics (ISMP, 2020)
- Is there a need for a follow up phone call?
- Any dose modifications or any cycle delays (UKONS, 2010)
- Is there any change to medication?
- Is there a history of an adverse event on a previous cycle?
- Disease monitoring and any other test/s as directed by the supervising Consultant (NCCP OAM protocols²)
- At defined intervals the following should be checked (specific to each drug) and reviewed as satisfactory prior to proceeding with treatment (NCCP OAM protocols).
- HCG (pregnancy) test completed (if applicable)

¹ Allergies should be checked prior to prescribing and not required at each review (JR)

² <https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/oral-anti-cancer-medicines/>

- FBC, renal and liver profile
- Calcium level (JR)
- Serum glucose
- Blood pressure
- Urinalysis
- Thyroid function tests
- ECG if clinically indicated or if history of cardiac problems
- Total cholesterol and triglycerides
- Other _____

Standard 5.3: Patients to have contact post commencement of cycle 1 of treatment.

Features of a service meeting this standard should include:

- 5.3.1 Follow up phone call by pharmacist to patient after 72 hours of commencing cycle 1 (Calabrese et al, 2015).
- 5.3.2 High risk patients to have a phone call made to them days 8 and 10 (Deluche et al, 2020).
- 5.3.3 All patients to have a phone call made to them days 15 & 30 (Deluche et al, 2020).
 - Contact or visits are beneficial especially in the 1st 2 cycles to improve treatment and symptoms experience (Vidall, 2010¹; Oakley et al, 2014; Wong et al, 2016).

Standard 5.4: Clinical SACT services should provide a 24-hour telephone advice service for patients prescribed oral therapies, with appropriately trained nursing, medical or pharmacy staff handling queries (UKONS, 2010; Deluche et al, 2020).

Features of a service meeting this standard should include:

- 5.4.1 All patients/families to be provided with 24hr/7 day contact details of the hospital at commencement of cycle 1 of OAM.
- 5.4.2 All patients should have access to advice from an appropriately qualified healthcare professional with experience in cancer treatment in the hospital (UKONS 2010).

¹ Vidall (2010) used home visits to monitor patients

Standard 5.5: Information strategies should be developed to standardise communication between hospitals, community pharmacists and GPs (NCCP, 2018) and primary care nurses and GPs require precise and regular information (UKONS, 2010).

Features of a service meeting this standard should include:

- 5.5.1 Adherence to standard 2.2 will facilitate this with community pharmacists.
- 5.5.2 Letter to GP to be sent if patient condition changes, medication changes or the dose changes. If there is stable care for a period of time beyond 6 months then a GP letter should be sent to provide an update (JR).
- 5.5.3 After the final cycle of a treatment course, the records for each patient should include the following (UKONS, 2010) and communicated to GP (JR):
 - Whether the course was completed or not.
 - If the course was not completed or if the planned dose was reduced, the reasons for cessation or reduction.
 - For completed courses of non-adjuvant treatment, a reference to the response.
 - The plans for on-going review and support.

Appendix 2: Microsoft® Excel® Data collection tool (using UID)

Excel Tab 1: Data collected (demographics):

- Gender.
- Age.
- Primary cancer.
- OAM specific regime/medication.
- OAM cycle (for first assessment on pilot).
- Aim of treatment (i.e. curative/palliative).
- Date of cancer diagnosis.
- Date of most recent recurrence.

Excel Tab 2 Data collected (geographical):

- Distance (in kilometres) travelled for OAM assessment.
- Methods of transport (i.e. car/bus).
 - Own or other driver (if car used).
 - Time off work for other driver (if has to be driven)-measured in minutes for round trip.
- Amount spent in parking (if car used).
- Amount spent for travel (if public transport used).
- Closest primary care centre to patient's home (measured in kilometres)
- Closest community hospital to patient's home (measured in kilometres)

Excel Tab 3 Data collected (appointment details):

- Type of assessment (i.e. physical assessment or virtual assessment).
- Did they attend on time (i.e. cancel/did not attend/attended on time).
 - Extent of time in delay (measured in minutes from appointment time) if relevant.
 - Reason for delay in attending if relevant.
- Patient reviewed at correct interval as per protocol.
 - Extent of interval delay (measured in days) if relevant.
 - Reason for any interval delay.
- Waiting time to be reviewed by ANP Oncology (measured in minutes from appointment time).
- Total time spent with ANP (measured in minutes)
 - Specific time (measured in minutes) performing observations, measuring weight and performing toxicity assessment (could be done by other nursing grades).
 - Specific time (measured in minutes) communicating outcome of assessment to General Practitioner and/or community pharmacist.
- Outcome of review (i.e. proceed with OAM/hold OAM/discontinue OAM).
- Toxicities requiring referral to hospital for medical review.
 - Description of toxicity (if relevant).
- Toxicities/problem requiring admission
 - Description of toxicity/problem if relevant.
- Adverse events previous cycle.

- Description of adverse events if relevant
- Near misses previous cycle.
 - Description of near misses if relevant.
- Clinical incident previous cycle.
 - Description clinical incident if relevant.
- Number of ad hoc queries from General Practitioner in previous cycle.
- Number of ad hoc queries from Community pharmacist in previous cycle.
- Number of ad hoc queries from other HCP in previous cycle.
- Number of ad hoc queries from patient in previous cycle.
- Number of ad hoc queries from family in previous cycle.

Appendix 3: EORTC OUT-PATSAT7

© QLQ-OUT-PATSAT7 Copyright 2017 EORTC Quality of Life Group. All rights reserved.

YOUR EXPERIENCE OF OUTPATIENT CARE

EORTC OUT-PATSAT7

We are interested in your MOST RECENT experience of the care received in the outpatient setting in this clinic. Please answer all the questions yourself by circling the number that best applies to you. There are no 'right' or 'wrong' answers

In the outpatient setting in this hospital, how would you rate services and care organisation, in terms of:

	Poor	Fair	Good	Very good	Excellent
34. The opportunity to see the same caregivers when you come to the outpatient clinic?	1	2	3	4	5
35. The ease of arranging medical appointments at convenient times?	1	2	3	4	5
36. The waiting time before obtaining a medical appointment?	1	2	3	4	5
37. The ease of communicating with the hospital services from home?	1	2	3	4	5
38. The information provided about what you should/should not do after you leave your hospital appointment?	1	2	3	4	5
39. The information on who to contact if you are worried after you leave your hospital appointment?	1	2	3	4	5
40. The provision of follow-up by the different caregivers (doctors, nurses, physiotherapists, psychologists, etc.) after treatment?	1	2	3	4	5

Appendix 4: Health Economic Analysis Questionnaires

DATE: _____

ID:

**A community-based ANP-led integrated oncology care model for adults
receiving oral anticancer medication**

HEALTH ECONOMICS QUESTIONNAIRE

PLEASE TAKE SOME TIME TO FILL OUT THE QUESTIONNAIRE.

IT WILL TAKE ABOUT 10 MINUTES TO COMPLETE.

THE QUESTIONNAIRE HAS 4 SECTIONS: A, B, C and D.



In this questionnaire we are trying to find out how **you are feeling** and about the **costs** to you arising from your cancer diagnosis and your current treatment. **Your answers are important** because they will give people in the Health Service an idea of how much the treatment costs you. Apart from the researchers, other nurses and doctors and health care professionals involved in your care **will not have access** any of the information you provide.

The information you give will be **completely confidential** and you will **not be identified** in any way. The detail you provide will be **held securely** as protected information.

Please try and answer every question. If you are not sure or cannot remember the exact details, please give the best answer you can. If you **do not want to answer a question** for whatever reason, then just leave it blank.

THANK YOU VERY MUCH FOR YOUR TIME AND HELP

SECTION A:	Your general health status
SECTION B:	Your private expenses
SECTION C:	Your general health service usage
SECTION D:	General demographic information

SECTION A: YOUR GENERAL HEALTH STATUS

We are interested in how you feel about **your health today**.

EQ-5D-5L

Under each heading, please tick the ONE box that best describes your health TODAY

Mobility

- I have no problems in walking about
- I have slight problems in walking about
- I have moderate problems in walking about
- I have severe problems in walking about
- I am unable to walk about

SELF-CARE

- I have no problems washing or dressing myself
- I have slight problems washing or dressing myself
- I have moderate problems washing or dressing myself
- I have severe problems washing or dressing myself
- I am unable to wash or dress myself

Usual Activities (e.g., work, study, housework, family or leisure activities)

- I have no problems doing my usual activities
- I have slight problems doing my usual activities
- I have moderate problems doing my usual activities
- I have severe problems doing my usual activities
- I am unable to do my usual activities

Pain/Discomfort


- I have no pain or discomfort
- I have slight pain or discomfort
- I have moderate pain or discomfort
- I have severe pain or discomfort
- I have extreme pain or discomfort

Anxiety/Depression

- I am not anxious or depressed
- I am slightly anxious or depressed
- I am moderately anxious or depressed
- I am severely anxious or depressed
- I am extremely anxious or depressed

- We would like to know how good or bad your health is TODAY
- This scale is numbered 0 to 100
- 100 means the best health you can imagine
- 0 mean the worst health you can imagine
- Mark an X on the scale to indicate how your health is TODAY
- Now, please write the number you marked on the scale in the box below.

YOUR HEALTH TODAY=



SECTION B: YOUR PRIVATE EXPENSES

This information gives the research team an understanding of the financial costs of having cancer and needing cancer treatment.

Q1. We are interested in whether you have missed **work** or arrived late/left early to attend the oncology clinic appointment with the ANP in the **last 3 months**. **If you have**, please fill in the section below and if not then move onto question 2.

TYPE	FREQUENCY		
YOUR TIME in the last 3 months			
In the last 3 months , did you miss work to attend appointments ?	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> Not applicable (NA)	If yes , how many days did you miss? (Please estimate) <hr/> If yes , how many hours did you miss? (Please estimate)	
In the last 3 months , did you arrive late or leave early from work because you were unwell ?	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NA	If yes , how many times did this happen? (Please estimate)	
In the last 3 months , did you miss work because you were unwell ?	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NA	If yes , how many days did you miss? (Please estimate)	

Q2. We would like to know what the **financial costs** are for your healthcare. Please tick which of the following you have to **pay for** and give us an estimate of how much you have had to pay over the last **3 months**. **If you have costs**, please fill in the section below and if not then move onto question 3.

YOUR EXPENSES		
Have you incurred any ' out of pocket ' expenses (spent any money)? If you do have , please estimate your expenses in relation to the categories listed below. If not , please leave blank		
<input type="checkbox"/> GP Visit	€ _____.00	In the last 3 months
<input type="checkbox"/> Blood tests	€ _____.00	
<input type="checkbox"/> Other, please specify _____ _____ _____ _____ _____ _____ _____	€ _____.00 € _____.00 € _____.00 € _____.00 € _____.00 € _____.00 € _____.00	

Q3. We are interested in whether **you spend any money** attending your appointments with the ANP for your oral anti-cancer treatment. **If you do**, please estimate how much you pay **per appointment**. If you don't have any costs, then move on to question 4.

YOUR CARE COSTS	Attending for blood tests	Attending the ANP oncology clinic
Do you usually get someone to look after your children when you have an appointment?	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NA	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NA
If yes , how much do you usually pay them per appointment ?	€_____ .00	
Do you usually get someone to look after other relatives when you have an appointment?	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NA	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NA
If yes , how much do you usually pay them per appointment ?	€_____ .00	

Q4. We would like to know if there are any **travel expenses** for you to attend your **appointments** (e.g., petrol, bus fare etc.). **If you do**, please estimate how much **per appointment**.

YOUR TRAVEL COSTS	For blood tests	Attending the ANP oncology clinic
How many kilometres (km) do you have to travel from your home to your appointment?	_____ km	_____ km
How do you usually travel to these appointments? (e.g., bus, car, train, taxi, walk, other) Please list all that apply.		
If you drive how much do you usually spend on parking per appointment?	€_____ .00	€_____ .00
If you use public transport how much do you usually spend per appointment?	€_____ .00	€_____ .00
If you usually pay someone to take you to your appointment how much do you usually spend per appointment?	€_____ .00	€_____ .00

SECTION C: YOUR GENERAL HEALTH SERVICE USAGE

Q1. We are interested in whether you have used any of the **healthcare services** listed below in the last **3 months**. If you have used a service, please indicate how **many visits** or **how often** you used that service in the last **3 months**.

TYPE	FREQUENCY OF USE	
SERVICE	Have you used this service in the last 3 months ?	If YES , approximately how many visits or how often did you use this service?
Outpatient clinic	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Dietician Services	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Optician Services	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Audiology (hearing) service	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
General Practitioner (GP)	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Practice Nurse (In GP practice)	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Public Health Nurse	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Physiotherapy Services	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Occupational Therapist	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Social Worker	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Psychological or Counselling Services	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick)	Visits: Number ____ Phone calls: Number ____
Other	<input type="checkbox"/> YES <input type="checkbox"/> NO → (Please tick and identify the service)	_____ _____

SECTION D: YOUR DEMOGRAPHICS

We are interested in learning more about you. This information is helpful to the research team but is not essential for you to complete.

1. What is your **gender**? Male Female Other
2. What is your **age** today? _____ years
3. Do you **smoke**? Yes No
4. Do you have a **medical card**? Yes No
5. Do you have a **GP visit card**? Yes No
6. Do you have **private health insurance**? Yes No
 - a. If **yes**, how much do you pay for this? €_____.00 per year or €_____.00 per month
7. What is your current **employment status**?

Full time employee	Part time employee	Unemployed
Student	Self-employed	Homemaker
Retired	Other If other, please specify: _____	
8. What is the highest level of **education** have you completed?

Primary level	Second level	Third level
---------------	--------------	-------------
9. What is your current **marital status**?

Married	Co-habiting	Widowed
Separated or divorced	Single or never married	
10. What is your **ethnic or cultural** background?

Irish	European (Non-Irish)	Other, please specify _____
-------	----------------------	-----------------------------

If you would like to provide feedback as to how you felt answering these questions, please do so:

Thank You!

You have now finished the questionnaire.

Thank you for taking the time to answer these questions.

If you would like any further information about this study, please contact


Janice Richmond or Mary Grace Kelly at

Letterkenny University Hospital on

0749123798 / 0749104642

National University of Ireland, Galway

Appendix 5: Proposed Model for OAM Care (Pilot)



References

American Association of Clinical Oncology (2013) ASCO/ONS Chemotherapy administration safety standards. Reference source: <https://www.ons.org/ascoons-chemotherapy-administration-safety-standards>. (Last accessed 09.02.2022).

Arain, M., Campbell, M. J, Cooper, C.L. et al (2010) What is a pilot or feasibility study? A review of current practice and editorial policy. BMC Medical Research Methodology, 10:67.

Balestroni G & Bertolotti, G. (2012) EuroQol-5D (EQ-5D): an instrument for measuring quality of life. Monaldi Arch Chest Diseases, 78(3):155-9. doi: 10.4081/monaldi.2012.121.

Battis, B., Clifford, L., Huq, M., Pejoro, E. & Mambourg, S. (2017). The impacts of a pharmacist-managed outpatient clinic and chemotherapy-directed electronic order sets for monitoring oral chemotherapy. Journal of Oncology Pharmacy Practice, 23, 582-590.

Brédart, A., Anota, A., Young, T., Tomaszewski, K. A., Arraras, J. I., Moura De Albuquerque Melo, H. & EORTC Quality of Life Group (2018). Phase III study of the European Organisation for Research and Treatment of Cancer satisfaction with cancer care core questionnaire (EORTC PATSAT-C33) and specific complementary outpatient module (EORTC OUT-PATSAT7). *European journal of cancer care*, 27(1), e12786.

Bugge, C., Williams, B., Hagen, S., Logan, J., Glazener, C., Pringle, S., & Sinclair, L. (2013). A process for Decision-making after Pilot and feasibility Trials (ADePT): development following a feasibility study of a complex intervention for pelvic organ prolapse. *Trials*, 14(1), 1-13.

Calabrese, E., Adams, L., Nestico, S., Giotis, A., Emmenegger, U. & Berry, S. (2015) An interprofessional model for the follow-up of patients receiving oral anti-cancer therapies. *Cancer Therapies*.

Calnan, M., Hutten, J., Tiljak, H., Saltman, R., Rico, A. & Boerma, W. (2006). The challenge of coordination: the role of primary care professionals in promoting integration across the interface. *Primary care in the driver's seat*, 85-104.

Cucinotta D, Vanelli M . (2020) WHO declares COVID-19 a pandemic. *Acta Biomed.* 2020;91(1):157–160. 32191675 10.23750/abm.v91i1.9397 7569573

Deluche, E., Darbas, T., Bourcier, K., Montangon, L., Bayard, G., Caille, E., Querrioux, J., Suchaud, C., Zabaleta, S. & Chaput, S. (2020). Prospective evaluation of an anti-cancer drugs management programme in a dedicated oral therapy center (DICTO programme). *Medical Oncology*, 37, 1-10.

Department of Health, Ireland (2017): National Cancer Strategy 2017-2026.
Reference source: <https://www.gov.ie/en/publication/a89819-national-cancer-strategy-2017-2026/>. (Last accessed 07.01.2022).

Greer, J.A., Amoyal, N., Nisotel, I, Fishbein, J. N., MacDonald, J., Stagl, J., Lennes, I., Temel, J. S., Safren, S. A. & Pirl, W. F. (2016) A Systematic Review of Adherence to Oral Antineoplastic Therapies. *Oncologist*. 2016; 21(3):354–376. 26921292 10.1634/theoncologist.2015-0405 4786357.

Hall, K. (2016). Oral Chemotherapy Program: Promoting Adherence Through

Process Improvement. *Journal of Oncology Navigation & Survivorship*, 7.

Hammond, L., Marsden, E., O'Hanlon, N., King, F., Henman, M. C. & Keane, C. (2012). Identification of risks associated with the prescribing and dispensing of oral anticancer medicines in Ireland. *International Journal of Clinical Pharmacy*, 34, 893-901.

Health Service Executive (2019) HSE Data protection policy. Reference source: <https://www.hse.ie/eng/gdpr/hse-data-protection-policy/>. (Last accessed 09.02.2022).

Herdman, Michael, Claire Gudex, Andrew Lloyd, M. F. Janssen, Paul Kind, David Parkin, Gouke Bonsel, and Xavier Badia. (2011). Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Quality of life research*, 20(10), 1727-1736.

House of the Oireachtas: Committee on the future of Healthcare: Sláintecare Report (2017). Reference Source <https://www.oireachtas.ie/en/committees/32/future-of-healthcare/>. (Last accessed 07.01.2022).

Institute for safe Medication practices (2020) Targeted Medication Safety Best

Practices for Hospitals. Reference source:

https://www.ismp.org/sites/default/files/attachments/2020-02/2020-2021%20TMSBP-%20FINAL_1.pdf (last accessed 07.01.2022)

Lancaster, G.A. (2015) Pilot and feasibility studies come of age. *Pilot and Feasibility Studies* 1 (1) DOI: <https://doi.org/10.1186/2055-5784-1-1>

Leung, M.S.T., Lin, S.G., Chow, J., & Harky, A. (2020). COVID-19 and Oncology: Service transformation during pandemic. *Cancer Medicine*, 9(19), 7161-7171.

Linedale, E.C., Mikocka-Walus, A., Gibson, P.R. & Andrews, J.M. (2020). The potential of integrated nurse-led models to improve care for people with functional gastrointestinal disorders: a systematic review. *Gastroenterology Nursing*, 43, 53-64.

Lopetegui, M., Yen, P.Y., Lai, A., Jeffries, J., Embi, P. & Payne, P. (2014). Time motion studies in healthcare: what are we talking about? *Journal of biomedical informatics*, 49, 292-299.

May, P., Laplant, K. & Mcgee, A. (2017). Practice model: Establishing and running an oral chemotherapy management clinic. *Asia-Pacific Journal of Oncology Nursing*, 4, 299-303.

Meier, J., Bergsbaken, J., Suzuki, S. (2018). Initiatives to improve safety of oral anticancer agents as delivered by community pharmacists. *European Medical Journal*. 2018;3(20):60–68.

Molassiotis, A., Liu, X.L. & Kwok, S.W. (2020). Impact of advanced nursing practice through nurse-led clinics in the care of cancer patients: A scoping review. *European Journal of Cancer Care*, e13358.

Muluneh, B., Schneider, M., Faso, A., Amerine, L., Daniels, R., Crisp, B., Valgus, J. & Savage, S. (2018). Improved adherence rates and clinical outcomes of an integrated, closed-loop, pharmacist-led oral chemotherapy management program. *Journal of oncology practice*, 14, e324-e334.

National Cancer Control Programme: Oral Anti-Cancer Medicines Model of Care Recommendations (2018). Reference source:

<https://www.gov.ie/en/publication/a89819-national-cancer-strategy-2017-2026/?referrer=http://www.health.gov.ie/wp-content/uploads/2017/07/National-Cancer-Strategy-2017-2026.pdf>. (Last accessed 23 October 2021).

National Cancer Institute: Targeted Cancer Therapies (2020). Reference source:

<https://www.cancer.gov/about-cancer/treatment/types/targeted-therapies/targeted-therapies-fact-sheet> (Last accessed 10 September 2021).

Oakley C., Lennan, E., Roe, H., Craven, O., Harrold, K. & Vidall C. (2010) Safe practice and nursing care of patients receiving oral anti-cancer medicines: a position statement from UKONS. Epub 2010 Jan 26. 2010; 4:177. Reference source: DOI: 10.3332/ecancer.2010.177. (Last accessed 19 January 2021).

Oncology Nursing Society (2016) Oral Adherence Toolkit Oncology Nursing Society; United States of America. Reference source:

https://www.ons.org/sites/default/files/ONS_Toolkit_ONLINE.pdf. (Last accessed 19 January 2021).

Paoletta, G.A., Boyd, A.D., Wirth, S.M., et al. (2018) Adherence to Oral Anticancer Medications: Evolving Interprofessional Roles and Pharmacist Workforce Considerations. *Pharmacy (Basel)*. 2018;6(1):23. Reference source: 29518017 10.3390/pharmacy6010023 5874562. (Last accessed 10 December 2020).

Randall, S, Crawford, T., Currie, J., et al. (2017) Impact of community-based nurse-led clinics on patient outcomes, patient satisfaction, patient access and cost effectiveness: A systematic review. *Int J Nurs Stud*. 2017;73:24–33. 28531549 10.1016/j.ijnurstu.2017.05.008

Richards, D.A. & Halberg, I.R (2015) *Complex interventions in health: an overview of research methods*. Routledge: England.

Richmond, J.P., Kelly, M.G., Johnston, A., Murphy, P. J., & Murphy, A. W. (2021). Current Management of Adults Receiving Oral Anti-Cancer Medications: A scoping Review. *European Journal of Oncology Nursing*, 102015.

Schwenkglens, M., & Matter-Walstra, K. (2016). Is the EQ-5D suitable for use in

oncology? An overview of the literature and recent developments. Expert review of pharmacoeconomics & outcomes research, 16(2), 207-219.

Shanyinde, M., Pickering, R. M., & Weatherall, M. (2011). Questions asked and answered in pilot and feasibility randomized controlled trials. BMC medical research methodology, 11(1), 1-11.

Skivington, K., Matthews, L., Simpson, S.A., Craig, P., Baird, J. & Blazeby, J.M., Boyd, K.A., Craig, N., French, D.P., McIntosh, E., Petticrew, M., Rycroft-Malone, J., White, M. & Moore, L. (2021) A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance BMJ 2021; 374 :n2061 doi:10.1136/bmj.n2061

Sung, H., Ferlay, J., Siegel, R.L., Laversanne, M., Soerjomataram, I., Jemal, A., & Bray, F. (2021). Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA: a cancer journal for clinicians, 71(3), 209-249.

Thompson, W. & McNamara, M. (2021) Constructing the advanced nurse

practitioner identity in the healthcare system: A discourse analysis. *Journal of Advanced Nursing*, 00 1-13. DOI: 10.1111/jan.15068.

Torrens, C., Campbell, P., Hoskins, G., Strachana, H., Wells, M., Cunningham, M., Bottonea, H., Polson, R., & Maxwell, M. (2020). Barriers and facilitators to the implementation of the advanced nurse practitioner role in primary care settings: A scoping review. *International journal of nursing studies*, 104, 1-21.
<https://doi.org/10.1016/j.ijnurstu.2019.103443>.

Vidall, C. (2010). Providing community oral chemotherapy services. *European Journal of Cancer Care*, 19, 29-34.

Vogler, S.A., & Lightner, A.L. (2020). Rethinking how we care for our patients in a time of social distancing during the COVID-19 pandemic. *Journal of British Surgery*, 107(8), 937-939.

United Kingdom Oncology Nursing Society (UKONS) (2010) Safe practice and nursing care of patients receiving oral anti-cancer medicines: a position statement from UKONS cancer.org/en/journal/article/177-safe-practice-and-nursing-care-of-

patients-receiving-oral-anti-cancer-medicines-a-position-statement-from-ukons

Reference source: <https://doi.org/10.3332/ecancer.2010.177>. Last accessed 29 January 2021.

Wong, S.F., Bounthavong, M., Nguyen, C. P. & Chen, T. (2016). Outcome assessments and cost avoidance of an oral chemotherapy management clinic. *Journal of the National Comprehensive Cancer Network*, 14, 279-285.

Wood, L. (2012) A review on adherence management in patients on oral cancer therapies. *Eur J Oncol Nurs*. 2012; 16(4):432–438. 22051845
10.1016/j.ejon.2011.10.002.