Title Assessing the transportability of clinical prediction models for cognitive impairment

using causal models

Jana Fehr*[1,2], Marco Piccininni[3,4], Tobias Kurth[3], Stefan Konigorski*[1,2,5] for the Alzheimer's
Disease Neuroimaging Initiative†

1 Digital Engineering Faculty, University of Potsdam, Germany

2 Digital Health and Machine Learning, Hasso-Plattner-Institute, Potsdam, Germany

3 Institute of Public Health, Charité – Universitätsmedizin Berlin, Berlin, Germany

4 Center for Stroke Research Berlin, Charité – Universitätsmedizin Berlin, Berlin, Germany

5 Hasso Plattner Institute for Digital Health at Mount Sinai, Icahn School of Medicine at

Mount Sinai, New York, NY, USA.

11
12   * Corresponding authors: Jana Fehr & Stefan Konigorski
13   Digital Engineering Faculty, University of Potsdam, Germany
14   Prof.-Dr.-Helmert-Str. 2-3
15   14482 Potsdam, Germany
16   e-Mail: jana.fehr@hpi.de; stefan.konigorski@hpi.de
17

†Data used in preparation of this article were obtained from the Alzheimer's Disease

Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators

within the ADNI contributed to the design and implementation of ADNI and/or provided data

but did not participate in the analysis or writing of this report. A complete listing of ADNI

investigators can be found at: https://adni.loni.usc.edu/wp-

content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Abstract

Background: Machine learning models promise to support diagnostic predictions, but may not perform well in new settings. Selecting the best model for a new setting without available data is challenging. We aimed to investigate the transportability by calibration and discrimination of prediction models for cognitive impairment in simulated external settings with different distributions of demographic and clinical characteristics.

Methods: We mapped and quantified relationships between variables associated with cognitive impairment using causal graphs, structural equation models, and data from the ADNI study. These estimates were then used to generate datasets and evaluate prediction models with different sets of predictors. We measured transportability to external settings under guided interventions on age, APOE ε4, and tau-protein, using differences between internal and external settings measured by calibration metrics and area under the receiver operating curve (AUC).

Results: Calibration differences indicated that models predicting with causes of the outcome were more transportable than those predicting with consequences. AUC differences indicated inconsistent trends of transportability between the different external settings. Models predicting with consequences tended to show higher AUC in the external settings compared to internal settings, while models with parents or all variables showed similar AUC.

Conclusions: We demonstrated with a practical prediction task example that predicting with causes of the outcome results in better transportability compared to anti-causal predictions measured by calibration differences. We conclude that calibration differences are more suitable than AUC differences to assess model transportability to external settings.

Keywords:

Alzheimer's Disease, clinical risk prediction, DAG, causality, transportability

Introduction

Dementia is the second leading cause of death globally,[1] and more than 55 million people

currently have dementia. Detecting dementia at an early stage of cognitive impairment is

essential to give affected individuals adequate care and eventually administer disease-

modifying treatments.[2] In recent years, several machine learning (ML) models have been

proposed to support clinical decision making by predicting the diagnosis of Alzheimer's

disease (AD) and cognitive impairment.[3–8] The models were developed with data from

different cohorts and included different predictor variables, such as image-derived brain

volumetric measures, cognitive test results, or demographic predictors. One obstacle for

deploying such prediction models in clinical practice is that they might not generalize well

when being transported (i.e., being applied) to other settings (e.g., in another hospital or

regions with different patient demographics). One reason for reduced transportability may be

that ML models learn non-causal associations between input and output variables, which

might be different in external settings.[9,10] This scenario can especially occur when models

predict a diagnosis based on clinical consequences of the disease (e.g., when prediction is

in the anti-causal direction).[11–13] For prospective applications, end-users face the challenge

of finding the most transportable model to their setting where data has not yet been

collected.

Causality research established two approaches to improve transportability for prediction

models. First, causal relationships can be incorporated in prediction models *a priori* for

learning relationships that are more stable across settings and can therefore avoid

systematic failures in external settings.[10,14–17] To this aim, directed acyclic graphs (DAGs) are

a useful tool to map assumed causal relationships between variables, represent differences

and commonalities between settings,[18,19] and select variables for transportable health

prediction tasks.[20–22] Second, the causal validity of learned relationships can be assessed

through guided interventions (also known as perturbations) on data distributions to simulate

differences between internal and external validation settings.[17,23,24] Research in

epidemiology and ML has adopted DAGs and interventions to develop transportable ML

models.[25–34] Piccininni et al. described the use of DAGs for selecting a single predictor in a

hypothetical clinical risk prediction model for AD.[25] They discussed that prediction models for

AD are more likely to transport well to different settings when the selected predictor is a

cause of AD and not a consequence. Rojas-Carulla et al. and Magliacane et al. applied

automatic hypothesis testing to determine a transportable predictor set across multiple

source domains.[30,31] Subbaswamy et al. used interventions to achieve that predictors do not

depend on unreliable parts of the data-generating process and thereby generalize to

unknown test data.[35] In another study, Singh et al. proposed a model predicting acute kidney

injury to ensure the fairness of predictions on unseen test data by applying DAGs and

interventions.[32] Steingrimsson et al. demonstrated an approach to assess transportability to

external settings where the outcome variable has not yet been measured using inverse-odds

weights.[36] These and other works[37–39] measured transportability to unseen data by

comparing mean-squared error or discrimination error (e.g., area under the receiver

operating curve (AUC)) between internal and external validation settings. Following the work

of Van Calster et al., however, we argue that discriminatory metrics may not be a suitable

metric to assess external validity because predicted risks can be unreliable even if

algorithms have good discrimination, and instead suggest calibration as a suitable metric to

measure transportability.[40]

In this work, we aim to compare the transportability, measured by calibration and

discrimination, of models predicting cognitive impairment with different sets of predictors to

simulated external settings to add evidence for two questions: 1) how to construct

transportable ML models and 2) how to assess the transportability.

Methods

<u>Data source and data preprocessing</u>

Data used in the preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). The primary goal of ADNI has been to test whether medical imaging, biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early AD. For up-to-date information, see www.adni-info.org. The ADNI study acquired multiple longitudinal measurements from elderly subjects across more than 50 clinics in USA and Canada.[41,42] The ADNI study enrolled participants who have mild cognitive impairment or a diagnosis of early AD and elderly controls between the age of 55 and 90 years old. We used the ADNI data subset that was created for the TADPOLE grand challenge (https://tadpole.grand-challenge.org/Data/), which is also available at https://ida.loni.usc.edu. We selected individuals who had a diagnosis and clinical measurements at baseline (n=1,737) and considered baseline variables that have been reported to be related to AD and had less than 30% of missing entries. Detailed information on data preprocessing is provided in Supplementary Text 1. Missing data were imputed using the R package 'mice' with default settings, and three imputed datasets were generated. All numeric variables were normalized by z-transformation. We defined the outcome variable 'cognitive impairment' based on the final diagnosis variable in ADNI by considering participants with cognitive normal or subjective memory complaint as "cognitively normal" and participants with mild cognitive impairment or AD as "with cognitive impairment."

<u>DAG creation</u>

In DAGs, nodes represent variables, and directed edges represent direct causal relationships pointing from the direct cause to the effect.[18,43,44] We reviewed the scientific literature to identify causal relationships between variables in our dataset that are involved in

cognitive impairment and AD processes (Supplementary Table S1) and mapped them in a

first DAG (Supplementary Figure S1). Then, we tested if the generated DAG was a good fit

to the ADNI dataset, using conditional independence testing with the R package 'dagitty.'[45]

We reviewed those test results with low p-values and large point estimates, which indicated

a violation of the implied conditional independence. We added 13 causal connections

(Supplementary Text 2) according to the test results and our domain expertise to create the

final DAG (Figure 1).

Figure 1: Directed acyclic graph of variables related to cognitive status. Predictor variables are
marked in blue, and the outcome variable (cognitive status) in green. Directed arrows indicate
assumed causal relationships between variables. The included variables are: APOE ε4 (apoe4), age,
sex, education (educ), CSF-Aβ (aβ), history of alcohol abuse (alc), history of smoking behavior
(smok), Body Mass Index (bmi), history of hypertension (hypert), CSF-tau (tau), history of
cardiovascular events (cardio), cognitive status (cogn), hippocampus (hippo), ventricles (ventr),
intracranial volume (icv), FDG-PET (fdg), Mini-Mental State Exam score (mmse).

<u>Semi-synthetic data generation using structural equation models</u>

We fitted a structural equation model (SEM) using the three imputed ADNI datasets to

quantify the causal relationships specified in our DAG. The SEM was implemented using the

'sem' function in the R package 'lavaan' with default parameters.[46] For numeric endogenous

(dependent) variables, the function computes weighted least squares estimates. For

categorical endogenous variables, the function automatically uses a diagonally weighted

least squares estimator and assumes that a conditional normally distributed latent variable

underlies the categorical variable (and estimates the thresholds).

We then used the SEM parameter estimates (Supplementary Table S2) to generate six

semi-synthetic datasets with 10,000 individuals each: one for training, one for internal

validation, and four for external validation of ML models. We bootstrapped exogenous

(independent) variables (age, sex, and APOE ε4) 10,000 times without replacement from the

original data. We used those to generate the endogenous variables for training and internal

validation sets, using the linear equations from the SEM (Supplementary Figure S2). We

generate the four external validation sets implemented by interventions on the variables to

reflect different populations with

1) a younger mean age, compared to the original data (73 years $\Rightarrow$ 35 years, which is

similar to the global world population mean at 31[1]**),**

2) a younger mean age compared to original data, but higher compared to the first age

validation setting $\Rightarrow$ 65 years (referred to as "age2"),

3) lower prevalence of the APOE ε4 gene compared to the original data (46.9% $\Rightarrow$

5.0%), and

4) a different mechanism generating the endogenous variable tau-protein, measured in

cerebrospinal fluid.

For the external age-intervention data, we sampled the age variable from a normal

distribution with a mean age of 35 (and mean of 65 for age2 setting) and standard deviation

of 10 and bootstrapped APOE ε4 and sex. For the APOE ε4 intervention, we sampled from a

Bernoulli distribution with a 5% probability. For the tau intervention, we altered the

mechanism determining tau levels by intervening on the parameters estimated by the SEM

for the tau equation. In particular, we arbitrarily changed the intercept from the tau equation

from -0.5 to 0.9, increased the influence of age on tau from 0.37 to 0.9, increased the

influence of apoe4 from 0.57 to 0.8, increased the influence of hypertension from 0.14 to 0.9,

and reduced the influence of alcohol from 0.57 to 0.001.

---

[1] https://worldpopulationreview.com/country-rankings/median-age

<u>Prediction algorithms</u>

We applied logistic regression, lasso regression, random forest, and generalized boosted

regression (GBM) to predict the cognitive state of an individual as either cognitive normal or

with cognitive impairment. Logistic regression was performed using the glm function in the

'stats' R package. Lasso regression was implemented using the 'glmnet' R package.[47] The

lasso model was initialized with an optimized penalization hyperparameter obtained from a

grid-search with 10-fold cross validation that selected the value of lambda for minimum

deviance. The random forest is an ensemble of regression trees, which aims at improving

generalizability compared to a single regression tree.[48] Previous works demonstrated the

strengths of random forests for diagnostic prediction modelling of AD.[3–5] The random forest

algorithm was applied from the 'randomForest' R package, using 500 trees and $\sqrt{p}$ number

of variables randomly sampled at each split (as per default), where p is the number of

predictors. GBM implements boosting by adding regression trees sequentially with respect to

the error of the current tree ensemble. This boosting approach increases robustness and

generalizability compared to a single regression tree.[49–51] The GBM algorithm was applied

using the 'gbm' R package with 100 trees (as per default).

Based on the causal assumptions in our DAG, we defined four predictor sets that included

either all variables or only those which are direct causes of the outcome (defined as parent

nodes), or only direct consequences of the outcome (defined as children nodes), or only

exogenous variables (age, sex, and APOE ε4) (Table 1). Each ML model was trained and

validated with each predictor set. We performed 10,000 repetitions to generate the six

datasets (one for training, one for internal validation, and four for external validation) for

training and validating all prediction models.

We repeated this procedure three times, each time using the SEM parameters obtained from

one of the three imputed ADNI datasets. As a sensitivity analysis, we additionally ran 100

repetitions using hyperparameter tuning to minimize the deviance for the random forest

(tuned parameters: number of predictors sampled for splitting at each node from 1 to 5, and

the minimum size of terminal nodes of 1, 5 or 10) using generated training datasets with

1,000 observations.

Table 1: Predictor sets with corresponding lists of variable names.
The predictor variables comprised age, APOE ε4, sex, years of education, Body Mass Index (BMI),
history of hypertension, history of alcohol abuse, history of smoking behavior, history of
cardiovascular events, Cerebrospinal fluid (CSF) of tau and Amyloid β (Aβ), volumes measured from
brain magnetic resonance imaging (sum of left and right hippocampal volumes, sum of left and right
ventricle volumes, intracranial volume, fluorodeoxy-glucose-positron emission tomography (FDG-
PET)), and Mini Mental State Exam score (MMSE).

Predictor set	Variable names
All nodes	age, APOE ε4, sex, education, BMI, history of hypertension, history of alcohol abuse, history of smoking behavior, history of cardiovascular events, CSF-tau, CSF-Aβ, hippocampus, ventricles, intracranial volume, FDG-PET, MMSE
Parent nodes	age, APOE ε4, sex, education, BMI, history of cardiovascular events, CSF-tau, CSF-Aβ
Children nodes	hippocampus, ventricles, intracranial volume, FDG-PET, MMSE
Exogenous nodes	age, APOE ε4, sex

<u>Assessing transportability</u>

We calculated calibration metrics and the AUC discrimination performance for all prediction

models in the internal validation setting and in each external validation setting. Calibration

was measured using the Integrated Calibration Index (ICI)[52] and the calibration component

of a three-way decomposed Brier score.[53] Low ICI and Brier scores indicate better

calibration. The Brier calibration component (termed 'reliability') was obtained from the bias-

corrected 'BrierDecomp' function of the 'SpecsVerification' R package using quantile bins of

predicted probabilities in 10% steps. The AUC was obtained from the 'pROC' R package.

AUC values close to one indicate good discrimination performance.

We assessed the transportability between the internal setting and each external validation

setting by differences in calibration (ICI or Brier score) and in AUC. Differences of zero

indicate equal performance in both internal validation and external settings and, therefore,

good transportability. Negative calibration differences values indicate decreased calibration

from internal validation to the intervention setting and therefore decreased transportability.

AUC differences greater than zero indicate decreased discriminatory performance from

internal validation to the intervention setting and thus low transportability.

We calculated the median, 2.5%, and 97.5% percentiles for performance metrics across all

10,000 repetitions for each of the three imputed ADNI datasets. See Figure 2 for a summary

of our workflow. All prediction algorithms, data simulations, and data analyses were

implemented using R version 4.0.3.

Results

Figure 2: Our approach to assess the transportability of machine learning models predicting cognitive
impairment. Orange boxes mark the four general steps of our workflow. We first mapped knowledge
about cognitive impairment into a Directed Acyclic Graph (DAG) and quantified those using Structural
equation modelling (SEM) and data from the Alzheimer's Disease Neuroimaging Initiative (ADNI). The
estimates were used in linear equations to generate datasets for training, internal validation, and four
external validation datasets with interventions on age, APOE ε4, and tau. The age variable was
intervened two times (age and age2) by sampling from normal distributions with two different mean
age values (35 and 65). We trained four machine learning algorithms (logistic regression, lasso
regression, random forest, and generalized boosted regression) to predict cognitive impairment using
four sets of predictors. We measured transportability between internal and external settings using
calibration differences, measured by Integrated Calibration Index (ICI) and Brier score, and
differences in Area under the Receiver Operating Curve (AUC). Steps 3 to 4 (data synthesis and
model training and validation) were repeated 10,000 times for each of the three imputed ADNI
datasets.

Description of the participants' characteristics

The ADNI study, represented in TADPOLE, recorded a total of 1737 participants with a
diagnosis at baseline together with their demographic information (age, sex, and education),
behavioral information (smoking and alcohol abuse history), clinical measurements (BMI,
FDG-PET imaging, brain volumetric measurements with MR imaging, Aβ and tau protein
concentrations in cerebrospinal fluid (CSF), Minimental State Cognitive Exam (MMSE) and
medical history (history of hypertension and cardiovascular events) (Table 2). Among all
participants, 1214 (69.9%) were diagnosed with cognitive impairment, which comprised 872
(71.8%) individuals with mild cognitive impairment and 342 (28.7%) individuals with AD. A

total of 523 (30.1%) individuals was diagnosed as cognitive normal, including 106 (20.3%)

individuals with subjective memory complaint.

Table 2: Participant characteristics of ADNI dataset at baseline stratified by cognitive status. Numeric
variables are indicated with * and are given with median and 25-75% interquartile range (IQR). All
other variables are categorical variables with two categories, and the absolute number and the
column-wise percentage of the reference category are given. Absolute numbers of missing values
(Nmiss) are given. Abbreviations: Body Mass Index (BMI), Cerebrospinal fluid (CSF), Amyloid β (Aβ),
Mini Mental State Exam score (MMSE).

Variables	Cognitive normal n=523 (30.1%)	Cognitive impairment n=1214 (69.9%)	Total n=1737 (100%)
Age*	73.7 (70.5, 78.0)	74.0 (68.3, 79.3)	73.9 (69.2, 78.9)
APOE ε4	149 (28.6%) Nmiss: 2	660 (54.8%) Nmiss: 10	809 (46.9%) Nmiss:12
sex (male)	253 (48.4%)	704 (58.0%)	957 (55.1%)
education* (years)	16 (14, 18)	16 (14, 18)	16 (14, 18)
CSF-Aβ* (pg/ml)	1271 (820.8, 1734.0) Nmiss: 156	741.5 (559.1, 1130.3) Nmiss: 366	854.2 (596.2, 1395.5) Nmiss: 522
CSF-tau* (pg/ml)	214.3 (175.2, 287.8) Nmiss: 156	281.6 (210.4, 379.2) Nmiss: 366	257.8 (193.4, 349.7) Nmiss: 522
history of alcohol abuse	16 (3.1%) Nmiss: 6	27 (2.3%) Nmiss: 27	43 (2.5%) Nmiss: 33
history of smoking	115 (26.8%) Nmiss: 94	223 (23.2%) Nmiss: 254	338 (24.3%) Nmiss: 348
BMI*	28.6 (25.8, 32.5) Nmiss: 4	28.17 (25.5, 31.2) Nmiss: 5	28.31 (25.5, 31.6) Nmiss: 9
history of hypertension	130 (24.9%)	460 (38.0%) Nmiss: 5	590 (34.1%) Nmiss: 5
history of cardiovascular events	343 (65.6%)	835 (68.8%)	1178 (67.8%)
MMSE*	29.0 (29.0, 30.0)	27.0 (25.0, 29.0)	28.0 (25.0, 29.0)

<u>Semi-synthetic data generation</u>

The SEM estimated all parameters quantifying the causal relationships in the DAG in Figure

1. We reviewed the estimated parameters and found that many were in agreement with

existing neurological domain knowledge. For example, age had a positive coefficient and

therefore increased CSF-tau (0.37), the likelihood of hypertension (0.11), and the likelihood

of cardiovascular events history (0.13) (Supplementary Table S3). Some estimated

relationships, however, were controversial to domain knowledge. For example, increasing

age decreased CSF-Aβ (-0.14) and the likelihood of cognitive impairment (-0.13). The SEM

additionally indicated a small correlation between sex and age and between age and APOE

ϵ4.

We compared endogenous variable distributions between the original ADNI data and

generated validation datasets generated by SEM parameters from the first imputed dataset

(Supplementary Table S4) and found that the percentage of cognitive impairment was

similar between the internal validation set (70.0%, 2.5% and 97.5% percentiles [69.1, 71.0])

and the original ADNI data (69.9%). We further compared endogenous variable distributions

between internal and external datasets. Lowering the mean age of 73.8 years in the internal

validation setting to 35 years in the external setting slightly decreased the prevalence of

cognitive impairment from 70.0% to 68.3%, increased the smoking prevalence from 23.4% to

34.1% and alcohol abuse history from 1.5% to 46.5%, decreased the prevalence of

hypertension from 35.0% to 8.0% and previous cardiovascular events from 65.1% to 32.4%.

Intervening on age increased the mean of Aβ from 1076.6 to 1521.0 pg/ml, shrank the mean

of tau from 292.7 to 180.8 pg/ml, and increased the MMSE from 27.3 to 29.6, in comparison

to the internal validation data. The differences between the internal and external variable

distributions were smaller but followed the same trends when reducing the mean population

age only slightly from 73.8 to 65.0 in the age2 intervention setting.

In the APOE ε4 intervention, lowering the prevalence of the APOE ε4 gene from the internal

setting to the external setting decreased the prevalence of cognitive impairment from 70.0%

to 63.1%, increased the mean of CSF-Aβ from 1076.6 to 1291.2 pg/ml and decreased the

mean CSF-tau from 292.7 to 260.1 pg/ml.

In the tau intervention, altering the intercept and coefficients that determine tau levels

between the internal setting to the external setting increased the level of CSF-tau from 292.7

to 476.4. Also they increased the prevalence of cognitive impairment from 70.0% to 82.3%.

<u>Internal calibration and discrimination of the models</u>

We evaluated the internal validation performance measured by calibration and AUC metrics

(Figure 3, Supplementary Figure S3, Supplementary Figure S4, and Supplementary Table

S5) of all models. ICI scores were close to 0 for all logistic and lasso models (e.g., logistic

and lasso with parents as predictors 0.009 2.5% and 97.5% percentiles [0.004, 0.016]).

Random forest and GBM models had lower calibration (indicated by higher ICI values)

compared to logistic and lasso models (using parents as predictors, random forest 0.035

[0.024, 0.046], GBM 0.029 [0.019, 0.039]). Random forest models predicting with exogenous

variables (age, sex, and APOE ε4) had very low calibration (0.292 [0.277, 0.305]). Brier

scores were exactly zero or closer to zero in the internal validation setting compared to the

ICI scores, except for random forest models predicting with exogenous variables, which

showed the lowest calibration (0.088 [0.080, 0.094]) among all models. AUC indicated the

best performance for models predicting with all variables, and of those models, logistic

regression and lasso regression achieved the highest AUC (0.75 [0.73, 0.76]) compared to

random forest (0.73 [0.72, 0.74]) and GBM (0.71 [0.70, 0.73]). All models predicting with

exogenous variables had the lowest discriminatory performance of 0.5.

Figure 3: Model performance in the internal validation setting, measured by the integrated calibration
index (ICI), Brier score, and area under the receiver operating curve (AUC). Cognitive impairment was
predicted using logistic regression, lasso regression, random forest (rf), and generalized boosted
regression (gbm) prediction models. Models were trained either with all predictor variables, only
parent nodes (direct causes) of the outcome, only children nodes (consequences) of the outcome, or

with the exogenous variables age, sex, and APOE ε4 (apoe4). Depicted are the full distributions of
ICI, Brier scores, and AUC, smoothed with a Gaussian kernel density function and medians marked
with ◊. The displayed metrics were obtained from 10,000 repetitions of data generation and model
training on the first imputed dataset.
331

<u>Measuring transportability by calibration differences</u>

We compared the transportability of prediction models measured by calibration differences

between internal validation and intervention settings (Supplementary Table S6). For this, we

focus on logistic regression and lasso regression predicting with all variables, parents,

children, and exogenous variables, since they showed good calibration in the internal

setting. In all intervention settings, models predicting with parent nodes were more

transportable than those predicting with children nodes (Figure 4, Supplementary Figure S3,

and Supplementary Figure S4). Models predicting with parents had good transportability in

intervention settings, indicated by a similar calibration (thereby small calibration difference)

between the internal validation and intervention setting. For example, the median ICI

difference between the internal validation and age intervention for logistic regression was

very small (ICI -0.009 [-0.045, 0.006]). Models predicting with children variables had low

transportability in intervention settings, as indicated by negative calibration differences. For

example, logistic regression predicting with children had a median ICI difference between

the internal and age intervention setting of -0.300 [-0.322, -0.276], which was 33.3-fold lower

than predicting with parents. The largest difference in the median ICI differences between

parents and children was in the age intervention setting (logistic regression: 0.291), and the

smallest one was in the APOE ε4 setting (logistic regression: 0.008). Logistic regression

models predicting with all variables equally or less transportable compared to models

predicting with parent variables (e.g., age intervention: all predictors median ICI -0.031 [-

0.073, 0.002], parent predictors: -0.009 [-0.045, 0.006]; tau: all predictors -0.014 [-0.022, -

0.005], parents 0.000 [-0.007, 0.008]). Similar results were observed for the lasso regression

models. In the age2 and APOE ε4 intervention setting, the calibration differences were close

to zero for logistic and lasso models with all predictors and parent predictors. Using

exogenous variables as predictors had calibration differences close to zero, except in the tau

intervention setting. We found low transportability of logistic and lasso regression models

with all sets of predictors in the tau-intervention setting. Only the models predicting with

parent variable had zero calibration differences.

Random forest and GBM models also showed generally lower ICI when using children as

predictors compared to parents and higher transportability for parent predictors compared to

all predictors. However, random forest and GBM models predicting with parents had ICI

differences between internal and external validation differences far from zero. Brier scores

supported the same trends as the ICI (Supplementary Figures S4 and S5, Supplementary

Table S6).

<u>Measuring transportability by AUC differences</u>

When measuring transportability with AUC differences, we found inconsistent transportability

trends between the intervention settings. In the age-intervention setting, models predicting

with children variables also had lower transportability compared to models predicting with

parent predictors, indicated by a positive AUC difference between internal validation and

intervention setting (logistic regression with parent predictors: -0.01 [-0.04, 0.02], logistic

regression with children 0.02 [0.00, 0.04], Figure 5, Supplementary Figures S6 and S7,

Supplementary Table S6). In the age2 and tau intervention settings, however, logistic

regression models predicting with children increased their AUC by 0.03 compared to the

internal validation setting, whereas logistic regression models predicting with parent

predictors had AUC differences close to zero. AUC differences indicated similar

transportability close to zero for all predictors compared to parent predictors in all

intervention settings. Lasso regression, random forest, and GBM models indicated similar

trends as logistic regression models.

Figure 4: Transportability between internal validation and external settings, measured by the
difference of integrated calibration index (ICI). Four intervention test sets were created with 1)
reducing the population mean age from 73 to 35 years, 2) reducing the population mean age from 73
to 65 years (age2), 3) reducing the APOE ε4 allele frequency from 46.6% to 5.0%, and 3) changing
the SEM-parameters for generating the endogenous variable tau. Cognitive impairment was predicted
using logistic regression, lasso regression random forest (rf), and generalized boosted regression
(gbm) prediction models. Models were trained either with all predictor variables, only parent nodes
(direct causes) of the outcome, only children nodes (consequences) of the outcome, or exogenous
variables (exo) age, sex, and APOE ε4 allele frequency. Depicted are the full distributions of ICI
differences from 10,000 repetitions on the first imputed dataset, smoothed with a Gaussian kernel
density function and medians marked with ◊.

Figure 5: Transportability between internal validation and external settings, measured by the difference of area under the receiver operating curve (AUC). Four intervention test sets were created with 1) reducing the population mean age from 73 to 35 years, 2) reducing the population mean age from 73 to 65 years (age2), 3) reducing the APOE ε4 allele frequency from 46.6% to 5.0%, and 3) changing the SEM-parameters for generating the endogenous variable tau. Cognitive impairment was predicted using logistic regression, lasso regression random forest (rf), and generalized boosted regression (gbm) prediction models. Models were trained either with all predictor variables, only parent nodes (direct causes) of the outcome, only children nodes (consequences) of the outcome, or exogenous variables (exo) age, sex, and APOE ε4 allele frequency. Depicted are the full distributions of AUC differences from 10,000 repetitions on the first imputed dataset, smoothed with a Gaussian kernel density function and medians marked with ◊.

<u>Sensitivity analyses with other imputed datasets and hyperparameter tuning</u>

We performed sensitivity analysis by using the other two imputed dataset as input for the

SEM-model. We did not find any differences between model performance results (ICI, Brier

score, and AUC) between replicates that used three different imputed datasets as input

(Supplementary Figures S5, S7, S8-9).

Optimizing hyperparameter for random forest models did not meaningfully improve

calibration or AUC performance (Supplementary Figure S10). The observed trends for

transportability measured by ICI differences (i.e., parent predictors being more transportable

compared to children) remained for random forest in the age-interventions settings, but in

the tau-intervention setting, optimized random forest models predicting with children may

have equal or better ICI transportability (-0.019 [-0.044, 0.012]), compared to parents (-0.032

[-0.063, -0.003]).

Discussion

In this study, as a first contribution, we have presented a causal data generation approach

for assessing the transportability of prediction models for cognitive impairment in synthetic

external settings with different distributions of age, APOE ε4 allele frequency, and tau. As a

second contribution, we assessed transportability by comparing performance between

internal and external validation settings, measured by the discrimination performance (AUC)

as in most prior studies,[30–32,34] but also by calibration (ICI and Brier calibration component).

Both calibration metrics, ICI, and the Brier score confirmed the previous causal theory that

prediction models that use direct causes of the outcome for the prediction are generally

more transportable.[20,25] We showed that, under a specific set of interventions, calibration

performance remained stable when ML models predicted only with direct causes (parent

nodes) but was reduced when predicting with consequences (children nodes) of the

outcome 'cognitive impairment.' We found that this held true in all prediction models (logistic

regression, lasso regression, random forest, and GBM). Measuring transportability by AUC

differences indicated inconsistent trends for transportability and did not reflect causal theory.

Calibration measures the closeness between the average predicted probabilities and the

relative frequency of the outcome event and therefore reflects the actual capability of

predicting the state of the outcome variable. In contrast, the AUC depends on the ranking of

the predicted risks and measures the probability that a predicted risk for a randomly drawn

individual with the event is higher compared to a randomly drawn individual without the

event.[40] With our results, we reinforce the claim that besides discrimination, good calibration

is important to achieve clinically useful prediction models.[40] Using calibration metric

differences to measure the transportability of prediction models, however, requires good

calibration in the internal setting. It is possible that a model predicting with children variables

has better transportability than a model predicting with parent variables if the model with

children predictors outperforms the model with parent variables in the internal validation

setting.

In our simulation, we found that the random forest and the GBM algorithms were not well

calibrated in the internal setting. The miscalibration might have happened because we

generated data assuming linear relationships in the SEM, whereas random forest and GBM

are designed to capture non-linear relationships.[54]

Our study further showed that causal thinking is essential when selecting predictors for

clinical prediction models. Previously developed prediction models for dementia and AD

have used brain volumetric measures or cognitive assessment scores as predictors because

they reduced prediction errors.[5,55,56] Similarly to another work[57], we assumed that these

predictors (brain volumetric measures and cognitive test results) are consequences of the

cognitive outcome status and therefore predict in the anti-causal direction leading to reduced

transportability to external settings. Another work suggested that predictors derived from

medical images may often predict in the anti-causal direction as they depict the

consequences of a disease, which may raise a caveat towards transportability.[15]

Our application to assess transportability has limitations. First, it cannot be empirically

verified if DAGs map causal relationships correctly and if all relevant factors were included.

We only included observed variables (other than latent variables for factors), and it is likely

that there are unobserved variables involved in the causal process of cognitive impairment.

Strong domain expertise is crucial to build accurate DAGs.[43] Conditional independence tests

can test if there is evidence against a given DAG in a dataset.[12] We applied conditional

independence tests to add directed connections between variables, but unexplainable

violations were present. One study suggested that causal relationships should generally be

assumed to exist between any two variables and that they should only be omitted when

evidence is available.[22] We ensured that our assumptions in the DAG correctly represent the

data by using semi-synthetic data so that any possible misspecification of the DAG did not

affect the evaluation of the model transportability.

Second, we applied a SEM to the ADNI data to quantify the causal relationships in our DAG.

While SEMs are widely applied for this purpose,[16] their methodology has limitations when

using categorical variables.[58–60] In our application, we had seven categorical variables and

found a small correlation between sex and age and between age and APOE ε4. We believe

this correlation might stem from biased selection in the ADNI study, which we did not

consider in our DAG. Additionally, we found that some SEM parameter estimates were

controversial to domain knowledge. For example, the relationship between age and cognitive

impairment was estimated to be -0.13, whereas the prevalence of cognitive impairment is

known to increase with age. The incorrect SEM estimates may have altered the effects of

the interventions. For example, reducing the mean age from 73.9 to 35 only reduced the

prevalence of cognitive impairment by 2%. This was likely because the ADNI study included

only elderly (>55 years) participants, and the SEM may not have been able to estimate the

correct relationship across this limited age range. Third, it was not possible to evaluate our

results with real data from external settings. Obtaining observational data from specific

healthcare settings for external validation is often difficult due to data protection. Simulating

external data may therefore be inevitable for anticipating transportability.[61] We simulated

external validation data by intervening on specific variables (either age or APOE ε4, and tau)

at a time. These interventions simplify general distribution shifts between populations in real-

world applications where multiple variables can vary jointly. Fourth, our prediction models

had suboptimal discrimination performance, similar to other studies,[37] with the highest

achieved AUC of 0.75 in the internal validation setting. Optimizing hyperparameter did not

improve the performance. Better performance could be achieved by using multi-modal data

and deep neural networks[3,62].

Our approach to assess the transportability of models predicting cognitive impairment can be

extended to overcome the described limitations. Future work could integrate our causal data

generation approach with the work of Pölsterl et al. and include unobserved variables in the

DAG.[57] To ensure that the estimated parameters of causal relationships follow biological

laws, future refinements could include the SEM with prior distributions as implemented in the

blavaan R package[63] or explore alternative latent variable models. Future research could

adapt our approach to measure transportability to models using deep learning to other use-

cases and high-dimensional data and compare simulated data with real world data (i.e., with

a prediction model). For example, recent deep learning models predicted AD from structural

brain MR images.[7,64] Deep learning models, however, often suffer from poor calibration,[65]

whereas we suggested good calibration in the internal validation setting as a requirement for

assessing transportability using calibration metrics. New approaches to train calibrated deep

learning models offer solutions for better calibration.[66,67] Additional research to generate

synthetic medical images on the basis of causal models has emerged[68], leaving an exciting

open challenge to identify causal structures in complex data[68] and assess transportability to

new settings using interventions and synthetic images.

Conclusions

Actionable machine learning for health algorithms requires good transportability to new

settings but measuring transportability before deployment is challenging. We have used an

approach to assess the transportability of prediction models for predicting cognitive

impairment using a causal graph and semi-synthetic data to simulate different external

validation scenarios. We conclude that calibration metrics may be more suitable than AUC to

measure transportability to external settings but require good calibration performance in the

internal validation setting. Our results also contain an empirical illustration of the existing

theory that models predicting with causes of the outcome have better transportability than

those predicting with consequences of the outcome and can help to better select predictors

for prediction models. Future research can adapt our approach to other use cases and high-

dimensional data such as images and apply new interventions to simulate more realistic

external scenarios.

Declarations

This is not a human study. All data used in the study are from public databases. No new

human data is generated in the study.

Ethics, consent and permissions

Ethical review and approval and obtaining informed consent was deemed unnecessary for

the present study, because we performed secondary analyses of data obtained from the

ADNI study, which was approved by the Institutional Review Boards of all of the participating

institutions. All participants represented in the ADNI dataset provided informed written

consent as described in. [41,42] All methods were carried out in accordance with relevant

guidelines and national regulations according to the Declaration of Helsinki (consent for

research). More details can be found at adni.loni.usc.edu.

Consent for publication

Not applicable.

Availability of data and code

The TADPOLE and ADNI datasets analyzed during the current study are both available at

the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (https://ida.loni.usc.edu)

upon consenting to the data sharing agreement. The R code implementing the SEM, data

simulation, prediction models and transportability measurements is accessible at

https://github.com/JanaFe/AssessingTransportability.

Competing interests

TK reports outside the submitted work to have received research grants from the German Joint Committee and the German Ministry of Health. He further reports personal compensation from Eli Lilly and Company, Teva, TotalEnergies S.E. the BMJ, and Frontiers. MP reports having received partial funding for a self-initiated research project from Novartis Pharma. MP further reports being awarded a research grant from the Center for Stroke Research Berlin (private donations) for a self-initiated project. The declared funding sources did not influence the present research study. All other authors do not have any conflicts of interest.

Funding

This research is supported through the German Federal Ministry of Education and Research (BMBF) within the project 'Syreal' (Grant No. 01/S21069A). The funder had no role in study design, data analysis and interpretation, or writing of the report.

Author's Contributions

MP and SK conceptualized the study. JF implemented the methods in R, ran the analyses, and drafted a first version of the manuscript. JF, MP, SK interpreted the results. SK supervised the study. TK consulted the creation of the DAG. All authors provided critical input to the manuscript and approved the final version.

Acknowledgements

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's

References

1   GBD 2016 Neurology Collaborators VL, Nichols E, Alam T, Bannick MS, Beghi E, Blake
N *et al.* Global, regional, and national burden of neurological disorders, 1990-2016: a
systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Neurology*
2019; **18**: 459–480.

2   Sabbagh MN, Boada M, Borson S, Doraiswamy PM, Dubois B, Ingram J *et al.* Early
Detection of Mild Cognitive Impairment (MCI) in an At-Home Setting. *Journal of*
Prevention of Alzheimer's Disease 2020; **7**: 171–178.

3   Weiner MW, Veitch DP, Aisen PS, Beckett LA, Nigel J, Green RC *et al.* Recent
publications from the Alzheimer's Disease Neuroimaging Initiative: Reviewing progress
toward improved AD clinical trials. *Alzheimers Dementia* 2017; **13**: 1–85.

4   Sarica A, Cerasa A, Quattrone A. Random forest algorithm for the classification of
neuroimaging data in Alzheimer's disease: A systematic review. *Frontiers in Aging*
Neuroscience 2017; **9**: 1–12.

5   Moore PJ, Lyons TJ, Gallacher J. Random forest prediction of Alzheimer's disease using
pairwise selection from time series data. *PLoS ONE* 2019; **14**: 1–14.

6   Al-Amyn Valliani A, Ranti D, Oermann KE. Deep Learning and Neurology: A Systematic
Review. *Neurology and Therapy* 2019; **8**: 351–365.

7   Kang MJ, Kim SY, Na DL, Kim BC, Yang DW, Kim EJ *et al.* Prediction of cognitive
impairment via deep learning trained with multi-center neuropsychological test data.
BMC Medical Informatics and Decision Making 2019; **19**: 1–9.

8   Grueso S, Viejo-Sobera R. Machine learning methods for predicting progression from
mild cognitive impairment to Alzheimer's disease dementia: a systematic review.
Alzheimer's Research & Therapy 2021; **13**: 1–29.

9   Siontis GCM, Tzoulaki I, Castaldi PJ, Ioannidis JPA. External validation of new risk
prediction models is infrequent and reveals worse prognostic discrimination. *Journal of
Clinical Epidemiology* 2015; **68**: 25–34.

10 Steyerberg EW. *Clinical Prediction Models: A practical approach to development,
validation and updating. Second Edition.* Second Edi. Springer Nature: Cham, 2019.

11 Schölkopf B, Janzing D, Peters J, Sgouritsa E, Zhang K, Mooij JMOOIJ J. On Causal
and Anticausal Learning. .

12 Peters J, Janzing D, Schölkopf B. *Elements of causal inference: foundations and
learning algorithms.* 2018 doi:10.1080/00949655.2018.1505197.

13 Prosperi M, Guo Y, Sperrin M, Koopman JS, Min JS, He X *et al.* Causal inference and
counterfactual prediction in machine learning for actionable healthcare. *Nature Machine
Intelligence* 2020; **2**: 369–375.

14 Kilbertus N, Parascandolo G, Schölkopf B, De BM. Generalization in anti-causal
learning. *arXiv* 2018.https://arxiv.org/abs/1812.00524v1 (accessed 26 Jan2022).

15 Castro DC, Walker I, Glocker B. Causality matters in medical imaging. *Nature
Communications* 2020; **11**: 1–10.

16 Richens JG, Lee CM, Johri S. Improving the accuracy of medical diagnosis with causal
machine learning. *Nature communications* 2020; **11**: 3923.

17 Schölkopf B, Locatello F, Bauer S, Ke NR, Kalchbrenner N, Goyal A *et al.* Toward
Causal Representation Learning. *Proceedings of the IEEE* 2021; : 1–24.

18 Pearl J. Causal diagrams for empirical research. *Biometrika* 1995; **82**: 669–688.

19 Pearl J, Bareinboim E. Transportability of causal and statistical relations: A formal
approach. In: *Proceedings of the Twenty- Fifth National Conference on Artificial
Intelligence (AAAI 2011).* AAAI Press: Menlo Park, CA, 2011, pp 247–54.

20 Schölkopf B, Janzing D, Peters J, Sgouritsa E, Zhang K, Mooij J. On causal and
anticausal learning. *Proceedings of the 29th International Conference on Machine
Learning, ICML 2012* 2012; **2**: 1255–1262.

21 Pearl J, Bareinboim E. External validity: From do-calculus to transportability across
populations. *Statistical Science* 2014; **29**: 579–595.

22 Tennant PW, Murray EJ, Arnold KF, Berrie L, Fox MP, Gadd SC *et al.* Use of directed
acyclic graphs (DAGs) to identify confounders in applied health research: review and
recommendations. *International Journal of Epidemiology* 2021; **50**: 620–631.

23 Moons KGM, Kengne AP, Woodward M, Royston P, Vergouwe Y, Altman DG *et al.* Risk
prediction models: I. Development, internal validation, and assessing the incremental
value of a new (bio)marker. *Heart (British Cardiac Society)* 2012; **98**: 683–690.

24 Moons KGM, Kengne AP, Grobbee DE, Royston P, Vergouwe Y, Altman DG *et al.* Risk
prediction models: II. External validation, model updating, and impact assessment. *Heart
(British Cardiac Society)* 2012; **98**: 691–698.

25 Piccininni M, Konigorski S, Rohmann JL, Kurth T. Directed acyclic graphs and causal
thinking in clinical risk prediction modeling. *BMC medical research methodology* 2020;
20: 179.

26 Ganopoulou M, Kangelidis I, Sianos G, Angelis L. Prediction Model for the Result of
Percutaneous Coronary Intervention in Coronary Chronic Total Occlusions. *Proceedings
of the 3rd International Conference on Statistics: Theory and Applications* 2021; **2018**.
doi:10.11159/icsta21.129.

27 Gebremedhin AT, Hogan AB, Blyth CC, Glass K, Moore HC. Developing a prediction
model to estimate the true burden of respiratory syncytial virus (RSV) in hospitalised
children in Western Australia. *Scientific Reports* 2022; **12**: 1–12.

28 Sperrin M, Martin GP, Pate A, Van Staa T, Peek N, Buchan I. Using marginal structural
models to adjust for treatment drop-in when developing clinical prediction models. 2018.
doi:10.1002/sim.7913.

29 Dickerman BA, Dahabreh IJ, Cantos KV, Roger ·, Logan W, Lodi S *et al.* Predicting
counterfactual risks under hypothetical treatment strategies: an application to HIV.
European Journal of Epidemiology 123AD; **1**: 3.

30 Rojas-Carulla M, Schölkopf B, Turner R, Peters J. Invariant models for causal transfer
learning. *Journal of Machine Learning Research* 2018; **19**: 1–34.

31 Magliacane S, Van Ommen T, Claassen T, Bongers S, Mooij JM, Versteeg P. Domain
adaptation by using causal inference to predict invariant conditional distributions.
Advances in Neural Information Processing Systems 2018; **2018-Decem**: 10846–10856.

32 Singh H, Singh R, Mhasawade V, Chunara R. Fairness violations and mitigation under
covariate shift. *FAccT 2021 - Proceedings of the 2021 ACM Conference on Fairness,
Accountability, and Transparency* 2021; : 3–13.

33 Wald Y, Feder A, Greenfeld D, Shalit U. On Calibration and Out-of-domain
Generalization. In: *35th Conference on Neural Information Processing Systems
(NeurIPS 2021)*. 2021.

34 Subbaswamy A, Saria S. From development to deployment: dataset shift, causality, and
shift-stable models in health AI. *Biostatistics (Oxford, England)* 2020; **21**: 345–352.

35 Subbaswamy A, Schulam P, Saria S. Preventing failures due to dataset shift: Learning
predictive models that transport. *AISTATS 2019 - 22nd International Conference on
Artificial Intelligence and Statistics* 2020; **89**.

36 Steingrimsson JA, Gatsonis C, Li B, Dahabreh IJ. Transporting a prediction model for
use in a new target population. *American Journal of Epidemiology* 2022; : kwac128.

37 John LH, Kors JA, Fridgeirsson EA, Reps JM, Rijnbeek PR. External validation of
existing dementia prediction models on observational health data. *BMC Medical*
Research Methodology 2022; **22**: 311.

38 Li C, Chua J, Schwarzhans F, Husain R, Girard MJA, Majithia S *et al.* Assessing the
external validity of machine learning-based detection of glaucoma. *Sci Rep* 2023; **13**:
558.

39 Qin Y, Alaa A, Floto A, Schaar M van der. External validity of machine learning-based
prognostic scores for cystic fibrosis: A retrospective study using the UK and Canadian
registries. *PLOS Digital Health* 2023; **2**: e0000179.

40 Van Calster B, McLernon DJ, Van Smeden M, Wynants L, Steyerberg EW, Bossuyt P *et*
al. Calibration: The Achilles heel of predictive analytics. *BMC Medicine* 2019; **17**: 1–7.

41 Mueller SG, Weiner MW, Thal LJ, Petersen RC, Jack C, Jagust W *et al.* The Alzheimer's
disease neuroimaging initiative. *Neuroimaging Clin N Am* 2005; **15**: 869–77.

42 Petersen RC, Aisen PS, Beckett LA, Donohue MC, Gamst AC, Harvey DJ *et al.*
Alzheimer's Disease Neuroimaging Initiative (ADNI) Clinical characterization.
2010www.neurology.org.

43 Pearl J. *Causality: Models, reasoning and inference.* Cambridge University Press:
Cambridge, 2000.

44 Hernán MA, Robins JM. Causal Inference. *Causal Inference: What If* 2019; : 235–281.

45 Ankan A, Wortel IMN, Textor J. Testing Graphical Causal Models Using the R Package
"dagitty". *Current Protocols* 2021; **1**: 1–22.

46 Rosseel Y. Lavaan: An R package for structural equation modeling. *Journal of Statistical*
Software 2012; **48**. doi:10.18637/jss.v048.i02.

47 Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models
via Coordinate Descent. *Journal of Statistical Software* 2010; **33**: 1–22.

48 Breiman L. Random forests. *Machine Learning* 2001; **45**: 5–32.

49 Friedman JH. Greedy function approximation: A gradient boosting machine. *The Annals*
of Statistics 2001; **29**: 1189–1232.

50 Friedman JH. Stochastic gradient boosting. *Computational Statistics and Data Analysis*
2002; **38**: 367–378.

51 Hastie T, Tibshirani R, Friedman JH. 10. Boosting and Additive Trees. In: *The Elements*
of Statistical Learning. Springer: New York, 2009, pp 337–384.

52 Austin PC, Steyerberg EW. The Integrated Calibration Index (ICI) and related metrics for
quantifying the calibration of logistic regression models. *Statistics in Medicine* 2019; **38**:
4051–4065.

53 Ferro CAT, Fricker TE. A bias-corrected decomposition of the Brier score. *Quarterly*
Journal of the Royal Meteorological Society 2012; **138**: 1954–1960.

54   Hastie T, Tibshirani R, Friedman J. *The Elements of Statistical Learning. Data Mining,*
719   *Inference, and Prediction*. Second Edi. Springer Science + Business Media: Berlin,
720   2009.

721   55   Lebedev AV, Westman E, Van Westen GJP, Kramberger MG, Lundervold A, Aarsland D
722   *et al.* Random Forest ensembles for detection and prediction of Alzheimer's disease with
723   a good between-cohort robustness. *NeuroImage: Clinical* 2014; **6**: 115–125.

724   56   Guest F, Kuzma E, Everson R, Llewellyn DJ, David Llewellyn CJ. Identifying key
725   features for dementia diagnosis usingmachine learning. *Alzheimer's & Dementia* 2020;
726   **16**: e046092.

727   57   Pölsterl S, Wachinger C. Estimation of Causal Effects in the Presence of Unobserved
728   Confounding in the Alzheimer's Continuum. In: *Information Processing in Medical*
729   *Imaging*. Springer International Publishing, 2021, pp 45–57.

730   58   Sass DA, Schmitt TA, Marsh HW. Evaluating Model Fit With Ordered Categorical Data
731   Within a Measurement Invariance Framework: A Comparison of Estimators. *Structural*
732   *Equation Modeling* 2014; **21**: 167–180.

733   59   Bandalos DL. Relative Performance of Categorical Diagonally Weighted Least Squares
734   and Robust Maximum Likelihood Estimation. *Structural Equation Modeling* 2014; **21**:
735   102–116.

736   60   DiStefano C, Morgan GB. A Comparison of Diagonal Weighted Least Squares Robust
737   Estimation Techniques for Ordinal Data. *Structural Equation Modeling* 2014; **21**: 425–
738   438.

739   61   Frangi AF, Tsaftaris SA, Prince JL. Simulation and Synthesis in Medical Imaging. *IEEE*
740   *Trans Med Imaging* 2018; **37**: 673–679.

741   62   Qiu S, Miller MI, Joshi PS, Lee JC, Xue C, Ni Y *et al.* Multimodal deep learning for
742   Alzheimer's disease dementia assessment. *Nature Communications* 2022; **13**.
743   doi:10.1038/s41467-022-31037-5.

744   63   Merkle EC, Rosseel Y. Blavaan: Bayesian Structural Equation Models Via Parameter
745   Expansion. *Journal of Statistical Software* 2018; **85**. doi:10.18637/jss.v085.i04.

746   64   Nigri E, Ziviani N, Cappabianco F, Antunes A, Veloso A. Explainable Deep CNNs for
747   MRI-Based Diagnosis of Alzheimer's Disease. *arXiv*
748   2020.http://arxiv.org/abs/2004.12204.

749   65   Guo C, Pleiss G, Sun Y, Weinberger KQ. On Calibration of Modern Neural Networks.
750   *arXiv* 2017.

751   66   Thiagarajan JJ, Venkatesh B, Anirudh R, Bremer PT, Gaffney J, Anderson G *et al.*
752   Designing accurate emulators for scientific processes using calibration-driven deep
753   models. *Nature Communications* 2020; **11**: 1–10.

754   67   Kumar A, Sarawagi S, Jain U. Trainable calibration measures for neural networks from
755   kernel mean embeddings. *35th International Conference on Machine Learning, ICML*
756   *2018* 2018; **6**: 4378–4389.

757   68 Pawlowski N, Castro DC, Glocker B. Deep structural causal models for tractable
758   counterfactual inference. In: *34th Conference on Neural Information Processing Systems*
759   *(NeurIPS 2020)*. 2020.

760
761