Viral cultures, Polymerase Chain Reaction Cycle Threshold Values and

Viral Load Estimation for SARS-CoV-2 Infectious Potential Assessment

in Hematopoietic Stem Cell and Solid Organ Transplant Patients: A

Systematic Review.

Tom Jefferson[1], Elizabeth A. Spencer[2], John M. Conly[7], Elena C. Rosca[3] , Susanna Maltoni[4], Jon

Brassey[5], Igho J. Onakpoya[1], David H. Evans[6], Carl J. Heneghan[2], Annette Plüddemann[2*].

[1]Department for Continuing Education, University of Oxford, UK

[2]Centre for Evidence-Based Medicine, Nuffield Department of Primary Care Health Sciences,

University of Oxford, UK

[3]Victor Babes University of Medicine and Pharmacy, Timisoara, Romania

[4]Division of Research and Innovation, IRCCS Azienda Ospedaliero-Universitaria di Bologna,

Bologna, Italy

[5]Trip Database Ltd, Newport, UK

[6]Li Ka Shing Institute of Virology and Dept. of Medical Microbiology & Immunology, University

of Alberta, Canada

[7]Departments of Medicine, Microbiology, Immunology & Infectious Diseases, and Pathology &

Laboratory Medicine, Synder Institute for Chronic Diseases and O'Brien Institute for Public

Health, Cumming School of Medicine, University of Calgary and Alberta Health Services,

Calgary, Canada.

*Corresponding author

Abstract

Background: Organ transplant recipients are at increased vulnerability to SARS-CoV-2 due to

immunosuppression and may pose a continued transmission risk especially within hospital

settings. Detailed case reports including symptoms, viral load and infectiousness, defined by the

presence of replication-competent viruses in culture, provide an opportunity to examine the

relationship between clinical course, burden and contagiousness, and provide guidance on

release from isolation.

Objectives: We performed a systematic review to investigate the relationship in transplant

recipients between serial SARS-CoV-2 RT-PCR cycle threshold (Ct) value or cycle of

quantification value (Cq), or other measures of viral burden and the likelihood and duration of

the presence of infectious virus based on viral culture including the influence of age, sex,

underlying pathologies, degree of immunosuppression, and/or vaccination on this relationship.

Methods: We searched LitCovid, medRxiv, Google Scholar and WHO Covid-19 databases, from 1

November 2019 until 31 December 2021. We included studies reporting relevant data for

transplantees with SARS-CoV-2 infection: results from serial RT-PCR testing and viral culture

data from the same respiratory samples. We assessed methodological quality using five criteria,

and synthesised the data narratively and graphically.

Results: We included 10 case reports and case series reporting on 38 transplantees. We

observed a relationship between proxies of viral burden and likelihood of shedding replication-

competent SARS-CoV-2. Two individuals shed replication-competent viruses over 100 days after

infection onset. Lack of standardisation of testing and reporting platforms precludes

establishing a definitive viral burden cut-off. However, most transplantees stopped shedding

competent viruses when the RT-PCR cycle threshold was above 30 despite differences across

platforms.

Conclusions

Viral burden is a reasonable proxy for infectivity when considered within the context of the

clinical status of each patient. Standardised study design and reporting are essential to

standardise guidance based on an increasing evidence base.

56   **Keywords**

57   COVID-19; SARS-CoV-2; transmission; organ transplants; viral culture; polymerase chain

58   reaction; viral load; cycle threshold calibration; infectivity

Introduction

Hematopoietic stem cell transplant (HSCT) and solid organ transplant (SOT) recipients have significant immunosuppression, affecting both cellular and humoral immunity, and less favourable outcomes with Severe Acute Respiratory Virus Syndrome 2 (SARS-CoV-2) infection, due to the immunosuppression and/or to pre-existing comorbidities[1]. Immunosuppression associated with transplantation places patients at risk for prolonged carriage and shedding of several respiratory viruses[2]. However, identification of respiratory viral shedding, recently by reverse transcriptase polymerase chain reaction (RT-PCR), depending on the testing platform does not correlate with the presence of replication-competent virus[3]. Accordingly, we sought to perform a systematic review of RT-PCR testing and viral culture of SARS-CoV-2, focussing on people receiving solid organ or hematopoietic stem cell transplants, following our published protocol[4].

Our research questions were:

1. What is the correlation between serial SARS-CoV-2 RT-PCR cycle threshold (Ct) value or cycle of quantification value (Cq), or other measures of viral burden and the likelihood of producing replication-competent virus?

2. What is the likelihood and duration of the presence of infectious virus based on viral culture, among transplant recipients with SARS-CoV-2 infection?

3. What is the influence of age, sex, underlying pathologies and degree of immunosuppression on infectiousness of SARS-CoV-2?

4. What is the relationship of vaccination status on infectiousness with SARS-CoV-2?

We included studies reporting serial Cts from sequential RT-PCR testing or other measures of

viral burden such as RNA gene copies of respiratory samples (from nasopharyngeal or throat

specimens) along with viral culture data on the same samples, from patients about to receive a

transplant or who were post-transplant, with SARS-CoV-2 infection.

Methods

Search Strategy

We searched the following electronic databases: LitCovid, medRxiv, Google Scholar and the

WHO Covid-19 database from November 2019 until December 31, 2021. No language

restrictions were applied.

The literature search terms were: (coronavirus OR covid-19 OR SARS-CoV-2) AND

(immunosuppressed OR immunocompromised OR transplant OR immunosuppression OR

"immune deficient" OR HIV) AND (CPE OR "cytopathic effect" OR "Viral culture" OR "virus

culture" OR vero OR "virus replication" OR "viral replication" OR "cell culture" or "viral load" OR

"viral threshold" OR "log copies" OR "cycle threshold").

Screening

Four reviewers independently screened titles and abstracts to identify studies for consideration

of full text. Full text screening was performed in duplicate and disagreements arbitrated by a

fifth reviewer.

Inclusion criteria

We included studies reporting serial Cts from sequential RT-PCR testing, or RNA gene copies of

respiratory samples (nasopharyngeal, throat, sputum, bronchoalveolar lavage, endotracheal

tube secretions) AND viral culture data from the same samples from patients about to receive a

transplant or post-transplant with SARS-CoV-2 infection. We included primary studies provided

they reported sufficient information to extract quantitative data on the PCR testing and the

viral culture for each included individual. Studies that included transplant and non-transplant

patients were included if we could ascertain the results separately. Studies reported only in

poster or abstract form were excluded. Reviews were excluded but the reference lists screened

for potential relevant primary studies.

Exclusion criteria

We excluded studies using post-mortem samples only and non-respiratory samples only. We

did not include studies of non-transplant patients or those not attempting viral cultures.

Data extraction

One reviewer extracted data, which was independently checked by a second reviewer.

Disagreements were arbitrated by a third reviewer. Data were extracted on study type and

study characteristics, including population, setting, sampling and laboratory methods, clinical

information, prescribed treatments, vaccination status, laboratory findings, and clinical

outcomes. For three studies we sought clarification from the corresponding authors.

Quality assessment

We assessed the quality of included studies according to five criteria:

1. Were the criteria for diagnosing a case clearly reported and appropriate?

2. Was the reporting of patient/population characteristics including clinical symptoms,

treatments with degree of immunosuppression and outcomes adequate?

3. Was the study period, including follow-up, sufficient to adequately assess any potential

relationship between viral burden measures and likelihood of producing replication-

competent virus and the rise in neutralising antibodies? We defined sufficient as more

than one observation.

4. Were the methods used to obtain RT-PCR results replicable, generalisable and

appropriate? We considered that each study should establish the relationship between

their Ct values and the target gene copy number, using internal standards.

5. Were the methods used to obtain viral culture results replicable and appropriate? We

considered the methods used should, at a minimum, include a description of specimen

sampling and management, preparation, media and cell line used, exclusion of

contamination or co-infection (use of good controls and appropriate antibacterials and

antimycotics and possible use of gene sequencing if available), and results of inspection

of culture.

Data reporting and pooling

We reported study flow according to PRISMA reporting standards[5]. We reported study

characteristics including age, sex, clinical symptoms, treatments and events in the participants

in tabular form. We presented data on disease burden measures and on viral culture in tabular

form. For studies reporting more than one patient participant, data were extracted related to

each participant if available. We plotted median, interquartile ranges (IQRs) and outliers for

viral culture results in relation to the duration of symptoms, and individual study plots to day

120 of viral culture results and cycle thresholds.

We were unable to meta-analyse the data on PCR cycle counts/RNA log copies and viral culture,

due to a lack of detailed information on laboratory practices, assays and because of the

absence of internal controls in some studies, and heterogeneous sampling. We therefore

reviewed the studies narratively, and where possible presented the results graphically within

the limitations noted.

We analysed the relationship between cycle threshold, days of onset of symptoms and

likelihood of shedding replication-competent virus by presenting the data on a scatter plot

Results

The literature search identified 12,988 titles for screening. Of these 63 underwent full-text

review. A total of 53 studies were excluded after full-text analysis: reasons are reported in the

list of excluded studies (see Supplementary File: List of excluded studies)

The 10 included studies (Figure 1 reported data for 30 transplant patients (7 females and 23

males): renal (21), cardiac (5), bone marrow transplant (9), liver (2), bilateral lung (1). For one

study the number of males and females was not available for the relevant patient group . [6] The

38 patients were in seven countries: Saudi Arabia[7], France [8] [9], Germany [10] [11] [12], Austria [13],

Denmark [14], Canada [15] and the USA[6], and were aged between 26 and 77 years old.

A total of 36 were infected with SARS-CoV-2 post-transplant: 21 patients in 3 studies had had

kidney transplant [7 8 12], 5 patients in 4 studies had had a cardiac transplant[7 10 15 9], 1 previous

bone marrow transplant for multiple myeloma [14], 1 liver transplant [14] and eight hematopoietic

cell transplants[6]. Two patients were infected with SARS-CoV-2 and subsequently underwent

transplant: 1 liver transplant [11], 1 patient had bilateral lung transplantation after a SARS-CoV-2

infection that severely affected the lungs [13].

Typically, patients received [6]a mixture of antivirals, antibiotics, convalescent plasma and

immune suppressants, as reported in Table 1. The clinical course of COVID-19 varied widely

amongst the included patients, from mild COVID-19 related symptoms to severe pneumonia

and lung failure; no deaths were specifically reported for this group, although deaths were

reported for 4 aggregate patients in one study within 30 days of diagnosis[6]. Prescribed

treatments reflected the variation in severity.

Quality Assessment

Table 2 reports study quality based on five criteria. Three studies [15 9 12] met all five criteria.

Follow-up was judged adequate in all studies; in nine studies the reporting of patient

characteristics was sufficiently comprehensive[6 8 10 13 11 14 15 9 12] and clinical information was

missing for one study [7]. Case definition was missing or unclear in four studies[6, 7 10 14], and

methods for RT- PCR testing were unclear for three studies[6 10 13]. The methods used for viral

culture were unclear in four studies [7 10 13 11] and one study reported using a cell line that has

not typically been used to demonstrate SARS-CoV-2 growth - Buffalo green monkey kidney

(BGMK) cell line[8].

Results of the studies

The results are reported in Table 1 and Table 3. The clinical course of infection was highly

variable (Tables and 1 and 3, and Figures 2 and 3). The time from transplant to infection varied

from days to years (see for example Aydillo et al[6] and Rajakumar et al[15]). Sampling schedules

varied between studies, with no regular timetable of testing taking place, so results for PCR and

viral culture are available for different time points in a patient's clinical course and with

different gaps in time between samples being taken.

In response to our first study question on the correlation between proxy indicators of viral

burden and infectiousness, the data from Figures 2 and 3 and Table 3 indicate a correlation

between viral burden (measured as log copies or Cq/Ct) and probable infectiousness. The data

suggests that earlier symptom onset is related to the likelihood of shedding replication-

competent virus (Figure 2). The median time for a positive culture from onset of symptoms was

16 days (IQR 8 to 27; range 1-105, mean 19.7 days n = 59 cultures performed). The median for a

negative culture was 40 days, mean 40.2 days (IQR 22 to 60; range 1-119, n = 107 cultures

performed).

Five patients reported by Alshukairi et al were all culture-negative; this was in samples taken on

days 9,12, 17,18 and day 26 since symptom onset, respectively [7].

Seven stem cell transplant recipients from the Aydillo et al letter who had repeated culture

assessments fluctuated in and out of shedding replication-competent virus. One who had a

single culture attempt was positive with an estimated Ct of 17.5 [6]. Patient MSK-6 shed

replication competent virus until day 62 from symptom onset with an estimated Ct of 19. MSK-

6 showed the longest duration infectious transplant recipient from the Aydillo et al letter[6].

Eight kidney transplant patients described by Benotmane and colleagues had positive viral

cultures [8].

For six patients we could identify the duration of probable infectiousness, which ranged from 8

days (patient 5: day 1 to 8) to 32 days (patient 7: day 7 to 38). Four patients were infectious

with reported Cts> 30 based on the individual platforms that were used to perform the Cts.

Rajakumar et al[15] described two cardiac transplant patients: viral culture found replication-

competent virus in samples from one patient on day 16 and in samples from the other patient

on day 4 and repeatedly up to day 27, after which all viral cultures were negative[15].For each

patient, viral culture was negative (i.e. no replication-competent virus observed) in samples

with PCR cycle counts of over 25. Within the samples giving positive viral cultures, the PCR

results showed that the cycle threshold for the N gene was lower than for the E gene by an

average of 5.4 Ct values.

In the study by Niyonkuru et al, the duration of infectiousness in the two patients, as indicated

by replication-competent virus, was 8 and 9 days (Figure 3) [14]. A cardiac transplant patient

described by Tarhini[9] and colleagues tested culture-positive with a Ct of 23 on day 103; all other

viral cultures were negative from samples with PCR Cts of 18 to over 40 [9].

Weigang et al[12] described a kidney transplant patient who experienced three hospital

admissions. During the first one (day zero to day 72), 19 RT-qPCR tests were performed, and

alongside that viral culture was performed, showing 8/19 positive cultures (Ct values ranging

from 15 to 25) and 11/19 negative (Ct values from 25 to 30). The patient was culture positive

again on day 105 (Ct of 23). After re-admission at day 140 the patient was still RT-qPCR positive,

but with viral culture negative; he was treated for 10 days (days 141-149) with remdesivir.

Subsequently, negative RT-qPCR tests until day 189 and negative cultures suggested that the

infection had resolved[12].

A heart transplant patient described by Decker[10] and co-workers had a positive viral culture on

day 18 and day 21 with 6.2 and 6.5 log10 copies/ml. [10]

Although the dataset was limited, we observed an inverse relationship (Ct/Cq) or direct

relationship (log copies): the viral burden indicated by these methods correlated with

infectiousness, as shown by the ability to produce replication-competent virus in culture. The

presence of replication-competent virus reflects one of the highest grades of evidence

supporting the capability for forward transmission of SARS-CoV-2[16, 17].

The robustness of the correlation is difficult to assess because laboratory methods differ; it was

not possible to pool the data to produce a summary cut-off value for infectiousness, due to

these variations and due to varying time windows for sampling from patients (see Figure 2 and

Table 3).

In response to our second research question (on the likelihood and duration of infectiousness

among transplant recipients with SARS-CoV-2 infection) the data indicate that regardless of

differences in laboratory practices, observed prolonged shedding of replication-competent

virus is associated with alternating increases and decreases of viral burden over time, which in

some cases may be up to around 100 days [9] [12].

Figure 4 shows the relationship between cycle threshold, symptom onset (in days) and

likelihood of shedding replication competent virus. In Benotmane, five results were reported

with a CT of 30 or above for a positive culture. [8] However, the cell line used in this study is not

demonstrated to support SARS-CoV-2 growth. In the other studies, despite a minimum of 10 different

PCR platforms being used and different culture techniques, the culture results are insensitive to

cycle thresholds above 30. The viral load estimates are affected by administration of courses of

anti-viral treatment including remdesivir. See Figures 3a (Cts/Cqs) and 3b (log copies).

Responding to our third research question (the influence of age, sex, underlying pathologies

and degree of immunosuppression on infectiousness): at present the heterogeneity and limited

amount of the available data preclude answering this question.

We are unable to answer our fourth and final research question on the relationship of

vaccination status on infectiousness because no study reported on vaccination status for these

transplant patients.

Discussion

This review included 10 reports of studies using viral culture and RT-qPCR testing among 38 transplant patients with immunosuppressive treatment who experienced COVID-19 infection. The evidence indicates a relationship between indicators of viral burden (Ct, Cq or RNA log copies) and probable infectiousness as indicated by the presence of replication-competent virus. Gaps in the data remain due to variable methods and reporting and establishing summary estimates of the relationship has not been possible. The data show a long-term rise and fall of viral burden associated with the likelihood of infectiousness that in some transplant patients appears to be a sequential pattern of going in and out of infectiousness. Replication-competent virus was most commonly observed in samples with PCR Ct values under 25; one study was an exception to this by reporting viable virus at Ct>30, but the use of a cell line not typically used for SARS-CoV-2 isolation makes interpretation unclear[8]. The duration of viral RNA shedding was variable, with the longest duration reported at 105 days[12]. Our findings suggest a Ct of 30 or greater, regardless of the platform, may be used as a reasonable proxy to rule out infectious SARS-CoV-2 as there is a consistent correlation between a rising Ct value and likelihood of isolating replication-competent virus. Such a value would be useful to guide clinicians managing these difficult patients, particularly if there were repeated values in this range. Below a Ct of 30, clinicians may choose to repeat NP or throat swabs to assess the direction of the Ct values to allow a more dynamic assessment which taken in conjunction with the clinical status may faciltate decision making for isolation or antiviral treatment considerations.

Regarding our third review question on the influence of patient variables on the likelihood of

the presence of infectious SARS-CoV-2, the included studies showed substantial heterogeneity;

some had missing data or few cultures available, and meta-analysis or pooling was not possible.

Variability in the clinical course of SARS-CoV-2 infection among transplant recipients has been

reported, including observed prolonged viral shedding[18]. Antiviral drugs may impact on these

observations, especially symptoms and viral burden. [19]

Two well-designed studies on immunosuppressed patients, which we were unable to include

because disaggregated data solely for transplant patients were not fully available, support our

conclusions[6, 20]. While this review is limited to transplant patients, evidence suggests similar

prolonged viral cultures are found in immunosuppressed cancer patients. We plan to perform a

further review in this group analysing the type of cancer and the impact of immunotherapies on

viral culture findings.

The transplant patient population is of particular importance: clinicians need guidance as to

when to release the patient from quarantine or isolation, given the heavy burden of

immunosuppression. We have tried to narrow the uncertainty and offer some general guidance

as to when patients are unlikely to be shedding replication-competent virus, but clinical

assessment of each patient must inform that decision because each patient and setting is

different.

The strengths of this review are that we followed our published protocol, entailing rigorous

literature searches, double checked data extraction and quality assessment, and a high level of

clinical and epidemiological expertise input to deliberate the findings. We were also able to

include data from an additional 8 transplant recipients after extensive correspondence with the

study authors[6]. Limitations include the small number of studies with viral culture and serial viral

load estimates among transplant patients, high variability in study design and reporting and

impossibility to pool results due to the well-known variability in sensitivity across assays[21].

Case series are conventionally considered low in the evidence hierarchy, as they may entail

inherent bias in the selection of study participants and therefore have limited generalisability;

however, here they are essential in providing the detailed reports needed for this unusual

patient group. The case reports included here comprise some of the most detailed longitudinal

reports of this patient group for whom data are needed. The evidence base is limited, however,

by heterogeneous design and reporting within the studies with, for example, different

observation windows for reporting of viral burden and culturability or clinical characteristics of

patients.

In addition to providing appropriate care for the individual patient, ongoing transmission of

SARS-CoV-2 is a concern, and immunosuppressed individuals may pose a challenge by

experiencing prolonged carriage of the virus that could lead to forward transmission. Based on

our findings we would offer the following general guidance to clinicians:

Physicians who are experienced with these immunosuppressed patient populations should

work with public health to direct their isolation and quarantine requirements. Infectious

patients with immunosuppressive treatment following solid organ or stem cell transplantation

should be isolated until at least two consecutive respiratory specimens collected ≥24 hours

apart demonstrate a rising RT-PCR Ct (i.e. indicating diminishing viral burden). After discharge,

they should be closely followed up for SARS-CoV-2 infection for several weeks to months,

depending on the individual clinical scenario.

For obtaining data, standardisation of methods is needed: each laboratory should use

consistently applied platforms with suitable internal standards to calibrate the relationship

between Ct and genome copy in these patient populations.

Publication of results of case series or other longitudinal study should be reported in a

standardised format to avoid loss of data. We suggest observation windows should be within a

short range of 3 to 7 days during the acute periods post-transplantation and during periods of

rejection when higher doses of immunosuppressants are employed, depending on clinical

circumstances. Each observation window should include a summary of symptoms and

interventions, the reporting of PCR cycle threshold and, for samples with Ct below 30, attempts

at viral culture if available. Description of patients should include past medical histories and

details of treatments received. Observed drug interactions should be highlighted. Reasons for

admission, discharge and changes in isolation should be clearly reported. To investigate the

duration of viral shedding, studies should report the time between the first positive and the

first negative viral cultures.

With additional data gathering and standardisation of methods, it will be possible for transplant

physicians to develop evidence-based approaches to dealing with these patients for the benefit

of the patients and their families and the community at large.

Acknowledgements

We gratefully acknowledge the contributions of Drs Mini Kamboj and Jeroen van Kampen

who provided additional data from their studies and helped us to progress this work.

Funding

This work is supported by the National Institute for Health Research School for Primary

Care Research [Project 569] and by the University of Calgary.

Author contributions

TJ, CH and JC designed the study. JB performed the literature searches. JB, TJ, SM, ER and

ES, screened the studies for eligibility and performed data extraction. Additional expertise

on clinical and laboratory issues was given by DE, JC, SM and ER. CH generated the data

figures. All authors contributed to interpreting and writing up the results and conclusions.

Ethics declarations

Ethics approval and consent to participate:

This is a systematic review and meta-analysis. Therefore, ethical approval is not applicable.

Consent for publication:

Not applicable.

Availability of data and materials

All data generated or analysed during this study are included in this published article [and its

supplementary information files].

Competing Interests

TJ's competing interests are accessible at: https://restoringtrials.org/competing-interests-tom-

jefferson

CJH holds grant funding from the NIHR, the NIHR School of Primary Care Research, the NIHR

BRC Oxford and the World Health Organization for a series of Living rapid review on the modes

of transmission of SARs-CoV-2 reference WHO registration No2020/1077093. He has received

financial remuneration from an asbestos case and given legal advice on mesh and hormone

pregnancy tests cases. He has received expenses and fees for his media work including

occasional payments from BBC Radio 4 Inside Health and The Spectator. He receives expenses

for teaching EBM and is also paid for his GP work in NHS out of hours (contract Oxford Health

NHS Foundation Trust). He has also received income from the publication of a series of toolkit

books and for appraising treatment recommendations in non-NHS settings. He is Director of

CEBM and is an NIHR Senior Investigator.

DE holds grant funding from the Canadian Institutes for Health Research and Li Ka Shing

Institute of Virology relating to the development of Covid-19 vaccines as well as the Canadian

Natural Science and Engineering Research Council concerning Covid-19 aerosol transmission.

He is a recipient of World Health Organization and Province of Alberta funding which supports

the provision of BSL3-based SARS-CoV-2 culture services to regional investigators. He also holds

public and private sector contract funding relating to the development of poxvirus-based Covid-

19 vaccines, SARS-CoV-2-inactivation technologies, and serum neutralization testing.

JMC holds grants from the Canadian Institutes for Health Research on acute and primary care

preparedness for COVID-19 in Alberta, Canada and was the primary local Investigator for a

Staphylococcus aureus vaccine study funded by Pfizer for which all funding was provided only to

the University of Calgary. He is co-investigator on a WHO funded study using integrated human

factors and ethnography approaches to identify and scale innovative IPC guidance

implementation supports in primary care with a focus on low-resource settings and using drone

aerial systems to deliver medical supplies and PPE to remote First Nations communities during

the COVID-19 pandemic. He also received support from the Centers for Disease Control and

Prevention (CDC) to attend an Infection Control Think Tank Meeting. He is a member and Chair

of the WHO Infection Prevention and Control Research and Development Expert Group for

COVID-19 and a member of the WHO Health Emergencies Programme (WHE) Ad-hoc COVID-19

IPC Guidance Development Group, both of which provide multidisciplinary advice to the WHO

and for which no funding is received and from which no funding recommendations are made

for any WHO contracts or grants. He is also a member of the Cochrane Acute Respiratory

Infections Working Group.

References

422
423

1.   Jering, K.S., et al., *Excess mortality in solid organ transplant recipients hospitalized with COVID-19: A large-scale comparison of SOT recipients hospitalized with or without COVID-19.* Clin Transplant, 2022. **36**(1): p. e14492. 10.1111/ctr.14492.

2.   Andersen, K.M., et al., *Long-term use of immunosuppressive medicines and in-hospital COVID-19 outcomes: a retrospective cohort study using data from the National COVID Cohort Collaborative.* (2665-9913 (Electronic)).

3.   Fernández-Ruiz, M. and J.M. Aguado, *Severe acute respiratory syndrome coronavirus 2 infection in the stem cell transplant recipient - clinical spectrum and outcome.* Curr Opin Infect Dis, 2021. **34**(6): p. 654-662. 10.1097/qco.0000000000000790.

4.   Jefferson, T., et al., *Viral cultures, PCR Cycle threshold values and viral load estimation for COVID-19 infectious potential assessment in transplant patients: systematic review - Protocol Version 30 December 2021.* medRxiv, 2022: p. 2021.12.30.21268509. 10.1101/2021.12.30.21268509.

5.   Page, M.J., et al., *The PRISMA 2020 statement: an updated guideline for reporting systematic reviews.* BMJ, 2021. **372**: p. n71. 10.1136/bmj.n71.

6.   Aydillo, T., et al., *Shedding of Viable SARS-CoV-2 after Immunosuppressive Therapy for Cancer.* New England Journal of Medicine, 2020. **383**(26): p. 2586-2588. 10.1056/NEJMc2031670.

7.   Alshukairi, A.N., et al., *Test-based de-isolation in COVID-19 immunocompromised patients: Cycle threshold value versus SARS-CoV-2 viral culture.* International Journal of Infectious Diseases, 2021. **108**: p. 112-115. 10.1016/j.ijid.2021.05.027.

8.   Benotmane, I., et al., *Long-term shedding of viable SARS-CoV-2 in kidney transplant recipients with COVID-19.* American Journal of Transplantation, 2021. **21**(8): p. 2871-2875. https://doi.org/10.1111/ajt.16636.

9.   Tarhini, H., et al., *Long-Term Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infectiousness Among Three Immunocompromised Patients: From Prolonged Viral Shedding to SARS-CoV-2 Superinfection.* The Journal of Infectious Diseases, 2021. **223**(9): p. 1522-1527. 10.1093/infdis/jiab075.

10.   Decker, A., et al., *Prolonged SARS-CoV-2 shedding and mild course of COVID-19 in a patient after recent heart transplantation.* Am J Transplant, 2020. **20**(11): p. 3239-3245. 10.1111/ajt.16133.

11.   Niess, H., et al., *Liver transplantation in a patient after COVID-19 - Rapid loss of antibodies and prolonged viral RNA shedding.* Am J Transplant, 2021. **21**(4): p. 1629-1632. 10.1111/ajt.16349.

12.   Weigang, S., et al., *Within-host evolution of SARS-CoV-2 in an immunosuppressed COVID-19 patient: a source of immune escape variants.* medRxiv, 2021: p. 2021.04.30.21256244. 10.1101/2021.04.30.21256244.

13.   Lang, C., et al., *Lung transplantation for COVID-19-associated acute respiratory distress syndrome in a PCR-positive patient.* The Lancet Respiratory Medicine, 2020. **8**(10): p. 1057-1060. 10.1016/S2213-2600(20)30361-1.

14.   Niyonkuru, M., et al., *Prolonged viral shedding of SARS-CoV-2 in two immunocompromised patients, a case report.* BMC Infectious Diseases, 2021. **21**(1): p. 743. 10.1186/s12879-021-06429-5.

15.   Rajakumar, I.A.-O., et al., *Extensive environmental contamination and prolonged severe acute respiratory coronavirus-2 (SARS CoV-2) viability in immunosuppressed recent heart transplant recipients with clinical and virologic benefit with remdesivir.* (1559-6834 (Electronic)).

16. Jefferson, T., et al., *Transmission of Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-*
CoV-2) from pre and asymptomatic infected individuals. A systematic review. Clinical
Microbiology and Infection, 2021. https://doi.org/10.1016/j.cmi.2021.10.015.
17. Jefferson, T.H., C.; Spencer, E.; Brassey, J.; Pluddeman, A.; Onakpoya, I.; Evans, D.; Conly, J. A
Hierarchical Framework for Assessing Transmission Causality of Respiratory Viruses. PrePrints,
2021. 10.20944/preprints202104.0633.v1.
18. Marinelli, T., et al., *Prospective Clinical, Virologic, and Immunologic Assessment of COVID-19 in*
Transplant Recipients. Transplantation, 2021. **105**(10): p. 2175-2183.
10.1097/tp.0000000000003860.
19. Thornton, C.S., et al., *Prolonged SARS-CoV-2 infection following rituximab treatment: clinical*
course and response to therapeutic interventions correlated with quantitative viral cultures and
cycle threshold values. Antimicrob Resist Infect Control, 2022. **11**(1): p. 28. 10.1186/s13756-022-
01067-1.
20. van Kampen, J.J.A., et al., *Shedding of infectious virus in hospitalized patients with coronavirus*
disease-2019 (COVID-19): duration and key determinants. medRxiv, 2020: p.
2020.06.08.20125310. 10.1101/2020.06.08.20125310.
21. Bruce, E.A., et al., *Predicting infectivity: comparing four PCR-based assays to detect culturable*
SARS-CoV-2 in clinical samples. EMBO Mol Med, 2021: p. e15290. 10.15252/emmm.202115290.
485
486

504 **List of supplementary files.**

506 Supplementary file. Literature search strategy.

508 Supplementary file. List of excluded studies, with reasons.

510 **Table 1.** Characteristics of transplant patients in included studies.

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
Alshukairi 2021	Pt1: 34 yr F Pt2: 71 yr M Pt3: 75 yr M Pt4: 26 yr M	Pt1: cardiac, in 2014, tacrolimus, MMF, prednisolone Pt2: renal, in 2014,	Pt1: Severe pneumonia; Pt2: pneumonia	Pt1: high-flow nasal cannula Pt2: low-flow cannula

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
	Pt5: 38 yr F	tacrolimus, MMF, prednisolone, DM, HTN, CAD Pt3: renal, in 2014, tacrolimus, MMF and prednisolone, HTN Pt4: renal, in 2018, tacrolimus, MMF, prednisolone, DM Pt5: renal in 2014, tacrolimus, AZA, prednisolone, APS & hypothyroidism.	Pt3: pneumonia Pt4: pneumonia Pt5: upper RTI	Pt3: low-flow cannula Pt4: oxygen not required Pt5: no cannula
Aydillo 2020	8 ppts (no. of M and F and age N/A for this	Various antibiotics and immunosuppressants	Variable from mild to severe	Hydroxychloroquine, remdesivir,

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
	subset)			azythromicin, convalescent plasma
Benotmane I 2021	14 M, 2 F, median age 63.3 yrs. Pts with at least two positive for SARS-CoV-2 NP swabs (of which one collected at least 7 days after symptom onset) during the follow-up period were	16 kidney transplant recipients; median time from transplant 3.8 yrs. Antithymocite globulin: 8/16; anti-CD25: 8/16; tacrolimus: 10/16; ciclosporin: 3/16; MMF/MPA: 14/16; mTOR inhibitors: 2/16; steroids: 10/16;	All 16 hospitalized for symptomatic COVID-19 between 4 March and 15 April 2020	lopinavir/ ritonavir: 1/16; hydroxychloroq uine: 8/16; tocilizumab: 2/16

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
	eligible for inclusion	belatacept: 2/16		
Decker A 2020	62 yr M	Heart transplant Nov 2019; subsequently pneumonia and acute respiratory distress syndrome; intermittent renal replacement therapy. Cyclosporine A (target range 135 ± 30 ng/mL), mycophenolate mofetil 500 mg b.i.d., prednisone 10 mg q.d.	1 March 2020 onset of symptoms and +ve PCR; mild symptoms, no cardiorespiratory decline, several weeks +ve.	hydroxychloroquine (400mg b.i.d. + 200mg b.i.d.) from day 7 to 14

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
		+ cotrimoxazole and due to cytomegalovirus high-risk constellation (D + R−), ganciclovir for 4 months post-transplantation, then valganciclovir prophylaxis.		
Lang C 2020	44 yr old F with mild untreated psoriatic arthritis and idiopathic CD4 lymphocytopen	Bilateral lung transplant day 58 after admission for Covid-19. Subsequently, standard triple	Day 0 admitted with fever, cough, +ve NP RT-PCR. Day 6 to ICU & intubation; day	Immunoglobulins, tocilizumab & lopinavir. Day 6 to ICU & intubation; day

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
	ia	immunosuppression was initiated, including tacrolimus, mycophenolate mofetil, and steroids; also 6 additional treatment cycles of immunoabsorption and antithymocyte globulin.	13 ECMO. Day 52 preparation begun for transplant: immunoabsorption therapy; day 58 bilateral lung transplant performed. Transferred to non ICU ward day 121.	13 ECMO. Bilateral lung transplant
Niess 2020	56 yr M patient listed for liver transplantation with a Model	admission for liver transplant: 18/3/2020 positive to COVID-19 screening on		

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
	For End-Stage Liver Disease Score of 19 points due to cryptogenic cirrhosis and a history of hepatitis B	25/3/2021 after 31-32 days: PCR negative + seroconversion after 36 days of symptom onset and 21 days after seronconversion: transplant; tacrolimus (to target of 4 to 7 ng/ml) + steroids		
Niyonkuru M 2021	Pt1: 66 yr M, recent liver transplant Pt2: 70 yr M,	Pt1: fatigue and tachypnea, hospital admission, tested positive for SARS-CoV-2	Pt1 day 0 +ve test. Symptom onset day 5, diminished taste and	Pt1: In ICU, non invasive ventilation, IV dexamethasone 6 mg daily

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
	previous bone marrow transplant for multiple myeloma	Pt2: pacemaker. Lenalidomide 25 mg daily. elevated CRP concentration (130 mg/L). Nocardia Farcinica was detected in blood cultures and pus from abscesses on the left leg. Trans-esophageal echocardiography showed vegetation on the pacemaker electrode.	smell; day 12 additional symptoms; day 14 hospital admission. On day 12 from symptom onset transferred to the ICU for non invasive ventilation. Pt2 admitted on October 21 due to fever and elevated	and remdesivir: 200 mg the first day and 100 mg the following 4 days. Pt2: no COVID-19 treatment (asymptomatic)

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
			CRP concentration (130 mg/L) because of systemic infection due to pacemaker contamination; periodic screening during admission; positive by week 4, always asymptomatic of COVID-19.	

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
Rajakumar 2021	Two cardiac transplant recipients (<3 months post-transplant): Pt1: 56 yr F; history of dilated cardiomyopathy with end-stage heart failure, type-2 diabetes mellitus, hypothyroidism, osteoporosis, and anemia.	Pt1: orthotopic heart transplant with antithymocyte globulin induction & standard triple immunosuppressive therapy & was discharged 30 days later. Pt2: orthotopic heart transplant with antithymocyte globulin induction therapy, discharged 4 weeks later.	NP swabs (& saliva & clinical & environmental samples) were collected at regular intervals beginning shortly after admission. Pt 1: 5 days post-discharge, rehospitalized following community	Pt1: after day 15, corticosteroids & antimicrobials initiated for presumptive COVID-19 pneumonitis & superimposed bacterial pneumonia; day 21 O2 requirements increased significantly. Despite

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
	Pt2: 33 yr F with a history of end-stage heart failure 2ndary to congenital heart disease, liver cirrhosis & kidney disease.	Pt1 and Pt2: prednisone, tacrolimus, and mycophenolate mofetil, and standard prophylactic medications	exposure to COVID-19. Nasal stuffiness & discharge, sneezing, fatigue & cough developed on admission day 4; NP swab +ve for SARS-CoV-2. All symptoms except fatigue resolved by day 12.	discontinuation of MMF & reduction of tacrolimus, cultivatable viral loads increased in the NP & saliva; chest radiograph & clinical condition deteriorated; mechanical ventilation considered. 10-day course of

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
			Dyspnea, cough & hypoxia then developed on day 15; chest radiograph revealed new bibasilar interstitial infiltrates. Discharged day 44. Pt2: tested +ve for SARS-CoV-2 shortly after discharge,	remdesivir initiated day 27. Clinical condition & chest radiograph improved; O2 discontinued day 32; discharged day 44. Pt2: intubation. MMF dose was reduced; dexamethasone &

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
			readmitted 1 week later with progressive dyspnea & hypoxemia requiring intubation. Significant viable virus burdens were noted in the NP swab (& saliva, & a used face cloth) 16 days after	antimicrobials initiated. MMF was subsequently held & later restarted when an echocardiogram demonstrated a reduction in left ventricular systolic function, suggesting acute graft rejection.

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
			initial SARS-CoV-2 positivity. 5-day course remdesivir initiated followed by clinical recovery & discharge 7 days later.	Pt1: 10-day course of remdesivir from day 27 to day 37 from hospital admission; improvement and subsequent discharge on day 44 Pt2: 5-day course remdesivir,

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
				improvement and discharge 7 days later
Tarhini 2021	71yr European M	Cardiac transplant, also diabetes mellitus & chronic kidney disease; prednisone, mycophenolic acid, belatacept	15 April 2020 hospitalized for asthenia, dry cough, myalgia, and low-grade fever for 1 week. No dyspnea or O2 need. +ve on admission and day 14 with <10% Covid-19	At readmission, treated in ICU for cardiac decompensation due to underlying respiratory infection. [Unclear what happened after this; last test day 121 & culture +ve at

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
			involvement on CT scan. Discharged day 39 after clinical improvement, despite persistent +ve PCR at day 32 (21 Ct). 23 June readmitted with dry cough, dyspnea, and O2 requirement.	day 103, negative serology throughout]

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
Weigang 2021	58 yr M	kidney transplant, also CHD, arterial hypertension, hyperlipidemia, obesity; basiliximab (20mg, day 0 and day 4 post-transplantation) and prednisone (250mg at day 0, 125 mg day 1, 50mg day 2 to 5, 20mg day 6 to 10, then 15 mg/day). Additionally tacrolimus (10mg day 0, 8mg day 1,	March 2020, mild respiratory symptoms & tested positive for SARS-CoV-2 for over 145 days; discharged on day 72 for home quarantine; re-hospitalized at day 106 to 126 due to another kidney transplant	Ivermectin 4 day course (33mg/day, day 56 to 60); 10 day course remdesivir (200 mg on day 140, then 100 mg/daily, day 141 to 149)

Study ID	Participants (data extracted for transplant patients)	Transplant, immunosuppressive treatment & comorbidities	Clinical course of COVID-19 infection	COVID-19 treatments
		5.5mg day 2, 5mg day 3 and 4, then 4mg/day) and Mycophenolate mofetil (2000mg/ day). Maintenance immunosuppression consisted of tacrolimus (4 to 6mg/day), mycophenolate mofetil (2,000mg/day), and prednisone (10 to 20mg/day)	failure. Days 140 to 149 treated with remdesivir; subsequent PCR tests negative & virus isolation negative.	

Abbreviations:

APS = anti-phospholipid syndrome

AZA = azathioprine

CAD = coronary artery disease

DM = diabetes mellitus

HTN = hypertension

MMF = mycophenolate mofetil

NP = nasopharyngeal

Pt = patient

RTI = respiratory tract infection

RT-PCR = reverse transcriptase polymerase chain reaction

Table 2. Quality of included studies.

Study ID	Were the criteria for diagnosing a case clearly reported and appropriate?	Was the reporting of patient/ population characteristics adequate?	Was the study period, including follow-up, sufficient?	Were the methods used to obtain RT-PCR results replicable and appropriate?	Were the methods used to obtain viral culture results replicable and appropriate?
Alshukairi 2021	Unclear*	No**	Yes	Yes	Unclear
Aydillo 2020	Unclear	Yes	Yes	Unclear	Yes
Benotmane I 2021	Yes	Yes	Yes	Yes	No***
Decker A 2020	No	Yes	Yes	Unclear	Unclear
Lang C 2020	Yes	Yes	Yes	Unclear	Unclear
Niess 2020	Yes	Yes	Yes	Yes	Unclear
Niyonkuru M 2021	No*	Yes	Yes	Yes	Yes
Rajakumar 2021	Yes	Yes	Yes	Yes	Yes

Tarhini 2021	Yes	Yes	Yes	Yes	Yes
Weigang 2021	Yes	Yes	Yes	Yes	Yes

525   *case definition unclear, article reports positive RT-PCR, but Ct cut-off not reported.

526   ** data on clinical symptoms lacking

527   ***The cell line used was not one that is demonstrated to support SARS-CoV-2 growth.

528   Therefore the cell culture results are not reliable.

530   **Table 3.** PCR cycle counts/log copies and viral culture results of included studies.

Study ID	Symptoms, days reported	RT-PCR Cycle count/log copies results	Viral culture results (days)
Alshukairi 2021	NR no viral culture at admission/onset of symptoms	Pt1: D3: positive, Ct NA, D26: positive, 22.87 Pt2: D3: positive, 11.58; D17: positive, 23.12 Pt3: D6: positive, 8.82, D19: positive, 13.88 Pt4: D4: positive, 10.38; D12: positive, 27.57 Pt5: D1: positive, 2.8, D9: positive, 14.84	Pt1: D26: negative Pt2: D17: negative Pt3: D19: negative Pt4: D12: negative Pt5: D9: negative
Aydillo 2020 *	All symptomatic; variable severity	Pt1: D3: 22.5, D27: 34 Pt2: D2: 18, D27: 32 Pt4: D5: 19, D28: 18.5, D50: 27.5 Pt6: D18: 21, D27: 21.5, D33: 30.5, D38: 20, D45: 19.5, D52:	Pt1 D3: positive, D27: negative Pt2: D2: positive, D27, negative Pt4: D5: positive, D28: positive, D50: negative Pt6: D18: negative, D27: negative, D33: negative, D38:

Study ID	Symptoms, days reported	RT-PCR Cycle count/log copies results	Viral culture results (days)
		20, D62: 18.5	negative, D45: positive, D52: positive, D62: positive
		Pt8: D1: 22, D17: 21, D22: 22, D37: 35	Pt8: D1: positive, D17: positive, D22: negative, D37: negative
		Pt9: D4: 13, D8: 22, D46: 33.5, D56: 27, D82: 38	Pt9: D4: positive, D8: positive, D46: negative, D56: negative, D82: negative
		Pt13: D2: 21.5, D50: 31.5	Pt13: D2: positive, D50: negative
		Pt14: D8: 17.5	Pt14: D8: positive
Benotmane 2021			Pt1: D5, D14: negative;
			Pt2: D21, D29: negative;
			Pt3: D25, D39: negative
			Pt4: D2: positive, D9, D16, D35: negative
		Patient 6: D23: Ct > 30; D29: Ct > 35	Pt5: D1, D8: positive; D23, D30: negative

Study ID	Symptoms, days reported	RT-PCR Cycle count/log copies results	Viral culture results (days)
		Patient 7: D38: Ct > 35 Patient 8: D28: Ct > 30 Patient 11: D27: Ct > 30	Pt6: D6, D9, D23, D29: positive Pt7: D7, D38: positive Pt8: D1: negative, D8, D28: positive Pt9: D11: negative, D30: negative Pt10: D2, D8: negative Pt11: D11, D27: positive Pt12: D17, D27: negative Pt13: D2, D9: positive; D31: negative Pt14: D3, D19: negative Pt15: D4, D17: negative Pt16: D20: positive, D30: negative.
Decker 2020	Mild symptoms. Day 0: transient	PCR +ve on days 1, 5, 7, 11, 18, 21, 25, 28, 33, and still on day 35 PCR remained positive on day 35	Viral culture +ve at day 18 and day 21 post-onset of symptoms.

Study ID	Symptoms, days reported	RT-PCR Cycle count/log copies results	Viral culture results (days)
	episode of fever & sore throat ; day 7 temperature increase; no clinical symptoms after day 20.	with copy numbers similar to the onset of infection. Concurrent with the second onset of fever there was an increased viral load after day 7 that slowly returned to the level of infection onset.	
Lang 2020	Admitted with fever & cough becoming severe and life threatening leading to bilateral lung transplant.	Day 0, Ct 27; Day 17, Ct 21; Day 23, Ct 23; Day 19, Ct 32; Day 31, Ct 32; Day 36, Ct 29; Day 48, Ct 39; Day 53, Ct 34; Day 59, Ct 33; Day 61, PCR negative; Day 62, Ct 36; Day 62, PCR negative; Day 64, PCR negative; Day 65, Ct 36; Day 66, Ct 39; Day 69, Ct 39; Day 70, PCR negative; Day 72, PCR negative; Day 74, PCR negative;	Samples cultured from day 48 and day 65; both negative by cell culture.

Study ID	Symptoms, days reported	RT-PCR Cycle count/log copies results	Viral culture results (days)
		Day 76, PCR negative; Day 95, PCR negative;	
Niess 2020	Mild symptoms of malaise and a dry cough	day 0: positive but asymptomatic day 8, 12, 19, 22 - positive PCR day 31, 32 - negative PCR day 49, 55, 60 - positive PCR day 58, 63, 64 - negative PCR results	Positive pre-transplant PCRs were not confirmed by cell cultures 4 negative viral cell culture results from samples taken on days 49, 54, 60 and 69 (*)
Niyonkuru 2021	Pt1: fatigue and tachypnea, then ventilation required. Pt2:	Pt1: day 15 from 1st positive PCR/day 3 from symptoms onset: Cq 21.7; day 21 from 1st positive PCR/day 9 from symptoms onset: Cq 19.21; day 43 from 1st positive PCR/day 13 from symptoms onset Cq 35.45	Pt1: day 15 from 1st positive PCR/day 3 from symptoms onset viral culture at 61,277 PFU/swab; day 21 from 1st positive PCR/day 9 from symptoms onset viral culture at 256,410 PFU/swab;

Study ID	Symptoms, days reported	RT-PCR Cycle count/log copies results	Viral culture results (days)
	asymptomatic for COVID-19.	Pt2: day 9 from 1st positive PCR: Cq: 22.33; day 12 from 1st positive PCR Cq 22.57; day 14 from 1st positive PCR: Ct approx 33.5; day 18 from 1st positive PCR: PCR negative.	day 22 from 1st positive PCR/day 13 from symptoms onset viral culture negative. Pt2: day 9 from 1st positive PCR: culture positive with 11082 PFU/swab; day 12 from 1st positive PCR: culture positive with PFU 55944/swab; day 14 from 1st positive PCR: culture negative.
Rajakumar 2021		Pt 1: hospital admission, positive PCR day 4 post-symptom onset Day 4: N gene Ct 16, E gene Ct 19 Day 12: N gene Ct 13, E gene Ct 20	Pt1 Viral culture Day 4: positive Day 12: positive Day 20: positive Day 27: positive

Study ID	Symptoms, days reported	RT-PCR Cycle count/log copies results	Viral culture results (days)
		Day 20: N gene Ct 14, E gene Ct 19	Day 32: negative
		Day 27: N gene Ct 15, E gene Ct 21	Day 60: negative
		Day 32: N gene Ct 32, E gene Ct 28	Pt 2: Viral culture
		Day 60: N gene Ct 30, E gene Ct 35	Day 16: positive
			Day 22: negative
		Pt 2:	Day 25: negative
		Day 16: N gene Ct 14, E gene Ct 20	Day 42: negative
		Day 22: N gene Ct 25, E gene Ct 33	Day 51: negative
		Day 25: N gene Ct 25, E gene 29	
		Day 42: N gene Ct 29, E gene Ct 37	
		Day 51: N gene Ct 37, E gene Ct 39	

Study ID	Symptoms, days reported	RT-PCR Cycle count/log copies results	Viral culture results (days)
Tarhini 2021	Severe infection requiring intensive care	Day 6: Ct=25, Day 16: Ct=24 Day 32: Ct=20 Day 78: Ct=34 Day 80: Ct=24 Day 91: Ct=28 Day 103: Ct=23 Day 109: Ct=18 Day 120: Ct=26 Day 132: Ct=40+ (negative) Day 136: Ct=40+ (negative) (*)	Day 6: culture negative Day 16: culture negative Day 80: culture negative Day 91: culture negative Day 103: culture positive Day 111: culture negative Day 120: culture negative (*)
Weigang 2021	Mild respiratory symptoms for over 145 days	38 PCR tests: days 0, 6, 10, 14, 16, 21, 24, 27, 31, 34, 38, 42, 45, 46, 49, 52, 56, 58, 63, 65, 71, 105, 113, 115, 119, 122, 126, 140, 143, 145, 146, 149, 150,	27 cell culture tests, days: 0, 6, 10, 14, 16, 21, 24, 27, 31, 34, 38, 42, 45, 49, 52, 56, 58, 63, 65, 71, 105, 106, 113, 115, 119, 140, 154; 27 results: -, +, +, +, -,

Study ID	Symptoms, days reported	RT-PCR Cycle count/log copies results	Viral culture results (days)
		154, 161, 167, 174, 189. 38	+, +, +,+, +, -, -, -, -, -, -, -, -, +, -, -, -, -, -, -, (i.e. positive on
		Ct values: 25, 16, 19, 19, 25, 23, 18, 26, 20, 17, 31, 27, 27, 50, 27, 28, 25, 28, 27, 29, 30, 23, 34, 29, 31, 31, 36, 26, 29, 34, 39, 45, 45, 34, 45, 45, 45, 45.	days 6, 10, 14, 21, 24, 27, 31, 34, 105)

(*) Numbers have been read from a figure in the published article and may not be an accurate

estimate.

Ct = cycle threshold

D = day

NR = not reported

NA = not available

Pt = patient

RT-PCR = reverse transcriptase polymerase chain reaction

Figure 1. PRISMA flow chart of study screening for inclusion.

Records identified through database searching
n = 12,987

Additional records identified through other sources
n= 1 through correspondence

Records screened for duplicates
n = 12,988

Records excluded
Duplicate of included study n=22

Records screened for inclusion:
title and abstract screening
n = 12,966

Excluded on title and abstract screen
No relevant data n = 12,903

Full-text articles assessed for eligibility
n = 63

Excluded on full text screen
No relevant data n = 53
(no transplant ppts n = 30;
no cell culture data n = 9;
no NP PCR cycle count data n = 9;
review n = 6)

Studies included in qualitative synthesis
n = 10

Studies included in quantitative synthesis (meta-analysis)
n = 0

Figure 2. SARS-CoV-2 culture results in transplant patients from days of symptom onset.

Culture positive (n=59 samples): median 16 (IQR 8-27, Range 1-105 days, mean 19.7 days
Culture negative (n=107 samples): median 40 (IQR 22-60), Range 1-119 days, mean 43.2 days

Figure 3a. Timings of positive culture results in Transplant Patients by duration of symptoms and Ct results
Alshukairi [7], Benotmane [8], Rajakumar [15] & Niyonkuru [14] (days 1- 42)

Figure 3b. Timings of positive culture results in Transplant Patients by duration of symptoms and Ct results
Aydillo [6] (day 1-90), Tarhini [9] & Weigang [12] (day 1-120)

🟥 Negative Culture

🟩 Positive Culture
  Number equals RT-PCR Cycle Threshold

Figure 3c. Timings of positive culture results in Transplant Patients by duration of symptoms and log10 copies/ml results
Decker [10] (days 1-36)

	patient	D0	D1	D2	D3	D4	D5	D6	D7	D8	D9	D10	D11	D12	D13	D14	D15	D16	D17	D18	D19	D20	D21	D22	D23	D24	D25	D26	D27	D28	D29	D30	D31	D32	D33	D34	D35	D36
Decker 2020 log10 copies/ml, > 4 log10 deemed +ve	1		4.5				4		6				6.2							6.2			6.5				5			4.5						4		4

■ Negative Culture
■ Positive Culture
Number equals log copies/ml

Figure 4. Relationship between cycle threshold, symptom onset (in days) and likelihood of shedding replication competent virus