Faster post-malnutrition weight gain during childhood is associated with risk of non-
communicable disease in adult survivors of severe malnutrition

Debbie S. Thompson[1], Kimberley McKenzie[1], Charles Opondo[2], Michael S. Boyne[3], Natasha Lelijveld[4,5],
Jonathan Wells[6], Tim J. Cole[6], Kenneth Anujuo[4], Mubarek Abera[7], Melkamu Berhane[7], Marko Kerac[4,5],
CHANGE Study Collaborators Group, Asha Badaloo[1]*

[1]Caribbean Institute for Health Research, The University of the West Indies, Kingston, Jamaica
[2]Department of Medical Statistics, Faculty of Epidemiology & Population Health, London
School of Hygiene & Tropical Medicine, London, UK
[3]Department of Medicine, The University of the West Indies, Kingston, Jamaica
[4]Department of Population Health, Faculty of Epidemiology and Population Health, London
School of Hygiene and Tropical Medicine, London, UK
[5]Centre for Maternal, Adolescent & Reproductive Child Health (MARCH), London School of
Hygiene & Tropical Medicine, London, UK
[6]UCL Great Ormond Street Institute of Child Health, London, UK
[7]Jimma University, Ethiopia
*CHANGE Study collaborators group: Stephen A. Wootton, Albert Koulman, Amir Kirolos,
Gerard Bryan Gonzales, Jonathan Swann, Abena S. Amoah.

Abstract

Background: Nutritional rehabilitation during severe malnutrition (SM) aims to rapidly restore

body size and minimize poor short-term outcomes. We hypothesized that too rapid weight gain

during and after treatment might however predispose to cardiometabolic risk in adult life.

Methods: Weight and height during hospitalization and one year post-hospitalization were

abstracted from hospital records of children who survived SM. Six definitions of post-malnutrition

weight gain/growth were analysed as continuous variables, quintiles and latent classes in age-sex

and minimum weight-for-age z-scores-adjusted regression models against adult anthropometry,

body composition (DEXA), blood pressure, blood glucose, insulin, and lipids.

Results: 60% of 278 participants were male, mean (SD) age 28.2 (7.7) years, mean (SD) BMI 23.6

(5.2) kg/m^2. Mean admission age for SM was 10.9 months (range 0.3-36.3 months) and 207/270

(77%) were wasted (weight-for-height z-score<-2). During childhood, mean rehabilitation weight

gain (SD) was 10.1(3.8) g/kg/day and 0.8(0.5) g/kg/day in the first year post-hospitalization.

Rehabilitation weight gain >12.9 g/kg/day was associated with higher adult BMI

(difference=0.5kg/m^2, 95%CI: 0.1-0.9, $p = 0.02$), waist circumference (difference=1.4cm, 95%CI:

0.4-2.4, p=0.005), fat mass (difference = 1.1kg, 95%CI: 0.2-2, p=0.02), fat mass index

(difference=0.32, 95%CI: -0.0001-0, p=0.05), and android fat mass (difference=0.09 kg, 95%CI:

0.01-0.2, p=0.03). Rehabilitation (g/month) and post-hospitalization (g/kg/month) weight gain

were associated with greater lean mass (difference = 0.7 kg, 95% CI: 0.1, 1.3, $p = 0.02$)

(difference=1.3kg, 95% CI: 0.3-2.4, p=0.015) respectively.

Conclusion: Rehabilitation weight gain exceeding 13g/kg/day was associated with adult adiposity

in young, normal-weight adult SM survivors. This raises questions around existing malnutrition

weight gain targets and warrants further studies exploring optimal post-malnutrition growth.

Keywords: Severe malnutrition, post malnutrition growth, rehabilitation weight gain, weight-

for-age z-scores, height-for-age z scores, weight-for-height z scores

Introduction

Severe malnutrition (SM) remains an important cause of mortality globally, and malnutrition contributes to the deaths of 500,000 children under the age of 5 years annually [1]. As global efforts improve the management of these children, alongside increased availability of ready to use therapeutic foods, the mortality from severe malnutrition is declining in some settings. However, while the short-term outcomes of severe childhood malnutrition may be improving, the long-term consequences are less clear. Although there is evidence of an association between prenatal undernutrition and later risk of chronic non-communicable diseases (NCDs) [2, 3], much of this evidence is from developed countries. Additionally, evidence of similar links between early postnatal malnutrition and later NCDs is recent and limited [4-6], and potential mechanisms are yet to be identified.

Most malnutrition treatment programs treat the attainment of normal weight-for-height as an indicator of recovery and aim for this to occur as rapidly as possible. To this end, energy dense feeds are usually provided during the rehabilitation stage of treatment of severe malnutrition; children are fed increasing amounts of energy and protein-enriched feeds to achieve a normal body weight. The recommended energy intake during the three day transition period is 100-135 kcal/kg/day [7] followed by intake of 150-225 kcal/kg/d during the rehabilitation ("catch-up growth") period, the goal being a weight gain of $\geq$ 10 g/kg/day as recommended by the World Health Organization (WHO) [8]. However, it is unclear whether better short-term outcomes are occurring at the expense of longer-term outcomes. The thrifty phenotype hypothesis proposes that poor infant growth is a risk factor for the subsequent development of type 2 diabetes and the metabolic syndrome due to the effects of poor early-life nutrition [9]. In such "metabolically thrifty" individuals, faster weight gain in the plastic developmental period of infancy could pose an additional risk for later NCDs [10].

Many countries with high levels of early childhood malnutrition also have high burdens of obesity

and NCDs including hypertension, type 2 diabetes and cardiovascular disease [11]. Data from the

WHO indicate that NCDs are the leading causes of death and disability globally, killing more than

three in five people worldwide and responsible for more than half the global burden of disease

[12]. NCDs also cause and perpetuate poverty while hindering economic development in low- and

middle-income countries (LMICs) as they struggle with the economic burden of providing

treatment for individuals with NCDs, thus recapitulating the cycle.

This study aims to explore rehabilitation and post-hospitalization weight gain and growth in

children who were hospitalized with severe malnutrition, and to investigate the associations

between weight gain during and after rehabilitation and adult obesity, body composition and NCD

risk. We hypothesize that faster rates of growth during nutritional rehabilitation are associated with

later NCD risk.

Methods

Setting: We conducted secondary analysis of data collected from members of the LION (Long-

term Implications of Nutrition) Cohort. The LION Cohort was retrospectively assembled and from

individuals who were admitted to the Tropical Metabolism Research Unit (TMRU) ward at the

University Hospital of the West Indies between the years 1963-1993 with a diagnosis of severe

malnutrition. Members of the cohort were followed up and as adults and were extensively

characterised in 2008-2012 in the Jamaica Marasmus and Kwashiorkor Adult Survivors

(JAMAKAS) Study (**Figure 1**) [4, 13-15]. All adult severe malnutrition survivors followed up in

the JAMAKAS Study for whom childhood hospital admission records were available were

included in this analysis. The Mona Campus Research Ethics Committee of the University of the

West Indies approved the study (CREC-MN. 204 20/21) and all participants provided written informed consent.

During hospitalization for severe malnutrition, treatment began with a "stabilisation" phase during which a diet was offered to provide energy for weight maintenance while treating acute complications such as infections, mineral and fluid imbalances. Since the 1960s, metabolism research has informed development of treatment guidelines in the TMRU [16] which led to a gradual standardisation of the approach by about the mid-1970s [17]. Since then, the treatment regime has been similar to the guidelines by WHO [18]. With clinical improvement, resolution of oedema and improved appetite and affect, the "rehabilitation" phase occurred, during which children were fed increasing amounts of energy and protein-enriched feeds to attain a body weight similar to WHO (1981) guidelines [18]. The final phase of treatment, referred to as "recovery", involved weaning the child to an age-appropriate diet before discharge from hospital. Children were discharged having attained weight-for-height $\geq$ 90% (NCHS) and were followed up as outpatients for up to two years post-hospitalization. During this period, they were fed a home-based diet.

Participant Characteristics: Childhood data abstracted from hospital records included admission weight, height and mid-upper arm circumference (MUAC) as well as weight, height/length and MUAC during hospitalization and one year post-discharge from hospital. Adult data were collected at the time of the JAMAKAS Study and included a detailed medical and drug history, anthropometry, blood pressure, body composition (by dual-energy x-ray absorptiometry, GE Lunar Prodigy), fasting glucose, insulin and serum lipids as described previously [19].

Exposure variables: We derived six definitions of post-malnutrition growth rate (PMGr) based on available data and established assessment methods (**Table 1**). The first three definitions (PMGr 1-

3) were derived from in-patient weight and age data starting at the time of minimum weight, which

was usually at admission or after oedema had resolved and continuing until the time of maximum

weight during treatment. The definitions respectively utilized the daily change in weight-for-age

z-score (WAZ) (using WHO 2006 growth standards) (PMGr1), g/kg of body weight (PMGr2) and

weight in grams (PMGr3) (**Table 1**), all common measures of growth in malnutrition programs.

Height-for-age z-score (HAZ) was not used during this period as height gain was small. Due to

individual variations in the attainment of maximum weight, weight gain was expressed 'per day'.

The remaining three definitions (PMGr 4-6) represented weight gain and height gain from the time

of hospital discharge until one year post-discharge. For these definitions, WAZ (PMGr4), g/kg

(PMGr 5) and HAZ (PMGr 6) were expressed as rate of change per month (**Table 1**).

Table 1: Definitions of post malnutrition growth (PMGr)

PMGr class	Abbreviated title	Definition	Time
1	Δ WAZ/day	Average change in WAZ per day	Date of minimum weight to date of maximum weight during hospitalization
2	g/kg/day	Average change in grams per kilogram of body weight per day	
3	g/day	Average change in grams per day	
4	Δ WAZ/month	Average change in WAZ per month	Date from hospital discharge to one year post-hospitalization
5	g/kg/month	Average change in grams per kilogram of body weight per month	
6	Δ HAZ/month	Average change in HAZ per month	

WAZ- weight-for-age z-scores, HAZ-height-for-age z-scores

Outcome variables included BMI, waist circumference, fat mass, fat mass index, % fat mass,

android fat, android: gynoid (AG) fat ratio (android fat divided by gynoid fat; a pattern of body fat

distribution associated with an increased risk for metabolic syndrome in healthy adults), lean mass,

lean mass index, systolic blood pressure, diastolic blood pressure, fasting glucose, fasting insulin,

LDL-cholesterol, triglyceride, and homeostatic model assessment for insulin resistance (HOMA-

IR). Body composition (height-adjusted lean mass and fat mass) was assessed by dual-energy X-

ray absorptiometry (DEXA).

Lean mass index based on DEXA was calculated using the formula: lean mass/height^2, and

similarly fat mass index based on DEXA was calculated using the formula: fat mass/height^2, each

in BMI units of kg/m^2. HOMA-IR was calculated using the formula:

148   HOMA-IR = fasting insulin (microU/L) x fasting glucose (mmol/L)/22.5.

Statistical Analyses: To determine the relationship between PMGr and adult body size, body

composition and NCD risk factors, we conducted separate linear regression analyses for each

outcome adjusted for age and sex and (as an additional model) minimum WAZ (i.e., WAZ at the

time of minimum weight). Additionally, all outcome measures of body composition and blood

pressure were adjusted for adult height. Adult BMI was treated as an outcome variable and not

adjusted for in the analyses as it was considered to be on the causal pathway between post

malnutrition weight gain and the outcomes. Each PMGr definition was evaluated as a continuous

variable, and we reported differences in continuous outcomes per unit faster growth depending on

the definition of PMGr. Based on the six definitions of post malnutrition growth, participants were

also grouped into quintiles which were used in age- and sex-adjusted regression analyses as ordinal

variables. Quintiles 1 (Q1) and 5 (Q5) represented the slowest and fastest rates of growth

respectively, and age- and sex-adjusted regression analyses were conducted comparing NCD risk

across the growth rate quintiles. Finally, a latent class analysis (LCA) was conducted to identify

qualitatively different growth pattern subgroups of the study population across 5 time points, viz.,

admission, time at minimum weight, time at maximum weight, discharge from hospital and one

year post-hospitalization. The LCA was fitted using a generalised structural equation model with

Gaussian-distributed outcomes at each of the timepoints included in the analysis. Model fit was

assessed using Akaike Information Criteria (AIC); latent classes were extracted from the model

with the lowest AIC, and which yielded reasonably-sized classes. These LCA-predicted classes

were plotted against anthropometric, body composition and clinical markers of NCD risk to

facilitate interpretation. Stata version 16.0 (StataCorp LLC, College Station, Texas, USA) was

used to conduct the statistical analyses and p-values ≤ 0.05 were considered statistically

significant.

Results

There were 1,366 children in the original TMRU/UHWI cohort, 47 of whom died during treatment

for malnutrition and the remaining 1,289 were discharged alive. The JAMAKAS study traced 729

adult survivors from the original cohort and enrolled 316 of them to the study. Subsequently, this

current 'CHANGE' (Child malnutrition & adult NCD: Generating evidence on mechanistic links

to inform future policy/practice) Study identified 278 adult participants for whom childhood

admission records were available for inclusion to this study. **Figure 1** summarises the recruitment

of the adult survivors of severe malnutrition.

We analysed data for 278 adults (60% male) who were treated for severe malnutrition as children.

Mean age at admission was 10.9 months, 237/270 (88%) were stunted (HAZ < -2), 207/270 (77%)

and 185/214 (87%) were wasted (WHZ <-2 and MUAC <12.5 cm respectively) and 182/278 (65%)

had oedematous malnutrition. As adults, mean age (SD) was 28.2 (7.7) years, and mean BMI (SD)

was 23.6 (5.2) kg/m^2 (**Table 2**).

Figure 1: Flow chart detailing recruitment of adult survivors of severe malnutrition (*n* = 278). "Unable to participate" includes adult survivors of severe malnutrition who were unavailable because of migration (*n* = 53), illness (*n* = 19), refusal (*n* = 14), or pregnancy (*n* = 30). TMRU, Tropical Metabolism Research Unit; UHWI, University Hospital of the West Indies; JAMAKAS, Jamaica Marasmus and Kwashiorkor Adult Survivors.

Table 2: Clinical characteristics of 278 survivors of severe malnutrition as children and as adults.

Children	Characteristics	At Admission N = 278	At Discharge n= 276	At 1 year PH n = 85
	Age (months)	11.9 (0.33 -36)	13.7 (3 - 42)	26.1 (15- 45)
	% Male	(167) 60.1 %	60.1%	56.5%
	% Oedema	(182) 65.5%	(0) 0%	(0) 0%
	Weight (kg)	5.4 (1.5), 278	7.3 (1.8), 276	9.8 (1.6), 85
	Height (cm)	64.6 (7.0), 270	66.5 (7.2), 195	80.1 (5.6), 81
	MUAC (cm)	10.8 (1.6), 214	12.9 (1.2), 33	14.(1.5), 18
	% Underweight (WAZ<-2)	92	63	41
	% Stunted (HAZ<-2)	88	88	59
	% Wasted (MUAC < 12.5cm)	87	27	17
	% Wasted (WHZ < -2)	77	5	15
	Days since admission	0	55 (13 - 251)	412 (343 - 588)

	Characteristics	All N = 278	Male n = 167	Female n= 111
Adults	Years since discharge	27.2 (16-51)	27.3 (16-51)	26.5 (17-46)
	Age (years)	28.2 (7.7)	28.5 (7.9)	27.7 (7.4)
	Weight (kg)	66.1 (14.9)	65.5 (11.7)	67 (18.7)
	Height (cm)	167.5 (9.0)	171.1 (7.6)	162 (7.8)
	BMI (kg/m^2)	23.6 (5.2)	22.3 (3.4)	25.4 (6.7)
	Waist circumference (cm)	78.8 (12.7)	76.3 (8.9)	82.6 (16.2)
	Waist-hip ratio	0.83 (0.06)	0.83 (.06)	0.82 (0.07)
	Diastolic blood pressure (mmHg)	71 (11)	71 (11)	71 (12)
	Systolic blood pressure (mmHg)	114 (12)	116 (12)	112 (13)
	Lean mass (kg)	48.4 (10.4)	54.4 (7.5)	39.3 (6.9)
	Lean mass index (kg/m^2)	17.1 (2.6)	18.5 (2.0)	14.9 (1.8)
	Fat mass (kg)	14.4 (13.2)	8.0 (7.3)	24.2 (14.1)
	Fat mass index (kg/m^2)	5.3 (5.0)	2.7 (2.5)	9.2 (5.3)
	% Fat	21.2 (15.5)	11.8 (8.6)	35.4 (12.5)
	Android fat mass (kg)	1.1 (1.2)	0.64 (0.68)	1.9 (1.4)
	Android fat %	24.1 (17.5)	14.4 (11)	38.8 (15)
	Android-gynoid fat ratio	0.9 (0.22)	0.9 (0.20)	0.9 (0.24)
	Fasting blood glucose (mmol/L)[‡]	4.6 (0.5)	4.7 (0.4)	4.5 (0.5)
	Fasting serum insulin (uIU/mL)[‡†]	4.7 (3.6)	3.3 (3.2)	6.3 (3.4)
	HOMA-IR[†]	0.9 (0.7)	0.7 (0.7)	1.3 (0.7)
	LDL-cholesterol[†] (mmol/L)	2.1 (0.7)	2.0 (0.6)	2.3 (0.7)
	Triglyceride[†] (mmol/L)	0.7 (0.4)	0.7 (0.3)	0.7 (0.4)

N = 278. PH = post-hospitalization. Z-scores based on WHO 2006 and 2007 references. Data are presented as means (± SDs) or (range). (‡, ‡†, †) n = 95, 83 and 82 respectively.

Subsequent analyses were based on 273 participants during rehabilitation and 84 participants post-hospitalization as some participants had missing maximum weight or discharge weight data. In 273 participants, mean weight gain during rehabilitation was, according to the three definitions,

210 0.07 WAZ/day, 10.1 g/kg/day and 62 g/day (**Table 3**). Mean (SD) change in WAZ during

211 rehabilitation was 2.2 (0.97). In the post-hospitalization phase, weight gain was much lower; mean

212 (SD) weight gain was 0.8 (0.5) g/kg/day and mean (SD) change in WAZ/ day was 0.002 (0.004)

213 for 84 participants. Each PMGr definition was grouped in quintiles (**Table 3 and Figure 2**)

214 **Table 3**: Descriptive data for definitions of post-malnutrition growth (PMGr)
215

PMGr Class	Description	Mean (SD)	Range
	Δ WAZ/ day	0.07 (0.03)	-0.0025, 0.20
1	Q1	0.03 (0.01)	-0.00, 0.04
(n = 273)	Q2	0.05 (0.00)	0.04, 0.06
	Q3	0.06 (0.00)	0.06, 0.07
	Q4	0.08 (0.00)	0.07, 0.09
	Q5	0.11 (0.02)	0.09, 0.20
	g/kg/day	10.1 (3.8)	0.32 - 27
2	Q1	5.5 (1.3)	0.3, 7.0
(n = 273)	Q2	7.9 (0.6)	7.0, 8.9
	Q3	9.7 (0.5)	8.9, 10.7
	Q4	11.8 (0.6)	10.7, 12.9
	Q5	15.7 (2.9)	12.9, 27.4
	g/day	61.6 (25.3)	2.5 - 189
3	Q1	31.8 (8.0)	2.5, 41.0
(n = 273)	Q2	46.9 (3.1)	41.4, 51.7
	Q3	58.6 (3.9)	52.0, 64.8
	Q4	71.6 (5.1)	65.0, 81.0
	Q5	99.9 (20.5)	81.1, 188.7
	Δ WAZ/ month	0.05 (0.1)	-0.22 - 0.46
4	Q1	-0.09 (0.06)	-0.22, -0.03
(n = 84)	Q2	-0.01 (0.01)	-0.03, 0.01
	Q3	0.05 (0.02)	0.01, 0.08
	Q4	0.11 (0.02)	0.08, 0.15
	Q5	0.21 (0.08)	0.15, 0.46
	g/kg/month	24.9 (15.6),	-0.5 - 7.1
5	Q1	5.4 (8.0)	-12.9, 14.4
(n = 84)	Q2	16.9 (2.4)	14.6, 20.9
	Q3	24.5 (2.2)	21.4, 28.0
	Q4	32.0 (3.3)	28.1, 37.2
	Q5	47.2 (12.9)	37.8, 90.2
	Δ HAZ/ month	0.10 (0.12)	-0.08 - 0.54
6	Q1	-0.04 (0.03)	-0.08, 0.01
(n = 47)	Q2	0.04 (0.01)	0.02, 0.06
	Q3	0.09 (0.01)	0.07, 0.11
	Q4	0.14 (0.02)	0.12, 0.17
	Q5	0.29 (0.11)	0.18, 0.54

216 PMGr - post malnutrition growth rate, Q – quintile: 1- slowest growth to 5- fastest
217 growth, WAZ - weight-for-age z-scores, HAZ - height-for-age z-scores

Figure 2: Mean growth rates of growth across four time periods according to the six PMGr definitions. A to C: PMGr1 to 3 (*rehabilitation*) and D to F: PMGr4 to 6 (*post-hospitalization*). Each PMGr definition was split into quintiles where Q1= slowest growth, and Q5 = fastest growth. PMGr at admission acted as comparator for the later timepoints.

Associations between rehabilitation weight gain and adult NCD risk

The three measures of rehabilitation weight gain (n = 273) were associated with adult NCD risk in

regression models adjusted for age, sex and minimum WAZ as follows:

PMGr1 as ΔWAZ/day was associated with waist circumference (difference=53cm, 95% CI: 6,

100, $p = 0.03$) and lean mass (difference=37.0 kg, 95% CI: 7, 67, $p = 0.02$) (**Supplementary Table**

1). Those with fastest weight gain had higher BMI (difference $= 0.4$kg/m^2, 95% CI: 0.02, 0.8, $p =$

0.04), waist circumference (difference $= 1.2$ cm, 95% CI: 0.3, 2, $p = 0.01$), lean mass (difference

$= 0.8$ kg, 95% CI: 0.2, 1.4, $p = 0.01$), fat mass (difference $= 0.8$ kg, 95% CI: 0.01, 1.7, $p = 0.048$)

and % fat mass (difference $= 0.85$, 95% CI: 0.02, 1.7, $p = 0.04$) than those with slowest weight

gain (**Supplementary Table 2, Figure 3**). Associations with waist circumference and % fat mass

remained significant after further adjusting for adult height. Visual examination of the box plots

(**Figure 3A, 3C, 3K, 3N**) suggested that only ΔWAZ ≥ 0.09/day was associated with measures of

adult adiposity and when further exploratory analyses (excluding Q5) were conducted, ΔWAZ $<$

0.09/day was not associated with adult adiposity (p-values > 0.2) (data not shown). In the fastest

weight gainers (Q5) in comparison to the slowest weight gainers (Q1), ΔWAZ/day was associated

with BMI (difference $= 2.1$ kg/m2, 95% CI: 0.3, 3.8, $p = 0.02$), waist circumference (difference $=$

6.5 cm, 95% CI: 2, 11, $p = 0.002$), waist-hip ratio (difference $= 0.03$, 95% CI: 0.003, 0.05, $p =$

0.03), fat mass (difference $= 4.0$ kg, 95% CI: 0.2, 8, $p = 0.04$) and android fat mass (difference $=$

0.36 kg, 95% CI: 0.01, 0.7, $p = 0.05$) (data not shown).

PMGr2 as weight gain in g/kg/day was associated with waist circumference (difference $= 0.43$

cm, 95% CI: 0.1, 0.8, $p = 0.02$) and lean mass (difference $= 0.28$ kg, 95% CI: 0.05, 0.5, $p = 0.02$)

(**Supplementary Table 1**). Those with fastest weight gain had higher BMI (difference $= 0.5$ kg/m^2,

95% CI: 0.1, 0.9, $p = 0.02$), waist circumference (difference $= 1.4$ cm, 95% CI: 0.4, 2.4, $p = 0.005$),

fat mass (difference = 1.1 kg, 95% CI: 0.2, 2, p = 0.02), fat mass index (difference = 0.32 kg/m2,

95% CI: -0.0001, 0.6, p = 0.05), % fat mass (difference = 1.1, 95% CI: 0.2, 2, p = 0.02), android

fat mass (difference = 0.09 kg, 95% CI: 0.01, 0.2, p = 0.03) and % android fat mass (difference =

1.2, 95% CI: 0.1, 2, p = 0.03) than those with slowest weight gain (**Supplementary Table 2,**

Figure 3). The associations with waist circumference, % fat mass, android fat mass and % android

fat mass remained significant after further adjusting for height (data not shown). Examination of

the box plots (**Figure 3B, 3D, 3L, 3M, 3O, 3P, 3Q**) suggested that only weight gain ≥ 12.9

g/kg/day was associated with measures of adult adiposity and when further exploratory analyses

(excluding Q5) were conducted, rehabilitation weight gain <12.9 g/kg/day was not associated with

adult adiposity (p-values > 0.3) (data not shown).

PMGr3 as weight gain in g/day was associated with lean mass (difference = 0.04 kg, 95% CI:

0.01, 0.1, p = 0.02) (**Supplementary Table 1**). Those with fastest weight gain had greater waist

circumference (difference = 1.2 cm, 95% CI: 0.2, 2, p = 0.01) and lean mass (difference = 0.7 kg,

95% CI: 0.1, 1.3, p = 0.02) than those with slowest weight gain (**Supplementary Table 2, Figure**

3). Examination of the box plots (**Figure 3E**) suggested that rehabilitation weight gain ≥ 81 g/day

was associated with adult waist circumference and when further exploratory analyses were

conducted, weight gain < 81 g/day was not associated with adult waist circumference (p-values >

0.3) (data not shown).

There was no association between any measure of rehabilitation weight gain (i.e., PMGr1, PMGr2,

and PMGr3) and adult blood pressure, blood glucose, insulin, or lipids (**Supplementary Tables 1**

and 2).

Associations between post-hospitalization weight gain and height gain and adult NCD risk.

Post-hospitalization weight gain (n = 84) and height gain (n = 47) were associated with NCD risk

in regression models adjusted for age, sex and minimum WAZ as follows:

PMGr4 as WAZ/month was associated with lean mass index (difference = 4.6 kg/m^2, 95% CI:

0.6, 9, p = 0.03) (**Supplementary Table 1**). Those with fastest weight gain had a higher lean mass

index (difference = 0.4 kg/m^2, 95% CI: 0.04, 0.7, p = 0.03) than those with the slowest weight gain

(**Supplementary Table 2, Figure 3**).

PMGr5 as g/kg/month was associated with lean mass index (difference = 0.03 kg/m^2, 95% CI:

0.01, 0.06, p = 0.02) (**Supplementary Table 1**). The fastest weight gainers had higher lean mass

(difference = 1.3 kg, 95% CI: 0.3, 2.4, p = 0.02) and lean mass index (difference = 0.4 kg/m^2, 95%

CI: 0.1, 0.7, p = 0.03) and a lower AG ratio (difference = -0.03, 95% CI: -0.07, -0.001, p = 0.05)

than the slowest weight gainers (**Supplementary Table 2, Figure 3**).

PMGr6 as height gain in ΔHAZ/month was not associated with NCD outcomes and there was no

difference between the fastest and slowest growers in terms of the NCD outcomes.

There was no association between any measure of post-hospitalization weight gain or growth (i.e.,

PMGr4, PMGr5, and PMGr6) and adult blood pressure, blood glucose, insulin, or lipids

(Supplementary Tables 1 and 2).

Figure 3: Association of NCD outcomes with PMGr definitions. Age, sex and minimum weight WAZ-adjusted linear regression results (coefficient, p-value): **A to B**- body mass index: PMGr1 (B = 0.4, p = 0.04), PMGr2 (B = 0.5, p = 0.02); **C to E**- waist circumference: PMGr1 (B = 1.2, p = 0.01), PMGr2 (B = 1.4, p = 0.01), and PMGr3 (B = 1.2, p = 0.01); **F to H**- lean mass: PMGr1 (B = 0.8, p = 0.01), PMGr3 (B = 0.7, p = 0.02) and PMGr5 (B = 1.3, p = 0.02); **I to J**- lean mass index: PMGr4 (B = 0.35, p = 0.03) and PMGr5 (B = 0.43, p = 0.01); **K to L-** fat mass: PMGr1 (B = 0.84, p = 0.048) and PMGr2 (B = 1.1, p = 0.01); **M**- fat mass index: PMGr2 (B = 0.32, p = 0.05); **N to O-** fat mass percent: PMGr1 (B = 0.85, p = 0.04) and PMGr2 (B = 1.1, p = 0.02); **P**- android fat mass: PMGr2 (B = 0.09, p = 0.03); **Q**- android fat mass %: PMGr2 (B = 1.2, p = 0.03); **R**- android-gynoid ratio: PMGr5 (B = -0.03, p = 0.045).

Latent class analysis

Latent class analysis (LCA) identified a model with 6 interpretable classes of at least 21 children each which had the lowest AIC (3159) (**Figure 4**). Class 1 (10%) consisted of the children who were most underweight on admission (mean WAZ < -6) and with the least weight gain during rehabilitation. Class 2 (21%) had the second most underweight children with the second lowest rehabilitation weight gain. Class 3 (30%) were children with the second highest rehabilitation weight gain. Class 4 (18%) had the highest rehabilitation weight gain. Class 5 (13%) had the second highest WAZ at admission and the third highest rehabilitation weight gain. Class 6 (8%) were the children who were least underweight on admission. All classes have a similarly steep level of catch-up during rehabilitation, except class 4. After discharge, class 1, with the greatest deficits, continued to catch up, and there was a pattern of decreasing catch-up growth up the classes, until class 6 where "catch down" was observed (**Figure 4).**

Age and sex-adjusted regression analyses against the NCD outcomes were carried out with LCA class as a categorical variable and Class I being the reference group. In children in Class 5 (of which 89% had oedematous malnutrition), change in WAZ was associated with greater adult fat mass (difference = 5.2kg, SE = 2.6; 95% CI: 0.1, 10, p = 0.045) and android fat mass (difference = 0.52 kg, SE = 0.24; 95% CI: 0.04, 0.99, p = 0.03) compared to children in Class 1. However, these associations were no longer significant after adjusting for minimum WAZ (data not shown).

Figure 4: Latent class analysis of patterns of weight-for-age z-score from admission for severe malnutrition to 1 year post-hospitalization in 278 adults. Classes ranked by lowest WAZ upon admission: 1, 2, 3, 4, 5, 6, and by highest weight gain during rehabilitation: 4, 3, 6, 5, 2, 1. PH- post hospitalization

Discussion

This study of adult survivors of severe malnutrition in childhood demonstrated that, during the rehabilitation phase of malnutrition treatment, weight gain above certain thresholds is associated with risk factors for NCDs (i.e., greater BMI, waist circumference, fat mass and android fat mass). Weight gain during the first year after hospitalization was inversely associated with android-gynoid fat ratio. Additionally, in adult survivors, faster weight gain during and after admission for severe malnutrition was associated with lean mass which is typically associated with lower NCD risk. It is important to note that these associations are independent of weight-for-age at the time of hospitalization and adult height. Neither rehabilitation weight gain nor post-hospitalization weight gain was associated with adult blood pressure or blood glucose, insulin, or lipids.

Post-malnutrition growth and adult adiposity

Weight gain targets during treatment of severe malnutrition have been in place since the 1970s, with the recommendation that a desirable rehabilitation weight gain exceeds 10g/kg/day [20]. We have demonstrated in this group of adult survivors of severe malnutrition that rehabilitation weight gain that exceeded specified thresholds was associated with adult adiposity. Measures of adiposity (i.e., BMI, waist circumference, fat mass and android fat mass) appeared similar across the first four quintiles of weight gain (defined as ΔWAZ, g/kg/day and g/day), whereas the fifth quintile showed greater adiposity. Thus, each unit increase in weight gain does not appear to increase NCD risk in the same way, and there is a threshold effect of rehabilitation weight gain on adult adiposity. This suggests that weight gain at or above specific thresholds (i.e., 13 g/kg/day, ΔWAZ > 0.09/day, 81g/day) could impute greater adult adiposity.

Faster rehabilitation weight gain was associated with greater adult BMI (an indicator of excess or inadequate weight) which itself is an independent risk factor for cardiovascular disease. Weight

gain above the previously specified thresholds was also associated with increased fat mass, and

this excess adipose tissue contributes to the production of proinflammatory cytokines which are

also associated with increased risk of cardiovascular disease. Importantly, faster rehabilitation

weight gain was associated with fat mass index, which has been shown to be a more complete

tool for diagnosing obesity than both BMI and fat mass as it differentiates fat and muscle mass of

the total body weight [21].   Furthermore, fat accumulation in the android compartment is

specifically known to be associated with increased cardiovascular and metabolic risk [22] and

studies in younger populations have also demonstrated that android fat is more closely related to

metabolic risk factors [23]. This tendency for some adult survivors of severe malnutrition to

accumulate visceral fat may be a consequence of adipose tissue maldevelopment which includes

altered body composition and increased visceral fat [24]. Rapid postnatal infant weight gain has

been shown to be associated with increases in both visceral and abdominal subcutaneous adipose

tissue, as well as total adiposity and the risk of obesity in middle adulthood [25]. Additionally,

increased visceral adipose tissue is linked to high triglycerides and low HDL cholesterol levels

[26]. However, while the tendency to store fat abdominally might be a persisting response to

adverse conditions and growth failure in fetal life and infancy, we have shown this tendency to

store fat in the abdominal compartment to be independent of minimum weight-for age z-score

during hospitalization. Therefore, the association between post-malnutrition weight gain in

childhood and adult android fat in this group of young, normal-weight severe malnutrition

survivors could signal an increased risk of clinical NCDs as they age.

It is also important to point out that although our findings indicate statistically significant

associations across the range of weight and body composition markers of NCD risk, these

individuals have not exceeded established thresholds for adverse outcomes such as obesity. So,

while faster weight gain was associated with higher adult BMI, only 11% of these participants

were obese (i.e., BMI >30 kg/m^2). Additionally, several studies have reported a lower

susceptibility to visceral adipose tissue deposition in blacks compared to whites [27, 28].

Therefore, although it is unclear whether the levels of visceral fat in these study participants are

sufficient to impute cardiometabolic risk, the possibility cannot be ruled out.

Our study produced the unexpected finding of an inverse association between post-hospitalization

weight gain (g/kg/month) and android-gynoid percent fat ratio (AG), an important predictor of

metabolic and cardiovascular disease risk. Indeed, in over 1,800 participants with a mean age of

35 years, AG was reported to have a greater association with cardiometabolic dysregulation than

android fat, gynoid percent fat or BMI [29]. While our findings are counterintuitive, they further

emphasize the complexity of these associations which may depend not only on the extent and rate

of weight gain, but also the age of the child at the time of weight gain, and other factors outside

the scope of this study.

Post-malnutrition growth and adult lean mass

Lean mass in adult survivors of severe malnutrition was associated with both rehabilitation and

post-hospitalization weight gain. While greater lean mass is generally associated with reduced

NCD risk, the association of lean mass with health appears complex [30]. Although lean mass

incorporates muscle mass, which is widely considered to protect against diabetes, high levels of

lean mass have been associated with higher blood pressure [31]. Additionally, it is possible that

in this group, lean mass is accreted in tandem with fat mass, with the benefit of higher lean mass

being obscured by the deleterious effects of fat (particularly visceral fat) accumulation.

The role of minimum WAZ

It is important to establish the role of minimum weight-for-age in these analyses as it has been

reported that the most underweight children tend to gain more weight during nutritional

rehabilitation. In our participants, however, most of the reported associations between both

rehabilitation and post-hospitalization weight gain and adult adiposity were independent of

minimum WAZ. However, in one subgroup of participants identified by latent class analysis,

minimum WAZ appears to be influential. Therefore, the relative importance of rehabilitation

weight gain versus admission WAZ in predicting NCD risk in these adult survivors may be

dependent on how weight gain is defined, and there could be an interplay between these two factors

influencing the association between childhood weight gain and adult adiposity.

Post malnutrition growth rate.

Rapid weight gain in infancy has been defined as a weight-for-age Z score gain greater than 0.67

between birth and 1.5 years [32]. Others have defined it as a change in WAZ > 0.67 over an

unspecified time period [33, 34]. Our study participants had a mean change in WAZ during

rehabilitation (mean duration 37 days) that was more than 3 times the + 0.67 cut-off. Our statistical

observation of associations between a one-unit change in WAZ per day (ΔWAZ/day) and very

large increases in lean mass and waist circumference must be interpreted with the caution that such

an increase in WAZ is biologically implausible. Nevertheless, our findings call into question

current guidelines which also recommend that a weight gain of >10 g/kg/day during rehabilitation

is desirable, as some of our participants who gained more than 10/g/kg/day during rehabilitation

were shown to have greater adiposity as adults. In India, WAZ and g/kg/day disagreed

substantially as methods for measuring growth velocity in more than a third of preterm children

studied [35]. How the two measures agree in children born at term is unclear, however, among our

participants the two measures were similarly associated with measures of adult adiposity but not

lean mass.

In summary, all three measures of in-treatment rehabilitation weight gain were associated with at

least one measure of adult adiposity; the same was not true of the measures of post-hospitalization

weight gain. While z-score is considered the most appropriate descriptor of malnutrition, among

our participants rehabilitation weight gain in g/kg/day had the most consistent associations with

adult NCD risk factors and has the additional advantage of being more applicable in clinical

settings. Nevertheless, we were unable to draw conclusions regarding the most appropriate proxy

for post-malnutrition weight gain from this study.

Strengths and Limitations

The strengths of the study include the uniqueness of the cohort and the availability of detailed

anthropometric and body composition measures in children and adults. The low mortality

(approximately 4%) minimizes the risk of survivor bias which limits some other malnutrition-

survivor studies [36]. We acknowledge however that many post-hospitalization factors occurring

in childhood, including the home diet and intercurrent illnesses, could have influenced the

observed associations. Additionally, data relating to adult dietary intake, physical activity, and co-

morbid conditions, all of which may confound the observed associations, were not evaluated.

Finally, the participants in this study were Afro-Caribbean and the findings may be different in

other ethnic groups and in other settings where other environmental and dietary conditions apply.

Conclusion

In this unique cohort of adult Afro-Caribbean severe malnutrition survivors, we demonstrated

statistically significant associations between rehabilitation and post-hospitalization weight gain

and adult NCD risk that were independent of admission weight and adult height. While the

implications of some of the findings are mixed, we have presented evidence of a risk of increased

BMI and adiposity, especially abdominal adiposity, in these adults whose rehabilitation weight

gain exceeded certain thresholds. Our findings raise the need to further explore the existing

guidelines relating to weight gain targets during malnutrition treatment in childhood and

underscore the need for further characterisation of optimal post-malnutrition weight gain in an

interventional trial.

Acknowledgements

We gratefully acknowledge the men and women who took part in the study. We also recognize

support from the members of the CHANGE Study Collaborators' Group.

Financial Support Statement

This work was supported by the UK Medical Research Council (grant number MR/V000802/1).

The initial project was supported by the New Zealand Health Research Council Grant 09/052,

Developmental Adaptation to an Obesogenic Environment Program.

References

1. Black, R.E., et al., *Maternal and child undernutrition and overweight in low-income and middle-income countries.* Lancet (London, England), 2013. **382**(9890): p. 427-451.

2. Nobili, V., et al., *Low birth weight and catch-up-growth associated with metabolic syndrome: a ten year systematic review.* Pediatr Endocrinol Rev, 2008. **6**(2): p. 241-7.

3. Darlow, B.A., J. Martin, and L.J. Horwood, *Metabolic Syndrome in Very Low Birth Weight Young Adults and Controls: The New Zealand 1986 VLBW Study.* J Pediatr, 2019. **206**: p. 128-133.e5.

4. Tennant, I.A., et al., *Impaired cardiovascular structure and function in adult survivors of severe acute malnutrition.* Hypertension, 2014. **64**(3): p. 664-71.

5. Thompson, D.S., et al., *Childhood severe acute malnutrition is associated with metabolic changes in adulthood.* JCI Insight, 2020. **5**(24).

6. Gonzalez-Barranco, J., et al., *Effect of malnutrition during the first year of life on adult plasma insulin and glucose tolerance.* Metabolism, 2003. **52**(8): p. 1005-11.

7. WHO, *Guideline: Updates on the Management of Severe Acute Malnutrition in Infants and Children.* 2013.

8. WHO, *Pocket Book of Hospital Care for Children: Guidelines for the Management of Common Childhood Illnesses.* 2nd Edition ed. Severe acute malnutrition. 2013, Geneva: WHO.

9. Hales, C.N. and D.J. Barker, *The thrifty phenotype hypothesis.* Br Med Bull, 2001. **60**: p. 5-20.

10. Boyne, M.S., et al., *Cardiometabolic Risk in Marasmus and Kwashiorkor Survivors*, in *Handbook of Famine, Starvation, and Nutrient Deprivation: From Biology to Policy*, V. Preedy and V.B. Patel, Editors. 2017, Springer International Publishing: Cham. p. 1-23.

11. Miranda, J.J., et al., *Understanding the rise of cardiometabolic diseases in low- and middle-income countries.* Nat Med, 2019. **25**(11): p. 1667-1679.

12. WHO. *Noncommunicable Diseases.* 2021 13 April 2021 [cited 2021; Available from: https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases.

13. Francis-Emmanuel, P.M., et al., *Glucose metabolism in adult survivors of severe acute malnutrition.* J Clin Endocrinol Metab, 2014. **99**(6): p. 2233-40.

14. Campbell CP, R.D., Badaloo AV, Gluckman PD, Martinez C, Gosby A, Simpson SJ, Osmond C, Boyne MS, Forrester TE, *Developmental contributions to macronutrient selection: a randomized controlled trial in adult survivors of malnutrition. .* Evol Med Public Health, 2016(1): p. 158.69.

15. Sheppard, A., et al., *Molecular evidence for differential long-term outcomes of early life severe acute malnutrition.* EBioMedicine, 2017.

16. Waterlow, J.C., A. Tomkins, and S.M. Grantham-McGregor, *Protein-energy malnutrition.* 1992: Edward Arnold, Hodder and Stoughton.

17. Picou, D., U.o.t.W.I.T.M.R. Unit, and U.o.t.W.I.D.o. Paediatrics, *Malnutrition and Gastroenteritis in Children: A Manual for Hospital Treatment and Management.* 1978: Caribbean Food and Nutrition Institute.

18. *The Treatment and management of severe protein-energy malnutrition.* 1981, World Health Organization: Geneva.

19. Thompson, D.S., et al., *Nonalcoholic Fatty Liver Disease in Nonobese Subjects of African Origin Has Atypical Metabolic Characteristics.* Journal of the Endocrine Society, 2019. **3**(11): p. 2051-2063.

20. Waterlow, J.C., M. Golden, and J. Patrick, *Protein energy malnutrition: Treatment.* 1978. p. 49-71.

21. Alpízar, M., et al., *Fat Mass Index (FMI) as a Trustworthy Overweight and Obesity Marker in Mexican Pediatric Population.* Children (Basel), 2020. **7**(3).

22. Kwon, H., D. Kim, and J.S. Kim, *Body Fat Distribution and the Risk of Incident Metabolic*
Syndrome: A Longitudinal Cohort Study. Scientific Reports, 2017. **7**(1): p. 10955.

23. Sari, C.I., et al., *Android Fat Deposition and Its Association With Cardiovascular Risk Factors in*
Overweight Young Males. Front Physiol, 2019. **10**: p. 1162.

24. Nakano, Y., *Adult-Onset Diseases in Low Birth Weight Infants: Association with Adipose Tissue*
Maldevelopment. J Atheroscler Thromb, 2020. **27**(5): p. 397-405.

25. Demerath, E.W., et al., *Rapid postnatal weight gain and visceral adiposity in adulthood: the Fels*
Longitudinal Study. Obesity (Silver Spring), 2009. **17**(11): p. 2060-6.

26. Scheuer, S.H., et al., *Abdominal Fat Distribution and Cardiovascular Risk in Men and Women*
With Different Levels of Glucose Tolerance. J Clin Endocrinol Metab, 2015. **100**(9): p. 3340-7.

27. Katzmarzyk, P.T., et al., *Racial differences in abdominal depot-specific adiposity in white and*
African American adults. Am J Clin Nutr, 2010. **91**(1): p. 7-15.

28. Hoffman, D.J., et al., *Comparison of visceral adipose tissue mass in adult African Americans and*
whites. Obes Res, 2005. **13**(1): p. 66-74.

29. Okosun, I.S., J.P. Seale, and R. Lyn, *Commingling effect of gynoid and android fat patterns on*
cardiometabolic dysregulation in normal weight American adults. Nutrition &Amp; Diabetes,
2015. **5**: p. e155.

30. Wells, J.C.K. and M.K. Shirley, *Body composition and the monitoring of non-communicable*
chronic disease risk. Glob Health Epidemiol Genom, 2016. **1**: p. e18.

31. Grijalva-Eternod, C.S., D.A. Lawlor, and J.C. Wells, *Testing a capacity-load model for*
hypertension: disentangling early and late growth effects on childhood blood pressure in a
prospective birth cohort. PLoS One, 2013. **8**(2): p. e56078.

32. Zhou, J., et al., *Rapid Infancy Weight Gain and 7- to 9-year Childhood Obesity Risk: A Prospective*
Cohort Study in Rural Western China. Medicine (Baltimore), 2016. **95**(16): p. e3425.

33. Penny, M.E., M.M. Jimenez, and R.M. Marin, *Early rapid weight gain and subsequent overweight*
and obesity in middle childhood in Peru. BMC Obes, 2016. **3**: p. 55.

34. Lin, Q., et al., *Combined effects of weight change trajectories and eating behaviors on childhood*
adiposity status: A birth cohort study. Appetite, 2021. **162**: p. 105174.

35. Simon, L., et al., *Neonatal growth velocity of preterm infants: The weight Z-score change versus*
Patel exponential model. PLoS One, 2019. **14**(6): p. e0218746.

36. Lelijveld, N., et al., *Chronic disease outcomes after severe acute malnutrition in Malawian*
children (ChroSAM): a cohort study. Lancet Glob Health, 2016. **4**(9): p. e654-62.