Open label phase I/II clinical trial and predicted efficacy of SARS-CoV-2 RBD protein vaccines SOBERANA 02 and SOBERANA Plus in children

Authors: Rinaldo Puga-Gómez[1,2], Yariset Ricardo-Delgado[1], Chaumey Rojas-Iriarte[3], Leyanis Céspedes-Henriquez[4], Misleidys Piedra-Bello[1], Dania Vega-Mendoza[1], Noelvia Pestana Pérez[1], Beatriz Paredes-Moreno[5], Meiby Rodríguez-González[5], Carmen Valenzuela-Silva[6], Belinda Sánchez-Ramírez[7], Laura Rodríguez-Noda[5], Rocmira Pérez-Nicado[5], Raul González-Mugica[5], Tays Hernández-García[7], Talía Fundora-Barrios[7], Martha Dubet Echevarría[8], Juliet María Enriquez-Puertas[8], Yenicet Infante Hernández[8], Ariel Palenzuela-Díaz[9], Evelyn Gato-Orozco[9]; Yanet Chappi-Estévez[10], Julio Cesar Francisco-Pérez[11], Miladi Suarez Martinez[4,] Ismavy C. Castillo-Quintana[5], Sonsire Fernandez-Castillo[5], Yanet Climent-Ruiz[5], Darielys Santana-Mederos[5], Yanelda García-Vega[7], María Eugenia Toledo-Romani[12], Yury Valdés-Balbín[5], Dagmar García-Rivera[5*], Vicente Vérez-Bencomo[5*] and SOBERANA Research Group.

1. "Juan Manuel Marquez" Pediatric Hospital, Av. 31, Marianao, Havana 11400, Cuba
2. Central Clinic "Cira García". St. 20, # 4101, Miramar, Playa, La Habana 11300, Cuba
3. Policlinic "5 de Septiembre"; Av. 7th, Santa Fé, Playa, La Habana, 11600, Cuba
4. Policlinic "Carlos J. Finlay"; Av, 51 and 124, Marianao, La Habana 11400, Cuba
5. Finlay Vaccine Institute. Av. 21 ·#19810, Atabey, Playa, Havana 11600, Cuba
6. Cybernetics, Mathematics and Physics Institute. 15th Street #55, Vedado, Plaza de la Revolución, La Habana 10400, Cuba.
7. Centre of Molecular Immunology. Av 15th. and 216 Street, Siboney, Playa, Havana 11600, Cuba.
8. National Civil Defense Research Laboratory. San José de Las Lajas, Mayabeque 32700, Cuba
9. Centre for Immunoassays. 134 St and 25, Cubanacán, Playa, La Habana, 11600 Cuba
10. National Clinical Trials Coordinating Center. Av. 5th and 62 Street, Miramar, Playa, Havana 11300, Cuba.
11. Pediatric Hospital "Borrás-Marfán". St. 17 #. 801, Vedado, La Habana, 10400. Cuba
12. "Pedro Kourí" Tropical Medicine Institute. Av "Novia del Mediodía", Kv 6 1/2, La Lisa, Habana, 11400, Cuba

Soberana Research Group:

María Elena Mesa-Herrera[1], Yarmila García-Cristiá[1], Leonor Verdecia-Sánchez[1], Misleidys Piedra-Bello[1], Dania Vega-Mendoza[1], Rafael del Valle Rodríguez[1], Yudalvies Oquendo-de la Cruz[1], Noelvia Pestana-Pérez[1], Daysi Álvarez-Montalvo[1], Randy Grillo-Fortún[1], Liset López-González[1], Omaida Fonte Galindo[4], Yeseni Reyes-González[4], Ana Beatriz González-Álvarez[4], Linet Gorrita-Mora[4], Rodrigo Valera-Fernández[5], Ivis Ontivero-Pino[5], Marisel Martínez-Pérez[5], Esperanza Cabellero-Gonzalez[5], Aniurka Garcés-Hechavarría[5], Dayle Martínez-Bedoya[5], Maite Medina-Nápoles[5], Yeney Regla Domínguez-Pentón[5], Yadira Cazañas-Quintana[7], Thais Fundora Barrios[7], Diana R. Hernández Fernández[7], Gretchen Bergado-Báez[7], Ivette Orosa-Vazquez[7], Franciscary Pi-Estopiñan[7], Marianniz Díaz-Hernández[7], Otto Cruz-Sui[8], Enrique Noa-Romero[8], Arilia García-López[10], Sandra Rivadereira Muro[10], Gerardo Baro-Roman[9]

Abstract:

Objectives: To evaluate heterologous scheme in children 3-18 y/o using two SARS-CoV-2 r-RBD protein vaccines.

Methods: A phase I/II open-label, adaptive and multicenter trial evaluated the safety and immunogenicity of two doses of SOBERANA02 and a heterologous third dose of SOBERANA Plus in 350 children 3-18 y/o in Havana Cuba. Primary outcomes were safety (in phase I) and safety/immunogenicity (in phase II) measured by anti-RBD IgG ELISA, molecular and live-virus neutralization tests and specific T-cells response. An immunobridging and prediction of efficacy were additional analysis.

Results: Local pain was the unique adverse event with >10% frequency, none was serious or severe. Two doses of SOBERANA 02 elicited humoral immune response similar to natural infection; the third dose increased significantly the response in all children, similar to that achieved in vaccinated young adults and higher than in convalescents children. The neutralizing titer was evaluated in a participant´s subset: all had neutralizing antibodies vs. alpha and delta; 97.9% vs. beta. GMT was 173.8 (CI 95% 131.7; 229.5) vs. alpha, 142 (CI 95% 101.3; 198.9) vs. delta and 24.8 (CI 95% 16.8; 36.6) vs. beta. An efficacy over 90% was estimated.

Conclusion: Heterologous scheme was safe and immunogenic in children 3-18 y/o.
Trial registry: https://rpcec.sld.cu/trials/RPCEC00000374

Keywords: COVID-19; SARS-CoV-2; conjugate vaccine; pediatric vaccine, heterologous immunization scheme; subunit vaccine

Introduction

Children protection against COVID-19 is pivotal for controlling virus dissemination and reducing disease incidence. COVID-19 cases and hospitalizations among children and adolescents, firstly driven by the delta variant and recently by omicron, have risen sharply even in countries with high adult vaccination coverage (Delahoy et al., 2021; Elliott et al., 2021). This context has accelerated the clinical trials of anti-SARS-CoV-2 vaccines for children (Xia et al., 2021; Han et al., 2021; Ali et al., 202; Frenck et al., 2021; Walter et al., 2021; Wallace et al., 2021).

For more than 30 years the Finlay Vaccine Institute has produced tetanus toxoid-conjugated vaccines, applied to children worldwide; their safety has been extensively proved through hundreds of millions of doses (Verez-Bencomo et al., 2004; Huang and Wu., 2010). SOBERANA 02 was designed as an anti SARS-CoV-2 recombinant RBD conjugated to tetanus toxoid (Valdes-Balbin et al., 2021a; 2021b). It is the unique conjugate vaccine in WHO´s vaccine pipeline (WHO COVID-19, 2021). T-cell epitopes present in tetanus toxoid were expected to promote RBD specific B- and T-cell memory, and high affinity, longstanding RBD IgG antibodies.

SOBERANA 02 has proved its safety, immunogenic and efficacy in adults 19-80 y/o; after two doses, its efficacy was 71%. Combined with a third dose of SOBERANA Plus (recombinant RBD dimer vaccine) in a three-dose heterologous scheme, efficacy increased to 92.4% (Toledo-Romani et al., 2021a; 2021b; 2021c). On August 2021, the Cuban Regulatory Authority granted their emergency use authorization in adults, being extensively applied nationally for preventing COVID-19 in Cuba (CECMED, 2021).

Here, we reported the first results of open label phase I-II clinical trial in children 3-18 years old to evaluate the safety and immunogenicity of two doses of SOBERANA 02 and a third dose of SOBERANA Plus. We avoided a placebo-controlled trial in this age group due to ethical concerns (Dal-Ré and Caplan., 2021); alternatively, a recommended comparison or immunobridging with adult´s immunogenicity was established (FDA, 2021) and the clinical efficacy was estimated based on immunological results.

Method

Study design

We designed a phase I/II study, open-label, adaptive and multicenter to evaluate the safety, reactogenicity and immunogenicity of two doses of SOBERANA02 and a heterologous third dose of SOBERANA Plus in children (3-11 y/o) and teenager (12-18 y/o). Two interim analysis would decide interruption/continuation of the study, depending on serious adverse events (AEs) during phase I.

Phase I was conceived as a two-step, incorporating firstly 25 children 12-18 y/o (sequence 1). The first interim report (no serious AE) seven days after vaccinated these children allowed incorporating 25 children aged 3-11 (sequence 2 of phase 1) and started the phase II in 12-18 y/o (n=150). A second interim report seven days after sequence 2 (no serious AE) allowed starting phase II in children 3-11 y/o (n=150). (Figure 1).

Phase I was conducted at "Juan Manuel Marquez" Pediatric Hospital in Havana; phase II was conducted in policlinics "5 de Septiembre" and "Carlos J. Finlay" in Havana, Cuba. Clinical sites were certified by the National Immunization Program.

Participants

Children were recruited at community level across the primary health system. They were included following physical examination, parent interview and, for phase I, clinical laboratory assays. Key inclusion criteria were weight-height nutritional assessment; physical examination without alterations; clinical laboratory results within the range of reference values (only phase I) and microbiology laboratory tests. Key exclusion criteria were any acute infection, previous or current history of SARS CoV-2 infection and being a contact of a positive COVID-19 case. A detailed description of selection criteria appears in Supplementary Material I.

Ethical issues

The trial was approved by the Ethical Committee at the "Juan Manuel Marquez" Pediatric Hospital and endorsed by the Cuban National Pediatric Group. The Cuban National Regulatory Agency (Centre for State Control of Medicines and Medical Devices, CECMED) approved the trial (June10th, 2021, Authorization Reference: 05.010.21BA).

Independent Data Monitoring Committees formed by five external members specialized on paediatric clinical practice, immunology and statistics were in charge of two interim analyses during phase I.

The trial was conducted according to Helsinki's Declaration, Good Clinical Practice and the Cuban National Immunization Program. During recruitment, the medical investigators provided to the parents, both orally and written, all information about the vaccine and potential risks and benefits. Written informed consent was obtained from both parents; children ≥over 12 y/o should assent. The decision to participate was not remunerated.

The National Clinical Trials Coordinating Centre (CENCEC) was responsible for monitoring
the trial in terms of adherence to the protocol, Good Clinical Practice and data accuracy.

Trial registry: RPCEC00000374 (Cuban Public Registry of Clinical Trials and WHO
International Clinical Registry Trials Platform) (IRCT, 2021)

Products under evaluation

SOBERANA 02 (RBD chemically conjugated to the carrier protein tetanus toxoid) and
SOBERANA Plus (RBD dimer), adjuvated in alumina hydroxide were produced at the Finlay
Vaccine Institute and the Centre for Molecular Immunology, in Havana, Cuba, under GMP
conditions. Both are subunit vaccines based SARS-CoV-2 RBD, sequence Arg319-Phe541
produced in genetically modified CHO cells. Formulations are detailed in Supplementary
Material II-Table S1.

Product batches used: SOBERANA 02 (E1002S02X, E1002S02); SOBERANA Plus
(E1001SP).

Procedures

RT-PCR SARS-CoV-2 was performed to all participants at least 72 hours before each dose.
Participants with negative PCR results received the vaccine by intramuscular injections in the
deltoid region. *Immunization schedule*: two doses of SOBERANA 02 and a heterologous third
dose of SOBERANA Plus 28 days apart (immunization on days 0, 28, 56). After each
immunization, participants were on-site evaluated during one hour. Medical controls visits
were planned at 24, 48 and 72 hours, and on days 7, 14 and 28 after each dose. AEs were
parents-registered daily. Serum samples were collected on day 0 (before vaccination) and 14
days after the second and the third dose (days 42 and 70). Peripheral blood mononuclear cells
were obtained before vaccination and after the third dose (day 70) in a participants' subset of
45 children randomly selected in each age subgroup.

Outcomes

Primary outcomes. Phase I: occurrence of serious AEs, measured daily during 28 days after
each dose. Phase II: % of subjects with seroconversion ≥ 4 (fold increase of IgG anti-RBD
respect to pre-vaccination level), on days 42 and 70.

Secondary outcomes. Phase II: Occurrence of serious AEs, measured during 28 days after each
dose. Phase I and Phase II: Solicited local and systemic AEs, measured during 7 days after each
dose; unsolicited AEs, measured during 28 days after each dose; neutralizing antibody titres
(on days 42 and 70, on a sample subset), inhibition of RBD-hACE2 interaction (on days 42,
70). Outcomes are detailed in Supplementary Material III)

Safety assessment

Pain, erythema, swelling, induration and temperature were local-solicited AEs at the injection
site. Fever $\geq 38\,^{\circ}$C, low-grade fever ($<38\,^{\circ}$C), general discomfort and rash were systemic-
solicited AEs. Any other events were parent-recorded throughout the 28 days follow-up
period after each dose. Clinical laboratory test included pre-vaccination and post-vaccination
biochemical serum analysis (only during phase I).

AEs were classified as serious or not serious; severity—according to Brighton Collaboration
definition and the Common Terminology Criteria for AE version 5·0—was: mild when AE
was transient (not interfering with activities), moderate (when caused mild to moderate
limitation in activity), or severe (limitating activity). AEs were reviewed for causality and
classified according to WHO: inconsistent causal association to immunization, consistent
causal association to immunization, undetermined, unclassifiable (WHO, 2018).

 Immunogenicity assessment

A quantitative ultramicro ELISA (UMELISA SARS-CoV-2 anti- RBD, Centre for Immunoassay, Havana, Cuba) determined the concentration of anti-RBD IgG antibodies in sera, expressed as AU/mL. A competitive ELISA determined the inhibitory capacity of antibodies for blocking the RBD-hACE2 interaction, expressed as % inhibition and molecular inhibitory titer ($mVNT_{50}$). Conventional virus neutralization titer ($cVNT_{50}$) vs. D614G strain was evaluated in a subset of samples randomly selected from the individuals with seroconversion after two doses (on day 42 n=123) and after three doses (n=131). Other subset of samples (n=48) were randomly selected for neutralizing tests vs alpha, beta, and delta compared with D614G, after the third dose (on day 70). RBD-specific T-cell response producing IFN- γ and IL-4 were quantified by enzyme-linked immunospot (ELISpot) assay using human IFN-γ ELISpot[PLUS] HRP kit (Mabtech, Sweden) and human IL-4 ELISpot[plus] HRP kit (Mabtech, Sweden) following the manufacturer's instructions. A detailed description of immunological assays is described in Supplementary Material IV. All immunological evaluations were performed by external laboratories, except T cell response.

Children Convalescent Serum Panel

A Cuban children convalescent serum panel (CCCSP) was made with sera from 82 patients (3-18 y/o) recovered from COVID-19: five from children with moderate symptomatic disease, 29 with mild disease and 48 asymptomatic; concerning prevalent variant of concern, 21 from delta wave and 61 from pre-delta wave when D614G and beta variants predominated in Havana. Convalescent children were attended by specialized doctors; their parents gave written consent for studying their immunological status after natural infection allowing their use for epidemiological research. This panel was characterized by anti-RBD IgG concentration (UA/ml), inhibition of RBD-hACE2 interaction (% of inhibition and molecular neutralization titre) and virus neutralization titre ($cVNT_{50}$) as described.

Statistical analysis

For phase I, the calculation of sample size was done considering a two-sided 95% confidence interval for one proportion with a width equal to 0.09 to estimate a serious AE rate <1%. For phase II a similar method was used to estimate a seroconversion around 50%, with a lower bound of the confidence interval >30% (hypothesis work) and a dropout of 20%. This resulted in a sample size of 350 subjects (including subjects from phase I).

Safety and reactogenicity endpoints are described as frequencies (%). Quantitative demographic characteristics are reported as mean, standard deviation (SD), median, interquartile range, and range. Seroconversion rate for anti-RBD IgG antibodies (≥4-fold increase in antibody concentration over baseline) was expressed as % with the 95% confidence interval and seroconversion index as median and interquartile range. Anti-RBD IgG concentration and % of inhibition of RBD-hACE2 interaction were expressed as median and interquartile range; mVNT50 and cVNT50 were expressed as geometric mean titer (GMT) and 95% confidence intervals (CI).

Spearman´s rank correlation was used to assess relationships among techniques used to evaluate the immune response. ROC curve was used to choose the most appropriate cut-off for humoral tests regarding the $cVNT_{50.}$ The Student t-Test or the Wilcoxon Signed-Rank Test were used for before-after statistical comparison.

A stopping criterion due to unacceptable toxicity (frequency >1%) was evaluated iteratively, with a Bayesian algorithm. As an additional analysis, a non-inferiority of the immune response in 3-11 and 12-18 y/o participants compared with that in 19-39 y/o participants (from not

concurrent phase I and phase II studies (Toledo-Romani et al., 2021a; 2021b) was assessed, using the geometric mean ratio of SARS-CoV-2 50% neutralizing titers (cVNT$_{50}$). The geometric mean ratio and two-sided 95% confidence intervals were calculated by exponentiating the mean difference of the logarithms of the titers and the corresponding confidence intervals (based on the Student's t distribution). The non-inferiority criterion was met, since the lower boundary of the two-sided confidence interval for the geometric mean ratio was >0.67 (FDA, 2021).

A prediction of clinical efficacy was estimated using the regression linear model reported for seven vaccines based on immunogenicity data (Khoury et al 20211; Earle et al., 2021). Statistical analyses were done using SPSS version 25.0; STATISTICA version 12.0, R version 3.2.4, EPIDAT version 3.1 and Prism GraphPad version 6.0. An alpha signification level of 0.05 was used.

Results.

Figure 1 and Table 1 describe the study design and demographic characteristics of the participants. From 11th June to 14th July 2021, 426 children (3-18 y/o) were recruited, 350 that accomplished selection criteria were included, and 306 completed the study. There was a balanced ratio on sex and ethnicity; mean age was 11.3 years (SD 4.5).

Phase I started by vaccinating 25 children 12-18 y/o with SOBERANA 02; the first interim analysis done seven days after vaccination indicated absence of serious AEs. In consequence, the trial proceeded to phase I, sequence 2, incorporating 25 children aged 3-11; the second interim analysis showed no serious AE. The trial proceeded to phase II, vaccinating 150 children 12-18 y/o and 150 children 2-11 y/o with SOBERANA 02 first dose.

Figure 1. Flow chart: recruitment, inclusion, vaccination and follow up of 3-18 years old children in phase I/II trial.

During the vaccination scheme, 86 children (53.1%) suffered at least one AE; frequency was higher (60%) in teenagers than in young children (46.3%). Severe and serious vaccine-associated AEs (VAAE) did not occur (Table 2). Local AE predominated; the most common was local pain (47.7%), all others had frequencies <5%; only 1.1% reported fever. More than

>90% of AE were classified as mild and lasted 72 hours or less, and 88.5% were associated with vaccination (Table 3, Supplemental Material V-Table S2). AEs were more frequent after the first dose than after the second and the third dose (Figure 2, Supplemental Material V-Table S3). Few unsolicited adverse events were recorder (Supplemental Material V-Table S4). No clinically relevant changes were observed in hematology and blood chemistry analyses (Supplemental Material V-Table S5)

Table 1: Demographic characteristics of subjects included in the clinical trial

	Age groups		
	3-11 years	**12-18 years**	**Total**
N	175	175	350
Sex			
Female	80 (45.7%)	83 (47.4%)	163 (46.6%)
Male	95 (54.3%)	92 (52.6%)	187 (53.4%)
Skin colour			
White	122 (69.7%)	116 (66.3%)	238 (68.0%)
Black	9 (5.1%)	11 (6.3%)	20 (5.7%)
Multiracial	44 (25.1%)	48 (27.4%)	92 (26.3%)
Age (years)			
Mean (SD)	7.4 (2.5)	15.1 (2.1)	11.3 ± 4.5
Median (IQR)	8.0 (5.0)	15.0 (4.0)	11.5 ± 7.0
Range	3; 11	12;18	3-18
Weight (kg)			
Mean (SD)	29.4 (10.1)	54.7 (9.0)	42.0 ± 15.9
Median (IQR)	27.5 (14.0)	55.0 (13.0)	43.0 ± 27.7
Range	13.0; 58.0	32.0; 80.0	13.0; 80.0
Height (cm)			
Mean (SD)	129.1 (17.2)	164.3 (9.6)	146.7 ± 22.5
Median (IQR)	131.0 (26.0)	164.0 (13.0)	151.0 ± 34.0
Range	92; 172	142; 190	92-190
BMI (kg/m^2)			
Mean (SD)	17.0 (2.0)	20.2 (2.3)	18.6 ± 2.7
Median (IQR)	16.7 (2.7)	19.9 (3.8)	18.3 ± 4.1
Range	13.2; 22.8	14.6; 25.5	13.2-25.5

Data are n (%) unless otherwise specified. Mean (SD)=Mean ± Standard Deviation. Median (IQR)=Median ± Interquartile Range. BMI=Body mass index. Range=(Minimum; Maximum)

Table 2: General characteristics of adverse events

	Age groups		
	3-11 years	**12-18 years**	**Total**
N	175	175	350
Subjects with some AE	81 (46.3%)	105 (60.0%)	186 (53.10%)
Subjects with some VAAE	76 (43.4%)	101 (57.7%)	177 (50.6%)
Subjects with some Serious AE	0 (0.0%)	1 (0.6%)	1 (0.3%)
Subjects with some Serious VAAE	0 (0.0%)	0 (0.0%)	0 (0.0%)
Subjects with some Severe AE	0 (0.0%)	0 (0.0%)	0 (0.0%)
Subjects with some Severe VAAE	0 (0.0%)	0 (0.0%)	0 (0.0%)
Total Adverse Events	141	182	323
Mild AE	135 (95.7%)	167 (91.8%)	302 (93.5%)
Moderate AE	6 (4.3%)	15 (8.2%)	21 (6.5%)
Severe AE	0 (0.0%)	0 (0.0%)	0 (0.0%)
Serious AE	0 (0.0%)	1 (0.5%)*	1 (0.3%) *
Local AE	118 (83.7%)	147 (80.8%)	265 (82.0%)
Systemic AE	23 (16.3%)	35 (19.2%)	58 (18.0%)
VAAE	126 (89.4%)	160 (87.9)	286 (88.5%)
Serious VAAE	0 (0.0%)	0 (0.0%)	0 (0.0%)
Severe VAAE	0 (0.0%)	0 (0.0%)	0 (0.0%)

Data are n (%). AE=Adverse Event. VAAE=Vaccine-Associated Adverse Event;
* Serious AE: Dengue required hospitalization.

Table 3: Frequency of solicited adverse events

	3-11 years	12-18 years	Total
		Age groups	
N	175	175	350
Subjects with some AE	81 (46.3%)	105 (60.0%)	186 (53.10%)
Subjects with solicited local AE			
Any	74 (42.3%)	98 (56.0%)	172 (49.1%)
Local pain	69 (39.4%)	98 (56.0%)	167 (47.7%)
Swelling	9 (5.1%)	2 (1.1%)	11 (3.1%)
Local warm	4 (2.3%)	0 (0.0%)	4 (1.1%)
Erythema	5 (2.9%)	1 (0.6%)	6 (1.7%)
Induration	5 (2.9%)	1 (0.6%)	6 (1.7%)
Subjects with solicited systemic AE			
Any	5 (2.9)	4 (2.3)	9 (2.6)
General discomfort	1 (0.6%)	3 (1.7%)	4 (1.1%)
Fever ($\geq 38\,°C$)	2 (1.1)	1 (0.6)	3 (0.9)
Low-grade fever ($<38\,°C$)	4 (2.3)	1 (0.6)	5 (1.4)

Data are n (%) unless otherwise specified. AE=Adverse Event

Figure 2. Frequency of solicited adverse events after each doses, by age group and overall.

Before vaccination, 97.4 % of children were negative for anti-RBD antibodies. Two doses of
SOBERANA 02 induced seroconversion in 95% participants (CI 95% 92.7; 97.6),
accomplishing with the study hypothesis: more than 50% of seroconversion with

lower boundary of the two-sided 95% confidence interval of >0.3). By age subgroup,
seroconversion was 99.4% (CI 95% 96.5; 99.9) in children 3-11 y/o and 91.8% (CI 95% 86.4;
95.6) in 12-18 y/o. Global seroconversion index was 28.3 (42.6 in children 3-11 y/o, 25.3 in
12-18 y/o); the median anti-RBD IgG was 57.0 UA/mL (25^{th}-75^{th} percentile 29.8; 153.4),
being higher in younger children (93.3 UA/mL; 25th-75th percentile 39.0; 214.4). Specific
antibody response was higher than the elicited by natural infection, evaluated in Cuban

321   children convalescent panel (anti-RBD IgG median 11.5; 25^{th}-75^{th} percentile 5.3; 24.2). The
322   heterologous third dose increased seroconversion to 100% and seroconversion index to 154.5;
323   anti-RBD IgG titers also increased significantly (p<0.005) to 325.7 UA/mL (25^{th}-75^{th}
324   percentile 141.5; 613.8) (Table 4).

325
326
327

Table 4: Humoral immune response induced after two doses of SOBERANA 02 and a third heterologous doses with SOBERANA Plus.

		Age groups						CCCSP
		3-11 years		12-18 years		Total		
		Post-2nd dose	Post-3rd dose	Post-2nd dose	Post-3rd dose	Post-2nd dose	Post-3rd dose	
	N	159	156	159	150	318	306	82
Anti-RBD IgG seroconversion rate	N (%)	157/158 (99.4)	155/155 (100.0)	146/159 (91.8)	150/150 (100%)*	303/317 (95.6)	305/305 (100.0)*	N.D
	95% CI	96.5; 99.9	99.6; 100.0	86.4; 95.6	99.6; 100.0	92.7; 97.6	99.8; 100.0	
Anti-RBD IgG AU/mL	Median	93.3	329.8*	49.9	325.6*	57.0	325.7*	11.5
	25th-75th	39.0; 214.4	162.9; 685.6	22.3; 90.7	108.0; 555.8	29.8; 153.4	141.5; 613.8	5.3; 24.2
Seroconversion index	Median	42.6	155.4*	25.3	154.3*	28.3	154.5*	N.D
	25th-75th	19.5; 97.0	75.4; 260.2	11.0; 45.7	50.2; 266.3	14.9; 69.8	67.2; 260.9	
RBD:hACE2 Inh%	Median	78.3	92.5*	60.1	92.2*	67.4	92.4*	20.8
	25th-75th	49.4; 88.9	88.8; 93.4	36.7; 79.7	87.7; 93.6	42.1; 86.9	88.3; 93.5	10.9; 40.8
mVNT$_{50}$	GMT	274.1	1418.3*	143.7	1116.2*	198.5	1261.2*	35.2
	95% CI	216.8; 346.5	1180.0; 1704.7	115.0; 179.5	924.0; 1348.3	168.4; 233.9	1105.5; 1438.8	25.3; 48.9
cVNT$_{50}$ vs D614G	N	63	66	60	65	123	131	70
	GMT	28.4	181.6*	24.4	137.9*	26.4	158.4*	9.2
	95% CI	18.5; 43.6	120.6; 273.3	17.6; 40.0	101.8; 186.9	20.2; 34.5	123.0; 204.0	6.8; 12.5

T0 or baseline anti-RBD IgG 1.95 (25th-75th percentile 1.95; 1.95) for both age subgroups and total. ND: Not determined

Anti-RBD IgG seroconversion rate: % of subjects with seroconversion (CI 95%).

Seroconversion index: increase of IgG concentration respect to baseline (median; 25th-75th percentile)

AU/mL=anti-RBD IgG concentration expressed in arbitrary units/mL.

RBD:hACE2 INH%: RBD:hACE2 inhibition % at a dilution 1/100.

mVNT$_{50}$: molecular virus neutralization titer; as serum dilution inhibiting 50% of RBD:hACE2 interaction.

cVNT$_{50}$: conventional live-virus neutralization titer.

GMT=Geometric Mean Titer. 95% CI= 95% Confidence Interval; 25th-75th=25-75 percentile.

* p<0.005 versus Post - 2nd dose McNemar test (anti-RBD IgG seroconversion %), Wilcoxon Signed Ranks test (anti-RBD IgG AU/mL, RBD:hACE2 Inh%) or Paired Student t test (mVNT$_{50}$, cVNT$_{50}$, log-transformed). CCCSP=Cuban children convalescent serum panel

The capacity of anti-RBD IgG for blocking RBD- hACE2 interaction after two doses of SOBERANA 02 was 67.4 % (25^{th}-75^{th} percentile 42.1; 86.9) and the $mVNT_{50}$ was 198.5 (CI 95% 168.4; 233.9); both increased significantly (p<0.005) after the third dose to 92.4 % (25^{th}-75^{th} percentile 88.3; 93.5) and 1261 (CI 95% 1105,5; 1438.8) respectively. These values were higher among the youngers (3-11 y/o) after the second dose, but were similar in both age subgroups after the third dose. The mVNT50 of antibodies was higher after both two and three doses than after natural infection (Table 4).

After two doses of SOBERANA 02 the neutralizing titer versus D614G variant was higher (GMT 26.4 CI 95% 20.2; 34,5) than the children convalescent panel value (GMT 9.2 CI 95% 6.8; 12.5); and the third dose significantly (p<0.005) boosted the response to GMT 158.4 (CI 95% 123.0; 204.0). The neutralizing titer versus the variants alpha, beta and delta was evaluated in 48 children; 100% had neutralizing antibodies vs. alpha and delta; 97.9% vs. beta (Table 5). GMT was 173.8 (CI 95% 131.7; 229.5) vs. alpha, 142 (CI 95% 101.3; 198.9) vs. delta and 24.8 (CI 95% 16.8; 36.6) vs. beta (a 2.2-fold decrease for delta and 7.0-fold decrease for beta compared to D614G).

Table 5. Conventional live-virus neutralization titers against SARS-CoV-2 variants alpha, delta and beta.

		D614G	Alpha	Delta	Beta
$cVNT_{50}$	N	48	48	48	48
	GMT	173.8	142.0	76.8*	24.8*
	95% CI	131.7; 229.5	101.3; 198.9	54.8; 107.7	16.8; 36.6

Sera from 48 children vaccinated with complete schedule (two doses SOBERANA 02 + one dose SOBERANA Plus, 28 days apart) were evaluated against D614G, alpha, delta and beta variants. $cVNT_{50}$=conventional live-virus neutralization titer. GMT=Geometric Mean Titer. 95% CI= 95% Confidence Interval. * p<0.005 Paired Student t test ($cVNT_{50}$, log-transformed) respect to D614G variant.

There was a good correlation among all humoral immunological variables, with coefficients greater than 0.7 (Supplemental Material VI-Table S6). Predictive cut-off for attaining $cVNT_{50}$

over 50 was estimated by ROC curve as: 192.2 AU/ml for IgG concentration, 87.1% for the inhibition of RBD:hACE2 and 427 for $mVNT_{50}$ (Supplemental Material, Table S7, Figure S1)

RBD-specific T cell response in a subset of 45 participants fully vaccinated was determined by measuring IFN-γ and IL-4 expression in peripheral blood mononuclear cells. The number of IFN-γ and IL-4 secreting cells was statistically higher (p<0.001) than their baseline levels (Figure 3), suggesting a mixed Th1/Th2 profile. IFN-γ expression was higher in older children while IL-4 was higher in younger children, evidencing differences in the balance Th1/Th2.

Figure 3. IFN-γ- and IL-4-secreting cells in peripheral blood mononuclear cells stimulated with RBD. Children 3-11 (N=24) and 12-18 years old (N=21) received two doses (on days 0, 28) of SOBERANA 02 and a heterologous third dose (on day 56) of SOBERANA Plus. p value represents the statistic differences as indicated.

The safety and immune response in children was compared with all young adults aged 19-39 vaccinated during Phase I and Phase II studies (FDA, 2021) (Supplemental Material VII-Table S8). Safety profile was similar in children respect to young adults (Supplemental Material VII-Table S9, Figure 2). An immunobridging analysis was performed for anti-RBD

IgG, mVNT50 and cVNT50 between children and young adults. IgG elicited after two doses
of SOBERANA 02 was 57.0 UA/ml (25th-75th percentile 29.8; 153.4), while for young adults
it was 46.4 (25th-75th percentile 17.4; 108.8); after the heterologous third dose of
SOBERANA Plus these values increased to 325.7 (25th-75th percentile 141.5; 613.8) in
children and 228.0 (25th-75th percentile 95.8; 394.3) in young adults. $mVNT_{50}$ was 198.5 (IC
95% 168.4; 233.9) in children after the second dose and 1261.2 (IC 95% 1105.5; 1438.8) after
the third; in young adults were 94.9 (IC 95% 75.0; 120.2) and 503.7 (IC 95% 432.6; 586.6)
after two and three doses (Figure 4). We found significant difference (p<0.05) for IgG and
$mVNT_{50}$ between children and young adults, with higher values in the children both after the
second and the third dose. Viral neutralization titers after the second dose were measured at
different time points in children and young adults (on day 42 in children and day 56 in adults)
making their comparison only approximate.

The non-inferiority analysis was performed by cVNT50, following FDA recommendation
(FDA, 2021). After three doses (on day 70), cVNT50 in children was 158.4 (IC 95% 123.0;
204.0) and 122.8 (80.2; 188.0) for young adults (n=43, data available) (Figure 4). The
immune response in 3-18 y/o, as well as 3-11 y/o and 12-18 y/o was non-inferior to that
observed in 19-39 y/o young adults. The geometric mean ratio of the neutralizing geometric
mean titer in 3-18 y/o participants respect to young adults 14 days after third dose was 1.25
(CI 95% 0.77; 2.02) (Table 6), which met the non-inferiority criterion (i.e., a lower boundary
of the two-sided 95% confidence interval of >0.67). In summary, both age groups (3-11 y/o
and 12-18 y/o) met the non-inferiority criterion

Figure 4: Immunobridging comparison of humoral immune response elicited in children (3-18 y/o) respect to young adults (19-39 y/o from Phase I and II clinical trial), after two doses of SOBERANA 02 (day 42) and a third dose of SOBERANA Plus (day 70). A) anti-RBD IgG median (25th-75th percentile); B) mVNT$_{50}$ GMT (CI 95%); C) cVNT$_{50}$ GMT (CI 95%). Bleeding was on day 42 and 70 (14 days after second or third dose), except for cVNT50 adults after second dose was on day 56. Mann-Whitney U test (anti-RBD IgG AU/mL) or Student t test (mVNT$_{50}$, cVNT$_{50}$, log-transformed).

408 Table 6: Immunobridging of $cVNT_{50}$ in children and young adults after heterologous scheme
409 (two doses of SOBERANA 02 and a third heterologous doses of SOBERANA Plus)

Age group	No. of participants	$cVNT_{50}$ GMT (95% CI)	Geometric mean ratio (95% CI) vs. 19 to 39 y/o
19-39 y/o	43	127.0 (89.6; 179.80)	--
3-18 y/o	131	158.4 (123.0; 204.0)	1.25 (0.77; 2.02)
3-11 y/o	66	181.6 (120.6; 273.3)	1.43 (0.80; 2.54)
12-18 y/o	65	137.9 (101.8; 186.9)	1.08 (0.68; 1.73)

GMT and two-sided 95% confidence intervals were calculated by exponentiating the mean logarithm of the titers and the corresponding confidence intervals (based on the Student's t distribution). The geometric mean ratio and two-sided 95% confidence intervals were calculated by exponentiating the mean difference of the logarithms of the titers (in child/adolescents cohorts minus the 19-39-year-old cohort) and the corresponding confidence intervals (based on the Student's t distribution). The non-inferiority criterion was met, since the lower boundary of the two-sided confidence interval for the geometric mean ratio was greater than 0.67.

Based on immunogenicity data of vaccinated children and the immune response to natural
infection (children convalescent panel), a prediction of clinical efficacy was estimated
through a regression linear model. By using $cVNT_{50}$ is the predictive variable, estimated
efficacy is 91.3% (CI 95% 84.6; 95.1) after two doses and 97.4 % (IC 95% 91.5; 99.2) after
three doses (Figure 5). This study was conducted in Havana during delta wave; after receiving
the first dose 29 children (partially vaccinated) (8.2%) of 350 children became PCR-positive,
this value dropped down to 13 (4.0%) out of 321 children after completing the two-dose
scheme) and no one after the third dose but the follow-up period after third dose was only 28
days.

Figure 5. Prediction of clinical efficacy in children from correlation between antibody responses and efficacy rate. Panels displays correlation of cVNT$_{50}$ neutralization and ratios, respectively for seven vaccines. The y-axis is estimated log risk ratio reported on the vaccine efficacy scale. The x-axis is log ratio of the peak geometric mean neutralization at 7-28 days post-vaccination, relative to human or children convalescent sera.

Discussion.

This study describes, for a first time, the safety and immunogenicity of SOBERANA 02 in children 3-18 y/o, after two doses schedule and heterologous third doses with SOBERANA Plus. The frequency of local and systemic AE was 49 and 2.6% respectively, lower than

mRNA COVID-19 (Frenck et al., 2021; Ali et al., 2021). After two doses schedule,
BNT162b2 vaccines reported 86% and 66% of children (aged 12-15) with local and systemic
AEs; mRNA-1273 reported 94.2% and 68.3% (aged 12-17 y/o) respectively (Frenck et al.,
2021; Ali et al., 2021). In our study, local pain was reported in 51.4% of children (12-18 y/o)
after first dose, 17% after the second and 17.3% after the third one. BNT162b2 and mRNA-
1273 vaccine in adolescents reported 86 % and 94.2% with local pain after first dose; 79%
and 92.4% after the second, respectively (Frenck et al., 2021; Ali et al., 2021). SOBERANA
02 and SOBERANA Plus caused general discomfort (the most frequent systemic AE) only in
1.7% of children 12-18 y/o, while mRNA vaccines provoked fatigue, headache, chills, muscle
pain or fever between 10-68.5% of adolescents (Frenck et al., 2021; Ali et al., 2021). In
children aged 3-11 local pain was the unique EA with frequency >10% during this study;
children aged 5-11 vaccinated with BNT162b2 reported local pain (74%), redness (19%),
swelling (15%), fatigue (39%) and headache (28%) (Walter et al., 2021). Myocarditis and
pericarditis have been reported in adolescent after mRNA COVID-19 vaccination (Oster et
al., 2021; Marshall et al., 2021); these AEs were not observed here.

The comparison between humoral immune response elicited by vaccination respect to those
elicited by natural infection has been useful for the development of several anti-SARS-CoV-2
vaccine (Keech et al., 2020; Yang et al., 2021). Two shots of SOBERANA 02 every 28 days
in children induced a robust humoral response, with higher levels of antibodies but similar
neutralizing capacity compared with those elicited by natural infection. The third dose with
SOBERANA Plus boosted the levels of antibodies and also the neutralizing capacity,
surpassing the immune response in convalescent children, similar than occurred during
clinical trials in adults (Toledo-Romani et al., 2021a; 2021b).

The induction of specific T-cell response is critical for protection of viral infections. The
heterologous three doses schedule in children developed a balanced activation of IFN-γ and
IL-4-secreting cells from PBMC, indicating a mixed Th1/Th2 response, as reported in adults
after the same vaccination scheme (Toledo-Romani et al., 2021a).

The SARS-CoV-2 variants of concern alpha, beta, delta and recently omicron have been
modifying the pandemic landscape worldwide (Fontane et al., 2021). Here, we reported the
capacity of anti-RBD antibodies for neutralizing alpha, beta and delta variant, with a fold-
reduction of 2.2 for delta and 7.0 for beta respect to D614G similar to we found in adults
(Toledo-Romani et al., 2021b). Official and public communication from the "Pedro Kourí"
Tropical Medicine Institute refers the capacity of adults vaccinated with SOBERANA 02 +
SOBERANA Plus for neutralizing omicron variant (Portal-Miranda 2022; Carles JM, 2022).

Recently, the Pasteur Institute of Iran informed that Pasteurcovac, the Iranian brand of
SOBERANA 02, is 100% effective against omicron variant in adults (Biglari 2022).
472
We conducted this clinical trial during the delta wave, the worst period of Cuban epidemic
(Rodriguez; 2021); in such a context and due to ethical reasons, a placebo-controlled clinical
trial was not ethical. Lacking a control group, two analytical tools complemented the study:
immunobridging with the immune response in young adults (FDA, 2021); and prediction of
clinical efficacy based on immunological response (Khoury et al., 20211; Kristen et al.,
2021). Firstly, we found a non-inferior response for the ratio of SARS-CoV-2 50%
neutralizing titers after full three doses scheme in 3-11 and 12-18 year-old participants
relative to 19-to-39-year-old as reference population (no concurrent). The comparison met the
non-inferiority criterion with ratio 1.43 (CI 95% 0.8 to 2.54) for 3-11 years old and 1.08 (CI
95% 0.68-1.73) for 12-18-year-old, accomplishing with the recommended (FDA, 2021).
Similar analysis has been reported by BNT162b2 and mRNA-1273 vaccines but using 19-25
years old as reference population (Walter et al., 2022; Ali et al., 2021). We considered young
adults upon 39 year as immunocompetent age (Lopez-Sejas 2016; Ventura et al., 2017; Thapa
and Farber 2019) allowing us wide-ranging the reference population for having more $cVNT_{50}$
data available.
488
Secondly, the prediction of clinical efficacy based on immunological response have been used
for other vaccines (Khoury et al., 20211; Kristen et al., 2021). Using this model, we
anticipated a clinical efficacy in adults aged 19-80 between 58% and 87% (for two doses
scheme) and between 81% and 93% for the three-dose scheme (Toledo-Romani et al., 2021b).
These results were confirmed during a phase III clinical trial reporting a 71% of efficacy for
the two-dose schedule of SOBERANA 02 and 92.4% for the heterologous three-dose
schedule (Toledo-Romani et al., 2021c). Here, the model predicts 89.3-91.3% clinical
efficacy after two doses of SOBERANA 02 and 95.1-97.4% after the third doses of
SOBERANA Plus in children.
Children vaccination as 2 y/o is key for controlling the pandemic, cutting the transmission and
reducing the emergence of new VOCs (Petersen and Buchy, 2021). The safety and
immunological results reported here supported the emergency use authorization of
SOBERANA 02 and SOBERANA Plus as heterologous scheme for children between 2-18
y/o. A massive immunization campaign started on 5[th] September 2021; fully vaccinating 1.8
millions of Cuban children (96% of Cuban 2-18 y/o population (Augustin, 2022). These

results support public health vaccination strategies, providing children as young as two years
a safe and effective vaccine scheme to prevent COVID-19.
506
507

508 *Funding*
509 This work was supported by the Finlay Vaccine Institute, BioCubaFarma and the National
510 Funds for Sciences and Technology from the Ministry of Science, Technology and
511 Environment (FONCI-CITMA-Cuba, contract 2020-20).

512
Declaration of Interests
The authors R.P.G, Y.R.D, C.R.I, L.C.H. M.P.B, D.V.M, N.P.P, C.V.S, A.P.D, E.G.O, Y.C.E,
J.C.F.P, M.S.M, M.D.E, J.M.E.P, Y.I.H, M.E.T.R declare that they have no known
competing financial interests or personal relationships that could have appeared to influence
the work reported in this paper.

The authors B.P.M, M.R.G, B.S.R, L.M.R.N, R.P.N, R.G.M, T.H.G, T.F.B, I.C.Q, S.F.C,
Y.C.R, D.S.M, Y.G.V, Y.V.B, D.G.R, V.V.B work at Finlay Vaccine Institute or the Centre
of Molecular Immunology, institutions that develop and manufacture the vaccine candidates
but they haven't received an honorarium for this paper

B.S.R., S.F.C., Y.C.R, L.R.N., D.S.M., Y.V.B., D.G.R. and V.V.B., have filed patent
applications related to the vaccine SOBERANA 02.
524

525 Acknowledgments

526 We especially thanks all the parents and children for participated in the clinical trial. We
527 recognize the contribution of all the medical and nurse staff at clinical site. We thank Dr. Lila
528 Castellanos for scientific advice.

529
530 References
531

1. Ali K, Berman G, Zhou H, Deng W, Faughnan V, Coronado-Voges M, et al.
Evaluation of mRNA-1273 SARS-CoV-2 Vaccine in Adolescents. N Engl J Med 2021;
385: 2241-51. DOI: 10.1056/NEJMoa2109522

2. Augustin Ed . Cuba leads the world in vaccinating children as young as two against
Covid. The Guardia. February 1st, 2022.
https://www.theguardian.com/world/2022/feb/01/cuba-leads-world-vaccinating-
children (Accessed February 17th, 2022)
3. Avci FY, Li X, Moriya T, Kasper D. A mechanism for glycoconjugate vaccine
activation of the adaptive immune system and its implications for vaccine design.
Nature Medicine 2011;17:1602-10. https://doi.org/10.1038/nm.2535
4. Biglari A. Official says Pasteurcovac 100% effective against Omicrom variant. Islamic
Republic News Agency. https://en.irna.ir/news/84638920/Official-says-Pasteurcovac-
100-effective-against-Omicrom-variant 2022. (Accessed February 11th 2022)
5. Carles Zerquera JM. Cuban´s vaccines against Omicron. Rapid response to
BMJ 2022; 376 doi: https://doi.org/10.1136/bmj.o66; January 29th, 2022 Available in:
https://www.bmj.com/content/376/bmj.o66/rapid-responses (Accessed February 17th,
2022)
6. CECMED Cuban National Regulatory Agency. Resolution 144/2021. Authorization for
emergency use of SOBERANA02 in adults. https://www.cecmed.cu 2021 (Accessed
November 17th; 2021)
7. Dal-Ré R, Caplan A. Current COVID-19 vaccine trials in high-income countries: are
placebo-controlled trials ethical? Clin Microbiol Infect 2021;27:1565-67. doi:
10.1016/j.cmi.2021.08.005
8. Delahoy M, Ujamaa D, Whitaker M, O'Halloran A, Anglin O, Burns E, et al.
Hospitalizations Associated with COVID-19 Among Children and Adolescents —
COVID-NET, 14 States, March 1, 2020–August 14, 2021. MMWR Morb Mortal Wkly
Rep 2021;70:1255–60. doi: 10.15585/mmwr.mm7036e2
9. Earle K, Ambrosino D, Andrew Fiore-Gartland; Golblatt D, Gilbert PB, et. al. Evidence
for antibody as a protective correlate for COVID-19 vaccines. Vaccine 2021;39:4423–
28. https://doi.org/10.1016/j.vaccine.2021.05.063
10. Elliott P, Bodinier B, Eales O, Wang H, Haw D, Elliott J, et al. Rapid increase in
Omicron infections in England during December 2021: REACT-1 study. Science 2022;
doi: 10.1126/science.abn8347
11. FDA. Vaccines and Related Biological Products Advisory Committee Meeting. June
10, 2021. Licensure and Emergency Use Authorization of Vaccines to Prevent
COVID-19 for Use in Pediatric Populations. https://www.fda.gov/advisory-
committees/advisory-committee-calendar/vaccines-and-related-biological-products-

advisory-committee-june-10-2021-meeting-announcement. 2021 (Accessed September
14th, 2021)
12. Fontanet A, Autran B, Lina B, Kieny MP, Abdool Karim SS, Sridhar D. SARS-CoV-2
variants and ending the COVID-19 pandemic. Lancet 2021; 397: 952–54. doi:
https://doi.org/10.1016/S0140-6736(21)00370-6
13. Frenck R, Klein N, Kitchin N, Gurtman A, Absalon J, Lockhart S et al. Safety,
Immunogenicity, and Efficacy of the BNT162b2 Covid-19 Vaccine in Adolescents. N
Engl J Med 2021;385: 239–50. doi: 10.1056/NEJMoa2107456
14. Han B, Song Y, Li C, Yang W, Ma Q, Jiang Z et al. Safety, tolerability, and
immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy
children and adolescents: A double-blind, randomised, controlled, phase 1/2 clinical
trial. Lancet Infect. Dis 2021;21:1645-53. https://doi.org/10.1016/S1473-
3099(21)00319-4
15. Huang YL, Wu CY. Carbohydrate-based vaccines: challenges and opportunities.
Expert Rev Vaccines 2010;9:1257–74 doi: 10.1586/erv.10.120
16. IRCT International Register Clinical Trials. Identifier RPCEC00000374. Phase I-II
study, sequential during phase I, open-label, adaptive and multicenter to evaluate the
safety, reactogenicity and immunogenicity of a heterologous two-dose schedule of the
prophylactic anti-SARS- CoV-2 vaccine candidate, FINLAY-FR- 2 and a dose of
FINLAY-FR-1A, in Cuban children and adolescents. (COVID-19).
https://rpcec.sld.cu/en/trials/RPCEC00000374-En (Accessed December 1st, 2021)
17. Keech C, Albert G, Cho I, Robertson A, Reed P, Neal S, et al. Phase 1-2 Trial of a
SARS-CoV-2 Recombinant Spike Protein Nanoparticle. N Engl J Med 2020;383:
2320-32. doi: https://doi.org/10.1056/NEJMoa2026920
18. Khoury DS, Cromer D, Reynaldi A, Schlub T, Wheatley A, Juno J, et. al. Neutralizing
antibody levels are highly predictive of immune protection from symptomatic SARS-
CoV-2 infection. Nature Medicine 2021; 27:1205–11. https://doi.org/10.1038/s41591-
021-01377-8
19. Lopez-Sejas N, Campos C, Hassouneh F, Sanchez-Correa B, Tarazona R, Pera A,
Solana R. Effect of CMV and Aging on the Differential Expression of CD300a,
CD161, T-bet, and Eomes on NK Cell Subsets. Front Immunol 2016;7:1-13. doi:
10.3389/fimmu.2016.00476
20. Marshall M, Ferguson ID, Lewis P, Jaggi P, Gagliardo C, Stewart JC. Symptomatic
acute myocarditis in 7 adolescents after Pfizer-BioNTech COVID-19 vaccination.
Pediatrics 2021;148:2-11. doi e2021052478

21. Oster ME, Shay DK, Su JR, Gee J, Creech B, Broder KR et al. Myocarditis cases reported after mRNA-Based COVID-19 vaccination in the US from December 2020 to August 2021. JAMA 2022;327:331-340. doi:10.1001/jama.2021.24110

22. Petersen E, Buchy P. Vaccination against SARS-CoV-2 should be included in childhood vaccination programs. Int. J Infectious Diseases 2021;106:429–430 doi: https://doi.org/10.1016/j.ijid.2021.04.082

23. Portal-Miranda JA. Las vacunas cubanas generan anticuerpos ante ómicron: Ese éxito no puede llevarnos a la confianza. http://www.cubadebate.cu/opinion/2022/01/26/ 2022 (Accessed January 26th, 2022)

24. Rodriguez A. Virus slams Cuba as it races to roll out its new vaccines. ABC News, July 21th, 2021. https://abcnews.go.com/Health/wireStory/virus-slams-cuba-races-roll-vaccines-78953663 (Accessed February 17th, 2022)

25. Thapa P, Farber D. The Role of the Thymus in the Immune Response. Thorac Surg Clin, 2019; 29:123-131 doi: 10.1016/j.thorsurg.2018.12.001

26. Toledo-Romani ME, Garcia-Carmenate M, Valenzuela-Silva C, Baldoquin-Rodriguez W, Martínez-Pérez M, Rodríguez-Gonzalez MC, et al. Efficacy and Safety of SOBERANA 02, a COVID-19 conjugate vaccine in heterologous three doses combination. MedRxiv 2021c https://doi.org/10.1101/2021.10.31.21265703

27. Toledo-Romani ME, García-Carmenate M, Verdecia-Sánchez L, Pérez-Rodríguez S, Rodriguez-González M, Valenzuela-Silva C. et al. Safety and immunogenicity of anti-SARS CoV-2 conjugate vaccine SOBERANA 02 in a two-dose or three-dose heterologous scheme in adults: Phase IIb Clinical Trial. MedRxiv 2021b https://doi.org/10.1101/2022.01.01.21268271

28. Toledo-Romani ME, Verdecia-Sánchez L, Rodriguez-González M, Rodríguez-Noda L, Valenzuela-Silva C, Paredes-Moreno et al. Safety and immunogenicity of anti-SARS CoV-2 vaccine SOBERANA 02 in homologous or heterologous scheme. MedRxiv 2021a doi: https://doi.org/10.1101/2021.11.14.21266309

29. Valdes-Balbin Y, Santana-Mederos D, Paquet F, Fernandez S, Climent Y, Chiodo F et al. Molecular Aspects Concerning the Use of the SARS-CoV-2 Receptor Binding Domain as a Target for Preventive Vaccines. ACS Cent Sci 2021;7:757−67. https://doi.org/10.1021/acscentsci.1c00216

30. Valdes-Balbin Y, Santana-Mederos D, Quintero L, Fernandez S, Rodriguez L, Sanchez-Ramírez B et al. SARS-CoV-2 RBD-Tetanus toxoid conjugate vaccine induces a strong neutralizing immunity. ACS Chem Biol 2021;16:1223–33. https://doi.org/10.1021/acschembio.1c00272

31. Ventura MT, Casciaro M, Gangemi S, Buquicchio R. Immunosenescence in aging:
between immune cells depletion and cytokines up-regulation. Clin Mol Allergy 2017;
15:21 doi: https://doi.org/10.1186/s12948-017-0077-0
32. Verez-Bencomo V, Fernandez-Santana V, Hardy E, Toledo ME, Rodriguez MC,
Heynngnezz L et. al. A synthetic conjugate polysaccharide vaccine against
Haemophilus influenzae type b. Science 2004;305: 522-525 doi:
10.1126/science.1095209
33. Wallace M, Woodworth KR, Gargano JW, Scobie HM, Blain M, Moulia D et al. The
Advisory Committee on Immunization Practices' interim recommendation for use of
Pfizer-BioNTech COVID-19 vaccine in adolescents aged 12–15 years—United States,
May 2021. MMWR Morb Mortal Wkly Rep 2021;70:749–52.
https://doi.org/10.15585/mmwr.mm7020e1
34. Walter EB, Talaat KR, Sabharwal C, Gurtman A, Lockhart S, Paulsen GC et al.
Evaluation of the BNT162b2 Covid-19 Vaccine in Children 5 to 11 Years of Age. N
Engl J Med 2022;386:35-46 doi: 10.1056/NEJMoa2116298
35. WHO COVID-19 – Landscape of novel coronavirus candidate vaccine development
worldwide. https://www.who.int/publications/m/item/draft-landscape-of-covid-19-
candidate-vaccines, 2021 (accessed December 29th, 2021)
36. World Health Organization. Causality Assessment of an Adverse Event Following
Immunization (AEFI). 2nd Edition. Geneva: WHO; 2019. ISBN: 9789241516990
https://www.who.int/publications/i/item/causality-assessment-aefi-user-manual-2019
2019 (Accessed June 17th, 2021)
37. Xia S, Zhang YT, Wang Y, Wang H, Yang Y, Fu-Gao G et al. Safety and
immunogenicity of an inactivated COVID-19 vaccine, BBIBP-CorV, in people
younger than 18 years: a randomised, double-blind, controlled, phase 1/2 trial. Lancet
Infect Dis 2022;22:196-208. doi: 10.1016/S1473-3099(21)00462-X
38. Yang S, Li Y, Dai L, Wang J, He P, Li C, et al. Safety and immunogenicity of a
recombinant tandem-repeat dimeric RBD-based protein subunit vaccine (ZF2001)
against COVID-19 in adults: two randomised, double-blind, placebo-controlled, phase 1
and 2 trials. Lancet Infect Dis. 2021; 21:1107-11. doi: https://doi.org/10.1016/S1473-
3099(21)00127-4
670